

Phylogeny and biogeography of some Cretaceous spatangoid echinoids with special emphasis on taxa from the Western Interior Seaway

Authors: Byrum, Steven, and Lieberman, Bruce S.

Source: Journal of Paleontology, 95(3) : 613-623

Published By: The Paleontological Society

URL: <https://doi.org/10.1017/jpa.2020.102>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Phylogeny and biogeography of some Cretaceous spatangoid echinoids with special emphasis on taxa from the Western Interior Seaway

Steven Byrum¹ and Bruce S. Lieberman^{2*} 

¹Florida Museum of Natural History, University of Florida, Gainesville, Florida 32611, USA <s.byrum@ufl.edu>

²Department of Ecology and Evolutionary Biology and Biodiversity Institute, University of Kansas, Lawrence, Kansas 66045, USA <blieber@ku.edu>

Abstract.—Members of the echinoid order Spatangoida, a highly diverse and abundant marine invertebrate clade, were important denizens of the Cretaceous Western Interior Seaway (WIS), an epicontinental seaway that divided North America in two during an interval of greenhouse conditions between roughly 100 and 65 million years ago. A phylogenetic analysis of spatangoids was conducted using a character matrix of 32 characters from 21 species. Species that occur in the WIS were considered comprehensively, and species from other regions such as South America, Europe, and North Africa were also incorporated into the analysis. Phylogenetic patterns retrieved are largely congruent with preexisting family-level classifications; however, species within several genera, especially *Hemiaster* and *Heteraster*, need to be reassigned so that classification better reflects phylogeny. The genera *Washitaster* and *Heteraster* are closely related, as are *Mecaster*, *Palhemiaster*, and *Proraster*; *Pliotoxaster*, *Macraster*, and *Hemiaster*; and *Micraster* and *Diplodetus*. Biogeographic patterns were also considered using the phylogeny, and several episodes of vicariance and range expansion were identified. These were possibly related to some of the various major episodes of sea-level rise and fall during the Cretaceous. In particular, Valangian–mid-Aptian regressions may have caused vicariance within *Heteraster* and *Washitaster* while other early spatangoid vicariance may be related to regressions during the late Aptian–early Cenomanian. Further, vicariance caused by regressions during the mid-Cenomanian–Maastrichtian may have driven diversification within *Micraster* and *Diplodetus*. Last, transgressions during the late Aptian–early Cenomanian seem to have spurred prominent range expansions in *Mecaster* and *Hemiaster*.

Introduction

Echinoids are a highly diverse group of marine invertebrates known from a plethora of modern and ancient habitats (Smith, 1984; Smith and Jeffery, 2000; Mooi, 2001; Kroh and Mooi, 2018; Arachchige et al., 2019). The group experienced a sharp diversity decline at the end of the Paleozoic, when the majority of stem-group echinoids went extinct after the end-Permian mass extinction (Thompson et al., 2018). However, their diversity rebounded in the Mesozoic era with the appearance of irregular echinoids (Saucède et al., 2007). Echinoids were abundant denizens of the shallow benthic communities of the southern parts of the Western Interior Seaway (WIS), in what is today the southwestern United States (e.g., Texas and New Mexico) and northern Mexico (Sonora and Coahuila). This is where the WIS opened into the much larger North Atlantic/West Tethys for much of its duration (Blakey, 2014). Many of the first echinoids found in this region were described in various nineteenth-century studies (e.g., Roemer, 1849, 1852; Clark, 1893). Subsequent works include descriptions by Clark (1915) and

Cooke (1946, 1953, 1955) and publications focused on taxa from Texas (Adkins, 1928; Whitney and Kellum, 1966; Smith and Rader, 2009; Morgan, 2016; Thompson, 2016). Echinoids have been frequent subjects of phylogenetic analysis (relatively recent publications in this area include Kroh and Smith, 2010; Mooi et al., 2014; Koch et al., 2018; Thompson et al., 2018).

Among the irregular echinoids, the most diverse and abundant are the heart urchins of the order Spatangoida, which inhabit all oceans and a wide array of habitats (Stockley et al., 2005). Spatangoids are deposit-feeding burrowers that radiated in the Cretaceous and survive into the present (Mortensen, 1950; Fischer, 1966; Smith, 1984; Mooi, 2001; Smith and Wright, 2008, 2012; Smith and Kroh, 2011; Kroh and Mooi, 2018). They have also been incorporated into phylogenetic analyses that considered both living and fossil taxa (Villier et al., 2004; Stockley et al., 2005; Saucède et al., 2007; Kroh and Smith, 2010; Kroh et al., 2014). The goal of the present study is to focus on species-level phylogenetic relationships of Cretaceous spatangoids, with special emphasis on taxa from the WIS, although it was not possible to consider to any great extent the diverse range of taxa present in the Mississippi Embayment, an important and related biogeographic region (see Zachos, 2017), as taxa in this region primarily radiated too late to be

* Corresponding Author

Table 1. Taxa considered in phylogenetic analysis, relevant material examined, and geographic occurrence; KUMIP = Division of Invertebrate Paleontology, Biodiversity Institute, University of Kansas, Lawrence, USA; NMMNH = New Mexico Museum of Natural History, Albuquerque, USA; STIB = Universität Bonn, Germany; USNM = United States National Museum of Natural History, Washington, D.C., USA; UT = Non-Vertebrate Paleontology Laboratory, University of Texas, Austin, USA. For geographic occurrence, 0 = Western Europe; 1 = North Africa; 2 = South America; 3 = the WIS.

| Taxa | Relevant material examined | Stage | Geographic occurrence |
|--|---|---------------|-----------------------|
| <i>Toxaster retusus</i> (Lamarck, 1816) | UT 11704 | Valanginian | 0,1 |
| <i>Heteraster oblongus</i> (Brongniart, 1821) | UT 14575 | Valanginian | 0 |
| " <i>Heteraster</i> " <i>texanus</i> (Roemer, 1849) | KUMIP 370536, 370548–370556, UT BEG 34165, STIB Texas 157 | Albian | 3 |
| <i>Washitaster riovistae</i> (Adkins, 1920) | UT BEG 21491–21493 | Albian | 3 |
| <i>Washitaster wenoensis</i> (Adkins, 1920) | UT BEG 21496 | Albian | 3 |
| <i>Macraster elegans</i> (Shumard, 1853) | KUMIP 420253, 490608, 490611, UT BEG 21947, 34161 | Albian | 3 |
| <i>Mecaster fourneli</i> (Agassiz and Desor, 1847) | UT 14285 | Turonian | 1,2 |
| <i>Micraster schroederi</i> Stolley, 1892 | UT 53845 | Campanian | 0 |
| <i>Pliotoxaster comanchei</i> (Clark, 1915) | USNM 103893, UT BEG 21264, BEG 21270 | Albian | 3 |
| <i>Proraster dalli</i> (Clark, 1891) | USNM 19114, UT 83036 | Albian | 3 |
| <i>Proraster humphreysanus</i> (Meek and Hayden, 1857) | USNM 331 | Maastrichtian | 3 |
| <i>Proraster bexari</i> (Clark, 1915) | USNM 8330 | Albian | 3 |
| <i>Palhemaster calvini</i> (Clark, 1915) | UT BEG 21268, BEG 21487 | Albian | 3 |
| <i>Hemiasster bufo</i> (Brongniart, 1822) | | Cenomanian | 0 |
| <i>Hemiasster cranium</i> Cooke, 1946 | USNM 103890 | Albian | 3 |
| " <i>Hemiasster</i> " <i>estepi</i> Lucas, 2000 | NMMNH P-26501, 26508, 26515 | Albian | 3 |
| " <i>Hemiasster</i> " <i>batnensis</i> (Coquand, 1862) | UT 14252, 83026 | Cenomanian | 2,3 |
| " <i>Hemiasster</i> " <i>texanus</i> (Roemer, 1849) | UT BEG 34771, STIB Texas 156 | Santonian | 3 |
| <i>Diplodetus americanus</i> Stephenson, 1941 | USNM 76285, UT 83001 | Maastrichtian | 3 |
| <i>Diplodetus parvistella</i> Schluter, 1900 | | Maastrichtian | 0 |
| <i>Linthia variabilis</i> Slocum, 1909 | | Maastrichtian | 3 |

relevant to the present analysis. The present study also uses the phylogenetic perspective to consider aspects of the biogeographic relationship between the WIS and other parts of the Cretaceous world.

Materials and methods

Taxonomic sampling.—Twenty-one taxa were selected for phylogenetic analysis, comprising representatives of the "Toxasteridae," Hemiassteridae, and Micrasteridae, the largest and most morphologically diverse clades within the Spatangoida (Smith and Kroh, 2011). Due to the large number of spatangoid species, not all could be considered in the phylogenetic analysis. Thus, only Cretaceous species, and not Cenozoic and modern forms, were analyzed. All well-preserved Cretaceous WIS species for which at least 75% of the characters used in the phylogenetic analysis could be coded were considered. Additional Cretaceous spatangoid species from outside the WIS were selected using the criteria that: (1) type specimens or quality photographs of these needed to be accessible, (2) species needed to be defined such that at least 75% of characters could be coded, and (3) all major Cretaceous biogeographic regions should be represented. This meant that species that were the subject of more recent systematic treatments were better represented in the analysis. Taxa subjected to phylogenetic analysis and relevant material examined are listed in Table 1. We follow the classification scheme of Smith and Kroh (2011) as it has been widely utilized, although an updated classification has recently been presented by Kroh (2020).

Characters.—Character and character state descriptions were developed via detailed comparative examination of test features. The preexisting literature was also considered to

identify characters that had previously been used to diagnose groups and infer relationships among groups. Villier et al. (2004), Smith and Stockley (2005), and Kroh and Smith (2010) were especially helpful in this regard. In total, 32 characters were identified (Appendix), including a combination of qualitative and quantitative characters. Quantitative characters were measured for coding using the program ImageJ; character states are noncontinuous relative to each other, and multistate characters were treated as unordered (Table 2). A variety of means for coding characters exist, and we have followed the precepts and principles presented by Wiley and Lieberman (2011) for identifying hypotheses of homologies and coding them. No explicit models of character weighting were used, but different morphological systems varied in the number of characters they contained. Further, some characters may display more homoplasy than others; thus, different characters may provide varying degrees of phylogenetic resolution.

Phylogenetic methods.—A parsimony analysis was implemented using the matrix in Table 2 via PAUP 4.0 (Swofford, 2003), with *Toxaster retusus* (Lamarck, 1816) employed as the outgroup; this taxon has been previously treated as an earliest representative of the group by Stockley et al. (2005). (Nexus files utilized are provided as Supplemental Data.) Data were subjected to a branch-and-bound search (Hendy and Penny, 1982) to find the most parsimonious tree(s). A bootstrap analysis was conducted using a heuristic search in PAUP 4.0 to determine support for each node appearing in the most parsimonious tree(s) using 100 bootstrap replicates in a stepwise search that employed five random replicated addition sequences per bootstrap replication with one tree held. Jackknife analysis was also used, again using 100 replicates in a stepwise search that employed five random addition

Table 2. Character state distributions for taxa used in phylogenetic analysis. ? = missing data.

| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 | 34 | 35 | |
|---|---|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|--|
| <i>Toxaster retusus</i> | 0 | 0 | 1 | 0 | 0 | 0 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | ? | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 2 | 1 | 2 | 2 | 0 | | | |
| <i>Heteraster oblongus</i> | 0 | 0 | 1 | 0 | 1 | 1 | 0 | 1 | 1 | 1 | 2 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 2 | 2 | 1 | | | |
| " <i>Heteraster</i> " <i>texasus</i> | 0 | 0 | 1 | 0 | 0 | 1 | 0 | 1 | 1 | 1 | 2 | 1 | 2 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 1 | 0 | 2 | 1 | 1 | | | |
| <i>Washitaster riovistae</i> | 0 | 0 | 1 | 0 | 0 | 0 | 1 | 1 | 1 | 1 | 2 | 1 | 2 | 0 | 0 | 0 | 1 | 0 | 0 | 1 | 1 | 1 | 0 | 1 | 1 | 1 | 0 | 1 | 0 | 2 | 1 | 2 | | | | |
| <i>Washitaster wenoensis</i> | 0 | 0 | 1 | 0 | 0 | 1 | 0 | 1 | 1 | 1 | 2 | 1 | 2 | 0 | 0 | 0 | 0 | 1 | ? | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | ? | ? | 1 | 0 | 2 | | |
| <i>Macraster elegans</i> | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 2 | 0 | 1 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 2 | 1 | 2 | 2 | 0 | | | |
| <i>Mecaster fourmeli</i> | 1 | 0 | 2 | 0 | 0 | 0 | 1 | 0 | 0 | 3 | 1 | 0 | 3 | 0 | 0 | 1 | 0 | 0 | 1 | 1 | 1 | 1 | 1 | 0 | 1 | 1 | 1 | 1 | 0 | 2 | 2 | 0 | | | | |
| <i>Micraster schroederi</i> | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 2 | 0 | 0 | 2 | 1 | 0 | 0 | 0 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 2 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | | | | |
| <i>Pliotaxaster comanchei</i> | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 2 | 0 | 0 | 2 | 0 | 1 | 0 | 0 | 1 | 1 | 1 | 0 | 0 | 1 | 0 | 1 | 1 | 1 | 1 | 1 | 2 | 2 | 0 | | | | |
| <i>Proraster dalli</i> | 1 | 0 | 1 | 1 | 2 | 0 | 1 | 1 | 0 | 2 | 2 | 0 | 3 | ? | 0 | 2 | 0 | 0 | 1 | ? | ? | 1 | 1 | 1 | 2 | 0 | 0 | ? | ? | 2 | 2 | 1 | | | | |
| <i>Proraster bexari</i> | 0 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 0 | 3 | 1 | 0 | 3 | 0 | 0 | 2 | 0 | ? | 1 | 1 | ? | 1 | 1 | 0 | 1 | 0 | 0 | ? | ? | 2 | 1 | 1 | | | | |
| <i>Proraster humphreysanus</i> | 0 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 0 | 3 | 2 | 0 | 3 | 0 | 0 | 2 | 0 | 0 | 1 | 1 | 0 | 1 | 1 | ? | 0 | 0 | 1 | 1 | 1 | 2 | 1 | 2 | | | | |
| <i>Palhemiaster calvini</i> | 0 | 1 | 1 | 0 | 0 | 0 | 1 | 0 | 0 | 3 | 0 | 0 | 3 | 0 | 0 | 1 | 0 | ? | 1 | 1 | 0 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 2 | 1 | 0 | | | | |
| <i>Hemiaster bufo</i> | 0 | 1 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 3 | 1 | 0 | 3 | 0 | ? | 2 | 0 | 1 | 1 | 1 | 0 | 1 | 0 | 1 | 1 | 0 | 1 | 1 | 1 | 2 | 1 | 0 | | | | |
| <i>Hemiaster cranium</i> | 0 | 1 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 3 | 1 | 0 | 3 | 0 | 0 | 1 | 0 | 0 | 1 | 1 | 0 | 1 | 0 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 0 | 1 | | | | |
| " <i>Hemiaster</i> " <i>batnensis</i> | 0 | 2 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 3 | 1 | 0 | 3 | 0 | 0 | 2 | 0 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 1 | 1 | 1 | 1 | 1 | 2 | 2 | 0 | | | | |
| " <i>Hemiaster</i> " <i>estepi</i> | 0 | 2 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 2 | 1 | 0 | 2 | 0 | 0 | 2 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 1 | 1 | 2 | 0 | 1 | | | | |
| " <i>Hemiaster</i> " <i>texasus</i> | 1 | 1 | 1 | 0 | 0 | 0 | 1 | 0 | 1 | 3 | 1 | 0 | 3 | 0 | 0 | 2 | 0 | 0 | 1 | 0 | 1 | 1 | 0 | 0 | 1 | 1 | 1 | 2 | 1 | 2 | 1 | 0 | | | | |
| <i>Linthia variabilis</i> | 0 | 1 | 2 | 0 | 1 | 0 | 1 | ? | 0 | 3 | 0 | ? | 3 | 0 | 0 | 2 | 0 | ? | 1 | 1 | ? | 1 | 1 | 1 | 1 | 1 | 1 | ? | ? | ? | ? | | | | | |
| " <i>Diplodetus</i> " <i>americanus</i> | 0 | 2 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 2 | 0 | 0 | 2 | 1 | 0 | 1 | 0 | 1 | 1 | 1 | 0 | 1 | 0 | 1 | 1 | 0 | 1 | 1 | 1 | 2 | 1 | 0 | | | | |
| <i>Diplodetus parvistella</i> | 0 | 1 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 1 | 0 | ? | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 1 | ? | 1 | 1 | 0 | 1 | 1 | 0 | ? | ? | 0 | 0 | 0 | | | | |

sequences per replication, with the percentage of characters removed value chosen at 10%, which equates to three characters, and one tree held. For both support tests, groups were retained that were compatible with the 50% majority rule consensus tree. In addition, a Bremer branch support analysis (Bremer, 1994) was conducted to calculate node support using the difference between the length of the most parsimonious tree and the largest tree in which a node is maintained.

Finally, a Bayesian analysis was conducted using MrBayes 3.2.7, again using *Toxaster retusus* as an outgroup, to explore how well supported the various nodes in the parsimony tree were. The original file was exported from the simplified nexus file provided as Supplemental Data. No partitioning of the data was implemented as there were no a priori reasons for differentiating among character types or morphological systems. A standard MKV model was employed without a particular starting tree and performed on the CIPRES cluster. The analysis ran for 10 million generations and sampled every 1,000 generations. The posterior probability values for nodes retrieved in the analysis were then compared with those in the parsimony tree (see Cole, 2019). Note that there has been extensive discussion about whether a Bayesian or a parsimony approach is better justified for the analysis of morphological character data, and consideration of this topic goes substantially beyond the scope of this contribution. We apply these different methods here only for purposes of considering commonality of results achieved across different methods. The interested reader is referred to discussions by Wiley and Lieberman (2011) and Wright (2019) for greater amplification of the various issues involved with applying different phylogenetic methods to morphological data.

Biogeographic analysis.—Biogeographic patterns were inferred by first substituting the area of occurrence of the species for a given species name. The areas utilized were Western Europe, North Africa, South America, and the WIS (Fig. 1), and these represent major areas of endemism for echinoids during the Late Cretaceous (see e.g., Smith, 1984, 1992; Rosen and Smith,

1988; Néraudeau and Mathey, 2000). Biogeographic states were then mapped to ancestral nodes using the modified version of Fitch parsimony optimization for multistate characters, which treats all transformations between different character states as unordered (described by Lieberman, 2000). This method minimizes the assumptions about the way biogeographic change happens between different regions. Transitions between ancestral and descendant nodes or ancestral and descendant terminals that involved range contraction were treated as potential instances of vicariance, and equivalent transitions involving range expansion were treated as potential instances of dispersal following Lieberman (2000).

Repositories and institutional abbreviations.—Type, figured, and other specimens examined in this study are deposited in the following institutions: Division of Invertebrate Paleontology, Biodiversity Institute, University of Kansas (KUMIP); New Mexico Museum of Natural History (NMMNH); United States National Museum of Natural History (USNM); and Non-Vertebrate Paleontology Laboratory, University of Texas, Austin (UT and UT BEG). High-quality images of type specimens were also provided by the Universität Bonn (STIB).

Systematic paleontology

Order Spatangoida Agassiz, 1840

Family Hemiasteridae Clark, 1917

Genus *Hemiaster* Desor in Agassiz and Desor, 1847

Type species.—*Spatangus bufo* Brongniart, 1822, Late Cretaceous, Cenomanian, France.

"*Hemiaster*" *texasus* (Roemer, 1849)

Figure 2.1

1849 *Hemiaster texasus* Roemer, p. 77, 393.

1953 *Hemiaster texasus*; Cooke, p. 33, pl. 13, figs. 1–4 (see for more complete synonymy).




Figure 1. Paleogeographic reconstruction for Late Cretaceous ~90 Ma; © 2016 Colorado Plateau Geosystems Inc.

- 1955 *Hemiaster texanus*; Cooke, p. 109, pl. 29, figs. 5–10.
 1991 *Mecaster texanum*; Smith in Smith and Bengtson, p. 13.
 2013 *Mecaster texanum*; Oliveira et al., p. 11, figs. 7A–C, 9A–I.
 2017 *Mecaster texanus*; Manso and Souza-Lima, p. 225, figs. 5, 6.

Holotype.—STIB Texas 156, New Braunfels, Texas, Late Cretaceous, Santonian.

Materials.—STIB Texas 156; UT BEG 34771, from the Austin Formation, Late Cretaceous, Santonian, Hondo River, 4.0 miles northwest of Hondo, Medina County, Texas, USA.

Remarks.—“*Hemiaster*” *texanus* does not share a sister group relationship with *Hemiaster bufo* (Brongniart, 1822), but the phylogenetic analysis presented herein suggests that they are relatively closely related. Further, it has been assigned to




Figure 2. “*Hemiaster*” *texanus*: (1) STIB Texas 156; and “*Heteraster*” *texanus* (STIB Texas 157) (2). Scale bars = 1 cm.

Hemiaster previously (e.g., Cooke, 1953) and is closely related to other taxa traditionally assigned to *Hemiaster* (e.g., *H. cranium* Cooke, 1946; “*Hemiaster*” *batnensis* (Coquand, 1862), see Berndt, 2003; and “*Hemiaster*” *estepi* Lucas, 2000). Thus, rather than creating a new monotypic genus or lumping the species in with other genera, “*Hemiaster*” *texanus* is assigned to a paraphyletic genus following the ‘within quotes’ convention of Wiley (1979). Photographs of the holotype had not been previously published, and thus one is provided herein.

Genus *Proraster* Lambert, 1895

Type species.—*Schizaster atavus* Arnaud in Cotteau, 1883; Late Cretaceous, Campanian, near Juillaguet (now Boisé-la-Tude), Charente, and Mensignac, Dordogne, France.

Proraster humphreysanus (Meek and Hayden, 1857)
Figure 3.2

1857 *Hemiaster?* *humphreysanus* Meek and Hayden, p. 147.
1953 *Hemiaster humphreysanus*; Cooke, p. 30, pl. 12, figs. 1–4 (see for more complete synonymy).
2017 *Hemiaster humphreysanus*; Hook and Cobban, p. 2.

Holotype.—USNM PAL 331, Cedar Creek Anticline, Montana, Pierre Shale, Late Cretaceous, Maastrichtian.

Materials.—USNM PAL 331.

Remarks.—This species is referred to *Proraster* on the basis of its close relationship to *Proraster dalli* (Clark, 1891) in the phylogenetic analysis and is a new taxonomic combination. Cooke (1953) had posited that these species were closely

similar. However, since the type species of *Proraster* could not be included in the phylogenetic analysis, it is possible that the generic assignment of *Proraster humphreysanus* could change upon subsequent study. Photographs of the holotype had not been previously published, and thus one is provided herein.

Family Micrasteridae Lambert, 1920

Genus *Diplodetus* Schlüter, 1900

Type species.—*Diplodetus brevistella* Schlüter, 1870; Late Cretaceous, Darup, Westphalia, Germany.

“*Diplodetus*” *americanus* (Stephenson, 1941)
Figure 3.1

1941 *Micraster (Plesiaster) americanus* Stephenson, p. 69, pl. 7, figs. 1–4.

1953 *Micraster americanus*; Cooke, p. 38, pl. 15, figs. 10–13.

2000 *Diplodetus americanus*; Jagt, p. 288.

Holotype.—USNM PAL 76285, six miles east of Castroville, Texas, cut in San Antonio road in Bexar County, ‘Navarro group,’ Late Cretaceous, Maastrichtian.

Materials.—USNM 76285; UT 83001, cut in San Antonio road in Bexar County, Corsicana Marl, Navarro group, Late Cretaceous, Maastrichtian.

Remarks.—“*Diplodetus*” *americanus* does not share an exclusive sister group relationship with *Diplodetus parvistella* (Schlüter, 1900), but the phylogenetic analysis presented herein suggests that they are relatively closely related. Given that *Diplodetus parvistella* and the type of the genus, *Diplodetus brevistella* Schlüter, 1870, are closely related (e.g.,




Figure 3. (1) “*Diplodetus*” *americanus* (USNM PAL 76285). (2) *Proraster humphreysanus* (USNM PAL 331). Scale bars = 1 cm.

Smith and Kroh, 2011), and to avoid creating a new monotypic genus, “*Diplodetus*” *americanus* is referred to a genus that is designated as paraphyletic following the within-quotes convention of Wiley (1979), though other taxonomic rearrangements are also possible. However, since the type species of *Diplodetus* could not be included in the phylogenetic analysis, it is possible that the generic assignment of “*Diplodetus*” *americanus* could change upon subsequent study. Photographs of the holotype have not been published for several decades, and thus one is provided herein.

Family Toxasteridae Lambert, 1920

Genus *Heteraster* d’Orbigny, 1855

Type species.—*Spatangus oblongus* Brongniart, 1821; Late Cretaceous, Valanginian–Hauterivian, France.

“*Heteraster*” *texanus* (Roemer, 1849)

Figure 2.2

- 1849 *Toxaster texanus* Roemer, p. 393.
 1928 *Heteraster texanus*; Adkins, p. 55, pl. 3, fig. 1.
 1946 *Enallaster texanus*; Cooke, p. 231, pl. 33, figs. 9–12 (see for more complete synonymy).
 1955 *Enallaster texanus*; Cooke, p. 102.
 2004 *Heteraster texanus*; Villier et al., p. 267.
 2009 *Enallaster texanus*; Smith and Rader, p. 9.
 2014? *Heteraster texanus*; Silva-Martinez et al., p. 388, fig. 6e–g.
 2016 *Heteraster texanus*; Thompson, p. 100.
 2017 *Heteraster texanus*; Melo et al., p. S120, fig. 7.
 2019 *Heteraster texanus*; Martha et al., p. 246.

Holotype.—STIB Texas 157, Fredricksburg, Texas, Late Cretaceous, Albian.

Materials.—KUMIP 370536, 370548–370556 (detailed locality information is available for all KUMIP specimens at <https://biodiversity.ku.edu/invertebrate-paleontology/collections/collections-search>); STIB Texas 157; UT BEG 34165.

Remarks.—“*Heteraster*” *texanus* does not share an exclusive sister group relationship with *Heteraster oblongus* (Brongniart, 1821), but the phylogenetic analysis presented herein suggests that they are relatively closely related. Further, the species has traditionally been assigned to *Heteraster* (e.g., see Villier et al., 2004, Thompson, 2016; Yavari et al., 2016). Thus, rather than creating a new monotypic genus, or lumping the species within *Washitaster* Lambert, 1927, “*Heteraster*” *texanus* is assigned to a paraphyletic genus following the within-quotes convention of Wiley (1979). Photographs of the holotype had not been previously published, and thus one is provided herein.

Results

Phylogenetic analysis.—Two separate parsimony analyses were run. The first included all taxa except for *Linthia variabilis* Slocom, 1909, which was coded as missing for 25% of the

character states. This analysis resulted in a strict consensus of four most parsimonious trees of length 111, consistency index (CI) of 0.41, and retention index (RI) of 0.60 (Fig. 4), with one character parsimony-uninformative. Then, *Linthia variabilis* was added and a branch-and-bound analysis performed again. (Both nexus files utilized are provided as Supplemental Data.) This resulted in a strict consensus of 83 trees of length 114, a CI of 0.40, and an RI of 0.60 (Fig. 5). Overall, the topology resembles that of Figure 4, but resolution decreased, possibly due to the high amount of missing data for *Linthia variabilis*. In terms of tree support statistics, four nodes of the cladogram in Figure 4 were supported by bootstrap values greater than 60, with 15 nodes supported by jackknife values greater than 60 (and seven of these values were above 80). Bremer support was found for four nodes in the tree (Fig. 4). The tree in Figure 4, with the characters used in phylogenetic analysis mapped to the tree, is provided as a figure in Supplemental Data. The Bayesian




Figure 4. Strict consensus of four most parsimonious trees of length 111 recovered in analysis, excluding relatively incomplete taxon *L. variabilis*. Numbers above node in bold are bootstrap support values, numbers below node in italics are jackknife support values, and numbers in normal font below node are Bremer support values. *Species occurring outside of the WIS.

analysis (not shown) of the matrix without *Linthia variabilis* contained five nodes with probabilities greater than 60, generally aligning with the results of the parsimony analysis but containing poor resolution of later nodes.

The analyses suggest several notable relationships relative to traditional groupings and previous analyses. Regarding traditional groupings, the genera *Hemiaster*, *Heteraster*, and *Proraster* are paraphyletic or even polyphyletic (see also Smith and Bengtson, 1991; Néraudeau, 1994). To make the classification more compatible with the phylogeny, several taxonomic changes were suggested (Fig. 4). For example, *Proraster* is treated herein as containing *Proraster dalli*, *Proraster humphreysanus*, and *Proraster bexari* (Clark, 1915) (Fig. 4). In addition, some taxa were assigned to paraphyletic genera using the within-quotes convention of Wiley (1979). These are “*Heteraster*” *texanus*; “*Hemiaster*” *batnensis*; “*Hemiaster*” *texanus*; “*Hemiaster*” *estepi*; and “*Diplodetus*” *americanus* (Fig. 4).




Figure 5. Strict consensus of 83 most parsimonious trees of length 114 recovered in analysis, including relatively incomplete *L. variabilis*. Numbers above node in bold are bootstrap support values, numbers below node in italics are jackknife support values, and numbers in normal font below node are Bremer support values. *Species occurring outside of the WIS.

At this time, this is preferable to establishing several new monotypic genera as not all relevant species could be considered in the present analysis. The phylogeny indicates that *Mecaster fourneli* (Agassiz and Desor, 1847) and *Palhemiaster* Lambert, 1916 are closely related, with *Proraster* the sister group of these; a paraphyletic set of species referable to *Hemiaster* and “*Hemiaster*” represent several successive lineages sister to these. In addition, the following pairs of genera are closely related: *Diplodetus* plus *Micraster* Agassiz, 1836; *Washitaster* plus *Heteraster*; and *Pliotoxaster* Fourteau, 1907 plus *Macraster* Roemer, 1888. Further, *Washitaster* plus *Heteraster* is the sister group to all spatangoids included here (excluding *Toxaster* Agassiz, 1840).

Biogeographic analysis.—Biogeographic patterns provide evidence for vicariance within the spatangoids (Fig. 6). For example, following an initial range expansion by the group from being distributed in Western Europe and North Africa to being distributed in these areas and the WIS during the Valanginian, there was subsequent vicariance between northern Africa and the WIS plus Western Europe during the Albian comprising *Heteraster* and *Washitaster*. There is further vicariance between Western Europe and the WIS within *Heteraster* and *Washitaster*. Another vicariance event occurred in the Campanian associated with the divergence of the WIS “*Diplodetus*” *americanus* and the Western European *Micraster* and *Diplodetus parvistella*. The rest of the diversification within the clade represents speciation occurring within the WIS, although at this time it could not be determined whether this involved smaller scale vicariant events within the WIS or actual sympatric differentiation.

There also appears to have been range expansion by spatangoids from the WIS into Europe, North Africa, and South America. For example, at some time between the Albian and the Cenomanian, there was a range expansion by the lineage leading to *Hemiaster bufo* (Fig. 6). There was also a range expansion from the WIS into South America during the Cenomanian associated with the origin of “*Hemiaster*” *batnensis*. Finally, sometime between the Albian and the Turonian, there was a range expansion from the WIS into North Africa and South America associated with the origin of *Mecaster fourneli*.

Discussion

Phylogenetic analysis.—In general, the phylogenetic results (Fig. 4) match various aspects of previous interpretations, and genera could be readily assigned to families as per the familial level classification presented by Smith and Kroh (2011). For example, *Micraster* and *Diplodetus* had been grouped into a monophyletic Micrasteridae due to the shared presence of an ethmophract apical disc and similarly patterned subanal fasciole (Smith and Stockley, 2005), although *Diplodetus* was found to be paraphyletic herein. Further, various authors (e.g., Smith and Bengtson, 1991; Stockley et al., 2005; Oliveira et al., 2013; Manso and Souza-Lima, 2017) have, either directly or on the basis of their taxonomic assignments, posited a relatively close relationship between *Mecaster* and *Hemiaster*. Some aspects of the phylogeny (Fig. 4) are also in agreement with the




Figure 6. Biogeographic distribution and ages of species mapped to tree shown in Figure 4 along with biogeographic distribution of ancestral nodes, calculated using modified version of Fitch parsimony described by Lieberman (2000), and minimum divergence age of nodes, determined using ghost lineage method described by Norell et al. (1992) and recently used by Guensburg et al. (2019). 0 = Western Europe; 1 = North Africa; 2 = South America; 3 = the WIS.

phylogeny presented by Stockley et al. (2005), although they included a much broader range of taxa. They also found *Mecaster* and *Hemiaster* to be closely related, although they did not retrieve a monophyletic Hemiasteridae. They also held *Toxaster* to be the earliest member of the spatangoids.

While the results from the analysis of Villier et al. (2004) cannot be directly compared with the analysis performed herein, as they focused on Early Cretaceous taxa and had fewer WIS representatives, as well as more representatives from outside of the WIS, some similarities emerge. In particular, Villier et al. (2004) also retrieved a clade consisting of *Heteraster oblongus*, “*Heteraster*” *texanus*, and *Washitaster riovistae* (Adkins, 1920) and found that a clade containing *Macraster elegans* was sister to two clades, one including Micrasteridae and the other including Hemiasteridae. They did not, however, find *Macraster* to be closely related to *Pliotoxaster comanchei* (Clark, 1915) (treated as *Palhemiaster* in their study).

Biogeographic patterns.—Smith (1984) summarized general patterns of echinoid biogeography and argued that during the Cretaceous, the present-day Mediterranean region (then Tethys) was a biodiversity hotspot for echinoids. The pattern of substantial within-region speciation in the WIS recovered herein suggests that the WIS might also have served as a biodiversity hotspot, at least among spatangoids. The recovery of episodes of range expansion between the WIS and South America matches patterns identified by Néraudeau and Mathey (2000) in their study of the biogeography of the Cretaceous of the present-day South Atlantic region. However, it is important to note that not all species of spatangoids could be considered herein. Changing the species sampled, or adding species, could potentially change the biogeographic character optimizations, which might alter the nature or number of episodes of range expansion, or vicariance; moreover, many species that undoubtedly existed are not preserved in the fossil record. Each of these will affect our ability to reconstruct biogeographic patterns (Turner et al., 2009). Thus, the biogeographic conclusions presented herein focus only on the available data.

Another relevant biogeographic study is that of Rosen and Smith (1988), who used parsimony analysis of endemism (PAE) on Campanian–Maastrichtian echinoids and found the WIS (in particular, a joint region composed of Texas and Mexico) to be the sister region to North Africa plus Europe. This matches the episodes of vicariance and range expansion between the WIS and Western Europe recovered herein. Further, Smith (1992) analyzed biogeographic patterns in Cenomanian echinoids using distance-based methods, cladistic biogeography, and PAE. The PAE indicated Brazil and Angola were closely related biogeographically and sister to the Texas region. Smith’s (1992) cladistic biogeographic analysis indicated that Texas could be the sister area of Western Europe and North Africa. Again, this matches some of the episodes posited herein for vicariance among spatangoids.

It is clear that sea-level changes would have had a major impact on the biogeography and evolution of echinoids (Smith and Rader, 2009). Indeed, the WIS evolved significantly during the time interval during which the taxa considered occurred. Multiple studies have been published on Cretaceous sea-level changes (e.g., Hallam, 1992; Haq, 2014). Prominent sea-level changes are associated with the following transgressive–regressive cycles seen in the North American Cretaceous carbonate platform series: the Coahuilan (Valanginian–mid-Aptian); the Comanchean (late Aptian–early Cenomanian); and the Gulfian (mid-Cenomanian–Maastrichtian) (Scott, 1993; Smith and Rader, 2009). The results from the biogeographic analysis presented herein suggest a possible connection between specific transgressive–regressive cycles and patterns of echinoid vicariance and range expansion. The regressions during the Coahuilan may coincide with the vicariance event that led to the diversification of the WIS “toxasterids,” including *Heteraster* and *Washitaster* (Fig. 6); the regression during the Comanchean may have led to the vicariance associated with the divergence at the base of all spatangoids, excluding the “toxasterids” (Fig. 6); another sea-level drop during the Gulfian may coincide with Campanian vicariance in the Micrasteridae (Fig. 6).

In addition, sea-level rise associated with the Comanchean could account for the range expansions associated with *Mecaster fourneli*, *Hemiaster bufo*, and “*Hemiaster*” *batnensis* (Fig. 6). However, a broader range of marine taxa will need to be considered to determine whether these are patterns that consistently emerge in other groups, thereby indicating biogeographic congruence and an overarching earth history control (e.g., Lieberman, 2000). If, instead, they do not, the patterns may be related to specific aspects of spatangoid ecology. At this time, it is not possible to determine the precise paleogeographic features that served as the specific geographic barriers that caused the episodes of vicariance when sea levels fell.

Acknowledgments

Thanks to L. Appleton and L. Boucher for their help providing access to collections at UT; M. Florence and K. Hollis for their help providing access to specimens and arranging the loan of material at USNM; and N. Ridgwell and S. Lucas for their help providing access to specimens and arranging the loan of material at NMMNH. Thanks to G. Heumann for providing photographs of holotypes at STIB, to T. Ewin for providing photographs of specimens at The Natural History Museum, London, and to J. Kimmig, who assisted with collections-related matters at KUMIP. B. Atkinson provided discussions on Bayesian phylogenetic approaches. C. Beard and L. Smith discussed various aspects of this research and commented on an earlier version of this manuscript. Thanks to A. Kroh, C. Souto, L. Zachos, and R. Mooi for comments on an earlier version of this manuscript. This research was made possible in part by a Panorama grant from the Biodiversity Institute, University of Kansas, and National Science Foundation grant DBI-1602067. In addition, this material is based on work supported while working at the National Science Foundation.

Accessibility of supplemental data

Data available from the Dryad Digital Repository: <https://doi.org/10.5061/dryad.6m905qfz2>.

References

- Adkins, W.S., 1920, The Weno and Paw Paw Formation of the Texas Comanchean: University of Texas Bulletin No. 1856, p. 101–117.
- Adkins, W.S., 1928, Handbook of Texas Cretaceous fossils: University of Texas Bulletin No. 2838, 384 p.
- Agassiz, L., 1836, Prodrôme d’une monographie des radiaires ou échinodermes: Mémoires de la Société des Sciences Naturelles de Neuchâtel, v. 1, p. 168–199.
- Agassiz, L., 1840, Description des échinodermes fossiles de la Suisse; seconde partie, Cidarides: Mémoires de la Société des Sciences Naturelles de Neuchâtel, v. 4, 107 p.
- Agassiz, L., and Desor, P.J.E., 1847, Catalogue raisonné des familles, des genres, et des espèces de la classe des échinodermes: Annales des Sciences Naturelles, Troisième Série, Zoologie, v. 7, p. 129–168.
- Arachchige, G.M., Jayakody, S., Mooi, R., and Kroh, A., 2019, Taxonomy and distribution of irregular echinoids (Echinoidea: Irregularia) from Sri Lanka: Zootaxa, v. 4541, p. 1–100.
- Berndt, R., 2003, Cenomanian echinoids from southern Jordan: Neues Jahrbuch für Geologie und Paläontologie, Monatshefte, v. 2003, p. 73–90.
- Blakey, R., 2014, Paleogeography and Paleotectonics of the Western Interior Seaway, Jurassic–Cretaceous of North America. http://www.search-and-discovery.com/pdfz/documents/2014/30392blakey/ndx_blakey.pdf.html [May 2019].
- Bremer, K., 1994, Branch support and tree stability: Cladistics, v. 10, p. 295–304.
- Brongniart, A., 1821, Sur les Caractères Zoologique des Formations: avec l’Application de ces Caractères à la Détermination de quelques Terrains de Craie: Paris, Madame Huzard, 38 p.
- Brongniart, A., 1822, Description Géologique des Environs de Paris: Paris, Chez G. Dufour et E. d’Ocagne Libraires, 475 p.
- Clark, H.L., 1917, Hawaiian and other Pacific echini: Memoirs of the Museum of Comparative Zoology, v. 46, p. 81–284.
- Clark, W.B., 1891, Revision of the Cretaceous Echinoidea of North America: Johns Hopkins University Circular, v. 10, no. 87, p. 75–77.
- Clark, W.B., 1893, The Mesozoic Echinodermata of the United States: Bulletin of the United States Geological Survey, v. 97, 207 p.
- Clark, W.B., 1915, The Mesozoic and Cenozoic Echinodermata of the United States: Monographs of the United States Geological Survey, v. 54, p. 9–100.
- Cole, S., 2019, Phylogeny and evolutionary history of diplobathrid crinoids (Echinodermata): Palaeontology, v. 62, p. 193–237.
- Cooke, C., 1946, Comanche echinoids: Journal of Paleontology, v. 20, p. 193–237.
- Cooke, C., 1953, American Upper Cretaceous Echinoidea: United States Geological Survey Professional Paper No. 254-A, p. 1–44.
- Cooke, C., 1955, Some Cretaceous Echinoids from the Americas: United States Geological Survey Professional Paper No. 264-E, p. 87–112.
- Coquand, H., 1862, Géologie et paléontologie de la région sud de la province de Constantine: Mémoires Société libre d’émulation de la Provence, Marseille, v. 2, 320 p.
- Cotteau, G., 1883, Échinides Jurassiques, Crétacés, et Tertiaires du sud-ouest de la France: Académie de La Rochelle, Société des Sciences Naturelles, 198 p.
- d’Orbigny, A., 1855, Paléontologie Française. Terrains Crétacés: Echinodermes, Échinoides irréguliers, v. 6, p. 97–272.
- Fischer, A., 1966, Order Spatangoida, in Moore, R.C., ed., Treatise on Invertebrate Paleontology, Part U, Echinodermata, Echinozoa-Echinoidea. Boulder, Colorado, and Lawrence, Kansas, Geological Society of America and University of Kansas Press, p. 367–695.
- Fourteau, R., 1907, Contribution à l’étude des échinides fossiles de la Craie Supérieure: Bulletin de l’Institut Égyptien, ser. 4, v. 6, no. 3, p. 140–175.
- Guensburg, T.E., Sprinkle, J., Mooi, R., Lefebvre, B., David, B., Roux, M., and Derstler, K., 2019, *Athenacrinus* n. gen. and other early echinoderm taxa inform crinoid origin and arm evolution: Journal of Paleontology, v. 94, p. 311–333.
- Hallam, A., 1992, Phanerozoic Sea-level Changes: New York, Columbia University Press, 266 p.
- Haq, B., 2014, Cretaceous eustasy revisited: Global and Planetary Change, v. 113, p. 44–58.
- Hendy, M., and Penny, D., 1982, Branch and bound algorithms to determine minimal evolutionary trees: Mathematical Biosciences, v. 59, p. 277–290.
- Hook, S.C., and Cobban, W.A., 2017, *Mecaster batnensis* (Coquand, 1862), a late Cenomanian echinoid from New Mexico, with a compilation of Late Cretaceous echinoid records in the Western Interior of the United States and Canada: Acta Geologica Polonica, v. 67, p. 1–30.
- Jagt, J.W.M., 2000, Late Cretaceous–early Paleogene echinoderms and the K/T boundary in the southeast Netherlands and northeast Belgium, Part 4, Echinoids: Scripta Geologica, v. 121, p. 181–375.
- Koch, N.M., Coppard, S.E., Lessios, H.A., Briggs, D.E.G., Mooi, R., and Rouse, G.W., 2018, A phylogenomic resolution of the sea urchin tree of life: BMC Evolutionary Biology, v. 18, p. 189.
- Kroh, A., 2020, Phylogeny and classification of echinoids, in Lawrence, J.M., ed., Sea Urchins: Biology and Ecology (fourth edition): Amsterdam, Elsevier, p. 1–17.
- Kroh, A., and Mooi, R., 2018, WoRMS Echinoidea: World Echinoidea Database. <http://www.marinespecies.org/echinoidea/> [May 2020].
- Kroh, A., and Smith, A.B., 2010, The phylogeny and classification of post-Palaeozoic echinoids: Journal of Systematic Palaeontology, v. 8, p. 147–212.
- Kroh, A., Lukender, A., and Gallemí, J., 2014, *Absurdaster*, a new genus of basal atelostomate from the Early Cretaceous of Europe and its phylogenetic position: Cretaceous Research, v. 48, p. 235–249.
- Lamarck, J.B.P.A. De M. De, 1816, Histoire Naturelle des Animaux sans Vertèbres, Présentant les Caractères, Généraux et Particuliers de ces Animaux, leur Distribution, leur Classes, leur Familles, leur Genres et la Citation Synonymique des Principales Espèces qui s’y Rapportent, Volume 3 (first edition): Paris, Chez Paul Renouard, 586 p.
- Lambert, J., 1895, Essai d’une monographie du genre *Micraster* et notes sur quelques échinides, in Grossouvre, A. de, Recherches sur la Craie Supérieure, pt. 1, fasc. 1: Paris, Imprimerie Nationale, p. 149–164.
- Lambert, J., 1916, Sur l’existence de l’étage Valanginien et sur l’oscillation barremienne dans l’Aube et dans l’Yonne: Mémoires de la Société Académique de l’Aube, v. 80, p. 19–94.
- Lambert, J., 1920, Sur quelques genres nouveaux d’échinides: Mémoires de la Société Académique d’Agriculture, des Sciences, Arts et Belles-lettres du Département de l’Aube, ser. 3, v. 55, p. 145–174.

- Lambert, J., 1927, Considérations sur les échinides de las Comanche Séries du Texas: Bulletin de la Société Géologique de France, ser. 4, v. 26, p. 263–278.
- Lieberman, B.S., 2000, Paleobiogeography: New York, Plenum Press/Kluwer Academic Publishers, 208 p.
- Lucas, S.G., 2000, A new species of the echinoid *Hemiaster* from the Lower Cretaceous (Albian) of southwestern New Mexico: New Mexico Museum of Natural History and Science Bulletin, v. 16, p. 101–106.
- Manso, C.L. de C., and Souza-Lima, W., 2017, *Mecaster texanus* (Echinoidea) from Turonian-Coniacian strata in the Sergipe-Alagoas Basin, Brazil: Carnets de Géologie, v. 17, p. 221–234.
- Martha, S.O., Taylor, P.D., and Rader, W.L., 2019, Early Cretaceous cyclostome bryozoans from the early to middle Albian of the Glen Rose and Walnut formations of Texas, USA: Journal of Paleontology, v. 93, p. 244–259.
- Meek, F.B., and Hayden, F.V., 1857, Descriptions of new species and genera of fossils collected by Dr. F.V. Hayden in Nebraska Territory, under the direction of Lieut. G.K. Warren, U.S. Topographical Engineers, with some remarks on the Cretaceous and Tertiary formations of the Northwest and the parallelism of the latter with those of other portions of the United States and Territories: Proceedings of the Philadelphia Academy of Natural Science, v. 9, p. 117–148.
- Melo, A.M., Sánchez, B.E.B., Marín, F.A.S., and Figueras, A.L., 2017, Equinoideos irregulares (Orden Spatangoida) del Cretácico de México: Revista de Biología Tropical, v. 65, suppl. 1, p. S113–S136.
- Mooi, R., 2001, Not all written in stone: interdisciplinary syntheses in echinoderm paleontology: Canadian Journal of Zoology, v. 79, p. 1209–1231.
- Mooi, R., Kroh, A., and Srivastava, D.K., 2014, Phylogenetic re-evaluation of fossil and extant micro-echinoids with revision of *Tridium*, *Cyamidia*, and *Lenicymidia* (Echinoidea: Clypeasteroidea): Zootaxa, v. 3857, no. 4, p. 501–526.
- Morgan, W.M., 2016, Collector's Guide to Texas Cretaceous Echinoids: Atglen, Schiffer, 160 p.
- Mortensen, T., 1950, A Monograph of the Echinoidea V (1). Spatangoida 1: Copenhagen, C.A. Reitzel, 432 p.
- Néraudeau, D., 1994, Hemiasterid echinoids (Echinodermata: Spatangoida) from the Cretaceous Tethys to the present-day Mediterranean: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 110, no. 3–4, p. 319–344.
- Néraudeau, D., and Mathey, B., 2000, Biogeography and diversity of South Atlantic Cretaceous echinoids: implications for circulation patterns: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 156, no. 1–2, p. 71–88.
- Norell, M., Novacek, M., and Wheeler, Q., 1992, Taxic origin and temporal diversity: the effect of phylogeny, in Extinction and Phylogeny: New York, Columbia University Press, p. 89–118.
- Oliveira, J., Manso, C.L. de C., Andrade, E.J., and Souza-Lima, W., 2013, O gênero *Mecaster* (Echinodermata: Spatangoida) do Cretáceo superior da Formação Jandaíra, Bacia Potiguar, Nordeste do Brasil: Scientia Plena, São Cristóvão, v. 9, no. 8, p. 1–17.
- Roemer, F., 1849, Texas: Mit besondere Rücksicht auf Deutsche Auswanderung und die Physischen Verhältnisse des Landes nach eigener Beobachtung Geschildert: Bonn, A. Marcus, 464 p.
- Roemer, F., 1852, Die Kreidebildungen von Texas und ihre Organischen Einschlüsse: Bonn, A. Marcus, 100 p.
- Roemer, F., 1888, *Macraster*, eine neue Spatangoiden-Gattung aus der Kriede von Texas: Neues Jahrbuch für Mineralogie, Geologie und Paläontologie, v. 1888, p. 191–195.
- Rosen, B., and Smith, A., 1988, Tectonics from fossils? Analysis of reef-coral and sea-urchin distributions from late Cretaceous to Recent, using a new method: Geological Society, London, Special Publications, v. 37, p. 275–306.
- Saucède, T., Mooi, R., and David, B., 2007, Phylogeny and origin of Jurassic irregular echinoids (Echinodermata: Echinoidea): Geological Magazine, v. 144, p. 333–359.
- Schlüter, C., 1870, Neue fossil Echiniden: Sitzungsberichte der Niederrheinischen Gesellschaft für Natur- und Heilkunde in Bonn: Bericht über Zustand der Gesellschaft während des Jahres, 1869, 132 p.
- Schlüter, C., 1900, Ueber einige Kreide-Echiniden: Zeitschrift der deutschen geologischen Gesellschaft, v. 52, p. 360–379.
- Scott, R.W., 1993, Cretaceous carbonate platform, U.S. Gulf Coast, in Simo, J.A.T., Scott, R.W., and Masse, J.-P., eds., Cretaceous Carbonate Platforms: Tulsa, American Association of Petroleum Geologists, p. 97–109.
- Shumard, B.F., 1853, Description of the species of Carboniferous and Cretaceous fossils collected, in Marcy's Natural History of the Red River of Louisiana. Appendix E: The 32d Congress, 2d Session, Executive Document No. 54, p. 197–211.
- Silva-Martínez, L.E., Blanco-Piñón, A., León-González, J.A.D., and Rodríguez-Vela, H., 2014, Equinoideos del Cretacio Tardío del Norte de Coahuila, Mexico: Boletín de la Sociedad Geológica Mexicana, v. 66, p. 377–395.
- Slocum, A.W., 1909, New echinoids from the Ripley group of Mississippi: Field Museum of Natural History Publication 134, Geology Series, v. 4, no. 1, p. 1–16.
- Smith, A.B., 1984, Echinoid Palaeobiology: London, Allen and Unwin, 190 p.
- Smith, A.B., 1992, Echinoid distribution in the Cenomanian: an analytical study in biogeography: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 92, no. 3–4, p. 263–276.
- Smith, A.B., and Bengtson, P., 1991, Cretaceous echinoids from north-eastern Brazil: Fossils and Strata, v. 31, p. 1–88.
- Smith, A.B., and Jeffery, C.H., 2000, Maastrichtian and Palaeocene Echinoids: A Key to World Faunas: London, The Palaeontological Association, 406 p.
- Smith, A.B., and Kroh, A., 2011, The Echinoid Directory—The Natural History Museum. <http://www.nhm.ac.uk/our-science/data/echinoid-directory/> [August 2019]
- Smith, A.B., and Rader, W., 2009, Echinoid diversity, preservation potential and sequence stratigraphical cycles in the Glen Rose Formation (early Albian, Early Cretaceous), Texas, USA: Palaeobiodiversity and Palaeoenvironments, v. 89, no. 1–2, p. 7–52.
- Smith, A.B., and Stockley, B., 2005, Fasciole pathways in spatangoid echinoids: a new source of phylogenetically informative characters: Zoological Journal of the Linnean Society, v. 144, p. 15–35.
- Smith, A.B., and Wright, C.W., 2008, British Cretaceous echinoids. Part 8. Atelostomata, 2. Spatangoida (1): Monograph of the Palaeontographical Society, v. 162, p. 569–635.
- Smith, A.B., and Wright, C.W., 2012, British Cretaceous echinoids. Part 9. Atelostomata, 2. Spatangoida (2): Monograph of the Palaeontographical Society, v. 166, p. 635–754.
- Stephenson, L.W., 1941, The larger invertebrate fossils of the Navarro group of Texas: University of Texas Bulletin No. 4104, 641 p.
- Stockley, B., Smith, A.B., Littlewood, T., Lessios, H., and MacKenzie-Dodds, J., 2005, Phylogenetic relationships of spatangoid sea urchins (Echinoidea): taxon sampling density and congruence between morphological and molecular estimates: Zoologica Scripta, v. 34, p. 447–468.
- Stolley, E., 1892, Die Kreide Schleswig-Holsteins: Mittheilungen aus dem Mineralogischen Institut der Universität Kiel, v. 1, p. 191–384.
- Swofford, D., 2003, PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods): Sunderland, Sinauer Associates.
- Thompson, J., Hu, S.-X., Zhang, Q.-Y., Petsios, E., Cotton, L., Huang, J.-Y., Zhou, C.-Y., Wen, W., and Bottjer, D., 2018, A new stem group echinoid from the Triassic of China leads to a revised macroevolutionary history of echinoids during the end-Permian mass extinction: Royal Society Open Science, v. 5, 171548.
- Thompson, W., 2016, Fossil Echinoids of Texas: A Monograph of Fossil Sea Urchins: Terre Haute, Indiana, River Styx Publishing, 225 p.
- Turner, A.H., Smith, N.D., and Callery, J.A., 2009, Gauging the effects of sampling failure in biogeographical analysis: Journal of Biogeography, v. 36, p. 612–625.
- Villier, L., Néraudeau, D., Clavel, B., Neumann, C., and David, B., 2004, Phylogeny of Early Cretaceous spatangoids (Echinodermata: Echinoidea): Palaeontology, v. 47, p. 265–292.
- Whitney, M., and Kellum, L., 1966, Echinoids of the Glen Rose Limestone of Texas: Papers of the Michigan Academy of Science, Arts and Letters, v. 51, p. 241–263.
- Wiley, E. 1979, An annotated Linnaean hierarchy, with comments on natural taxa and competing systems: Systematic Biology, v. 28, p. 308–337.
- Wiley, E., and Lieberman, B.S., 2011, Phylogenetics: Theory and Practice of Phylogenetic Analysis (second edition): Hoboken, Wiley-Blackwell, 406 p.
- Wright, A.M. 2019, A systematist's guide to estimating Bayesian phylogenies from morphological data: Insect Systematics and Diversity, v. 3, p. 1–14.
- Yavari, M., Yadzi, M., Gahalavand, H., Kamyabi Shadan, H., Adabi, M.H., and Villier, L., 2016, Early Cretaceous toxasterid echinoid *Heteraster* from the high Zagros basin, south of Iran: Carnets de Géologie, v. 16, p. 615–632.
- Zachos, L.G., 2017, Paleocene echinoid faunas of the eastern United States: Journal of Paleontology, v. 91, p. 1001–1024.

Accepted: 29 October 2020

Appendix

Apical disc

1. Plating: genital plate 2 does not extend between the posterior genital plates, referred to as ethmophract (0) / genital plate 2 extends between posterior genital and ocular plates, referred to as ethmolytic (1).
2. Position on test: posterior of midpoint (0) / at midpoint (1) / anterior of midpoint (2).

Frontal ambulacra (III)

3. Width is much narrower than (less than two-thirds the size of) central width of paired petals (0) / width is much broader than (more than 1.3 times the size of) central width of paired petals (1) / width is approximately equal to width of central width of paired petals (2).
4. Frontal ambulacra depth shallow (0) / frontal ambulacra depth substantial (1).
5. Distal end of frontal ambulacra tapers to form a shallow 'notch' present at ambitus (0) / distal end of frontal ambulacra does not form notch at ambitus (1) / very deep notch present at ambitus (2).
6. Pore pairs of the frontal ambulacra are uniform and uniserial (0) / pore pairs are heterogenous and not uniform (1).
7. Tubercles located between pore pairs in the poriferous zone: absent (0) / present (1).

Anterior paired ambulacra

8. Angle axis of petals forms relative to anterior/posterior axis through the apical disc: 6–14° (0) / 18–22° (1).
9. Relative size of pairs of pores on both axes: equally developed (0) / not equally developed (1).
10. Termination of paired petals: subpetaloid (0) / divergent (1) / in linear branches (2) / closed (3).

Posterior paired ambulacra

11. Relative size of pairs of pores on both axes: equally developed (0) / not equally developed (1).
12. Angle axis of petals forms relative to anterior/posterior axis through the apical disc: 2–7° (0) / 10–11° (1) / more than 14° (2).
13. Termination of paired petals: subpetaloid (0) / divergent (1) / in linear branches (2) / closed (3).

Fasciole

14. Subanal fasciole: absent (0) / present (1).
15. Loosely developed fasciole that does not fully encircle the test (peripetalous parafasciole): absent (0) / present (1).
16. Fully developed peripetalous fasciole: absent (0) / partial (fascioles are well defined and patterned but do not fully encircle the test) (1) / complete (2).
17. Multiple fasciole bands on the test: absent (0) / present (1).

18. Ambulacral plates: decrease in size at peripetalous fasciole (0) / do not decrease in size such that ambulacra are not 'pinched' (1).

Interambulacra

19. Interambulacra on aboral side: flush or level with paired ambulacra (0) / raised above ambulacra (1).

Periproct

20. In posterior view of the test, the periproct: positioned at the midline (0) / positioned above the midline (1). Note, this character can be influenced by ontogeny since in many spatangoids the periproct will migrate 'downward' as plates are added, and this analysis focused on adult specimens.

Plastron

21. Plates 2a and 2b: asymmetrical (protamphisternous) (0) / symmetrical (mesamphisternous) (1).

Peristome

22. Labrum: does not project over the peristome (0) / projects over the peristome (1).
23. Peristome shape: rounded (0) / pentagonal (1).
24. Peristome: not surrounded by apparent rim (0) / surrounded by apparent rim (1).
25. Peristome orientation from aboral view: flat (0) / oblique (1) / strongly oblique (2).

Test shape

26. From an adapical view, posterior end shape: rounded (0) / truncate (1). Note that one needs to pay careful attention with this character and consider a number of specimens as aspects of test shape can vary within species, although no specific polymorphism was found in this instance.
27. Anterior end in lateral view: has flat angular slope (0) / has inflated curvature (1).

Labrum

28. Posterior termination of labral plate extends to: second ambulacral plate (0) / third ambulacral plate (1) / fourth ambulacral plate (2).
29. Sternal suture in aboral view: straight or lightly curved (0) / concave (1).

Measurements

30. Anterior petal length: length to ambitus ratio: 0.5–0.55 (0) / 0.65–0.68 (1) / 0.72–1.0 (2). Note that for characters 30–32 it was possible to find distinct breaks between the different states such that they are discontinuous.
31. Posterior petal length: length to ambitus ratio: 0.3–0.47 (0) / 0.51–0.60 (1) / 0.65–1.0 (2).
32. Anterior ambulacra length/posterior ambulacra length ratio: 0.9–1.3 (0) / 1.54–1.9 (1) / 2.5–2.8 (2).