

## **Tall fescue interseeding and postemergence herbicides for false-green kyllinga (*Kyllinga gracillima*) control in turfgrass**

Authors: Elmore, Matthew T., and Tuck, Daniel P.

Source: Weed Technology, 36(4) : 548-552

Published By: Weed Science Society of America

URL: <https://doi.org/10.1017/wet.2022.51>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Research Article

**Cite this article:** Elmore MT and Tuck DP (2022) Tall fescue interseeding and postemergence herbicides for false-green kyllinga (*Kyllinga gracillima*) control in turfgrass. Weed Technol. 36: 548–552. doi: [10.1017/wet.2022.51](https://doi.org/10.1017/wet.2022.51)

Received: 24 January 2022

Revised: 25 April 2022

Accepted: 27 June 2022

First published online: 4 July 2022

**Associate Editor:**

Sandeep Singh Rana, Bayer Crop Science

**Keywords:**

Acetolactate synthase; ALS inhibitor; integrated pest management; PPO inhibitor; protoporphyrinogen oxidase; seeding; sulfonylurea; turfgrass

**Nomenclature:**

Carfentrazone; glyphosate; halosulfuron-methyl; imazosulfuron; sulfentrazone; false-green kyllinga, *Kyllinga gracillima* Miq.; tall fescue, *Lolium arundinaceum* (Schreb.) Darbysh

**Author for correspondence:**

Matthew Elmore, Assistant Extension Specialist, Rutgers University, Department of Plant Biology, 59 Dudley Road, New Brunswick, NJ 08901.  
E-mail: [matthew.elmore@rutgers.edu](mailto:matthew.elmore@rutgers.edu)

© The Author(s), 2022. Published by Cambridge University Press on behalf of the Weed Science Society of America. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



# Tall fescue interseeding and postemergence herbicides for false-green kyllinga (*Kyllinga gracillima*) control in turfgrass

Matthew T. Elmore<sup>1</sup>  and Daniel P. Tuck<sup>2</sup> 

<sup>1</sup>Assistant Extension Specialist, Rutgers University, Department of Plant Biology, New Brunswick, NJ, USA and <sup>2</sup>Field Researcher III, Rutgers University, Department of Plant Biology, New Brunswick, NJ, USA

**Abstract**

False-green kyllinga is a problematic  $C_4$  perennial Cyperaceae species. Previous research examined herbicide efficacy but has not integrated nonchemical practices with herbicides. Replicate field experiments were conducted to evaluate late-summer postemergence applications of herbicides in combination with turf-type tall fescue interseeding for false-green kyllinga control. Seven herbicide treatments and a nontreated control formed a complete factorial, with tall fescue interseeding or no seeding (eight-by-two factorial). Six herbicide treatments consisted of single postemergence applications of halosulfuron-methyl ( $70 \text{ g ha}^{-1}$ ), imazosulfuron ( $420 \text{ g ha}^{-1}$ ), and sulfentrazone + carfentrazone ( $280 + 30 \text{ g ha}^{-1}$ ) applied 4 wk before tall fescue interseeding (WBS) and the day of interseeding. Glyphosate ( $220 \text{ g ae ha}^{-1}$ ) applied the day of interseeding was the seventh herbicide treatment. Tall fescue was interseeded in September, and false-green kyllinga control was evaluated the following summer. In combination with tall fescue interseeding, imazosulfuron and halosulfuron applied 4 WBS as well as glyphosate applied the day of interseeding controlled false-green kyllinga better than all other treatments. In July, imazosulfuron and halosulfuron treatments applied 4 WBS and combined with interseeding had 1% to 6% false-green kyllinga cover, respectively, compared to 28% and 62% cover, respectively, without interseeding. Interseeding alone controlled false-green kyllinga <50%. Imazosulfuron, halosulfuron, and sulfentrazone + carfentrazone applied the day of seeding severely injured emerging tall fescue seedlings and reduced turfgrass quality the following spring. Applying imazosulfuron or halosulfuron in late summer and interseeding turf-type tall fescue 4 wk later or applying glyphosate the day of seeding are effective strategies for postemergence false-green kyllinga control. Integrating herbicides and the cultural practice of tall fescue interseeding provided more false-green kyllinga control than either practice alone.

**Introduction**

False-green kyllinga is a problematic  $C_4$  perennial sedge (Cyperaceae) species in turfgrass systems (Bryson et al. 1997). It has been reported in 15 US states from Connecticut to North and South Carolina and recently as far west as Nebraska (Bryson et al. 1997, USDA 2018; R. Gaussoin, personal communication). False-green kyllinga produces a dense mat of rhizomes that resembles desirable turfgrass and make it difficult to control with postemergence herbicides (McElroy et al. 2005).

Acetolactate synthase (ALS)-inhibiting herbicides halosulfuron and imazosulfuron as well as the protoporphyrinogen oxidase (PPO) inhibitor sulfentrazone control false-green kyllinga in cool-season turf (Elmore et al. 2019). Sequential postemergence herbicide applications usually control kyllinga better than single applications (Elmore et al. 2019; Gannon et al. 2012; McElroy et al. 2005). Imazosulfuron ( $\geq 420 \text{ g ha}^{-1}$ ) is more effective for false-green kyllinga control than halosulfuron-methyl ( $70 \text{ g ha}^{-1}$ ), which is more effective than sulfentrazone ( $140 \text{ g ha}^{-1}$ ) (Elmore et al. 2019). Similarly, in warm-season bermudagrass [*Cynodon dactylon* (L.) Pers.] turf, Gannon et al. (2012) found that single or sequential applications of sulfentrazone at 140 to  $420 \text{ g ha}^{-1}$  controlled false-green kyllinga <60%. Elmore et al. (2019) found that two sequential applications of imazosulfuron at 420 or  $740 \text{ g ha}^{-1}$  controlled false-green kyllinga 100% at all three locations, but single applications at  $420 \text{ g ha}^{-1}$  were less effective at certain locations (Elmore et al. 2019). There is no research to date that integrates herbicides and nonchemical strategies for postemergence *Kyllinga* spp. control.

Interseeding cool-season turfgrass in late summer is a common cultural practice for turfgrass management (Reicher et al. 2000). Combining herbicide applications with the cultural practice of interseeding tall fescue controlled perennial weeds dallisgrass (*Paspalum dilatatum* Poir.) and bermudagrass better than either strategy alone (Brosnan and Breeden 2013; Elmore et al. 2013). Similarly, the biocontrol *Sclerotinia minor* combined with interseeding a mixture of cool-season turfgrasses controlled dandelion (*Taraxacum officinale* F.H. Wigg.) better than the biocontrol or

interseeding alone (Abu-Dieyeh and Watson 2007). When interseeding is not integrated with an herbicide or other weed management practice, control of perennial weeds is inconsistent and weed control is poor (Abu-Dieyeh and Watson 2007; Elford et al. 2008; Jensen et al. 2014, 2017; Larsen et al. 2004; Larsen and Fischer 2005; Miller and Henderson 2012).

Previous research evaluated only mid- to late-spring applications of postemergence herbicides for false-green *kyllinga* control and without complementary cultural practices (Elmore et al. 2019; Gannon et al. 2012; McElroy et al. 2005). Late summer or early fall is the optimum time to seed cool-season turfgrasses such as tall fescue (Reicher et al. 2000). Tall fescue is well adapted throughout regions where cool-season turfgrass is grown because of its improved drought tolerance, insect pest tolerance, and persistence under low fertility (Bonos and Huff 2013). Therefore, the objective of this research was to evaluate the effect of tall fescue interseeding in combination with late-summer herbicide applications for false-green *kyllinga* control. We propose that combining the cultural practice of interseeding turf-type tall fescue with herbicide applications will control false-green *kyllinga* better than either seeding or herbicides alone.

## Materials and Methods

### Site

Replicate field experiments were conducted adjacently from 2019 to 2020 and 2020 to 2021 at Rutgers Hort Farm No. 2 in North Brunswick, NJ (40.47°N, 74.42°W) on a sandy loam soil with a pH of 5.7, 3.6% organic matter, and cation exchange capacity of 12 mEq 100 g<sup>-1</sup>. The site was infested with false-green *kyllinga* by sodding turf from an infested golf course fairway at Woodlake Country Club in Lakewood, NJ (40.09°N, 74.17°W) in June 2018. The sod was approximately 80% false-green *kyllinga* cover at harvest, with annual bluegrass (*Poa annua* L.) and creeping bentgrass (*Agrostis stolonifera* L.) the minority species. Foramsulfuron (40 g ha<sup>-1</sup>) was applied in July 2019 to suppress annual bluegrass and creeping bentgrass. The site was >95% false-green *kyllinga* cover by August 2019. The site was maintained at a 7-cm height of cut using a walk-behind rotary mower every 5 to 10 d during the growing season and irrigated as needed based on visible wilt to optimize tall fescue and false-green *kyllinga* growth before and during the experiments.

### Treatments

Seven herbicide treatments and a nontreated control were arranged in a complete factorial with tall fescue interseeding or no interseeding, forming a two-by-eight factorial treatment design. Treatments were applied in a strip-plot randomized complete block design with three replications. Tall fescue interseeding was the strip-plot factor. Whole plots (1.2 by 3.0 m) received herbicide treatments, which were single applications of sulfentrazone + carfentrazone at 280 + 30 g ha<sup>-1</sup> (Dismiss NXT; FMC Corp., Philadelphia, PA), halosulfuron at 70 g ha<sup>-1</sup> (Sedgehammer; Gowan Company, Yuma, AZ), or imazosulfuron at 420 g ha<sup>-1</sup> (Celero; NuFarm Americas Inc., Alsip, IL). These three herbicides were applied singly either 4 wk before seeding (WBS) based on label-recommended seeding intervals or the day of seeding, totaling six treatments. The seventh herbicide treatment was glyphosate (Credit 41 Extra; Nufarm Inc., Alsip, IL) at 220 g ae ha<sup>-1</sup> applied the day of seeding. Imazosulfuron and halosulfuron were applied with a nonionic surfactant at 0.25% v/v (Activator 90; Loveland

Products Inc., Loveland CO), according to the product label. This is the maximum labeled single application rate of halosulfuron and sulfentrazone + carfentrazone in cool-season turf and the lowest labeled single-application rate of imazosulfuron (Anonymous 2020, 2021a, 2021b). The highest labeled rate of halosulfuron and sulfentrazone + carfentrazone was selected, as single applications of these herbicides controlled false-green *kyllinga* <60% in previous research, and they are good candidates to be improved by interseeding (Elmore et al. 2019). The same rationale was used to select the lowest label rate of imazosulfuron, which controlled false-green *kyllinga* 78% to 95% in previous research compared to 89% to 99% control from the highest imazosulfuron rate (740 g ha<sup>-1</sup>), (Elmore et al. 2019). The 4-WBS treatments were applied on August 19, 2019 and August 27, 2020. The day of seeding herbicide treatments were applied on September 16, 2019 and September 24, 2020. Herbicide treatments were applied with water carrier at 420 L ha<sup>-1</sup> through a single AI11004EVS nozzle (Teejet; Spraying Systems Co., Glendale Heights, IL) using a hand-held CO<sub>2</sub>-pressurized (300 kPa) sprayer. Treatments were irrigated with 10 mm of irrigation 24 h after application to allow sufficient time for foliar absorption and move the herbicide into the soil in a controlled manner to prevent herbicide movement between plots.

Tall fescue interseeding and associated cultural practices occurred on September 16, 2020 and September 24, 2021, respectively, the same day as “day of seeding” herbicide applications. To prepare a seedbed for interseeding, a strip plot (1.5 m wide) across each block was core-cultivated using reciprocating 2.2-cm diam hollow tines on a 4.0-cm spacing (Toro® ProCore 648; The Toro Company, Bloomington, MN), then verticut twice (Ryan Ren-o-Thin® Power Rake; Schiller Grounds Care, Johnson Creek, WI) to create 0.6-cm-deep slits every 1.5 cm. A turf-type tall fescue blend of ‘Da Vinci’, ‘Leonardo’, and ‘Michelangelo’ (Winning Colors Turf Type Tall Fescue Blend; LebanonTurf, Lebanon, PA) was seeded at 400 kg ha<sup>-1</sup> (360 kg pure live seed ha<sup>-1</sup>) using a shaker jar inside a frame to prevent wind-aided seed drift. After seeding, plots were raked by hand perpendicular to the verticut slits using a leaf rake and then lightly irrigated. Day-of-seeding herbicide treatments were applied shortly after the foliage dried. Beginning 24 h after the day-of-seeding herbicide application, the experiment was irrigated daily until seedling emergence.

### Site Management After Treatment

Fertilizer (12 N : 24 P<sub>2</sub>O<sub>5</sub> : 8 K<sub>2</sub>O; 49 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>; Country Club® MD, 40% N derived from methylene urea; LebanonTurf, Lebanon, PA) was applied to the experiment 7 d after tall fescue seeding. In addition, 45 kg N ha<sup>-1</sup> was applied to the entire site on April 18, 2020, June 23, 2020, May 11, 2021, and July 6, 2021 (25 N : 0 P<sub>2</sub>O<sub>5</sub> : 5 K<sub>2</sub>O; Country Club®, 50% N derived from methylene urea; LebanonTurf, Lebanon, PA) to maintain tall fescue and *kyllinga* vigor throughout the spring and summer. Dithiopyr (Dimension® 2EW; Dow AgroSciences LLC, Indianapolis, IN) was applied to the entire site at 560 g ha<sup>-1</sup> in March 2020 and 280 g ha<sup>-1</sup> in April 2021 for preemergence crabgrass (*Digitaria* spp.) control. Spring dithiopyr application is a standard industry practice. A higher dithiopyr rate was used in 2020 because of COVID-19-related uncertainties about labor and farm management. Crabgrass control was excellent in both study years. Dithiopyr does not affect false-green *kyllinga* emergence from rhizomes but may have prevented false-green *kyllinga* establishment from seed. Azoxystrobin + difenconazole (470 + 300 g ha<sup>-1</sup>; Briskway™; Syngenta Crop Protection, LLC, Greensboro, NC)

was applied in late June and late July each year to control brown patch disease (caused by *Rhizoctonia solani*) of tall fescue. Clopyralid (Lontrel®; Dow AgroSciences LLC, Indianapolis, IN) was applied at 280 g ae ha<sup>-1</sup> on July 2, 2020 and July 14, 2021 for broadleaf weed control. Clopyralid was selected, as it does not reduce false-green kyllinga vigor (personal observation, M. Elmore). Glyphosate (Credit 41 Extra; Nufarm Inc., Alsip, IL) was applied at 220 g ae ha<sup>-1</sup> to the noninterseeded strips on February 24, 2020 and March 10, 2021 (during false-green kyllinga dormancy) for winter annual weed control.

### Data Collection

False-green kyllinga control was evaluated visually on a 0 (no control) to 100% (complete control) scale relative to the nontreated, nonseeded control in June, July, and August following herbicide applications—specifically, June 3, July 15, and August 15 in 2020 and June 17, July 19, and August 9 in 2021. Line-intersect counts were conducted July 15, 2020 and July 19, 2021 using a 0.8- by 0.9-m grid with 90 intersects. The presence or absence of false-green kyllinga under each intersect was recorded. Intersect count data were transformed to be expressed as percent false-green kyllinga cover. Tall fescue seedling injury was visually estimated at 3 wk after tall fescue seeding in both years on a 0 (no seedling stunting, chlorosis, or necrosis) to 100% (complete necrosis) scale relative to the nontreated, interseeded control. To detect meaningful residual effects of tall fescue seedling injury the following year when injury symptoms were no longer visible and turfgrass cover differences were minor, turfgrass quality ratings were conducted in May, June, and July each year. Turfgrass quality was assessed only in the seeded strips using the National Turfgrass Evaluation Program 1 to 9 scale, where 1 equaled low-quality turf, 9 equaled high-quality turf, and 6 was considered the minimum acceptable level of turfgrass quality (Morris and Shearman 1998). Green cover, density, uniformity, and to a lesser extent, color, were the primary components considered when estimating turfgrass quality. The presence or absence of false-green kyllinga was not considered when evaluating turfgrass quality, as false-green kyllinga displays desirable turfgrass characteristics during the warmer months.

### Statistical Analysis

Model assumptions were tested through residual analysis (Shapiro-Wilk statistic) in addition to visual assessments of Q-Q plots and histograms in SAS (Statistical Analysis Software, Inc., Cary, NC), and no transformations were required. The control data and intersect counts were analyzed in a strip-plot randomized complete block design ( $P = 0.05$ ). Seedling injury and turfgrass quality response variables related only to tall fescue seeding; thus, these variables were analyzed in a single-factor randomized complete block design with herbicide as the only fixed effect. For visible control and tall fescue seedling injury, nontreated control data were removed prior to analysis. Data were pooled across experiment years. ANOVA was performed using the mixed-model procedure in SAS, and Fisher's protected LSD test ( $P = 0.05$ ) was used to compare means (Saxton 2010). Herbicide treatment and tall fescue seeding were fixed effects, whereas year and block nested within year were considered random effects (Blouin et al. 2011). Year was a random effect, as the objective of the experiments was to estimate treatment effects over broad inference spaces (Blouin et al. 2011; Moore and Dixon 2015). To correlate grid

intersect count and visible data, the CORR procedure was conducted in SAS.

## Results and Discussion

### False-Green Kyllinga Control

Main effects of herbicide and interseeding were significant ( $P < 0.001$ ) for all visible control and line-intersect count data but are not presented. Herbicide-by-interseeding main-effect interactions were significant ( $P < 0.05$ ) for all visible control and line-intersect count data and are presented in Table 1. In June, both imazosulfuron treatments, halosulfuron applied 4 WBS, and glyphosate controlled false-green kyllinga similarly ( $\geq 86\%$ ), regardless of interseeding treatment (Table 1). By July, imazosulfuron and halosulfuron applied 4 WBS, as well as glyphosate controlled false-green kyllinga  $\geq 94\%$  in combination with seeding compared to  $\leq 77\%$  control without interseeding; in August, these treatments controlled false-green kyllinga  $< 40\%$  without interseeding and  $\geq 90\%$  with interseeding. In combination with interseeding, glyphosate, both imazosulfuron treatments, and halosulfuron applied 4 WBS controlled false-green kyllinga better than sulfentrazone + carfentrazone in July and August. Tall fescue interseeding improved false-green kyllinga control from every herbicide treatment in August. It is possible interseeding effects are overestimated by this research, as false-green kyllinga cover was  $> 95\%$  in August and other turfgrass species and weeds were suppressed before and during the experiments. In the authors' experience it is common to observe false-green kyllinga cover  $> 90\%$  in managed turf during late summer. Interseeding alone controlled false-green kyllinga 69% in June but only 23% by August. This indicates that false-green kyllinga will outcompete tall fescue during the summertime, even when tall fescue growth is managed with fertilizer to encourage density and fungicides to prevent brown patch disease (caused by *Rhizoctonia solani*). In the absence of interseeding, among treatments applied 4 WBS, imazosulfuron was more effective than halosulfuron, which was more effective than sulfentrazone + carfentrazone; the relative efficacy of these three herbicides aligns with Elmore et al. (2019).

False-green kyllinga cover as determined by line-intersect count align with visible-control estimates. For all herbicide treatments, tall fescue interseeding reduced false-green kyllinga cover compared to the herbicide alone. In combination with interseeding, both imazosulfuron treatments, halosulfuron applied 4 WBS, and glyphosate resulted in less false-green kyllinga cover ( $< 10\%$ ) than all other treatments (27% to 99%). Interseeding alone resulted in 57% false-green kyllinga cover compared to 95% for the nontreated control. False-green kyllinga cover as determined by intersect counts and control determined visually in July were highly correlated ( $r = -0.95$ ;  $P < 0.001$ ).

### Tall Fescue Seedling Injury

Imazosulfuron, halosulfuron, and sulfentrazone + carfentrazone applied the day of interseeding injured tall fescue seedlings substantially ( $\geq 55\%$ ) at 3 wk after interseeding (Table 2). Similarly, McFadden et al. (2022) found that halosulfuron (70 g ha<sup>-1</sup>) applied 0 or 7 d before seeding reduced tall fescue and Kentucky bluegrass establishment. This contrasts with Li et al. (2015), who reported no injury from sulfentrazone applied the day of seeding buffalograss [*Buchloe dactyloides* (Nutt.) Engelm.] and suggests that buffalograss is more tolerant to sulfentrazone than tall fescue at seeding. Sulfentrazone + carfentrazone applied 4 WBS caused minor



**Table 1.** False-green kyllinga control and cover following single late-summer applications of postemergence herbicides alone or in combination with tall fescue interseeding. Herbicide treatments were applied singly 4 wk before tall fescue interseeding (WBS; August 19, 2019 and August 27, 2020) or the day of interseeding (September 16, 2019 and September 24, 2020). Data presented are pooled across experiment years<sup>d</sup>.

			False-green kyllinga control <sup>a</sup>			
Herbicide timing	Herbicide	Tall fescue seeding	June	July	August	False-green kyllinga cover <sup>b</sup>
%						
4 WBS	Imazosulfuron <sup>c</sup>	No	95 a	77 bc	37 de	28 de
		Yes	99 a	99 a	98 a	1 f
	Halosulfuron	No	86 abc	49 de	20 f	62 b
		Yes	96 a	95 a	90 a	6 f
	Sulfentrazone + carfentrazone	No	46 e	7 f	3 g	89 a
		Yes	87 ab	73 bc	56 b	31 de
Day of interseeding	Imazosulfuron	No	98 a	85 ab	54 bc	27 e
		Yes	100 a	98 a	94 a	6 f
	Halosulfuron	No	48 e	7 f	2 g	97 a
		Yes	73 bcd	48 e	21 f	52 bc
	Sulfentrazone + carfentrazone	No	17 f	0 f	0 g	99 a
		Yes	72 cd	55 de	41 cd	44 cd
	Glyphosate	No	94 a	65 cd	22 f	51 bc
		Yes	97 a	94 a	90 a	9 f
N/A	None	No	.	.	.	95 a
		Yes	69 d	40 e	23 ef	57 bc
	P value		***	***	***	*

<sup>a</sup>False-green kyllinga control on a 0 (no control) to 100% (complete control) scale as determined by visual estimates relative to the nontreated control.

<sup>b</sup>Cover was determined by converting line-intersect count data to percent false-green kyllinga cover. Intersect counts were conducted on July 15, 2020 and July 19, 2021.

<sup>c</sup>Imazosulfuron (420 g ha<sup>-1</sup>) and halosulfuron (70 g ha<sup>-1</sup>) treatments included a nonionic surfactant at 0.25% (v/v). Sulfentrazone + carfentrazone (280 + 30 g ha<sup>-1</sup>) and glyphosate (220 g ae ha<sup>-1</sup>) were applied without a surfactant.

<sup>d</sup>Abbreviations: N/A, not applicable; \*, significant at  $P \leq 0.05$ ; \*\*\*, significant at  $P \leq 0.001$ .

**Table 2.** Tall fescue seedling injury and turfgrass quality following single late-summer applications of postemergence herbicides pooled across experiment years. Herbicide treatments were applied singly 4 wk before tall fescue interseeding (WBS; August 19, 2019 and August 27, 2020) or the day of interseeding (September 16, 2019 and September 24, 2020)<sup>d</sup>.

			Visual turfgrass quality score <sup>a</sup>		
Herbicide timing	Herbicide	Tall fescue seedling injury <sup>b</sup>	May	June	July
		%	Scale of 1 to 9		
4 WBS	Imazosulfuron <sup>c</sup>	13 cd	6.2 cd	7.1 ab	6.1 a
	Halosulfuron	0 d	7.3 ab	7.0 b	6.6 a
	Sulfentrazone + carfentrazone	16 c	6.7 bc	7.2 ab	7.4 a
Day of interseeding	Imazosulfuron	72 a	3.8 f	4.4 d	4.0 b
	Halosulfuron	55 b	5.6 de	7.2 ab	6.9 a
	Sulfentrazone + carfentrazone	85 a	5.0 e	6.3 c	7.4 a
N/A	Glyphosate	0 d	7.5 a	7.6 ab	6.5 a
	None	.	7.8 a	7.7 a	7.6 a
	P value	***	***	**	***

<sup>a</sup>Visual turfgrass quality ratings were determined visually using the National Turfgrass Evaluation Program 1 to 9 scale where 1 equaled low-quality turf, 9 equaled high-quality turf, and 6 was the minimum acceptable level of turfgrass quality.

<sup>b</sup>Injury was determined on a 0 (no seedling stunting, chlorosis, or necrosis) to 100% (complete necrosis) scale as determined by visual estimates relative to the nontreated control.

<sup>c</sup>Imazosulfuron (420 g ha<sup>-1</sup>) and halosulfuron (70 g ha<sup>-1</sup>) treatments included a nonionic surfactant at 0.25% (v/v). Sulfentrazone + carfentrazone (280 + 30 g ha<sup>-1</sup>) and glyphosate (220 g ae ha<sup>-1</sup>) were applied without a surfactant.

<sup>d</sup>Abbreviations: \*\*, significant at  $P \leq 0.01$ ; \*\*\*, significant at  $P \leq 0.001$ .

(16%) injury. Imazosulfuron applied 4 WBS caused 13% injury, but this was not statistically different from halosulfuron applied 4 WBS or glyphosate, which caused no injury.

### Turfgrass Quality

Trends in tall fescue injury at 3 wk after interseeding were apparent in turfgrass quality the following year. In May, imazosulfuron, halosulfuron, and sulfentrazone + carfentrazone applied the day of interseeding reduced turfgrass quality to <6.0. Among treatments applied 4 WBS, imazosulfuron and sulfentrazone + carfentrazone reduced turfgrass quality in May compared to the nontreated control. By June, only imazosulfuron and sulfentrazone + carfentrazone applied the day of interseeding reduced turfgrass quality. In

July, only imazosulfuron applied the day of interseeding reduced turfgrass quality compared to the nontreated control. It is possible that the spring broadcast application of dithiopyr slowed tall fescue seedling recovery from autumn injury, but tall fescue plants in 4-WBS treatments were mature and not displaying signs of herbicide injury when the dithiopyr application was made. We did not observe any injury during the spring or summer following the dithiopyr application.

Seedling injury and subsequent turfgrass quality reductions observed from imazosulfuron, halosulfuron, and sulfentrazone + carfentrazone applied the day of interseeding aligns with product labels that restrict seeding within 4 wk of application (Anonymous 2020, 2021a, 2021b). The transient turf quality reduction during springtime caused by imazosulfuron applied 4

WBS is noteworthy, as the lowest labeled rate (420 g ha<sup>-1</sup>) was applied. Considering the imazosulfuron half-life of 21 to 70 d in aerobic soils (Mikata et al. 1996; Morrica et al. 2001, 2002), this turf quality reduction suggests substantial reductions in tall fescue establishment are possible at the maximum labeled rate of 740 g ha<sup>-1</sup>, even if the label recommendations of seeding no sooner than 4 wk after application are heeded. Future research should evaluate imazosulfuron safety to tall fescue and other cool-season turfgrass species commonly seeded to lawns. Halosulfuron should be included for comparison to this and previous research of McFadden et al. 2022. With a half-life of 7 to 28 d, halosulfuron is less persistent than imazosulfuron in aerobic, slightly acidic soils (Kuwatsuka and Yamamoto 1997).

Applying imazosulfuron (420 g ha<sup>-1</sup>) or halosulfuron (70 g ha<sup>-1</sup>) in late summer followed by interseeding turf-type tall fescue as soon as 4 wk later is an effective strategy for selective postemergence false-green kyllinga control in cool-season turfgrass; glyphosate applied the day of interseeding is an effective nonselective option. Herbicides or interseeding alone provided poor false-green kyllinga control the following August, nearly 1 yr after herbicide application. Integrating herbicides and cultural practices to provide a competitive turf sward is critical for effective postemergence false-green kyllinga control with single-herbicide applications. Future research should investigate tall fescue seeding in autumn followed by postemergence herbicide applications in late spring.

**Acknowledgments.** This research was supported by funding from the Rutgers Center for Turfgrass Science. The authors would like to thank Christian Pucci, Christine Kochmer, Brandon McNally, and Katie Diehl for their assistance with this research.

The authors declare no conflicts of interest.

## References

- Abu-Dieyeh MH, Watson AK (2007) Grass interseeding and a fungus combine to control *Taraxacum officinale*. *J Appl Ecol* 44:115–124
- Anonymous (2020) Dismiss NXT herbicide product label. FMC Corp. Philadelphia, PA: FMC. 6 p
- Anonymous (2021a) Celero herbicide product label. NuFarm Americas Inc. Alsip, IL: NuFarm. 25 p
- Anonymous (2021b) Sedghammer herbicide product label. Gowan Company. Yuma, AZ: Gowan. 7 p
- Blouin DC, Webster EP, Bond JA (2011) On the analysis of combined experiments. *Weed Technol* 25:165–169
- Bonos and Huff (2013) Cool-season grasses: biology and breeding. Pages 591–660 in Stier et al., eds, *Turfgrass: Biology, Use and Management*. Agronomy monograph 56. Madison, WI: Crop Sci Soc of America
- Brosnan JT, Breeden GK (2013) Bermudagrass control with topramezone and triclopyr. *Weed Technol* 27:138–142
- Bryson CT, Carter R, McCarty LB, Yelverton FH (1997) *Kyllinga*, a genus of neglected weeds in the continental United States. *Weed Technol* 11:838–842
- Elford EMA, Tardif FJ, Robinson DE, Lyons EM (2008) Effect of perennial ryegrass overseeding on weed suppression and sward composition. *Weed Technol* 22:231–239
- Elmore MT, Brosnan JT, Mueller TC, Horvath BJ, Kopsell DA, Breeden GK (2013) Seasonal application timings affect dallisgrass (*Paspalum dilatatum*) control in tall fescue. *Weed Technol* 27:557–564
- Elmore MT, Patton AJ, Tuck DP, Murphy JA, Carleo J (2019) False-green kyllinga control in cool-season turfgrass. *Weed Technol* 33:329–334
- Gannon TW, Yelverton FH, Tredway, LP (2012) Purple nutsedge (*Cyperus rotundus*) and false-green kyllinga (*Kyllinga gracillima*) control in bermudagrass turf. *Weed Technol* 26:61–70
- Jensen AMD, Bühler O, Kvalbein A, Aamlid T (2017) Evaluation of the occurrence of turfgrasses and weeds after repeated overseeding on fairways. *Int Turfgrass Soc Res J* 13:389–393
- Jensen AMD, Petersen KN, Aamlid T. (2014) Pesticide-free management of weeds on golf courses: current situation and future challenges. *Eur J Turfgrass Sci* 42:61–64
- Kuwatsuka DS, Yamamoto I (1997) Degradation of the herbicide halosulfuron-methyl in two soils under different environmental conditions. *J Pestic Sci* 22:282–287
- Larsen SU, Fischer J (2005) Turfgrass management and weed control on golf course fairways without pesticides. *Int Turfgrass Res Soc J* 10: 1213–1221
- Larsen SU, Kristofferson P, Fischer J (2004) Turfgrass management and weed control without pesticides on football pitches in Denmark. *Pest Manag Sci* 60:579–587
- Li L, Sousek MD, Reicher Z (2015) Herbicides applied at or shortly after seeding are effective for weed control in seedling buffalograss. *Crop Forag Turfgrass Mgmt* 1, doi: [10.2134/cftm2015.0150](https://doi.org/10.2134/cftm2015.0150)
- McElroy JS, Yelverton FH, Warren Jr. LS (2005) Control of green and false-green kyllinga (*Kyllinga brevifolia* and *K. gracillima*) in golf course fairways and roughs. *Weed Technol* 19:824–829
- McFadden D, Fry J, Keeley S, Hoyle J, Raudenbush Z (2022) Establishment of Kentucky bluegrass and tall fescue seeded after herbicide application. *Crop Forage Turfgrass Mgmt* 8, doi: [10.1002/cft2.20151](https://doi.org/10.1002/cft2.20151)
- Mikata K, Yamamoto A, Tashiro S (1996) Degradation of imazosulfuron in flooded soils. *J Pestic Sci* 21:171–177
- Miller NA, Henderson JJ (2012) Organic management practices on athletic fields: part 1. the effect on color, quality, cover, and weed population. *Crop Sci* 52:890–903
- Moore KJ, Dixon PM (2015) Analysis of combined experiments revisited. *Agron J* 107:763–771
- Morrica P, Fidente P, Seccia S, Ventrigilia M (2002) Degradation of imazosulfuron in different soils—HPLC determination. *Biomed Chromatography* 16:189–494
- Morrica P, Giordano A, Seccia S, Ungard F, Ventrigilia M (2001) Degradation of imazosulfuron in soil. *Pest Manag Sci* 57:360–365
- Morris KN, Shearman RC (1998) NTEP Turfgrass Evaluation Guidelines. National Turfgrass Evaluation Program Workshop. Beltsville, MD. Retrieved from <https://ntep.org/pdf/ratings.pdf>
- Reicher ZJ, Throssell CS, Weisenberger DV (2000) Date of seeding affects establishment of cool-season turfgrasses. *HortSci* 35:166–1169
- Saxton AM (2010) DANDA.sas: Design and analysis macro collection version 2.11. Knoxville, TN: University of Tennessee. Accessed: November 18, 2018
- [USDA] United States Department of Agriculture (2018) Plants Profile, Natural Resources Conservation Service. <https://plants.usda.gov/core/profile?symbol=KYGR>. Accessed: January 15, 2022