

Road orientation affects the impact of roads on wildlife

Authors: Mulero-Pázmány, Margarita, Rollán, Laura, D'Amico, Marcello, and González-Suárez, Manuela

Source: Wildlife Research, 50(1) : 39-46

Published By: CSIRO Publishing

URL: <https://doi.org/10.1071/WR21149>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Road orientation affects the impact of roads on wildlife

Margarita Mulero-Pázmány^{A,B,*} , Laura Rollán^C, Marcello D'Amico^{D,E}  and Manuela González-Suárez^F 

For full list of author affiliations and declarations see end of paper

***Correspondence to:**

Margarita Mulero-Pázmány
Department of Animal Biology, Facultad de Ciencias, University of Málaga, 29071 Málaga, Spain
Email: muleromara@uma.es

Handling Editor:

Tom Sullivan

ABSTRACT

Context. Understanding the factors determining the impacts of roads and how they fragment landscapes limiting the movement of animals, is key to implement efficient mitigation measures.

Aims. Here we investigate if road orientation in relation to limiting resources, a largely overlooked factor on road impact assessments, can influence the movement of animals within a landscape where water resources are spatially clustered. **Methods.** We evaluated movement by monitoring animal tracks on unpaved roads: two with a North–South orientation and two with an East–West orientation. **Key results.** Animals were more likely to follow roads leading to limiting resources (i.e. East–West orientation), confirming human linear structures can facilitate wildlife movements. Carnivores were more likely to follow roads with any orientation and for longer compared to ungulates, whereas ungulates followed roads mainly in the orientation of limiting resources. **Conclusions.** Road orientation affects how roads influence the movement of animals in landscapes where resources are distributed along a spatial gradient with different effects for ungulates and carnivores. **Implications.** The key implications of this work affect the planning and implementation of mitigation strategies and safety measures. Our results suggest road-crossing infrastructure and fences will be most important in roads traversing a gradient to allow wildlife movement while preventing collisions. For roads along a gradient, crossing structures may be less important, but fences or appropriate signage could be useful to prevent or warn drivers of animals travelling on the road.

Keywords: animal movement, habitat fragmentation, landscape connectivity, mitigation, road ecology, roadkill, track census, wildlife collisions.

Introduction

Roads fragment landscapes, contribute to ecosystem degradation and threaten biodiversity (Bennett 2017). These linear structures are expected to increase by more than 60% globally by 2050 (Meijer *et al.* 2018), therefore, understanding the factors determining their impact on wildlife before they are designed and built is crucial to minimise environmental impact and reduce human safety risks along with the financial implications of wildlife collisions (Díaz-Varela *et al.* 2011; Morelle *et al.* 2013; Rytwinski *et al.* 2016; Visintin *et al.* 2018; Ascensão *et al.* 2019a). The most studied road impacts for wildlife are roadkill and fragmentation, both consequences of an animal choice to cross or avoid a road (Grilo *et al.* 2012). Their interaction produces the so-called barrier effect, which contributes to subdivide and isolate animal populations (Grilo *et al.* 2012) eventually compromising their viability (Ceia-Hasse *et al.* 2017). How roads affect wildlife depends on which species are in an area, their characteristics, and their local abundance (Jacobson *et al.* 2016; Visintin *et al.* 2016; González-Suárez *et al.* 2018; Ascensão *et al.* 2019a; Duffett *et al.* 2020), as well as on traffic and road properties (e.g. road width, traffic speed) and the characteristics of the surrounding environment (Borkovcová *et al.* 2012; D'Amico *et al.* 2015; Visintin *et al.* 2016). While diverse factors related to road properties and surrounding environment have been studied, there is still a gap in the literature regarding how the spatial properties of roads, such as their orientation in relation with limiting resources for animals, can influence how roads impact wildlife.

Received: 12 October 2021

Accepted: 16 April 2022

Published: 14 July 2022

Cite this:

Mulero-Pázmány M *et al.* (2023)
Wildlife Research, **50**(1), 39–46.
doi:[10.1071/WR21149](https://doi.org/10.1071/WR21149)

© 2023 The Author(s) (or their employer(s)). Published by CSIRO Publishing.

This is an open access article distributed under the Creative Commons Attribution 4.0 International License (CC BY).

OPEN ACCESS

The objective of this work was to fill this gap evaluating how a largely overlooked aspect of road design, orientation, affects the impact of these linear structures on wildlife. To achieve this goal, we conducted a study within a Mediterranean ecosystem, in which water is a limiting resource that shows strong spatial aggregation. Road orientation can be important because limiting resources for animals, such as water, are often associated to geographical features (e.g. rivers, mountain chains or coastlines) that form landscape gradients. These gradients result in periodic movements of animals to and from resources that may occur across different spatio-temporal scales, from individual daily movements connecting water/refuge/food to large seasonal migrations to track fresh pastures. Therefore, we predict that in landscapes with a resource gradient, such as our study area, animals will respond differently to roads that run along the gradient compared to those that intersect it. In general, we expect that animals could either avoid completely, or alternatively use as movement corridors, roads that run along the gradient, while roads that intersect the gradient, i.e. blocking their access to the resources, will have to be regularly crossed but will rarely be used as travel corridors.

Additionally, we anticipate that different species groups will be influenced differently by road orientation. To test this possibility, we studied two groups of mammals: large herbivores and carnivores for which we predict different responses. Herbivores, namely ungulates, are dependent on primary productivity and regularly move across the landscape water gradient, but previous work suggest they may avoid roads (D'Amico *et al.* 2016). Conversely, carnivores' movements are less influenced by water and pasture resources and previous work suggests they can be attracted to roads and use these for travelling and marking (Whittington *et al.* 2011). Thus, we predict carnivores will use and travel on roads more often and be less affected by their orientation as their

movement across the landscape is not strongly directional. Large herbivores, which are those most often involved in deadly collisions (Bissonette *et al.* 2008), are expected to cross roads traversing the gradient and could either avoid or travel along those roads that link resources.

Our study provides new understanding regarding how wildlife is affected by road design, in particular orientation within the landscape. When designing a road, its orientation will largely be determined by the need to connect areas and relevant socioeconomic factors, yet, even if overall orientation is unlikely to be flexible, mitigation measures and safety guidance (e.g. speed limits and signage) could be adapted to reduce impact to wildlife and humans. Understanding how orientation influences the movement of wildlife can anticipate the probability of animals crossing roads or moving along them, behaviours which pose different challenges to wildlife and human safety. Here we offer insights to promote more effective mitigation strategies, with implications for sustainable road management and wildlife conservation.

Materials and methods

Study area

Our study was carried out in Doñana Biological Reserve, a management unit of 6794 ha with the highest level of environmental protection in the core of the Doñana Natural Area. The Doñana Natural Area (36°59'N, 6°26'W) comprises both National and Natural Parks, and is located in the Guadalquivir river estuary, in the Atlantic coast of South-western Spain (Fig. 1). The climate is Mediterranean sub-humid with marked seasons (i.e. hot dry summer and mild winter, with precipitations mainly during winter and spring). Doñana Biological Reserve hosts three main ecosystems: sandy dunes; Mediterranean shrubland/forest;




Fig. 1. Study area in Doñana National Park. Dark grey rectangles indicate the four sampled trails. The dashed black line indicates the ecotone, with water resources located towards East.

and marshland in the eastern area of the reserve (Fig. 1). During summer (June–September) most available water and pastures are concentrated along the ecotone between Mediterranean shrubland/forest and marshland, that runs mainly in a North–South direction within the Biological Reserve. Ungulate species in this area include wild boar *Sus scrofa*, feral cattle *Bos taurus*, red deer *Cervus elaphus*, feral horse *Equus caballus*, and fallow deer *Dama dama* (listed in approximate order of abundance; [Castroviejo 1993](#)). Carnivore species include red fox *Vulpes vulpes*, large grey mongoose *Herpestes ichneumon*, European badger *Meles meles*, feral dog *Canis lupus familiaris*, small-spotted genet *Genetta genetta*, feral cat *Felis silvestris catus*, Iberian lynx *Lynx pardinus*, Eurasian otter *Lutra lutra*, and European wildcat *Felis silvestris* (listed in approximate order of abundance; [Castroviejo 1993](#)). Doñana Natural Area has a heterogeneous road-network mostly composed by unpaved sandy roads that were built according to the transportation needs for humans (i.e. park management, historical reasons), with low traffic volume (~10 vehicles/day; [Román et al. 2010](#)).

Data collection and variables definition

Within Doñana Biological Reserve, we selected a section of 50 m length in each of four unpaved roads (with sandy soil) close to the ecotone between Mediterranean shrubland/forest and marshland. Two of the surveyed roads run North–South (2° and 4°) perpendicular to the landscape water gradient. The other two roads run East–West (82° and 89°) in parallel with landscape water gradient (Fig. 1).

During 10 days in June–July 2019 we monitored ungulate and carnivore tracks on the selected road sections daily. The day before starting the data collection, we swept the sand in each road section using a heavy piece of wood pulled by a 4 × 4 vehicle to create a clean and homogeneous surface. Then, each road section was swept with a broom daily after tracks were recorded to avoid double counting them on subsequent days. Every sampling day, soon after sunrise, we surveyed the four road sections to record tracks starting each day on a different road section. For each track found we recorded: the species (when possible) or *species group* (i.e. ungulates or carnivores), the angle of the track where entering the road section and the angle where exiting it (recorded with a compass), *track length* within the road section (measured in metres with a tape), and described the *speed* of the animal that made the track as gait type: walk, trot, gallop. For each road section, we also visually estimated and recorded the percentage of *shrub coverage* in the band of 3 m adjacent to the roads because vegetation may affect animal movement.

Because not all tracks could be identified to species, our analyses focused on the two *species groups* (ungulates and carnivores). Track *speed* was also reclassified in two levels for analyses: slow for tracks that only showed gait = walk,

and fast for all others. Track *movement type* was classified in two levels based on the entrance and exit angles and the road orientation: cross for tracks where the animal had entered the road from one side and exited from the other (regardless of track length) and follow for tracks of >3 m in length where entrance and exit occurred from the same side of the road or were beyond the 50 m section (entrance or exit angle were not recorded). We excluded nine tracks with a length <3 m that we could not clearly classify as ‘cross’ or ‘follow’. Finally, track *movement direction* was defined as the travel direction (e.g. North–South or South–North) within the road for tracks with movement type = follow.

Data analyses

We fitted logistic mixed effects regression models using the *glmer* function from the *lme4* package in R ([R Foundation for Statistical Computing 2013](#)) to:

1. Predict the *movement type* (cross vs follow) of each observed track as a function of three predictors: *road orientation*, *species group* and estimated *shrub coverage* in the adjacent terrain of each road.
2. Predict the *speed* (fast vs slow) of each observed track as a function of three predictors: *road orientation*, *species group* and *movement type*, (we expected animals crossing to move faster than those following roads).
3. Predict the *movement direction* for animals following roads (*movement type* = follow) for each *road orientation* (one model for each orientation) as function of two predictors: *species group* and *speed*.
4. Predict the *track length*, a proposed proxy for the duration of road travel, for animals following roads (*movement type* = follow) as a function of four predictors: *road orientation*, *species group*, *movement speed* and *shrub coverage*.

In models that included *road orientation* and *species group* as predictors, we tested for an interaction between these two predictors to explore differential responses. We report interaction terms if these were significant and otherwise report simpler additive models. In all fitted models we used observation *date* nested within *transect ID* (unique identifier for each of road sections) as a random factor to control for temporal and spatial non-independence of the track data.

Results

We analysed a total of 289 tracks (Table 1). Of these, 264 were made by ungulates including 24 by feral species (horses and cattle), 240 by wild species (wild boars, red and fallow deer) and 25 by carnivores (foxes, badgers, felids and other unidentified meso-carnivores). We found 132 tracks in

Table 1. Recorded tracks by animal species groups, movement type (cross vs follow) and road orientation.

Species group	Cross			Follow		
	East–West	North–South	Total	East–West	North–South	Total
Ungulates	42	114	156	82	26	108
Carnivores	2	5	7	6	12	18
Total	44	119	163	88	38	126

East–West roads and 157 in North–South roads. A total of 163 tracks crossed the road and 126 followed its trajectory.

Animals moved differently on roads with different orientations being over five times more likely to follow and three times less likely to move fast on East–West roads compared to those with North–South orientation (Table 2, Figs 2, 3). Species differed in their use of roads, with carnivores being over seven times more likely to follow roads than ungulates and having a tendency (not significant) to move slower. However, groups did not differentially respond to road orientation: the interactions between *road orientation* and *species group* were not significant for *movement type* or *speed*. Finally, roads were used differently depending on the adjacent terrain and animals were more likely to follow roads in areas with more shrub coverage (Table 2).

There were no differences in *movement direction* associated with *species group* or *movement speed* (Table 3). However, several factors influenced *track length*, and thus, presumably the time animals followed roads. In particular, tracks were longer in North–South roads, if they belonged to carnivores, and when left by animals moving faster (Table 4, Fig. 3).

Table 2. Coefficient estimates (odds ratios (OR)) for models testing factors that influence how animals move on road (movement type and speed).

Variables	OR	95% CI
Movement type: follow/cross		
Intercept	0.348	0.106–1.353
Road orientation: East–West	5.189	2.268–13.117
Species group: ungulates	0.139	0.034–0.398
Shrub coverage	1.037	1.024–1.053
Speed: fast/slow		
Intercept	1.195	0.250–5.007
Road orientation: East–West	0.266	0.055–0.932
Species group: ungulates	0.299	0.079–1.073
Movement type: follow	0.477	0.171–1.195

We report the odds ratios for best estimates (OR) and their 95% bootstrapped confidence intervals. Predictors with 95% CI non-overlapping with one are highlighted in bold. Landscape direction is East–West.

Discussion

As predicted, our results showed that the movement of animals in relation to roads varies depending on the orientation of these structures, at least in landscapes where resources are spatially distributed along a gradient. We also found that ungulates and carnivores make different use of these structures, which suggests these groups could face different roadkill risk and fragmentation impacts.

Animals were more likely to follow roads along the environmental gradient that could be used to move to and from a key resource, which in this study area is freshwater. In semi-arid environments, wildlife depends highly on limited water (Kihwele *et al.* 2020), and many species travel daily to and from water and associated resources such as fresh pastures (Kasiringua *et al.* 2017). Our findings suggest that the animals may be taking advantage of existing roads for these movements, at least in the dry season when water is scarce. Several studies have described wildlife using roads, including some of the species we detected (e.g. wild boars and red foxes) (Macdonald 1979; Boughton *et al.* 2019). Roads can offer opportunities for foraging and marking behaviour (Monclús *et al.* 2009; Toger *et al.* 2018) and facilitate movement because they often present a flatter surface without vegetation or snow (Whittington *et al.* 2005; Brown *et al.* 2006). Indeed, we found that animals were more likely to follow unpaved roads in areas surrounded by more shrub coverage where movement may be hampered and left longer tracks when travelling faster because when flatter, obstacle-free terrain may be most important. However, wildlife may not use roads if these are heavily used by humans. In fact, wildlife are more likely to take advantage of roads with low traffic volume (Brown *et al.* 2006; Boughton *et al.* 2019), which is the case for our study area (10 vehicles/day; Román *et al.* 2010). Similarly, wildlife can travel along heavily trafficked roads and associated roadsides during low-traffic periods, such as night-time for certain roads (Brown *et al.* 2006).

As hypothesised, the two species groups differed in their use of roads, with carnivores over seven times more likely to follow roads and doing so for longer compared to ungulates. Carnivore species have been shown to use roads for diverse purposes, including as travel corridors (Whittington *et al.* 2005), for territorial displays (Monclús *et al.* 2009), for hunting (Andersen *et al.* 2017) and




Fig. 2. Model predictions (median shown as black symbols and 95% CI as error bars) showing changes in the probability of movement type being 'follow' for roads with different orientations and for different species groups. Plots also display the observed occurrence of 'follow' movements (grey circles positioned at $y = 1$) and cross movements (grey circles at $y = 0$) for each road orientation and species group. The size of the grey circles is proportional to the number of observations (e.g. we observed more 'follow' movement on East-West than on North-South roads).




Fig. 3. Model predictions (median shown as small symbols with 95% CI error bars) showing the probability of movement speed being fast (left panel) and estimated track length for movement type = 'follow' (right panel) across species group and roads with different orientations. Plots also display the observed occurrence of fast movements (circles positioned at $y = 1$) and slow movements (circles at $y = 0$) in the left panel and the observed track lengths (right panel) for each road orientation and species group. The size of these circles is proportional to the number of observations. There were more slow than fast movements on ungulates following East-West roads.

scavenging (Barrientos *et al.* 2018). In our study area, red foxes are known to actively select roads for defecation, probably as a marking behaviour (Suárez-Esteban *et al.* 2013). Conversely, prey species, such as ungulates in our case study, are more likely to exhibit road-gap avoidance as part of their anti-predator behaviour (Laurance *et al.* 2004; Chen and Koprowski 2016). In agreement with a previous study in this area (D'Amico *et al.* 2016), we found that ungulates do sometimes use roads, with a tendency to follow East-West roads more than those oriented North-South. This suggests road use may be an acceptable risk when there is a benefit such as easier travel to and from limiting resources.

Previous studies have evaluated how road design and some related characteristics influence wildlife impacts. Road design depends on road alignment, profile and section (Garber and Hoel 2019). Regarding road alignment, both the presence of curves and road sinuosity can affect roadkill risk (Grilo *et al.* 2011; D'Amico *et al.* 2015). Similarly, regarding road profile, road slope can have an effect on roadkill probability (D'Amico *et al.* 2015; Kang *et al.* 2016). Many road features related to road section have been often considered as predictors of different road impacts, such as roadkill and fragmentation, for example road size and the presence of road shoulders or ditches (Matos *et al.* 2012; D'Amico *et al.* 2015; Jacobson *et al.* 2016). Our study is the

Table 3. Coefficient estimates for models testing factors that influence the *movement direction* in animals that follow roads (*movement type* = follow).

Variables	β /OR	95% CI
Direction in East–West roads		
Intercept	0.828	0.050–11.302
Species group: ungulates	0.894	0.121–5.417
Speed: slow	1.247	0.189–7.962
Direction in North–South roads		
Intercept	1.354	0.258–11.420
Species group: ungulates	0.447	0.036–2.376
Speed: slow	2.615	0.468–44.418

We report odds ratios (OR) and their 95% bootstrapped confidence intervals. Landscape direction is East–West.

Table 4. Coefficient estimates for models testing factors that influence the *track length* in animals that follow roads (*movement type* = follow).

Variables	β	95% CI
Intercept	39.548	30.223 to 49.686
Road orientation: East–West	–7.796	–13.835 to –1.370
Species group: ungulates	–11.893	–19.115 to –4.841
Speed: slow	–8.739	–15.544 to –1.278
Shrub coverage	0.006	–0.102 to 0.109

We report best estimates and their 95% bootstrapped confidence intervals. Relevant predictors with 95% CI non-overlapping with zero are highlighted in bold.

first assessment of how road orientation in relation to key resources affects animal movement, which can ultimately influence landscape connectivity and direct impacts to wildlife and humans, but our approach had some limitations. First, we did not monitor animal tracks outside roads; thus, we could not test whether the studied species used roads differently from the natural landscape. However, our study design allowed us to compare the use on roads running both across and along the environmental gradient, testing the hypothesis that road use by animals could differ when both orientations are available. Second, our study area is small (<1 km²) and did not allow us to confirm that the patterns found here will apply to larger geographical scales. Studies addressing the consequences of animal movement on landscape connectivity are usually performed at a larger scale (landscape or regional, up to global) involving high economic costs (e.g. tagging or radio-marking several individuals of the target species; [Blazquez-Cabrera et al. 2016](#)) or complex modelling efforts (e.g. [Ascensão et al. 2019b](#)). Here, we used a smaller-scale approach that can represent a valuable first step to assess landscape connectivity with a relatively small effort, both economically and analytically. Third, the low

traffic levels and the lack of pavement of the studied roads seem to facilitate animal movement along them, but effects may be different for busy or paved roads. Future research would be necessary to understand how different types of roads facilitate or prevent access of wildlife to spatially aggregated limiting resources and how this may affect landscape connectivity and ultimately the persistence of animal populations.

Our results have potential implications for the conservation and management of species and areas, and for human safety by helping in understanding the mechanisms potentially reducing wildlife-vehicle collisions. Roadkills occur when wildlife use roads, which we show can be influenced by road orientation, suggesting this aspect of road design should be considered as a factor affecting collision probability. Road orientation seems to influence particularly large ungulates, which due to their size pose the greatest risk to humans in case of a collision ([Conover 2019](#)). We recommend that mitigation measures should be prioritised for roads paralleling resource gradients as these may be preferably used by large ungulates to access limiting resources. Fenced roadsides could be designed to act as suitable corridors for ungulates, allowing for required movements to essential resources but limiting collision risks ([Jakes et al. 2018](#)). For roads that transverse the environmental gradient, wildlife road-crossing structures will be essential to ensure access to limiting resources. Finally, while road orientation may be primarily driven by human requirements (i.e. which areas need to be connected), whenever possible the construction of new roads in directional landscapes should consider road orientation in relation to key resources for wildlife to reduce the impact on animals movement and landscape connectivity.

Conclusion

The main finding of our study is that road orientation affects how these linear structures influence animal movement in landscapes where resources are distributed along a spatial gradient with distinct effects for ungulates and carnivores. When a road is planned and designed its orientation will be primarily determined by human needs and costs, although environmental impact assessments will also take into consideration aspects that we studied here such as animal movement and critical resources ([Stokes 2015](#); [Broniewicz and Ogrodnik 2020](#)). However, we consider the key implications of our work affect the planning and implementation of mitigation strategies and safety measures. Our results suggest road-crossing structures and fences will be most important for roads traversing a gradient to allow wildlife movement while preventing collisions. For roads along a gradient, crossing structures may be less important, but fences or appropriate signage could be useful to prevent or warn drivers of animals travelling on the road.

Future work will be necessary to expand on our research by covering different areas with varying resource gradients, representing different species, and comparing roads with higher levels of traffic and other features (e.g. paved, multiple lanes). Additional work should also consider overcoming some of the limitations of our study, by including off-road data on animal movement and covering a wider region. Despite its limitations, our study expands our knowledge of how largely overlooked aspects of how roads are designed can affect animal movement and ultimately landscape connectivity.

References

- Andersen GE, Johnson CN, Barmuta LA, Jones ME (2017) Use of anthropogenic linear features by two medium-sized carnivores in reserved and agricultural landscapes. *Scientific Reports* **7**, 11624. doi:10.1038/s41598-017-11454-z
- Ascensão F, Kindel A, Teixeira FZ, Barrientos R, D'Amico M, Borda-de-Água L, Pereira HM (2019a) Beware that the lack of wildlife mortality records can mask a serious impact of linear infrastructures. *Global Ecology and Conservation* **19**, e00661. doi:10.1016/j.gecco.2019.e00661
- Ascensão F, Mestre F, Barbosa AM (2019b) Prioritizing road defragmentation using graph-based tools. *Landscape and Urban Planning* **192**, 103653. doi:10.1016/j.landurbplan.2019.103653
- Barrientos R, Martins RC, Ascensão F, D'Amico M, Moreira F, Borda-de-Água L (2018) A review of searcher efficiency and carcass persistence in infrastructure-driven mortality assessment studies. *Biological Conservation* **222**, 146–153. doi:10.1016/j.biocon.2018.04.014
- Bennett VJ (2017) Effects of road density and pattern on the conservation of species and biodiversity. *Current Landscape Ecology Reports* **2**, 1–11. doi:10.1007/s40823-017-0020-6
- Bissonette JA, Kassir CA, Cook LJ (2008) Assessment of costs associated with deer–vehicle collisions: human death and injury, vehicle damage, and deer loss. *Human-Wildlife Conflicts* **2**, 17–27. doi:10.26077/ns32-mk60
- Blazquez-Cabrera S, Gastón A, Beier P, Garrote G, Simón MÁ, Saura S (2016) Influence of separating home range and dispersal movements on characterizing corridors and effective distances. *Landscape Ecology* **31**, 2355–2366. doi:10.1007/s10980-016-0407-5
- Borkovcová M, Mrtka J, Winkler J (2012) Factors affecting mortality of vertebrates on the roads in the Czech Republic. *Transportation Research Part D: Transport and Environment* **17**, 66–72. doi:10.1016/j.trd.2011.09.011
- Boughton RK, Allen BL, Tillman EA, Wisely SM, Engeman RM (2019) Road hogs: implications from GPS collared feral swine in pastureland habitat on the general utility of road-based observation techniques for assessing abundance. *Ecological Indicators* **99**, 171–177. doi:10.1016/j.ecolind.2018.12.022
- Broniewicz E, Ogródnik K (2020) Multi-criteria analysis of transport infrastructure projects. *Transportation Research Part D: Transport and Environment* **83**, 102351. doi:10.1016/j.trd.2020.102351
- Brown GP, Phillips BL, Webb JK, Shine R (2006) Toad on the road: use of roads as dispersal corridors by cane toads (*Bufo marinus*) at an invasion front in tropical Australia. *Biological Conservation* **133**, 88–94. doi:10.1016/j.biocon.2006.05.020
- Castroviejo J (1993) 'Memoria: Mapa del Parque Nacional de Doñana.' (Consejo Superior de Investigaciones Científicas (CSIC))
- Ceia-Hasse A, Borda-de-Água L, Grilo C, Pereira HM (2017) Global exposure of carnivores to roads. *Global Ecology and Biogeography* **26**, 592–600. doi:10.1111/geb.12564
- Chen HL, Koprowski JL (2016) Differential effects of roads and traffic on space use and movements of native forest-dependent and introduced edge-tolerant species. *PLoS ONE* **11**, e0148121. doi:10.1371/journal.pone.0148121
- Conover MR (2019) Numbers of human fatalities, injuries, and illnesses in the United States due to wildlife. *Human-Wildlife Interactions* **13**, 12. doi:10.26077/r59n-bv76
- D'Amico M, Román J, de los Reyes L, Revilla E (2015) Vertebrate road-kill patterns in Mediterranean habitats: who, when and where. *Biological Conservation* **191**, 234–242. doi:10.1016/j.biocon.2015.06.010
- D'Amico M, Périquet S, Román J, Revilla E (2016) Road avoidance responses determine the impact of heterogeneous road networks at a regional scale. *Journal of Applied Ecology* **53**, 181–190. doi:10.1111/1365-2664.12572
- Díaz-Varela ER, Vázquez-González I, Marey-Pérez MF, Álvarez-López CJ (2011) Assessing methods of mitigating wildlife–vehicle collisions by accident characterization and spatial analysis. *Transportation Research Part D: Transport and Environment* **16**, 281–287. doi:10.1016/j.trd.2011.01.002
- Duffett D, D'Amico M, Mulero-Pázmány M, González-Suárez M (2020) Species' traits as predictors of avoidance towards roads and traffic. *Ecological Indicators* **115**, 106402. doi:10.1016/j.ecolind.2020.106402
- Garber NJ, Hoel LA (2019) 'Traffic and highway engineering.' (Cengage Learning)
- González-Suárez M, Zanchetta Ferreira F, Grilo C (2018) Spatial and species-level predictions of road mortality risk using trait data. *Global Ecology and Biogeography* **27**, 1093–1105. doi:10.1111/geb.12769
- Grilo C, Ascensão F, Santos-Reis M, Bissonette JA (2011) Do well-connected landscapes promote road-related mortality? *European Journal of Wildlife Research* **57**, 707–716. doi:10.1007/s10344-010-0478-6
- Grilo C, Sousa J, Ascensão F, Matos H, Leitão I, Pinheiro P, Costa M, Bernardo J, Reto D, Lourenço R, Santos-Reis M, Revilla E (2012) Individual spatial responses towards roads: implications for mortality risk. *PLoS ONE* **7**, e43811. doi:10.1371/journal.pone.0043811
- Jacobson SL, Bliss-Ketchum LL, de Rivera CE, Smith WP (2016) A behavior-based framework for assessing barrier effects to wildlife from vehicle traffic volume. *Ecosphere* **7**, e01345. doi:10.1002/ecs2.1345
- Jakes AF, Jones PF, Paige LC, Seidler RG, Huijser MP (2018) A fence runs through it: a call for greater attention to the influence of fences on wildlife and ecosystems. *Biological Conservation* **227**, 310–318. doi:10.1016/j.biocon.2018.09.026
- Kang W, Minor ES, Woo D, Lee D, Park C-R (2016) Forest mammal roadkills as related to habitat connectivity in protected areas. *Biodiversity and Conservation* **25**, 2673–2686. doi:10.1007/s10531-016-1194-7
- Kasiringua E, Kopij G, Procheş Ş (2017) Daily activity patterns of ungulates at water holes during the dry season in the Waterberg National Park, Namibia. *Russian Journal of Theriology* **16**, 129–138. doi:10.15298/rusjtheriol.16.2.02
- Kihwele ES, Mchomvu V, Owen-Smith N, Hetem RS, Hutchinson MC, Potter AB, Olf H, Veldhuis MP (2020) Quantifying water requirements of African ungulates through a combination of functional traits. *Ecological Monographs* **90**, e01404. doi:10.1002/ecm.1404
- Laurance SGW, Stouffer PC, Laurance WF (2004) Effects of road clearings on movement patterns of understory rainforest birds in central Amazonia. *Conservation Biology* **18**, 1099–1109. doi:10.1111/j.1523-1739.2004.00268.x
- Macdonald DW (1979) Some observations and field experiments on the urine marking behaviour of the red fox, *Vulpes vulpes* L. *Zeitschrift für Tierpsychologie* **51**, 1–22. doi:10.1111/j.1439-0310.1979.tb00667.x
- Matos C, Sillero N, Argaña E (2012) Spatial analysis of amphibian road mortality levels in northern Portugal country roads. *Amphibia-Reptilia* **33**, 469–483. doi:10.1163/15685381-00002850
- Meijer JR, Huijbregts MAJ, Schotten KCGJ, Schipper AM (2018) Global patterns of current and future road infrastructure. *Environmental Research Letters* **13**, 064006. doi:10.1088/1748-9326/aabd42
- Monclús R, Arroyo M, Valencia A, de Miguel FJ (2009) Red foxes (*Vulpes vulpes*) use rabbit (*Oryctolagus cuniculus*) scent marks as territorial marking sites. *Journal of Ethology* **27**, 153–156. doi:10.1007/s10164-008-0098-8
- Morelle K, Lehaire F, Lejeune P (2013) Spatio-temporal patterns of wildlife–vehicle collisions in a region with a high-density road network. *Nature Conservation* **5**, 53–73. doi:10.3897/natureconservation.5.4634

- R Foundation for Statistical Computing (2013) 'R: a language and environment for statistical computing.' (R Foundation for Statistical Computing: Vienna, Austria)
- Román J, Barón A, Revilla E (2010) Evaluación de los efectos del tránsito a motor sobre especies y comunidades de interés en el Espacio Natural Doñana. Estación Biológica de Doñana CSIC, Seville, Spain.
- Rytwinski T, Soanes K, Jaeger JAG, Fahrig L, Findlay CS, Houlahan J, van der Ree R, van der Grift EA (2016) How effective is road mitigation at reducing road-kill? A meta-analysis. *PLoS ONE* **11**, e0166941. doi:10.1371/journal.pone.0166941
- Stokes J (2015) What transportation agencies need in environmental impact assessments and other reports to minimise ecological impacts. In 'Handbook of road ecology'. (Eds R van der Ree, DJ Smith, C Grilo) pp. 43–50. (Wiley-Blackwell)
- Suárez-Esteban A, Delibes M, Fedriani JM (2013) Barriers or corridors? The overlooked role of unpaved roads in endozoochorous seed dispersal. *Journal of Applied Ecology* **50**, 767–774. doi:10.1111/1365-2664.12080
- Toger M, Benenson I, Wang Y, Czamanski D, Malkinson D (2018) Pigs in space: an agent-based model of wild boar (*Sus scrofa*) movement into cities. *Landscape and Urban Planning* **173**, 70–80. doi:10.1016/j.landurbplan.2018.01.006
- Visintin C, van der Ree R, McCarthy MA (2016) A simple framework for a complex problem? Predicting wildlife–vehicle collisions. *Ecology and Evolution* **6**, 6409–6421. doi:10.1002/ece3.2306
- Visintin C, Golding N, van der Ree R, McCarthy MA (2018) Managing the timing and speed of vehicles reduces wildlife-transport collision risk. *Transportation Research Part D: Transport and Environment* **59**, 86–95. doi:10.1016/j.trd.2017.12.003
- Whittington J, St. Clair CC, Mercer G (2005) Spatial responses of wolves to roads and trails in mountain valleys. *Ecological Applications* **15**, 543–553. doi:10.1890/03-5317
- Whittington J, Hebblewhite M, DeCesare NJ, Neufeld L, Bradley M, Wilmschurst J, Musiani M (2011) Caribou encounters with wolves increase near roads and trails: a time-to-event approach. *Journal of Applied Ecology* **48**, 1535–1542. doi:10.1111/j.1365-2664.2011.02043.x

Data availability. The data that support this study are available at <https://doi.org/10.6084/m9.figshare.20032637.v1>.

Conflicts of interest. The authors declare no conflicts of interest.

Declaration of funding. MM work was funded by European Union 'NextGenerationEU' (Programa María Zambrano, Ministerio de Universidades, Spain) and Liverpool John Moores University, UK. MD was supported by a post-doctoral grant from Fundação da Ciência e Tecnologia (CEECIND/03798/2017). Universidad de Málaga, Spain, provided funding to publish this article as open access.

Acknowledgements. Logistic and technical support was provided by ICTS-RBD-CSIC, Ministry of Science and Innovation and co-financed by FEDER Funds, and Instituto de Investigación en Recursos Cinegéticos Universidad de Castilla la Mancha (Consejo Superior de Investigaciones Científicas). We thank Jose María Galán (Espacio Natural Doñana) for his help on animal track identification. Doñana National Park registered project 2018/18.

Author affiliations

^ADepartment of Animal Biology, Facultad de Ciencias, University of Málaga, 29071 Málaga, Spain.

^BSchool of Biological and Environmental Sciences, Liverpool John Moores University, 3 Byrom Street, Liverpool L3 3AF, UK.

^CCybertracker Conservation, Plaza del Comercio 2, 21750 El Rocio, Huelva, Spain.

^DCIBIO-InBIO, School of Agriculture, University of Porto and University of Lisbon, Tapada da Ajuda Campus, 1349-017 Lisbon, Portugal.

^EPresent address: Department of Conservation Biology, Doñana Biological Station CSIC, C/Américo Vespucio 26, 41092 Seville, Spain.

^FEcology and Evolutionary Biology, School of Biological Sciences, University of Reading, Reading RG6 6EX, UK.