

## Two new species of Papillacarus (Acari, Oribatida, Lohmanniidae) from China

Authors: Ren, Guoru, Yang, Maofa, Liang, Wenqin, and Xie, Lixia

Source: Systematic and Applied Acarology, 23(5) : 824-837

Published By: Systematic and Applied Acarology Society

URL: <https://doi.org/10.11158/saa.23.5.4>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Two new species of *Papillacarus* (Acari, Oribatida, Lohmanniidae) from China

GUORU REN<sup>1</sup>, MAOFA YANG<sup>1\*</sup>, WENQIN LIANG<sup>1,2</sup> & LIXIA XIE<sup>3</sup>

<sup>1</sup> Institute of Entomology, Guizhou University, Guizhou Provincial Key Laboratory for Agricultural Pest Management of the Mountainous Region, Guiyang, 550025, China

<sup>2</sup> Guizhou Provincial Center for Disease Control and Prevention, Guiyang, 550004, China

<sup>3</sup> College of Plant Protection, Shandong Agricultural University, Shandong Provincial Key Laboratory for Biology of Vegetable Diseases and Insect Pests, Taian, 271018, China

\* Corresponding author, E-mail: gdgally@126.com

### Abstract

Two new species of Lohmanniidae, *Papillacarus (Papillacarus) internus* sp. nov. from soil in Hainan Province, South China and *Papillacarus (Vepracarus) hexagonus* sp. nov. from rotten tree and soil in Chongqing City, Southwest China are described and illustrated. *Papillacarus (P.) internus* sp. nov. is most similar to *Papillacarus (P.) lienhardi* (Mahunka, 1997), however, it can be distinguished from the latter by the following characters: 29–30 pairs of additional neotrichal setae, seta  $c_1$  shorter than  $c_2$ , all genital setae setiform and smooth. *Papillacarus (V.) hexagonus* sp. nov. is most similar to *Papillacarus (V.) gueyeae* (Pérez-Íñigo, 1989), however, it can be distinguished from the latter by the following characters: 38 pairs of additional neotrichal setae, 7 pairs of subcapitula setae, epimeral formula 9-4-3-4.

**Key words:** Lohmanniidae, new species, systematics, morphology, China

### Introduction

In this paper we describe two new species of Lohmanniidae: *Papillacarus (Papillacarus) internus* sp. nov. from soil in Hainan Province, South China, and *Papillacarus (Vepracarus) hexagonus* sp. nov. from rotten tree and soil in Chongqing City, Southwest China.

The oribatid mite genus *Papillacarus* was proposed by Kunst (1959) with *Lohmannia murciooides aciculata* Berlese, 1905 as type species. Currently, *Papillacarus* comprises 37 species in the two subgenera *Papillacarus* and *Vepracarus*, which are distributed in tropical regions (Subías 2004, online version 2018). The main generic characters of *Papillacarus* are the following: genital plates with transverse suture, anal and adanal plates separated, preanal plate narrow, two pairs of adanal setae present, notogastral setae setiform, branched or ciliate. The main subgeneric characters of *Papillacarus (P.)* are the following: pygidial setae setiform, ciliate; weak additional neotrichal setae, no more than 30 pairs. The main subgeneric characters of *Papillacarus (V.)* are the following: pygidial setae branched or tree-shaped; strong additional neotrichal setae, more than 20 pairs (Balogh 1961; Wallwork 1962; Aoki 1965; Balogh & Balogh 1987, 1992). Prior to this study, eight species of the genus *Papillacarus* have been recorded in China (Wang & Hu 1990, Li *et al.* 1991, Chen *et al.* 2010a, Chen *et al.* 2010b, Chen & Yang 2011): *Papillacarus (Papillacarus) echinatus* Li *et al.* 1991, *Papillacarus (Vepracarus) hirsutus* Aoki, 1961, *Papillacarus (P.) ondriasi* Mahunka, 1974, *Papillacarus (P.) undirostratus* Aoki, 1965, *Papillacarus (V.) cruzae* Corpuz-Raros, 1979, *Papillacarus (V.) punctatus* Wang & Hu, 1990, *Papillacarus (V.) jinggangshanensis* Chen *et al.*

2010b, *Papillacarus* (V.) *konglinensis* Chen & Yang, 2011. Identification keys for some species of *Papillacarus* were presented by Balogh and Balogh (1987), Ermilov *et al.* (2012), Ermilov and Tolstikov (2015).

## Material and methods

For the material, see *Material examined* section of each species. The specimens were mounted in lactic acid on temporary cavity slides for measurement and illustration. The body measurements are presented in micrometers. The body length was measured in lateral view, from the tip of the rostrum to the posterior edge of the ventral plate. Notogastral width refers to the maximum width in dorsal aspect. Lengths of body setae were measured in lateral aspect. Formulae for leg setation are given according to the sequence trochanter-femur-genu-tibia-tarsus (famulus included). Formulae for leg solenidia are given in brackets according to the sequence genu-tibia-tarsus. General terminology used in this paper follows that summarized by Grandjean (1950), Norton (1977) for leg setal nomenclature, and Norton and Behan-Pelletier (2009). Morphological description of both species follows with Ermilov and Deharveng (2016), Ermilov (2017), Ermilov *et al.* (2017). Drawings were made with a camera lucida using an Olympus transmission light microscope “Olympus CX41”.

## Descriptions

### *Papillacarus* (*Papillacarus*) *internus* sp. nov.

(Figs 1–10)

#### Diagnosis


Body size 467–488×227–243. Rostrum rounded. Prodorsum and notogaster surface densely punctate. Prodorsal, notogastral and neotrichal setae setiform, with short cilia, and thin, smooth apex. Five transverse bands present on notogaster surface. Bothridial seta pectinate, with 12 to 13 branches. Epimeral setal formula: 7-4-3-4. Genital seta setiform, smooth. Adanal seta setiform, barbed unilaterally.

*Measurements.* Body length 472 (holotype), 467–488 (9 paratypes); body width 236 (holotype), 227–243 (9 paratypes).

*Integument.* Body yellow-brown. Surface of prodorsum, notogaster, subcapitulum, genital, adanal plates and legs densely punctate.

*Prodorsum* (Figs 1–3). Rostrum widely rounded. Rostral seta (*ro*, 52) setiform. All prodorsal setae setiform, with short cilia, and thin, smooth apex. Lamellar seta (*le*, 62), interlamellar seta (*in*, 70) longer than others, posterior exobothridial seta (*exp*, 65) setiform, smooth. Anterior exobothridial seta (*exa*, 59) barbed bilaterally. Bothridial seta (*bs*, 82) pectinate, with 12 to 13 branches on one side, and four barbs on the opposite side. Postbothridial transverse band ( $S_b$ ) between bothridia.

*Notogaster* (Figs 1–3). Five transverse bands ( $S_2$ – $S_6$ ) present on notogaster,  $S_2$  and  $S_4$  complete, while  $S_3$ ,  $S_5$  and  $S_6$  interrupted medially. Sixteen pairs of primary notogastral setae and 29–30 pairs of additional neotrichal setae present, all setiform, with short cilia bilaterally or unilaterally and smooth apex. Setae  $c_1$ ,  $d_1$ ,  $e_1$ , and  $f_1$  never reaching the margin of notogaster, seta  $c_1$  shorter than distance  $c_1$ – $d_1$ , setae  $d_1$ ,  $d_2$  lie on ( $d_1$ ) resp. inside of a duplication ( $d_2$ ) of the transverse band  $S_4$ . Neotrichal setae of two types: 3 pairs long (*m*, 45–48; *n*, 50–53; *r*, 58–60) and 26–27 pairs short (21–33). Lyrifissures *ia*, *im*, *ip*, *ih* distinct, *ia* laterally to seta  $c_3$ , *ip* anterior laterally to  $f_2$ , *ih* laterally to  $h_3$ , *ips* not distinct. Notogastral setal lengths:  $c_1 \approx h_1 = 41$  (39–42),  $c_2 \approx d_1 \approx e_1 \approx f_1 = 52$  (51–54),  $c_3 \approx d_2 \approx e_2 = 57$  (56–58),  $f_2 \approx h_2 = 63$  (62–64),  $d_3 \approx h_3 \approx p_1 \approx p_2 \approx p_3 = 69$  (68–71).



**FIGURE 1.** *Papillacarus (Papillacarus) internus* sp. nov., adult: dorsal view. Scale bar 50 µm.


**Gnathosoma** (Figs 2, 4–6). Subcapitulum longer than wide (116–122×92–97). Setae *h*, *m*<sub>1</sub> and *m*<sub>2</sub> (11–20) thickened in medio-basal part, barbed bilaterally, represented by four pairs: *a* (26), *h* (17), *m*<sub>1</sub> (11), *m*<sub>2</sub> (18). Adoral setae smooth: *or*<sub>1</sub> (23) more or less triangular, wide in proximal part, blunt-ended; *or*<sub>2</sub> (30) long, setiform, blunt-ended; *or*<sub>3</sub> (25) long, setiform, pointed-ended. Palp (49) with setal formula 0-1-0-1-10(+1ω). Chelicera (139) with two setae, seta *cha* (17) short, thorn-like, seta *chb* (27) long, setiform and barbed. Trägårdh's organs (*Tg*) triangular, tip slightly pointed.

**Epimeral region** (Fig. 2). Epimeral plates distinct, partly distinct ridges, epimeral setae setiform, setae *1a*, *2a*, *3a*, *4a*, *4b* and one pair of lateral setae of epimere I smooth (6–9); other setae barbed bilaterally. Epimeral formula 7-4-3-4.

**Anogenital region** (Figs 2–3). Ten pairs of genital setae, setiform and smooth, four setae in outer row and six setae in inner row, setae *g*<sub>2</sub>, *g*<sub>5</sub>, *g*<sub>7</sub>, *g*<sub>8</sub> longer (17–22), others setae shorter (8–10). Two pairs of anal setae, setiform, barbed unilaterally. Four pairs of adanal setae (*ad*, 56–72), setiform, barbed unilaterally. Lyrifissure *iad* distinct.

**Legs** (Figs 7–10). All legs with one simple claw each of which with small basoventral tooth. Famulus ε conical and short, posterior to solenidion *ω*<sub>1</sub>. Solenidion *ω*<sub>1</sub> on tarsus I, *ω*<sub>1</sub> and *ω*<sub>2</sub> on tarsus II, φ on tibia III thickened and blunt distally. Other solenidia setiform, with thinner tips. Formulae

of leg setation and solenidia: I 0-5-3(2)-4(1)-18(2), II 0-6-3(1)-4(1)-13(2), III 2-4-2(1)-3(1)-12(0), IV 2-3-2(1)-3(0)-12(0) (see Table 1).




**FIGURE 2.** *Papillacarus (Papillacarus) internus* sp. nov., adult: ventral view. Scale bar 50  $\mu\text{m}$ .

**TABLE 1.** Leg setation and solenidia of adult *Papillacarus (Papillacarus) internus* sp. nov.

Leg	Trochanter	Femur	Genu	Tibia	Tarsus
I	—	<i>d</i> , ( <i>l</i> ), <i>bv</i> "', <i>v</i> "	( <i>l</i> ), <i>dσ</i> ", <i>σ</i> '	<i>xt</i> 2, <i>xt</i> 1, <i>l</i> ', <i>v</i> ', <i>φ</i>	( <i>ft</i> ), ( <i>tc</i> ), ( <i>it</i> ), ( <i>p</i> ), ( <i>u</i> ), ( <i>a</i> ), <i>s</i> , <i>m</i> , <i>n</i> , ( <i>pv</i> ), <i>ε</i> , <i>ω</i> <sub>1</sub> , <i>ω</i> <sub>2</sub>
II	—	<i>d</i> , ( <i>l</i> <sub>1</sub> ), <i>l</i> <sub>2</sub> ", <i>bv</i> "', <i>v</i> "	( <i>l</i> ), <i>dσ</i>	<i>xt</i> 2, <i>xt</i> 1, <i>l</i> ', <i>v</i> ', <i>φ</i>	( <i>ft</i> ), ( <i>tc</i> ), ( <i>p</i> ), ( <i>u</i> ), ( <i>a</i> ), <i>s</i> , ( <i>pv</i> ), <i>ω</i> <sub>1</sub> , <i>ω</i> <sub>2</sub>
III	<i>l</i> ', <i>v</i> '	<i>d</i> , <i>l</i> <sub>1</sub> ', <i>l</i> <sub>2</sub> ', <i>ev</i> '	<i>d</i> , <i>l</i> ', <i>σ</i>	<i>d</i> , <i>l</i> ', <i>v</i> ', <i>φ</i>	( <i>ft</i> ), ( <i>tc</i> ), ( <i>p</i> ), ( <i>u</i> ), <i>a'</i> , <i>s</i> , ( <i>pv</i> )
IV	<i>l</i> ', <i>v</i> '	<i>d</i> , <i>l</i> ', <i>ev</i> '	<i>d</i> , <i>l</i> ', <i>σ</i>	<i>d</i> , <i>l</i> ', <i>v</i> '	( <i>ft</i> ), ( <i>tc</i> ), ( <i>p</i> ), ( <i>u</i> ), <i>a'</i> , <i>s</i> , ( <i>pv</i> )

Note: Roman letters for normal setae Greek letter for solenidia, except *ε* for filum, *dσ*-seta and solenidion coupled. Single prime (') marks setae on anterior and double prime ("') setae on posterior side of the given leg segment. Parentheses refer to a pair of setae.




**FIGURES 3–6.** *Papillacarus (Papillacarus) internus* sp. nov., adult: 3. lateral view; 4. chelicera, left; 5. subcapitulum, ventral view; 6. palp, left. Scale bar 100 µm (3), 25 µm (4, 5, 6).

#### Material examined

Holotype, Chengmai County (19°63'N, 110°3'E), Haikou City, Hainan Province, 9 Jan. 2016, Guoru Ren, in soil. Nine paratypes, same data as holotype. The holotype and nine paratypes are deposited in the Institute of Entomology, Guizhou University, Guiyang of China (GUGC).

#### Etymology

The new specific name “*internus*” is derived from the Latin “*internus*”, and refers to the notogastral setae  $d_1$  and  $d_2$  which are situated on and inside of the duplication of the transverse band  $S_4$ .




**FIGURES 7–8.** *Papillacarus (Papillacarus) internus* sp. nov., adult: legs, femur to tarsus, antiaxial view. 7. leg I, right; 8. leg II, right. Scale bar 50  $\mu$ m.

#### Remarks

The new species *Papillacarus (P.) internus* sp. nov. is most similar to *Papillacarus (P.) lienhardi* (Mahunka, 1997) from Brunei in having setiform prodorsal and notogastral setae.

However, the new species differs from the latter by the following characters: (1) 29–30 pairs of additional neotrichal setae (*versus* approximately 25 pairs *P. lienhardi*); (2) seta  $c_1$  shorter than  $c_2$  (*versus* seta  $c_1$  longer than  $c_2$ ); (3) five transverse bands,  $S_4$  complete (*versus* four transverse bands,  $S_4$  interrupted medially); (4) all genital setae setiform and smooth (*versus* all genital setae ciliate).

The new species *Papillacarus* (*P.*) *internus* sp. nov. is also similar to *Papillacarus* (*P.*) *gramenicus* (Bayartogtokh, 2010) from Mongolia in having setiform prodorsal and notogastral setae. However, the new species differs from the latter by the following characters: (1) 29–30 pairs of additional neotrichal setae (*versus* 13 pairs); (2) epimeral formula 7-4-3-4 (*versus* 8-4-4-4); (3) four pairs of subcapitular setae (*versus* five pairs).




**FIGURES 9–10.** *Papillacarus* (*Papillacarus*) *internus* sp. nov., adult: legs, femur to tarsus, antiaxial view. 9. leg III, left; 10. leg IV, left. Scale bar 50  $\mu\text{m}$ .

***Papillacarus (Vepracarus) hexagonus* sp. nov.**  
(Figs 11–20)

*Diagnosis*


Body size 455–472×210–228. Rostrum truncate, but medially convex. Prodorsum and the whole surface of notogaster covered with bacilliar papillae. Prodorsal, notogastral and neotrichal setae with branches, tree-shaped. Four transverse bands present on notogaster surface. Bothridial seta pectinate, with 9 to 10 branches. Epimeral setal formula: 9-4-3-4. Genital seta setiform or tree-shaped. Adanal seta branched.



**FIGURE 11.** *Papillacarus (Vepracarus) hexagonus* sp. nov., adult: dorsal view. Scale bar 50 µm.

**Measurements.** Body length 462 (holotype), 455–472 (15 paratypes); body width 213 (holotype), 210–228 (15 paratypes).

**Integument.** Body yellow-brown. Surface of Prodorsum, notogaster, subcapitulum, genital, adanal plates and legs papilliform.




**FIGURE 12.** *Papillacarus (Vepracarus) hexagonus* sp. nov., adult: ventral view. Scale bar 50 µm.

**Prodorsum** (Figs 11–13). Rostrum truncate, but medially convex. Rostral seta (*ro*, 30) bear several long branches on each side. Prodorsal surface papilliform in front of postbothridial transverse band and covered with fine spots posterior of this band. Lamellar seta (*le*, 26), interlamellar seta (*in*, 36), anterior exobothridial seta (*exa*, 32), posterior exobothridial seta (*exp*, 38) tree-shaped. Bothridial seta (*bs*, 69) pectinate, with 9 to 10 branches on one side, and three barbs on the opposite side. Postbothridial transverse band ( $S_b$ ) between bothridia.

**Notogaster** (Figs 11–13). Four transverse bands ( $S_2$ – $S_5$ ) present on notogaster, all interrupted medially. Sixteen pairs of primary notogastral setae and 38 pairs of additional neotrichal setae present, all with 5–7 branches, tree-shaped. Setae  $c_1$ ,  $c_2$ ,  $d_1$ ,  $e_1$ ,  $h_1$  and  $f_1$  never reaching margin of notogaster, setae  $d_1$ ,  $e_1$ , looking like hexagram in dorsal view. Lyrifissures *ia*, *im*, *ip*, *ih* distinct, *ia* laterally to seta  $c_3$ , *ip* laterally to  $f_2$ , *ih* anterior laterally to  $h_3$ , *ips* not distinct. Lengths of notogastral setae:  $c_1$ ,  $c_2$ ,  $d_1$ ,  $d_2$ ,  $e_1$ ,  $f_1$ ,  $h_1$ ,  $p_1$ ,  $p_2$  (13–25);  $c_3$ ,  $d_3$ ,  $e_2$ ,  $f_2$ ,  $h_2$ ,  $h_3$ ,  $p_3$  (31–48).


**Gnathosoma** (Figs 12, 14–16). Subcapitulum longer than wide (98–105×80–84). Surface of subcapitulum papilliform. Seven pairs of subcapitula setae, *a* (22–24) setiform, smooth, pointed-ended;  $m_1$  and  $m_4$  (18–23) setiform, barbed unilaterally;  $h_1$ ,  $h_2$ ,  $m_2$ ,  $m_3$  (12–17) setiform, barbed

bilaterally. Three pairs of smooth adoral setae:  $or_1$  (14) wide in proximal part, tapering up, blunt-ended;  $or_2$  (22) long, setiform, blunt-ended;  $or_3$  (17) long, setiform, pointed-ended. Palp (38) with setal formula 0-1-0-3-10(+1 $\omega$ ). Chelicera (134) with two setae, seta  $cha$  (4) short, thorn-like, seta  $chb$  (36) long, setiform and smooth. Trägårdh's organs ( $Tg$ ) triangular, rounded distally.




**FIGURES 13–16.** *Papillacarus (Vepracarus) hexagonus* sp. nov., adult: 13. lateral view; 14. palp, right; 15. subcapitulum, ventral view; 16. chelicera, left. Scale bar 50  $\mu$ m (13), 25  $\mu$ m (14, 15, 16).

*Epimeral region* (Fig. 12). Epimeral plates distinct, partly distinct ridges, epimeral I and II neotrichous, setae setiform or tree-shaped, epimeral formula 9-4-3-4, setae 1a, 2a, 3a, 4a, and one pair of lateral setae of epimere I setiform and smooth (6–10); other setae tree-shaped (11–26).



**FIGURES 17–18.** *Papillacarus (Vepracarus) hexagonus* sp. nov., adult: legs, femur to tarsus, antiaxial view. 17. leg I, right; 18. leg II, right; Scale bar 50 µm.

*Anogenital region* (Figs 12– 13). Ten pairs of genital setae, setiform or tree-shaped, four setae in outer row and six setae in inner row,  $g_7$ ,  $g_8$ ,  $g_{10}$  setiform and smooth (17–19), other setae tree-shaped (11–14). Two pairs of anal setae bearing several long branches unilaterally (*an*, 31–34). Four pairs of adanal setae with several long branches (*ad*, 33–42). Lyrifissure *iad* distinct.



**FIGURES 19–20.** *Papillacarus (Vepracarus) hexagonus* sp. nov., adult: legs, femur to tarsus, antiaxial view. 19. leg III, left; 20. leg IV, left. Scale bar 50  $\mu\text{m}$ .

**Legs** (Figs 17–20). Surface of leg femur papilliform. All legs with one simple claw each of which with small basoventral tooth. Famulus  $\varepsilon$  conical short, posterior to solenidion  $\omega_1$ . Solenidion  $\omega_1$  on tarsus I,  $\omega_1$  and  $\omega_2$  on tarsus II,  $\varphi$  on tibia III thickened and blunt distally. Other solenidia setiform, with thinner tips. Formulae of leg setation and solenidia: I 0-5-3(2)-4(1)-17(2), II 0-6-3(1)-4(1)-11(2), III 2-4-2(1)-3(1)-11(0), IV 2-3-2(1)-3(0)-12(0) (see Table 2).

**TABLE 2.** Leg setation and solenidia of adult *Papillacarus (Vepracarus) hexagonus* sp. nov.

Leg	Trochanter	Femur	Genu	Tibia	Tarsus
I	—	<i>d</i> , ( <i>l</i> ), <i>bv</i> ", <i>v</i> "	( <i>l</i> ), <u><i>dσ</i></u> ", <i>σ'</i>	<i>xt2</i> , <i>xt1</i> , <i>l'</i> , <i>v'</i> , <i>φ</i>	( <i>ft</i> ), ( <i>tc</i> ), <i>it'</i> , ( <i>p</i> ), ( <i>u</i> ), ( <i>a</i> ), <i>s</i> , <i>m</i> , <i>n</i> , ( <i>pv</i> ), <i>ε</i> , <i>ω</i> <sub>1</sub> , <i>ω</i> <sub>2</sub>
II	—	<i>d</i> , ( <i>l</i> <sub>1</sub> ), <i>l</i> <sub>2</sub> ", <i>bv</i> ", <i>v</i> "	( <i>l</i> ), <u><i>dσ</i></u>	<i>xt2</i> , <i>xt1</i> , <i>l'</i> , <i>v'</i> , <i>φ</i>	( <i>ft</i> ), ( <i>tc</i> ), ( <i>p</i> ), ( <i>u</i> ), ( <i>a</i> ), <i>s</i> , ( <i>pv</i> ), <i>ω</i> <sub>1</sub> , <i>ω</i> <sub>2</sub>
III	<i>l'</i> , <i>v'</i>	<i>d</i> , <i>l</i> <sub>1</sub> ', <i>l</i> <sub>2</sub> ', <i>ev'</i>	<i>d</i> , <i>l'</i> , <i>σ</i>	<i>d</i> , <i>l'</i> , <i>v'</i> , <i>φ</i>	( <i>ft</i> ), ( <i>tc</i> ), ( <i>p</i> ), ( <i>u</i> ), <i>a'</i> , <i>s</i> , <i>pv'</i>
IV	<i>l'</i> , <i>v'</i>	<i>d</i> , <i>l'</i> , <i>ev'</i>	<i>d</i> , <i>l'</i> , <i>σ</i>	<i>d</i> , <i>l'</i> , <i>v'</i>	( <i>ft</i> ), ( <i>tc</i> ), ( <i>p</i> ), ( <i>u</i> ), <i>a'</i> , <i>s</i> , ( <i>pv</i> )

Note: Roman letters for normal setae Greek letter for solenidia, except *ε* for famulus, *dσ*-seta and solenidion coupled. Single prime (') marks setae on anterior and double prime (") setae on posterior side of the given leg segment. Parentheses refer to a pair of setae.

#### Material examined

Holotype, Simian Mountain (28°34'N, 106°20'E), Chongqing City, 23 Sept. 2017, Guoru Ren, in rotten tree and soil. Fifteen paratypes, same data as holotype. The holotype and fifteen paratypes are deposited in the Institute of Entomology, Guizhou University, Guiyang of China (GUGC).

#### Etymology

The new specific name “hexagonus” is derived from the Latin “hexagonus”, and refers to the notogastral setae *d*<sub>1</sub> and *e*<sub>1</sub>, which look like a hexagram in dorsal view.

#### Remarks

The new species *Papillacarus (V.) hexagonus* sp. nov. is most similar to *Papillacarus (V.) gueyeae* (Pérez-Íñigo, 1989) described from Senegal, in having tree-shaped prodorsum and notogastral setae. However, the new species differs from the latter by the following characters: (1) 38 pairs of additional neotrichal setae (*versus* 46–50 pairs in *Papillacarus (V.) gueyeae*); (2) bothridial seta with 9 to 10 branches (*versus* 13 to 15 branches); (3) epimeral formula 9-4-3-4 (*versus* 8-5-3-3); (4) seven pairs of subcapitular setae (*versus* six pairs); (5) genital setae *g*<sub>4</sub>–*g*<sub>6</sub> tree-shaped; *g*<sub>10</sub> setiform and smooth (*versus* *g*<sub>4</sub>–*g*<sub>6</sub> setiform, smooth; *g*<sub>10</sub> tree-shaped).

The new species *Papillacarus (V.) hexagonus* sp. nov. is also similar to *Papillacarus (V.) cornutus* (Sarkar & Subías, 1984) described from India in having tree-shaped prodorsal and notogastral setae. However, the new species differs from the latter by the following characters: (1) body size 455–472×210–228 (*versus* 342–349×134–141 in *Papillacarus (V.) cornutus*); (2) body surface covered with polygonal network sculpture (*versus* without); (3) 38 pairs of additional neotrichal setae (*versus* 18 pairs); (4) genital setae *g*<sub>4</sub>–*g*<sub>6</sub> tree-shaped; *g*<sub>10</sub> setiform and smooth (*versus* all genital setae bilaterally barbed).

#### Acknowledgements

We would like to express our gratitude to Dr. Yong Chen (Institute of Entomology, Guizhou University, Guiyang, China) and Professor Roy A. Norton (State University of New York, Syracuse, U.S.A.) for providing literature. This project was supported by the Program of Ministry of Science and Technology of the People's Republic of China (2015FY210300); the Program of Excellent Innovation Talents, Guizhou Province, China (No. 20164022); the Funds for National Natural Sciences Foundation of China (No. 31501847), the key research and development program of Shandong Province (2016GNC110012).

## References

- Aoki, J. (1961) Beschreibungen von neuen Oribatiden Japans. *Japanese Journal of Applied Entomology and Zoology*, 5(1), 64–69.  
<https://doi.org/10.1303/jjaez.5.64>
- Aoki, J. (1965) Oribatiden (Acarina) Thailands. I. *Nature and Life in Southeast Asia*, 4, 129–193.
- Balogh, J. (1961) An outline of the family Lohmanniidae Berlese, 1916 (Acari: Oribatei). *Acta Zoologica Academiae Scientiarum Hungaricae*, 7(1–2), 19–44.
- Balogh, J. & Balogh, P. (1987) A new outline of the Family Lohmanniidae Berlese, 1916 (Acari: Oribatei). *Acta Zoologica Academicae Scientiarum Hungaricae*, 33(3–4), 327–398.
- Balogh, J. & Balogh, P. (1992) *The oribatid mites genera of the world. Vol. 1*. Budapest, Hungarian National Museum Press, 263 pp.
- Berlese, A. (1905) Acari nuovi. Manipulus III (Acari di Giava). *Redia*, 2, 10–32.
- Chen, Y. & Yang, M.F. (2011) Two new species of the family Lohmanniidae (Acari, Oribatida) from China. *International Journal of Acarology*, 37(5), 448–454.  
<https://doi.org/10.1080/01647954.2010.525524>
- Chen, J., Liu, D. & Wang, H.F. (2010a) Oribatid mites of China: a review of progress, with a checklist. *Zoosymposia*, 4, 186–224.
- Chen, Y., Yang, M.F. & Liang, W.Q. (2010b) A taxonomic study on the genus *Vepracarus* Aoki (Acari, Oribatida, Lohmanniidae), with description of a new species from China. *Acta Zootaxonomica Sinica*, 35(1), 35–37.
- Corpuz-Raros, L.A. (1979) Philippine Oribatei (Acarina) II. Family Lohmanniidae. *Kalikasan, Philippine Journal Biology*, 8(3), 315–334.
- Ermilov, S.G. (2017) A new species of *Lohmannia* (*Lohmannia*) (Acari, Oribatida, Lohmanniidae) from Vietnam, with supplementary description of *L. (Lohmannia) turcmenica* (Bulanova-Zachvatkina, 1960). *Systematic & Applied Acarology*, 22(2), 193–207.  
<https://doi.org/10.11158/saa.22.2.4>
- Ermilov, S.G. & Deharveng, L. (2016) Contribution to the knowledge of oribatid mites (Acari, Oribatida) of Vanuatu. *Systematic & Applied Acarology*, 21(5), 681–688.  
<https://doi.org/10.11158/saa.21.5.9>
- Ermilov, S.G. & Tolstikov, A.V. (2015) Report on oribatid mites (Acari, Oribatida) from Brazil, with description of a new species of the genus *Papillacarus*. *Acarina*, 23(1), 49–54.
- Ermilov, S.G., Anichkin, A.E. & Wu, D. (2012) Two new species of the genus *Papillacarus* (Acari: Oribatida: Lohmanniidae) from caves of Southern Vietnam. *Zootaxa*, 3593, 75–88.
- Ermilov, S.G., Hugo-Coetzee, E.A. & Khaustov, A.A. (2017) Contribution to the knowledge of oribatid mites of the family Lohmanniidae (Acari, Oribatida) from South Africa. *Systematic & Applied Acarology*, 22(5), 666–682.  
<https://doi.org/10.11158/saa.22.5.6>
- Grandjean, F. (1950) Etude sur les Lohmanniidae (Oribates, Acariens). *Archives de Zoologie experimentale et generale*, 87(2), 95–161.
- Kunst, M. (1959) Bulgarische Oribatiden (Acarina) III. *Acta Universitatis Carolina, Biologica*, 6(1), 51–74.
- Li, Y.R., Li, L.S. & Chen, H.N. (1991) A new species of Papillacarus from Chongqing, China (Oribatida: Lohmanniidae). *Acta Entomologica Sinica*, 34(4), 496–498.
- Mahunka, S. (1997) Oribatids from Brunei II (Acari: Oribatida). (*Acarologica Genavensis LXXXII*). *Revue Suisse de Zoologie*, 104(3), 661–700.  
<https://doi.org/10.5962/bhl.part.80015>
- Norton, R.A. (1977) A review of F. Grandjean's system of leg chaetotaxy in the Oribatei (Acari) and its application to the family Damaeidae. In: Dindal, D.L. (Editor), *Biology of oribatid mites*. SUNY College of Environmental Science and Forestry, Syracuse, pp. 33–61.
- Norton, R.A. & Behan-Pelletier, V.M. (2009) Oribatida. Chapter 15, In: Krantz, G.W. & Walter, D.E. (eds). *A manual of acarology 3rd edition*. Texas Tech University Press, Lubbock, pp. 430–564.
- Pérez-Íñigo, C. (1989) Première découverte du genre *Vepracarus* en Afrique continentale (Acari, Oribatei, Lohmanniidae). *Revue de Zoologie Africaine-Journal of African Zoology*, 103, 389–393.
- Subías, L.S. (2004) Listado sistemático, sinonímico y biogeográfico de los ácaros oribátidos (Acariformes: Oribatida) del mundo (excepto fósiles). *Graellsia*, 60 (número extraordinario), pp. 3–305. Online version accessed in January 2018, 605 pp.
- Wallwork, J.A. (1962) Some Oribatei from Ghana X. The family Lohmanniidae. *Acarologia*, 4(3), 457–487.
- Wang, X.Z. & Hu, S.H. (1990) Studies on the Chinese species of Lohmanniidae (Acari: Oribatida). *Contract Shanghai Institute Entomology*, 9, 127–140

Submitted: 25 Feb. 2018; accepted by Marut Fuangarworn: 19 Apr. 2018; published: 11 May 2018