

## **First mitochondrial genome of *Amblyomma triste* Koch, 1844 (Acari: Ixodidae): Evidence for studying species within the *A. maculatum* group**

Authors: Ossa-López, Paula A., Uribe, Juan E., Ramírez-Chaves, Héctor E., and Rivera-Páez, Fredy A.

Source: Systematic and Applied Acarology, 29(1) : 93-108

Published By: Systematic and Applied Acarology Society

URL: <https://doi.org/10.11158/saa.29.1.7>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Article

## First mitochondrial genome of *Amblyomma triste* Koch, 1844 (Acari: Ixodidae): Evidence for studying species within the *A. maculatum* group

PAULA A. OSSA-LÓPEZ<sup>1,2</sup>, JUAN E. URIBE<sup>3</sup>, HÉCTOR E. RAMÍREZ-CHAVES<sup>2,4</sup> & FREDY A. RIVERA-PÁEZ<sup>2\*</sup>

<sup>1</sup>Doctorado en Ciencias, Biología, Facultad de Ciencias Exactas y Naturales, Universidad de Caldas, Calle 65 No. 26-10, 170004, Manizales. Caldas, Colombia.

<sup>2</sup>Grupo de Investigación en Genética, Biodiversidad y Manejo de Ecosistemas (GEBIOME), Departamento de Ciencias Biológicas, Facultad de Ciencias Exactas y Naturales, Universidad de Caldas, Calle 65 No. 26-10, 170004, Manizales, Caldas, Colombia.

<sup>3</sup>Departamento de Biodiversidad y Biología Evolutiva, Museo Nacional de Ciencias Naturales (MNCN-CSIC), José Gutiérrez Abascal 2, 28006 Madrid, Spain.

<sup>4</sup>Centro de Museos, Museo de Historia Natural, Universidad de Caldas, Calle 58 No. 21-50, 170004, Manizales, Caldas, Colombia.

\*Corresponding author: fredy.rivera@ucaldas.edu.co

### Abstract

The tick species that comprise the *Amblyomma maculatum* group (Acari: Ixodidae) are widely distributed in the Nearctic and Neotropical regions, ranging from the United States of America (USA) to Argentina. This group includes three species: *Amblyomma maculatum*, *Amblyomma tigrinum*, and *Amblyomma triste*, which parasitize a high number of vertebrates, including domestic mammals and humans, where they are the main vectors of *Rickettsia parkeri* s.s. The identification, distribution, and validity of the three species within the group have been controversial and continue to raise questions. Previous studies have mostly focused on the analysis of partial nuclear and mitochondrial genes, highlighting the need for new integrative analyses that contribute to clarifying their systematics and ecology. In this investigation, we obtained the first mitochondrial genome of *A. triste* (14,808 bp), which, upon comparison with the reported mitochondrial genomes of *A. maculatum*, exhibits a genetic distance of 4.2%, providing new evidence for the validity of the former species. The order, composition, and structure of the mitogenome of *A. triste* are consistent with the characteristics reported for most Metastriata including taxa within *Amblyomma*, that count with several mitochondrial genomes published in the last few years. The attributes of the mitogenomes, such as the absence of paralogs and a small genome, confirm the utility in studies involving complexes of cryptic species as it also serves as a robust tool for inferring phylogenies. Furthermore, we reported new molecular markers at the mitochondrial level which can be used in phylogenetic studies for other tick species, especially those with controversial or challenging taxonomy. We highlight the need of sequencing the mitochondrial genome of *A. tigrinum*, which is part of the *A. maculatum* group, as well as experimental crosses of populations from different places in the Americas to obtain additional evidence for species recognition.

**Key words:** Metastriata, Pathogens, Public Health, Systematics, Ticks

### Introduction

Ticks (Acari: Ixodida) are obligate hematophagous ectoparasites capable of transmitting multiple pathogens such as bacteria, helminths, protozoa, and viruses (de la Fuente *et al.* 2008; Baneth 2014; Bezerra-Santos *et al.* 2022). Ixodida comprises three families: Argasidae (soft ticks), Ixodidae (hard

ticks), and Nuttalliellidae (Nava *et al.* 2017; Guglielmone *et al.* 2020; 2021; 2023). Currently, Ixodidae is the most diverse family with approximately 759 species and 15 genera, where *Amblyomma* is the third largest in terms of species number (136), surpassed by *Ixodes* and *Haemaphysalis* with 266 and 176 species, respectively (Guglielmone *et al.* 2023). *Amblyomma* species have received special attention due to their involvement in pathogen transmission (e.g., *Anaplasma*, *Ehrlichia*, *Hepatozoon*, *Rickettsia*) and by the presence of cryptic species (complexes or groups) in America, such as the *Amblyomma cajennense*, *Amblyomma maculatum*, and *Amblyomma ovale* (Dumler *et al.* 2001; Demoner *et al.* 2013; Rivera-Páez *et al.* 2017; Suwanbongkot *et al.* 2019), and the *Amblyomma testudinarium* and *Amblyomma marmoreum* complexes in Asia and Africa, respectively (Mohamed *et al.* 2022; Cotes-Perdomo *et al.* 2023b).

The *A. maculatum* group is widely distributed from the United States of America (USA), Argentina, Belize, Bolivia, Brazil, Chile, Colombia, Costa Rica, Ecuador, French Guiana, Guatemala, Honduras, Mexico, Nicaragua, Paraguay, Peru, Uruguay, and Venezuela (Mendoza-Uribe & Chávez-Chorocco 2004; Guzmán-Cornejo *et al.* 2006; Mertins *et al.* 2010; Abarca *et al.* 2012; Lado *et al.* 2018; Rivera-Páez *et al.* 2018; Ossa-López *et al.* 2022; Guglielmone *et al.* 2023). Currently, the phylogenetic position of the *A. maculatum* group within *Amblyomma* is unclear (Santodomingo *et al.* 2021; Cotes-Perdomo *et al.* 2023a). In addition, the three species of the *A. maculatum* group are morphologically and phylogenetically closely related, casting doubts on their identification and taxonomic recognition (Koch 1844; Mendoza-Uribe & Chávez-Chorocco 2004; Estrada-Peña *et al.* 2005; Guzmán-Cornejo *et al.* 2006; Mertins *et al.* 2010; Abarca *et al.* 2012; Paddock *et al.* 2015; Lado *et al.* 2018; Guglielmone *et al.* 2023). For example, some studies suggest the possibility that *A. maculatum* and *A. triste* are conspecific (Nava *et al.* 2017; Lado *et al.* 2018). Other species such as *A. tigrinum* always cluster into a well-defined monophyletic lineage (Lado *et al.* 2018), and differs from *A. maculatum* and *A. triste* in the adult morphology, and ecological preferences. However, the larvae and nymphs of these three species are extremely difficult to differentiate using morphological traits (Guglielmone *et al.* 2000; Mendoza-Uribe & Chávez-Chorocco 2004; Nava *et al.* 2007; 2017; Lado *et al.* 2018; Guglielmone *et al.* 2021; 2023).

For the *A. maculatum* group, Lado *et al.* (2018) reported four morphotypes with distinct geographical distribution: i) *A. triste* s.s., defined as morphotype I, including specimens from Argentina, southern Brazil, Paraguay, and Uruguay; ii) *A. maculatum* s.s., defined as morphotype II and including specimens from Colombia (Department of Santander), eastern United States of America (USA), Guatemala, Honduras, Mexico, Nicaragua, and Venezuela; iii) morphotype III, which presents a combination of morphological characters limiting the strict assignment to *A. maculatum* s.s. or to *A. triste* s.s.; this morphotype includes ticks from southern and southwestern USA (Arizona and Texas), and northern Mexico; and iv) morphotype IV, which also exhibits a combination of characters of *A. maculatum* s.s. and *A. triste* s.s., and is found in Chile, Ecuador, and Peru. Ossa-López *et al.* (2022) confirmed the wide distribution of *A. maculatum* s.s. (morphotype II) in Colombia and indicated that the sequences of the individual from the Department of Santander analyzed by Lado *et al.* (2018), consistently clustered with other samples from Colombia (from the departments of Caldas, Cundinamarca, and Tolima) instead of clustering with sequences from the USA.

Similarly, Ossa-López *et al.* (2022) confirmed the presence of *A. triste* s.s. (morphotype I) in Colombia using morphological and molecular data, therefore, two out of the three species within the *A. maculatum* group are found in the country, and *A. tigrinum* is currently excluded from the Colombian territory due to lack of verified records (Guglielmone *et al.* 2011; Guglielmone & Robbins 2018; Guglielmone *et al.* 2021; Ortiz-Giraldo *et al.* 2021; Guglielmone *et al.* 2023).

The medical and veterinary importance of *A. maculatum* is well-documented due to its role as the principal reservoir of *Hepatozoon americanum*, and together with *A. triste*, both tick species are

vectors of *Rickettsia parkeri* s.s. (Mathew *et al.* 1998; 1999; Paddock *et al.* 2004; Venzal *et al.* 2004; Paddock & Goddard 2015; Colombo *et al.* 2016; Romer *et al.* 2020). For that reason, more research on reproductive compatibility, distribution, taxonomy, and molecular affinities is needed to clarify the controversial specific taxonomy and to contribute to the understanding of their role in pathogen transmission (Mathew *et al.* 1998; 1999; Ewing *et al.* 2000; Paddock *et al.* 2004; Venzal *et al.* 2004; Colombo *et al.* 2016; Allerdice *et al.* 2020; Romer *et al.* 2020; Cuervo *et al.* 2021; Ossa-López *et al.* 2022; Guglielmone *et al.* 2023).

Mitochondrial genomes (mitogenomes) have proven to be important for the taxonomy, systematics, and population genetics of ticks, emerging as a valuable source of informative molecular markers (Wang *et al.* 2019; Uribe *et al.* 2020; Kneubehl *et al.* 2022; Cotes-Perdomo *et al.* 2023a). Mitochondrial genomes of 120 tick species have been used in phylogenetic analyses (Kelava *et al.* 2021), recovering controversial relationships at the family and genera levels. The absence of paralogous and small size make the mitogenomes a robust molecular marker for inferring phylogenetic hypotheses in studies of cryptic diversity (Zaharias *et al.* 2020; Cotes-Perdomo *et al.* 2023a; 2023b). In this study, sequencing techniques were combined to generate the first mitogenome of *A. triste*, aiming to evaluate its composition (content and organization) and establish its phylogenetic relationships and differentiation from other closely related species within the genus *Amblyomma*.

## Material and methods

### *Sample collection and DNA extraction*

In the year 2018, two male ticks from the *A. maculatum* group were found parasitizing a capybara (*Hydrochoerus hydrochaeris*, Rodentia, Caviidae) at the vereda Las Plumas, municipality of Arauca, Department of Arauca (6°36' 18"N; 70°31' 51"W), Colombia (Ossa-López *et al.* 2022). The collections were executed within the framework permit granted by the National Environmental Licensing Authority (ANLA) to the Universidad de Caldas as stipulated in resolution 02497 of December 31, 2018. The specimens were stored in cryogenic tubes with absolute alcohol and were morphologically identified following the keys of Kohls (1956); Estrada-Peña *et al.* (2005); Barros-Battesti *et al.* (2006); Nava *et al.* (2017); Lado *et al.* (2018) and Guglielmone *et al.* (2021; 2023). The specimens were confirmed as *A. triste*, primarily based on the presence of a thick and highly sclerotized spine and a thin and weakly sclerotized seta on the tibia of legs II-IV. In addition, the spiracular plates (peritreme) are almost oval, with short and wide dorsal projection, and longer spines on coxa IV (Ossa-López *et al.* 2022). The total DNA from each individual was extracted using the DNeasy Blood and Tissue kit (Qiagen) following the manufacturer's recommended protocols, with an extended incubation period (16 h) in lysis buffer at 56 °C.

### *Amplifying and sequencing*

The molecular confirmation was made with partial sequences of three mitochondrial genes, *12S*, *16S* and *cox1* (Ossa-López *et al.* 2022). DNA from a single tick was used for subsequent molecular processes and the sequencing of the complete mitogenome. The fragments were obtained by conventional PCR and sanger sequencing (Macrogen Inc., South Korea) (Supplementary Material 1). The sequences were analyzed using Geneious® Prime 2022.1 software (Kearse *et al.* 2012).

Two strategies were implemented for the amplification and sequencing of the mitogenome. Initially, two fragments that complete ~10kb (70% of the mitogenome) were amplified by long-range PCR and pooled together at equimolar concentration to prepare a Nextera XT DNA library, which was sequenced using Nova Seq 6000 150 PE (150 × 2 bp; 10 Gb/sample) in Illumina platform.

Subsequently, primers were designed along the missing region using as a reference the mitogenome of *A. maculatum* (MW719251; available in the National Center for Biotechnology Information – NCBI database) for primer-walking sequencing by Sanger (Supplementary Material 2).

#### *Assembly, annotation, and alignment*

For each raw file from Illumina sequencing, the quality was checked using FastQC (Andrews 2010). The cleaning, filtering, and adapter trimming of each raw file was made across Trimmomatic (Bolger *et al.* 2014). The partial *cox1* sequence was used as a scaffold for reference-guided assembly using the "Map to Reference(s)" option in Geneious® Prime 2022.1. Subsequently, the protein-coding (PCGs; coding sequences - CDSs), transference RNAs (tRNAs), and the two ribosomal RNAs (rRNAs) genes were identified using MITOS web server (Bernt *et al.* 2013). The annotation was manually verified with the ORFs searcher in Geneious Prime® 2023.2.1. The genes obtained from the *Amblyomma aureolatum* transcriptome from SRA repository in GenBank (SRR4301110; Martins *et al.* 2017) assembled *de novo* by Uribe *et al.* (2020) were included in our analysis, beside the following mitogenomes available in the NCBI: *Amblyomma cajennense* s.s. (OP901701; OP901707), *A. maculatum* (MW719251), *Amblyomma mixtum* (OP901702; OP901703), *Amblyomma ovale* (MT554102; MT554103), *Amblyomma patinoi* (OP901704), *Amblyomma sculptum* (NC\_032369; OP901706), *Amblyomma tonelliae* (OP901705), and *Dermacentor nitens* (NC\_023349) used as outgroup.

#### *Phylogenetic analyses*

The genes were aligned and filtered separately using MAFFT (Katoh & Standley 2013) and BMGE (Criscuolo & Gribaldo 2010). The CDSs + rRNAs filtered genes were concatenated in a matrix at the nucleotide level (Matrix-NT), which was the subject of phylogenetic analyses by probabilistic methods such as Bayesian Inference (BI; Rannala & Yang 1996; Yang & Rannala 1997) and Maximum Likelihood (ML; Felsenstein 1981). BI analyses were performed using MrBayes v3.2.7a (Ronquist *et al.* 2012), two parallel runs, 2000000 generations, in which the initial 25% of sampled data were discarded as burn-in; and ML analyses was performed in IQ-TREE (Nguyen *et al.* 2015) and the robustness was assessed using 1000 bootstrap pseudoreplicates. ModelFinder (Kalyaanamoorthy *et al.* 2017) was used to select the best evolutionary model based on the Bayesian Information Criterion (BIC; Schwarz, 1978): GTR+I+G+F for BI and GTR+F+I+G4 for ML. FigTree v. 1.4.3 (Rambaut 2007) was used to visualize all the phylogenetic trees. Genetic distances were estimated using the *p-distance* method with the MEGA 11 program (Tamura *et al.* 2021) and corroborated with the identity percentage matrix in Geneious Prime® 2023.2.1.


## **Results**

#### *Content and organization of the mitochondrial genome of Amblyomma triste*


The complete mitogenome of the male tick sequenced is available in GenBank as record OR497835. This molecule is completely circular, has a length of 14,808 bp, and it contains (i) 13 CDSs: *cox1-3*, *nad1-6*, *nad4L*, *atp6*, *atp8* and *cytb*; (ii) two rRNA genes: a small subunit ribosomal RNA gene (*rrnS* or *12S*), and a large subunit ribosomal RNA gene (*rrnL* or *16S*); (iii) 22 tRNAs; and (iv) two intergenic spaces: between *12S* and tRNA-Ile, and between tRNA-Leu and tRNA-Cys, corresponding to the non-coding regions 1 and 2 (NCR1 and NCR2) respectively (Figure 1; Table 1).

In the organization of the mitogenome there are two overlaps in the same strand between: *atp6* and *atp8*, and between *nad4* and *nad4L*; and six overlaps in different strands between: *nad2* and

tRNA-Trp; tRNA-Trp and tRNA-Tyr; tRNA-Tyr and *cox1*; tRNA-Arg and tRNA-Asn; tRNA-Asn and tRNA-Ser; and between tRNA-Glu and *nad1*. All the tRNAs vary in length from 54 to 68 nucleotides (Figure 1; Table 1). Prediction of the tRNA secondary structure showed that 20 of the tRNAs have the standard cloverleaf structure, while tRNA-S1 (trnS1) and tRNA-C (trnC) were missing the D-arm or DHU arm (Figure 2).



**FIGURE 1.** Circular and linear representation of the mitochondrial gene order of *Amblyomma triste*. Gene scaling is approximate, and the end of the arrow indicates the direction of the genes (the genes encoded in the H-strand or “+” and L-strand or “-”). All genes have standard nomenclature including the 22 tRNA genes, which are designated by a three-letter code for the corresponding amino acid. The CDS genes are in yellow, the rRNAs in red, the tRNAs in pinkish-purplish color, and the two potential control regions (CR), are non-coding in gray.



**FIGURE 2.** Predicted secondary structure of the mitochondrial tRNA genes of *Amblyomma triste* obtained using MITOS. A. trnS1(aga), lacks DHU arm; B. trnS2(tca), normal cloverleaf structure; C. trnC(tgc), lacks DHU arm.

**TABLE 1.** Mitochondrial genome organization of *Amblyomma triste*.


| Genes/regions                             | Positions and nt sequence lengths(bp) | Strand | Initiation and stop codons |
|-------------------------------------------|---------------------------------------|--------|----------------------------|
| tRNA-Met (M)                              | 1–64 (64)                             | H      |                            |
| <i>nad2</i>                               | 65–1030 (1194)                        | H      | ATT/TAA                    |
| tRNA-Trp (W)                              | 1030–1091 (62)                        | H      |                            |
| tRNA-Tyr (Y)                              | 1090–1151 (62)                        | L      |                            |
| <i>cox1</i>                               | 1144–2682 (1539)                      | H      | ATT/TAA                    |
| <i>cox2</i>                               | 2698–3370 (673)                       | H      | ATG/T                      |
| tRNA-Lys (K)                              | 3371–3435 (65)                        | H      |                            |
| tRNA-Asp (D)                              | 3436–3497 (62)                        | H      |                            |
| <i>atp8</i>                               | 3498–3653 (156)                       | H      | ATT/TAA                    |
| <i>atp6</i>                               | 3647–4309 (663)                       | H      | ATG/TAA                    |
| <i>cox3</i>                               | 4325–5102 (778)                       | H      | ATG/T                      |
| tRNA-Gly (G)                              | 5103–5162 (60)                        | H      |                            |
| <i>nad3</i>                               | 5163–5507 (345)                       | H      | ATT/TAA                    |
| tRNA-Ala (A)                              | 5516–5578 (63)                        | H      |                            |
| tRNA-Arg (R)                              | 5583–5640 (58)                        | H      |                            |
| tRNA-Asn (N)                              | 5640–5703 (64)                        | H      |                            |
| tRNA-Ser <sup>AGA</sup> (S1)              | 5702–5758 (57)                        | H      |                            |
| tRNA-Glu (E)                              | 5761–5823 (63)                        | H      |                            |
| <i>nad1</i>                               | 5803–6763 (961)                       | L      | ATT/T                      |
| tRNA-Leu <sup>UUA</sup> (L <sub>2</sub> ) | 6764–6826 (63)                        | L      |                            |
| <i>rrnL</i>                               | 6836–8035 (1200)                      | L      |                            |
| tRNA-Val (V)                              | 8039–8100 (62)                        | L      |                            |
| <i>rrnS</i>                               | 8101–8788 (688)                       | L      |                            |
| Non-coding region (NCR1)                  | 8789–9111 (323)                       |        |                            |
| tRNA-Ile (I)                              | 9112–9177 (66)                        | H      |                            |
| tRNA-Gln (Q)                              | 9180–9247 (68)                        | L      |                            |
| tRNA-Phe (F)                              | 9250–9308 (59)                        | L      |                            |
| <i>nad5</i>                               | 9313–10971 (1659)                     | L      | ATT/TAA                    |
| tRNA-His (H)                              | 10972–11035 (64)                      | L      |                            |
| <i>nad4</i>                               | 11040–12362 (1323)                    | L      | ATG/TAA                    |
| <i>nad4L</i>                              | 12356–12631 (276)                     | L      | ATG/TAA                    |
| tRNA-Thr (T)                              | 12638–12698 (61)                      | H      |                            |
| tRNA-Pro (P)                              | 12703–12764 (62)                      | L      |                            |
| <i>nad6</i>                               | 12771–13205 (435)                     | H      | ATT/TAA                    |
| <i>cytb</i>                               | 13210–14286 (1077)                    | H      | ATG/TAA                    |
| tRNA-Ser <sup>UCA</sup> (S2)              | 14295–14358 (64)                      | H      |                            |
| tRNA-Leu <sup>CUA</sup> (L1)              | 14361–14419 (59)                      | L      |                            |
| Non-coding region (NCR2)                  | 14421–14750 (330)                     |        |                            |
| tRNA-Cys (C)                              | 14753–14806 (54)                      | H      |                            |

### Genetic distances and phylogenetic analyses

The concatenated matrix with CDSs indicates that *A. triste* has a 95.8% of identity with *A. maculatum* (MW719251), a record provided by the Oklahoma State University Tick Rearing Facility, USA (Brenner & Raghavan, 2021; Table 2). *A. ovale* has also a 80.6% of identity with *A. aureolatum*, and the species of the *A. cajennense* complex between 76.1 and 81.9% (Table 2).

**TABLE 2.** Percent identity matrix of concatenated protein-coding genes (CDS).

| Species                                | 1    | 2    | 3    | 4    | 5    | 6    | 7    | 8    | 9    | 10   | 11   | 12   | 13   | 14 |
|----------------------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|----|
| 1 <i>A. triste</i> (This study)        | -    |      |      |      |      |      |      |      |      |      |      |      |      |    |
| 2 <i>A. maculatum</i> (MW71925)        | 95.8 | -    |      |      |      |      |      |      |      |      |      |      |      |    |
| 3 <i>A. aureolatum</i> (SRR4301110)    | 76.7 | 76.9 | -    |      |      |      |      |      |      |      |      |      |      |    |
| 4 <i>A. ovale</i> (MT554102)           | 77.6 | 77.2 | 80.6 | -    |      |      |      |      |      |      |      |      |      |    |
| 5 <i>A. ovale</i> (MT554103)           | 77.6 | 77.3 | 80.6 | 97.8 | -    |      |      |      |      |      |      |      |      |    |
| 6 <i>A. cajennense</i> s.s. (OP901701) | 78.1 | 78   | 76.2 | 76.7 | 76.7 | -    |      |      |      |      |      |      |      |    |
| 7 <i>A. cajennense</i> s.s. (OP901707) | 78   | 77.9 | 76.2 | 76.7 | 76.6 | 99.8 | -    |      |      |      |      |      |      |    |
| 8 <i>A. patinoi</i> (OP901704)         | 77.9 | 77.8 | 76.2 | 76.7 | 76.6 | 88.1 | 88.1 | -    |      |      |      |      |      |    |
| 9 <i>A. mixtum</i> (OP901702)          | 78.3 | 78   | 76.7 | 76.8 | 76.9 | 87.6 | 87.5 | 87.6 | -    |      |      |      |      |    |
| 10 <i>A. mixtum</i> (OP901703)         | 78.2 | 77.9 | 76.6 | 76.8 | 76.9 | 87.4 | 87.3 | 87.6 | 95.6 | -    |      |      |      |    |
| 11 <i>A. sculptum</i> (NC_032369)      | 78   | 77.6 | 76.2 | 76.4 | 76.3 | 85.7 | 85.6 | 85.3 | 85.5 | 85.7 | -    |      |      |    |
| 12 <i>A. sculptum</i> (OP901706)       | 78.1 | 77.6 | 76.1 | 76.4 | 76.3 | 85.6 | 85.6 | 85.3 | 85.5 | 85.7 | 99.4 | -    |      |    |
| 13 <i>A. tonelliae</i> (OP901705)      | 78.4 | 78   | 77   | 77.5 | 77.4 | 82   | 81.9 | 82.2 | 82.3 | 82.3 | 82   | 81.9 | -    |    |
| 14 <i>D. nitens</i> (NC_023349)        | 73.6 | 73.5 | 71.1 | 71.9 | 71.9 | 72.8 | 72.7 | 72.7 | 73.5 | 73.1 | 73.1 | 73.1 | 72.7 | -  |



**FIGURE 3.** Phylogenetic tree based on the concatenation of 13 CDS and two rRNA genes. The tree was inferred with the best-fit substitution model and the topology shown corresponds to Bayesian Inference (BI: MrBayes v3.2.7a) and Maximum Likelihood (ML: IQ-TREE). Branch support indicated bootstrap percentages (ML/BI). Accession codes are in square brackets and mitogenome obtained in this study are shown in bold.

The interspecific genetic distances between *A. maculatum* and *A. triste* from each of the genes (CDSs and rRNA) are shown in Table 3. The *nd3* and *nd5* genes exhibited the greatest divergence, with 5.4% and 5% respectively. The phylogenetic relationships of 14 tick species based on the

concatenated Matrix-NT, using Maximum Likelihood (ML) and Bayesian Inference (BI) analyses, are shown in Figure 3. The topologies of the trees using ML and BI were identical, recovering a highly supported (100%) clade constituted by the two species of the *A. maculatum* group (*A. maculatum* and *A. triste*). Another well supported (73/100% respectively for ML/BI) was constituted by taxa of the *A. ovale* group. Similarly, the taxa of the *A. cajennense* complex were recovered in a separated clade with a 100% of support in both ML/BI reconstructions (Figure 3).

**TABLE 3.** Pairwise distance (in percentage) of mitochondrial genes (CDS and rRNA) between *A. maculatum* and *A. triste*, according to the *p*-distance method.

|                               | Genes        | <i>A. maculatum</i> (MW71925) |
|-------------------------------|--------------|-------------------------------|
| <i>A. triste</i> (This study) | <i>atp6</i>  | 3.9                           |
|                               | <i>atp 8</i> | 4.3                           |
|                               | <i>cox 1</i> | 4.2                           |
|                               | <i>cox 2</i> | 3.0                           |
|                               | <i>cox 3</i> | 3.8                           |
|                               | <i>cybt</i>  | 3.7                           |
|                               | <i>nd1</i>   | 3.6                           |
|                               | <i>nd2</i>   | 3.7                           |
|                               | <i>nd3</i>   | 5.4                           |
|                               | <i>nd4</i>   | 4.1                           |
|                               | <i>nd4L</i>  | 3.3                           |
|                               | <i>nd5</i>   | 5.0                           |
|                               | <i>nd6</i>   | 3.4                           |
|                               | <i>12S</i>   | 2.1                           |
|                               | <i>16S</i>   | 2.9                           |

## Discussion

### *Content and organization of the mitochondrial genome of Amblyomma triste*

The mitogenome of *A. triste* obtained in the present study is the first one recorded for this species, and it is like the mitogenomes reported for other arthropods including ticks (Montagna *et al.* 2012; Liu *et al.* 2013; Cameron *et al.* 2014; Brenner & Raghavan 2021).

In ticks, mitogenomes exhibit size ranges between 14 to 16 kb, circular organization, double-stranded DNA, with 37 genes: 13 CDSs, 22 tRNAs genes, and two rRNA genes, as well as two control or non-coding regions (NCRs); the arrangement of the genes of is similar to that found in the majority of species of Ixodidae ticks (Burger *et al.* 2013; Simon & Hadrys 2013; de Lima *et al.* 2017; Li & Liang 2018; Wang *et al.* 2019; Kelava *et al.* 2021; Cotes-Perdomo *et al.* 2023a).

In the *A. triste* mitogenome, the start codons (ATT and ATG) for the CDSs correspond to the typical start codons found in insects, which are the same as those commonly adopted by ticks (ATN) (Liu *et al.* 2013; Wang *et al.* 2019; Uribe *et al.* 2020; Brenner & Raghavan 2021). Regarding the termination codons (TAA and T), the CDSs correspond to the termination codons, which in ticks are mainly TAA and TAG, but sometimes “T” or “TA” may be converted into a complete termination codon by polyadenylation after translation (Montagna *et al.* 2012; Liu *et al.* 2013; de Lima *et al.* 2017; Uribe *et al.* 2020; Brenner & Raghavan 2021; Chavatte & Octavia 2021).

The mitochondrial rRNA genes (*12S* and *16S*) of *A. triste* exhibit a complex functional structure with a relatively slow evolution rate; this is particularly important as these genes have long been used as markers in tick population genetics, and phylogenetic and systematic studies (Araya-Anchetta *et al.* 2015; Wang *et al.* 2019).

Tick mitogenomes contain unique copy of the *12S* and *16S* rRNA genes, and due to gene rearrangement, the position of the rRNA genes shifts, whereas the gene order and the location in the N strand remain unchanged; therefore, the combined use of CDSs and rRNA genes is recommended for taxonomic studies, particularly when involving cryptic species and complexes (Burger *et al.* 2014; Mans *et al.* 2019; Wang *et al.* 2019).

The size and structure of the tRNA genes in this study (54 to 68 nucleotides) are consistent with the ranges reported for other tick species (50 to 90 bp), and most tRNA genes have a complete cloverleaf structure (Liu *et al.* 2013; Wang *et al.* 2019; Chavatte & Octavia 2021), except for trnS1 and trn-C, which lacks DHU arm or D-arm. This is a common feature in most animal species, including ticks (Liu *et al.* 2013; Cameron *et al.* 2014; Williams-Newkirk *et al.* 2015; Wang *et al.* 2019; Chavatte & Octavia 2021). Mitochondrial tRNA-C secondary structures are variable among tick species with some missing D-arm and/or T-arm and some having standard cloverleaf structure (Burger *et al.* 2012; Montagna *et al.* 2012; Williams-Newkirk *et al.* 2015).

Similarly, tick mitogenomes may contain two non-coding regions (NCRs), as in *Metastrata*, or one as in *Ixodes* (non-Australasian *Prostrata*), and *Argasidae*. In both cases, these regions have a conserved location within the mitogenome (Montagna *et al.* 2012; Kelava *et al.* 2021). Alternatively, they may be specific to the group, for example, *Metastrata* and Australasian *Ixodes* (*Prostrata*) have two non-coding regions, but with different locations within the mitogenome (Montagna *et al.* 2012; Liu *et al.* 2022). Changes in the non-coding regions of the mitochondrial DNA suggest potential differences in replication/transcription mechanisms, as NCRs contain regulatory elements (Burger *et al.* 2012; Liu *et al.* 2013; Wang *et al.* 2019; Chavatte & Octavia 2021).

#### *Genetic distances and phylogenetic analyses*

Mitogenomes can provide a deeper understanding of tick systematics, allowing the resolution of phylogenetic relationships that are problematic when using partial genes (Wang *et al.* 2019; Kelava *et al.* 2021; Mohamed *et al.* 2022). In this study, the interspecific distances of CDSs+rRNA genes between *A. triste* and *A. maculatum* were 4.2%, similar to the reported by Cotes-Perdomo *et al.* (2023b) when comparing the complete genomes of two species of the *A. marmoreum* complex (*Amblyomma nuttalli* and *Amblyomma sparsum*) with an interspecific distance of 4.4%. In comparison with other *Amblyomma* species, the intra-specific distances found were: *A. ovale* (2.2%); *A. cajennense* s.s. (0.2%); *A. mixtum* (4.4%); *A. sculptum* (0.6%), divergence percentage is consistent with the findings reported by Cotes-Perdomo *et al.* (2023a) for the *A. cajennense* complex. For other tick genera, Reynolds *et al.* (2022) reported intra-specific divergences ranging from 5.8% to 10.6% for *Dermacentor albipictus*, suggesting the existence of a species complex, and recommend further evaluation of other genes (*nad1*, *nad2*, *nad5*, *cox1*, and *atp8*) that may contribute to highlighting the genetic differences.

Traditionally, the *A. maculatum* group has been studied using divergences of gene fragments such as *12S*, *16S*, and *cox1*, which have been the most commonly used and reported genes for hard ticks (Beati & Keirans 2001; Marrelli *et al.* 2007; Beati *et al.* 2013; Paternina *et al.* 2016; Rivera-Páez *et al.* 2017; 2018; Lado *et al.* 2018). Our results demonstrate that these genes do not exhibit the highest interspecific divergences. For that reason, further research involving mitogenomes or other genes that display greater genetic distances should be conducted. This could be useful for differentiation of cryptic species such as *A. cajennense* complex, *A. marmoreum* complex and *A. ovale* complexes.

As reported by Ossa-López *et al.* (2022), the genetic distances of each of the evaluated genes (1.5–1.8% for *12S*; 2.3–7.1% for *16S*, and 4.9–7.0% for *cox1*) obtained between *A. triste* and *A. maculatum* from Colombia are larger than those reported by Lado *et al.* (2018), although they are lower than those found between *A. tigrinum* compared to all morphotypes that comprise the *A. maculatum* group (3.9–4.3%). The phylogenetic results also confirmed differences between *A. maculatum* from USA and Colombia based on genetic distances of each of the evaluated genes (1.2–1.8 for *12S*; 1.5–5.8% for *16S*, and 3–3.7% for *cox1*).

Although this study emphasizes the divergence percentages for the CDSs + rRNA genes, with *12S* and *16S* showing low percentages of 2.1% and 2.9% respectively, in comparison to the CDSs: *nd3*, *nd5*, *atp8*, and *cox1* stand out with interspecific divergences of 5.4%, 5.0%, 4.3%, and 4.2% respectively. Three of these genes (*nd5*, *atp8*, and *cox1*) align with the recommendations of Reynolds *et al.* (2022), and should be explored in phylogenetic studies, using the primers designed in our study for *A. maculatum* (MW719251), which can be implemented for other populations or species (Supplementary Material 2).

Finally, there are still several questions and gaps in the systematics and diversity of hard ticks. This study shows that the *A. maculatum* group comprises at least two species in Colombia. We highlight the need to establish both intra- and inter-specific divergences among the species within the *A. maculatum* group, new information on the mitogenomes of the different morphotypes reported by Lado *et al.* (2018), which originate from diverse geographic areas, is needed. Also, obtaining and evaluating the mitogenome of *A. tigrinum*, is crucial for elucidating both intra- and interspecific divergences within the species that comprise the *A. maculatum* group, and to carry out experimental crosses to obtain additional evidence for species recognition.

## Acknowledgments

Ministerio de Ciencia, Tecnología e innovación of Colombia - Minciencias Program “Relación, distribución, taxonomía de especies de garrapatas asociadas a mamíferos silvestres en zonas endémicas de rickettsiosis en Colombia. Un acercamiento a la comprensión de la relación vectores patógenos-reservorios” (Code: 120385270267 and CTO 80740- 200-2021) – Project “Garrapatas asociadas a mamíferos silvestres en el departamento de Caldas: Diversidad, detección de patógenos y distribución (Code:71717)”. Minciencias for funding the PhD in Science-Biology of Paula Andrea Ossa López "Convocatoria del Fondo de Ciencia, Tecnología e Innovación del Sistema General de Regalías para la conformación de una lista de proyectos elegibles para ser viabilizados, priorizados y aprobados por el OCAD dentro del Programa de Becas de Excelencia cohorte 1–2019". We thank the Universidad de Caldas, Unidad Administrativa Especial de Salud de Arauca. Award “For Women in Science 2022” conducted in collaboration with L'Oréal, Minciencias, ICETEX and La Comisión Nacional de Cooperación con la UNESCO. Part of the laboratory work was conducted in and with the support of the Molecular Systematics Lab (<https://www.mncn.csic.es/en/investigación/servicios-cientifico-tecnicos/molecular-systematics-laboratory>) of MNCN-CSIC, José Gutiérrez Abascal, Madrid, Spain.

## Funding

This project was funded by the Vicerrectoría de Investigaciones y Posgrados - Universidad de Caldas proyecto “Morfología interna y marcadores moleculares en garrapatas (Acari: Ixodidae): una aproximación a las interacciones con pequeños mamíferos y sus patógenos” [code 0318322].

Ministerio de Ciencia, Tecnología e Innovación of Colombia - Minciencias Program “Relación, distribución, taxonomía de especies de garrapatas asociadas a mamíferos silvestres en zonas endémicas de rickettsiosis en Colombia. Un acercamiento a la comprensión de la relación vectores patógenos-reservorios”, granted by the (Code: 120385270267 and CTO 80740- 200-2021). Minciencias for funding the PhD in Science-Biology of Paula Andrea Ossa López "Convocatoria del Fondo de Ciencia, Tecnología e Innovación del Sistema General de Regalías para la conformación de una lista de proyectos elegibles para ser viabilizados, priorizados y aprobados por el OCAD dentro del Programa de Becas de Excelencia cohorte 1–2019". Part of this work was funded by Award “For Women in Science 2022” conducted in collaboration with L'Oréal, Minciencias, ICETEX and La Comisión Nacional de Cooperación con la UNESCO. Comunidad de Madrid for Atracción de Talento contract (REFF 2019-T2/AMB-13166) of JEU.

## References

- Abarca, K., López, J., Acosta-Jamett, G., Lepe, P., Soares, J.F. & Labruna, M.B. (2012) A third *Amblyomma* species and the first tick-borne rickettsia in Chile. *Journal of Medical Entomology*, 49, 219–222.  
<https://doi.org/10.1603/ME11147>
- Allerdice, M.E., Snellgrove, A.N., Hecht, J.A., Hartzer, K., Jones, E.S., Biggerstaff, B.J., Ford, S.L., Karpathy, S.E., Delgado-de la Mora, J., Delgado-de la Mora, D. & Licona-Enriquez, J.D. (2020) Reproductive incompatibility between *Amblyomma maculatum* (Acari: Ixodidae) group ticks from two disjunct geographical regions within the USA. *Experimental and Applied Acarology*, 82(4), 543–557.  
<https://doi.org/10.1007/s10493-020-00557-4>
- Andrews, S. (2010) FastQC: A Quality Control Tool for High Throughput Sequence Data [Online]. Available online at <http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>
- Araya-Anchetta, A., Busch, J.D., Scoles, G.A. & Wagner, D.M. (2015) Thirty years of tick population genetics: a comprehensive review. *Infection, Genetics and Evolution*, 29, 164–179.  
<https://doi.org/10.1016/j.meegid.2014.11.008>
- Baneth, G. (2014) Tick-borne infections of animals and humans: a common ground. *International Journal for Parasitology*, 44(9), 591–596.  
<https://doi.org/10.1016/j.ijpara.2014.03.011>
- Barros-Battesti, D.M., Arzua, M. & Bechara, G.H. (2006) *Carrapatos de importância médico veterinária da região neotropical: Um guia ilustrado para identificação de espécies*. São Paulo, Vox/ICTTD-3/Butantan. 223 pp.
- Beati, L. & Keirans, J.E. (2001) Analysis of the systematic relationships among ticks of the genera *Rhipicephalus* and *Boophilus* (Acari: Ixodidae) based on mitochondrial 12S ribosomal DNA gene sequences and morphological characters. *Journal of Parasitology*, 87, 32–48.  
[https://doi.org/10.1645/0022-3395\(2001\)087\[0032:AOTSRA\]2.0.CO;2](https://doi.org/10.1645/0022-3395(2001)087[0032:AOTSRA]2.0.CO;2)
- Beati, L., Nava, S., Burkman, E.J., Barros-Battesti, D., Labruna, M.B., Guglielmone, A.A., Cáceres, A.G., Guzmán-Cornejo, C., León, R., Durden, A.L. & Faccini, J.L. (2013) *Amblyomma cajennense* (Fabricius, 1787) (Acari: Ixodidae), the cayenne tick: phylogeography and evidence for allopatric speciation. *BMC Evolutionary Biology*, 13, 267.  
<https://doi.org/10.1186/1471-2148-13-267>
- Bernt, M., Donath, A., Jühling, F., Externbrink, F., Florentz, C., Fritzsche, G., Pütz, J., Middendorf, M. & Stadler, P.F. (2013) MITOS: improved de novo metazoan mitochondrial genome annotation. *Molecular Phylogenetics and Evolution*, 69, 313–319.  
<https://doi.org/10.1016/j.ympev.2012.08.023>
- Bezerra-Santos, M.A., de Macedo, L.O., Nguyen, V.L., Manoj, R.R., Laidoudi, Y., Latrofa, M.S., Beugnet, F. & Otranto, D. (2022) *Cercopithifilaria* spp. in ticks of companion animals from Asia: new putative hosts and vectors. *Ticks and Tick-borne Diseases*, 13(4), 101957.  
<https://doi.org/10.1016/j.ttbdis.2022.101957>
- Bolger, A.M., Lohse, M. & Usadel, B. (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, 30, 2114–2120.  
<https://doi.org/10.1093/bioinformatics/btu170>
- Brenner, A.E. & Raghavan, R. (2021) Complete mitochondrial genome sequence of the Gulf Coast tick (*Ambly-*

- omma maculatum*). *Microbiology Resource Announcements*, 10, e00431–21.  
<https://doi.org/10.1128/mra.00431-21>
- Burger, T.D., Shao, R. & Barker S.C. (2013) Phylogenetic analysis of the mitochondrial genomes and nuclear rRNA genes of ticks reveals a deep phylogenetic structure within the genus *Haemaphysalis* and further elucidates the polyphyly of the genus *Amblyomma* with respect to *Amblyomma sphenodonti* and *Amblyomma elaphense*. *Ticks and Tick-borne Diseases*, 4(4), 265–74.  
<https://doi.org/10.1016/j.ttbdis.2013.02.002>
- Burger, T.D., Shao, R., Beati, L., Miller, H. & Barker, S.C. (2012) Phylogenetic analysis of ticks (Acari: Ixodida) using mitochondrial genomes and nuclear rRNA genes indicates that the genus *Amblyomma* is polyphyletic. *Molecular Phylogenetics and Evolution*, 64(1), 45–55.  
<https://doi.org/10.1016/j.ympev.2012.03.004>
- Burger, T.D., Shao, R., Labruna, M.B. & Barker, S.C. (2014) Molecular phylogeny of soft ticks (Ixodida: Argasidae) inferred from mitochondrial genome and nuclear rRNA sequences. *Ticks and Tick-borne Diseases*, 5, 195–207.  
<https://doi.org/10.1016/j.ttbdis.2013.10.009>
- Cameron, S.L. (2014) Insect mitochondrial genomics: implications for evolution and phylogeny. *Annual Review of Entomology*, 59, 95–117.  
<https://doi.org/10.1146/annurev-ento-011613-162007>
- Chavatte, J.M. & Octavia, S. (2021) The complete mitochondrial genome of *Dermacentor (Indocentor) auratus* (Acari, Ixodidae). *Parasite*, 28, 6.  
<https://doi.org/10.1051/parasite/2021002>
- Colombo, V.C., Antoniazzi, L.R., Fasano, A.A., Beldoménico, P.M. & Nava, S. (2016) *Amblyomma triste* en simpatría con *Amblyomma tigrinum* (Acari: Ixodidae) en la provincia de Santa Fe, Argentina. *Medicina (Buenos Aires)*, 76, 304–306.
- Cotes-Perdomo, A.P., Nava, S., Castro, L.R., Rivera-Paéz, F.A., Cortés-Vecino, J.A. & Uribe, J.E. (2023a) Phylogenetic relationships of the *Amblyomma cajennense* complex (Acari: Ixodidae) at mitogenomic resolution. *Ticks and Tick-borne Diseases*, 14(3), 102125.  
<https://doi.org/10.1016/j.ttbdis.2023.102125>
- Cotes-Perdomo, A.P., Sanchez-Vialas, A., Thomas, R., Jenkins, A. & Uribe, J.E. (2023b) New insights into the systematics of the Afrotropical *Amblyomma marmoreum* complex (Acari, Ixodidae) and a novel *Rickettsia africae* strain using morphological and metagenomic approaches. *bioRxiv*, 2023.08.18.553479.  
<https://doi.org/10.1101/2023.08.18.553479>
- Crisuolo, A. & Gribaldo, S. (2010) BMGE (Block Mapping and Gathering with Entropy): a new software for selection of phylogenetic informative regions from multiple sequence alignments. *BMC Ecology and Evolution*, 10, 210.  
<https://doi.org/10.1186/1471-2148-10-210>
- Cuervo, P.F., Flores, F.S., Venzal, J.M. & Nava, S. (2021) Niche divergence among closely related taxa provides insight on evolutionary patterns of ticks. *Journal of Biogeography*, 48(11), 2865–2876.  
<https://doi.org/10.1111/jbi.14245>
- de la Fuente, J., Estrada-Pena, A., Venzal, J.M., Kocan, K.M. & Sonenshine, D.E. (2008) Overview: Ticks as vectors of pathogens that cause disease in humans and animals. *Frontiers in Bioscience*, 1(13), 6938–6946.  
<https://doi.org/10.2741/3200>
- de Lima, P.H.C., Barcelos, R.M., Klein, R.C., Vidigal, P.M.P., Montandon, C.E., Fabres-Klein, M.H., Dergam, J.A. & Mafra, C. (2017) Sequencing and comparative analysis of the *Amblyomma sculptum* mitogenome. *Veterinary Parasitology*, 30, 247, 121–128.  
<https://doi.org/10.1016/j.vetpar.2017.10.007>
- Demoner, L.deC., Rubini, A.S., Paduan, K.dosS., Metzger, B., de Paula Antunes, J.M., Martins, T.F., Mathias, M.I. & O'Dwyer, L.H. (2013) Investigation of tick vectors of *Hepatozoon canis* in Brazil. *Ticks and Tick-borne Diseases*, 4(6), 542–546.  
<https://doi.org/10.1016/j.ttbdis.2013.07.006>
- Dumler, J.S., Barbet, A.F., Bekker, C.P., Dasch, G.A., Palmer, G.H., Ray, S.C., Rikihisa, Y. & Rurangirwa, F.R. (2001) Reorganization of genera in the families Rickettsiaceae and Anaplasmataceae in the order Rickettsiales: unification of some species of *Ehrlichia* with *Anaplasma*, *Cowdria* with *Ehrlichia* and *Ehrlichia* with *Neorickettsia*, descriptions of six new species combinations and designation of *Ehrlichia equi* and ‘HGE agent’ as subjective synonyms of *Ehrlichia phagocytophila*. *International Journal of Systematic and Evolutionary Microbiology*, 51, 2145–2165.  
<https://doi.org/10.1099/00207713-51-6-2145>
- Ewing, S.A., Panciera, R.J., Mathew, J.S., Cummings, C.A. & Kocan, A.A. (2000) American canine hepatozoono-

- sis: an emerging disease in the New World. *Annals of the New York Academy of Sciences*, 916, 81–92.  
<https://doi.org/10.1111/j.1749-6632.2000.tb05277.x>
- Estrada-Peña, A., Venzal, J.M., Mangold, A.J., Cafrune, M.M. & Guglielmone, A.A. (2005) The *Amblyomma maculatum* Koch, 1844 (Acari: Ixodidae: Amblyomminae) tick group: diagnostic characters, description of the larva of *A. parvitarsum* Neumann, 1901, 16S rDNA sequences, distribution and hosts. *Systematic Parasitology*, 60(2), 99–112.  
<https://doi.org/10.1007/s11230-004-1382-9>
- Felsenstein, J. (1981) Evolutionary trees from DNA sequences: a maximum likelihood approach. *Journal of Molecular Evolution*, 17, 368–376.  
<https://doi.org/10.1007/BF01734359>
- Folmer, O., Black, M., Hoeh, W., Lutz, R. & Vrijenhoek, R. (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology*, 3, 294–299.
- Guglielmone, A.A., Mangold, A.J., Luciani, C.E., & Viñabal, A. (2000) *Amblyomma tigrinum* (Acari: Ixodidae) in relation to phytogeography of central-northern Argentina with notes on hosts and seasonal distribution. *Experimental and Applied Acarology*, 24, 983–989.  
<https://doi.org/10.1023/A:1010775528628>
- Guglielmone, A.A., Estrada-Peña, A., Mangold, A.J., Barros-Battesti, D.M., Labruna, M.B., Martins, J.R., Venzal, J.M., Arzua, M. & Keirans, J.E. (2003) *Amblyomma aureolatum* (Pallas, 1772) and *Amblyomma ovale* Koch, 1844 (Acari: Ixodidae): hosts, distribution and 16S rDNA sequences. *Veterinary Parasitology*, 113(3–4), 273–288.  
[https://doi.org/10.1016/S0304-4017\(03\)00083-9](https://doi.org/10.1016/S0304-4017(03)00083-9)
- Guglielmone, A.A., Nava, S. & Díaz, M. (2011) Relationships of south American marsupials (Didelphimorphia, Microbiotheria and Paucituberculata) and hard ticks (Acari: Ixodidae) with distribution of four species of Ixodes. *Zootaxa*, 3086, 1–30  
<https://doi.org/10.11646/zootaxa.3086.1.1>
- Guglielmone, A.A. & Robbins, R.G. (2018) *Hard ticks (Acari: Ixodida: Ixodidae) parasitizing humans: a global overview*. Cham, Switzerland, Springer international publishing, 314 pp.  
<https://doi.org/10.1007/978-3-319-95552-0>
- Guglielmone, A., Petney, T. & Robbins, R. (2020) Ixodidae (Acari: Ixodoidea): descriptions and redescrptions of all known species from 1758 to December 31, 2019. *Zootaxa*, 4871, 1–322.  
<https://doi.org/10.11646/zootaxa.4871.1.1>
- Guglielmone, A.A., Nava, S. & Robbins, R.G. (2021) *Neotropical Hard Ticks (Acari: Ixodida: Ixodidae)*. Cham, Switzerland, Springer International Publishing, XXV, 486 pp.  
<https://doi.org/10.1007/978-3-030-72353-8>
- Guglielmone, A.A., Nava, S. & Robbins, R.G. (2023) Geographic distribution of the hard ticks (Acari: Ixodida: Ixodidae) of the world by countries and territories. *Zootaxa*, 5251(1), 1–274.  
<https://doi.org/10.11646/zootaxa.5251.1.1>
- Guzmán-Cornejo, C., Perez, T.M., Nava, S. & Guglielmone, A.A. (2006) Confirmation of the presence of *Amblyomma triste* Koch, 1844 (Acari: Ixodidae) in Mexico. *Systematic & Applied Acarology*, 11, 47–50.  
<https://doi.org/10.11158/saa.11.1.5>
- Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K.F., von Haeseler, A. & Jermini, L.S. (2017) ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods*, 14, 587–589.  
<https://doi.org/10.1038/nmeth.4285>
- Katoh, K. & Standley, D.M. (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution*, 30, 772–780.  
<https://doi.org/10.1093/molbev/mst010>
- Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper, A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P. & Drummond, A. (2012) Geneious basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics*, 28(12), 1647–9.  
<https://doi.org/10.1093/bioinformatics/bts199>
- Kelava, S., Mans, B.J., Shao, R., Moustafa, M.A.M., Matsuno, K., Takano, A., Kawabata, H., Sato, K., Fujita, H., Ze, C., Plantard, O., Hornok, S., Gao, S., Barker, D., Barker, S.C. & Nakao, R. (2021) Phylogenies from mitochondrial genomes of 120 species of ticks: insights into the evolution of the families of ticks and of the genus *Amblyomma*. *Ticks and Tick-borne Diseases*, 12, 101577.  
<https://doi.org/10.1016/j.ttbdis.2020.101577>
- Koch, C.L. (1844) Systematische übersicht über die Ordnung der Zecken. *Archiv für Naturgeschichte*, 10, 217–

- Kneubehl, A.R., Muñoz-Leal, S., Filatov, S. de Klerk, D.G., Pienaar, R., Lohmeyer, K.H., Bermúdez, S.E., Suriyamongkol, T., Mali, I., Kanduma, E., Latif, A.A., Sarih, M., Bouattour, A., Pérez de León, A.A., Teel, P.D., Labruna, M.B., Mans, B.J. & Lopez, J.E. (2022) Amplification and sequencing of entire tick mitochondrial genomes for a phylogenomic analysis. *Scientific Reports*, 12, 19310.  
<https://doi.org/10.1038/s41598-022-23393-5>
- Kohls, G.M. (1956) Concerning the identity of *Amblyomma maculatum*, *A. tigrinum*, *A. triste* and *A. ovatum* of Koch, 1844. *Proceedings of the Entomological Society of Washington*, 58, 143–147.
- Lado, P., Nava, S., Mendoza-Urbe, L., Caceres, A.G., Delgado-de la Mora, J., Licona-Enriquez, J. D., Delgado-de la Mora, D., Labruna, M.B., Durden, L.A., Allerdice, M.E.J., Paddock, C.D., Szabó, M.P.J., Venzal, J.M., Guglielmone, A.A. & Beati, L. (2018) The *Amblyomma maculatum* Koch, 1844 (Acari: Ixodidae) group of ticks: Phenotypic plasticity or incipient speciation? *Parasites & Vectors*, 11, 610.  
<https://doi.org/10.1186/s13071-018-3186-9>
- Li, K. & Liang, A.P. (2018) Hemiptera mitochondrial control region: new sights into the structural organization, phylogenetic utility, and roles of tandem repetitions of the noncoding segment. *International Journal of Molecular Sciences*, 19, 1292.  
<https://doi.org/10.3390/ijms19051292>
- Liu, G.H., Chen, F., Chen, Y.Z., Song, H.Q., Lin, R.Q., Zhou, D.H. & Zhu, X.Q. (2013) Complete mitochondrial genome sequence data provides genetic evidence that the brown dog tick *Rhipicephalus sanguineus* (Acari: Ixodidae) represents a species complex. *International Journal of Biological Sciences*, 9(4), 361–369.  
<https://doi.org/10.7150/ijbs.6081>
- Liu, J., Yu, J., Yu, X., Bi, W., Yang, H., Xue, F., Zhang, G., Zhang, J., Yi, D., Ma, R., Zhou, Y., Lan, G., Gu, J., Wu, W., Li, Z. & Qi, G. (2022) Complete mitogenomes of ticks *Ixodes acutitarsus* and *Ixodes ovatus* parasitizing giant panda: Deep insights into the comparative mitogenomic and phylogenetic relationship of Ixodidae species. *Genes*, 13(11), 2049.  
<https://doi.org/10.3390/genes13112049>
- Mans, B.J., Featherston, J., Kvas, M., Pillay, K.A., Klerk, D.G., Pienaar, R., de Castro, M.H., Schwan, T.G., Lopez, J.E., Teel, P., Pérez de León, A.A., Sonenshine, D.E., Egekwu, N.I., Bakkes, D.K., Heyne, H., Kanduma, E.G., Nyangiwe, N., Bouattour, A. & Latif, A.A. (2019) Argasid and ixodid systematics: implications for soft tick evolution and systematics, with a new argasid species list. *Ticks and Tick-borne Diseases*, 10, 219–240.  
<https://doi.org/10.1016/j.tbbdis.2018.09.010>
- Mangold, A.J., Bargues, M.D. & Mas-Coma, S. (1998) Mitochondrial 16S rDNA sequences and phylogenetic relationships of species of *Rhipicephalus* and other tick genera among Metastrata (Acari: Ixodidae). *Parasitology Research*, 84(6), 478–484.  
<https://doi.org/10.1007/s004360050433>
- Marrelli, M.T., Souza, L.F., Marques, R.C., Labruna, M.B., Matioli, S.R., Tonon, A.P., Ribolla, P.E.M., Marinotti, O. & Schumakeret, T.T.S. (2007) Taxonomic and phylogenetic relationships between neotropical species of ticks from genus *Amblyomma* (Acari: Ixodidae) inferred from second internal transcribed spacer sequences of rDNA. *Journal of Medical Entomology*, 44, 222–228.  
<https://doi.org/10.1093/jmedent/44.2.222>
- Martins, L.A., Galletti, M.F., Ribeiro, J.M., Fujita, A., Costa, F.B., Labruna, M.B., Daffre, S. & Fogaça, A.C. (2017) The distinct transcriptional response of the midgut of *Amblyomma sculptum* and *Amblyomma aureolatum* ticks to *Rickettsia rickettsii* correlates to their differences in susceptibility to infection. *Frontiers in Cellular and Infection Microbiology*, 7, 129.  
<https://doi.org/10.3389/fcimb.2017.00129>
- Mathew, J.S., Ewing, S.A., Panciera, R.J. & Kocan, K.M. (1999) Sporogonic development of *Hepatozoon americanum* (Apicomplexa) in its definitive host, *Amblyomma maculatum* (Acarina). *Journal of Parasitology*, 85(6), 1023–1031.  
<https://doi.org/10.2307/3285663>
- Mathew, J.S., Ewing, S.A., Panciera, R.J. & Woods, J.P. (1998) Experimental transmission of *Hepatozoon americanum* Vincent-Johnson et al., 1997 to dogs by the Gulf Coast tick, *Amblyomma maculatum* Koch. *Veterinary Parasitology*, 80, 1–14.  
[https://doi.org/10.1016/S0304-4017\(98\)00189-7](https://doi.org/10.1016/S0304-4017(98)00189-7)
- Mendoza-Urbe, L. & Chávez-Chorocco, J. (2004) Ampliación geográfica de siete especies de *Amblyomma* (Acari: Ixodidae) y primer reporte de *A. oblongoguttatum* Koch, 1844 para Perú. *Revista Peruana de Entomología*, 44, 69–72
- Mertins, J.W., Moorhouse, A.S., Alfred, J.T. & Hutcheson, H.J. (2010) *Amblyomma triste* (Acari: Ixodidae): new

- North American collection records, including the first from the United States. *Journal of Medical Entomology*, 47, 536–542.  
<https://doi.org/10.1093/jmedent/47.4.536>
- Mohamed, W.M.A., Moustafa, M.A.M., Thu, M.J., Kakisaka, K., Chatanga, E., Ogata, S., Hayashi, N., Taya, Y., Ohari, Y., Naguib, D., Qiu, Y., Matsuno, K., Bawm, S., Htun, L. L., Barker, S.C., Katakura, K., Ito, K., Nonaka, N. & Nakao, R. (2022) Comparative mitogenomics elucidates the population genetic structure of *Amblyomma testudinarium* in Japan and a closely related *Amblyomma* species in Myanmar. *Evolutionary Applications*, 15, 1062–1079.  
<https://doi.org/10.1111/eva.13426>
- Montagna, M., Sasser, D., Griggio, F., Epis, S., Bandi, C. & Gissi, C. (2012) Tick-Box for 3'-End Formation of Mitochondrial Transcripts in Ixodida, Basal Chelicerates and *Drosophila*. *PLoS ONE*, 7(10), e47538.  
<https://doi.org/10.1371/journal.pone.0047538>
- Nava, S., Lareschi, M., Rebollo, C., Benítez Usher, C., Beati, L., Robbins, R.G., Durden, L.A., Mangold, A.J. & Guglielmone, A.A. (2007) The ticks (Acari: Ixodida: Argasidae, Ixodidae) of Paraguay. *Annals of Tropical Medicine and Parasitology*, 101, 255–270.  
<https://doi.org/10.1179/136485907X176319>
- Nava, S., Venzal, J.M.M., González-Acuña, D.G., Martins, T.F.F. & Guglielmone, A.A. (2017) Ticks of the southern cone of America: diagnosis, distribution, and hosts with taxonomy, ecology and sanitary importance. Elsevier, London, San Diego, 348 pp.
- Nguyen, L.T., Schmidt, H.A., von Haeseler, A. & Minh, B.Q. (2015) IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and Evolution*, 32, 268–274.  
<https://doi.org/10.1093/molbev/msu300>
- Norris, D.E., Klompen, J.S.H., Keirans, J.E. & Black, W.C. (1996) Population genetics of *Ixodes scapularis* (Acari: Ixodidae) based on mitochondrial 16S and 12S genes. *Journal of Medical Entomology*, 33, 78–89.  
<https://doi.org/10.1093/jmedent/33.1.78>
- Ortiz-Giraldo, M., Tobón-Escobar, W.D., Velásquez-Guarín, D., Usma-Marín, M.F., Ossa-López, P.A., Ramírez-Chaves, H.E., Carvajal-Agudelo, J.D. & Rivera-Páez, F.A. (2021) Ticks (Acari: Ixodoidea) associated with mammals in Colombia: a historical review, molecular species confirmation, and establishment of new relationships. *Parasitology Research*, 120(2), 383–394.  
<https://doi.org/10.1007/s00436-020-06989-6>
- Ossa-López, P.A., Robayo-Sánchez, L.N., Uribe, J.E., Ramírez-Hernández, A., Ramírez-Chaves, H.E., Cortés-Vecino, J.A. & Rivera-Páez, F.A. (2022) Extension of the distribution of *Amblyomma triste* Koch, 1844: Morphological and molecular confirmation of Morphotype I in Colombia. *Ticks and Tick-Borne Diseases*, 13, 101923.  
<https://doi.org/10.1016/j.ttbdis.2022.101923>
- Paddock, C.D. & Goddard, J. (2015) The Evolving Medical and Veterinary Importance of the Gulf Coast tick (Acari: Ixodidae). *Journal of Medical Entomology*, 52(2), 230–252.  
<https://doi.org/10.1093/jme/tju022>
- Paddock, C.D., Sumner, J.W., Comer, J.A., Zaki, S.R., Goldsmith, C.S., Goddard, J. & Ohl, C.A. (2004) *Rickettsia parkeri*: A newly recognized cause of spotted fever rickettsiosis in the United States. *Clinical Infectious Diseases*, 38, 805–811.  
<https://doi.org/10.1086/381894>
- Paddock, C.D. & Goddard, J. (2015) The Evolving Medical and Veterinary Importance of the Gulf Coast tick (Acari: Ixodidae). *Journal of Medical Entomology*, 52, 2, 230–252.  
<https://doi.org/10.1093/jme/tju022>
- Paternina, L.E., Verbel-Vergara, D. & Bejarano, E.E. (2016) Comparación y utilidad de las regiones mitocondriales de los genes 16S y COX1 para los análisis genéticos en garrapatas (Acari: Ixodidae). *Biomedical*, 36(2), 295–302.  
<https://doi.org/10.7705/biomedica.v36i2.3116>
- Rambaut, A. (2007) FigTree v1.4.3, A Graphical Viewer of Phylogenetic Trees. Available <http://tree.bio.ed.ac.uk/software/figtree>.
- Rannala, B. & Yang, Z. (1996) Probability distribution of molecular evolutionary trees: a new method of phylogenetic inference. *Journal of Molecular Evolution*, 43, 304–311.  
<https://doi.org/10.1007/BF02338839>
- Reynolds, S., Hedberg, M., Herrin, B. & Chelladurai, J.R.J.J. (2022) Analysis of the complete mitochondrial genomes of *Dermacentor albipictus* suggests a species complex. *Ticks and Tick-Borne Diseases*, 13(6), 102038.  
<https://doi.org/10.1016/j.ttbdis.2022.102038>

- Rivera-Páez, F.A., Sampieri, B.R., Labruna, M.B., da Silva, M.R., Martins, T.F. & Camargo-Mathias, M.I. (2017) Comparative analysis of germ cells and DNA of the genus *Amblyomma*: adding new data on *Amblyomma maculatum* and *Amblyomma ovale* species (Acari: Ixodidae). *Parasitology Research*, 116, 2883–2892. <https://doi.org/10.1007/s00436-017-5592-x>
- Rivera-Páez, F.A., Labruna, M.B., Martins, T.F., Perez, J.E., Castaño-Villa, G.J., Ossa-López, P.A., Gil, C.A., Sampieri, B., Aricapa-Giraldo, H.J. & Camargo-Mathias, M.I. (2018) Contributions to the knowledge of hard ticks (Acari: Ixodidae) in Colombia. *Ticks and Tick-Borne Diseases*, 9(1), 57–66. <https://doi.org/10.1016/j.ttbdis.2017.10.008>
- Romer, Y., Borrás, P., Govedic, F., Nava, S., Carranza, J. I., Santini, S., Armitano, R., & Lloveras, S. (2020) Clinical and epidemiological comparison of *Rickettsia parkeri* rickettsiosis, related to *Amblyomma triste* and *Amblyomma tigrinum*, in Argentina. *Ticks and Tick-Borne Diseases*, 11(4), 101436. <https://doi.org/10.1016/j.ttbdis.2020.101436>
- Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D.L., Darling, A., Höhna, S., Larget, B., Liu, L., Suchard, M.A. & Huelsenbeck, J.P. (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology*, 61, 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Santodomingo, A., Uribe, J.E., Lopez, G. & Castro, L.R. (2021) Phylogenetic insights into the genus *Amblyomma* in America, including the endangered species *Amblyomma albopictum*, *Amblyomma macfarlandi*, and *Amblyomma usingeri*. *International Journal of Acarology*, 47, 1–9. <https://doi.org/10.1080/01647954.2021.1933176>
- Schwarz, D. (1978) Estimating the dimension of a model. *Annals of Statistics*, 6(6), 461–464. <https://doi.org/10.1214/aos/1176344136>
- Simon, S. & Hadrys, H. (2013) A comparative analysis of complete mitochondrial genomes among Hexapoda. *Molecular Phylogenetics and Evolution*, 69, 393–403. <https://doi.org/10.1016/j.ympev.2013.03.033>
- Suwanbongkot, C., Langohr, I.M., Harris, E.K., Dittmar, W., Christofferson, R.C. & Macaluso, K.R. (2019) Spotted Fever Group Rickettsia Infection and Transmission Dynamics in *Amblyomma maculatum*. *Infection and Immunity*, 87(4), e00804–18. <https://doi.org/10.1128/IAI.00804-18>
- Tamura K., Stecher, G. & Kumar, S. (2021) MEGA 11: Molecular Evolutionary Genetics Analysis Version 11. *Molecular Biology and Evolution*, 38(7), 3022–3027. <https://doi.org/10.1093/molbev/msab120>
- Uribe, J.E., Nava, S., Murphy, K.R., Tarragona, E.L. & Castro, L.R. (2020) Characterization of the complete mitochondrial genome of *Amblyomma ovale*, comparative analyses and phylogenetic considerations. *Experimental and Applied Acarology*, 81(3), 421–439. <https://doi.org/10.1007/s10493-020-00512-3>
- Venzal, J.M., Portillo, A., Estrada-Peña, A., Castro, O., Cabrera, P.A. & Oteo, J.A. (2004) *Rickettsia parkeri* in *Amblyomma triste* from Uruguay. *Emerging Infectious Diseases*, 10, 1493–1495. <https://doi.org/10.3201/eid1008.030999>
- Wang, T., Zhang, S., Pei, T., Yu, Z. & Liu, J. (2019) Tick mitochondrial genomes: structural characteristics and phylogenetic implications. *Parasites & vectors*, 12(1), 451. <https://doi.org/10.1186/s13071-019-3705-3>
- Williams-Newkirk, A.J., Burroughs, M., Changavil, S.S. & Dash, G.A. (2015) The mitochondrial genome of the lone star tick (*Amblyomma americanum*). *Ticks and Tick-Borne Diseases*, 6, e793–e801. <https://doi.org/10.1016/j.ttbdis.2015.07.006>
- Yang, Z. & Rannala, B. (1997) Bayesian phylogenetic inference using DNA sequences: a Markov Chain Monte Carlo method. *Molecular Biology and Evolution*, 14, 717–724. <https://doi.org/10.1093/oxfordjournals.molbev.a025811>
- Zaharias, P., Pante, E., Gey, D., Fedosov, A.E. & Puillandre, N. (2020) Data, time and money: evaluating the best compromise for inferring molecular phylogenies of non-model animal taxa. *Molecular Phylogenetics and Evolution*, 42, 106660. <https://doi.org/10.1016/j.ympev.2019.106660>

*Submitted: 1 Sept. 2023; accepted by Xiao-Feng Xue: 24 Dec. 2023; published: 29 Jan. 2024*