

## **Socioeconomic Predictors of Intestinal Parasitic Infections Among Under-Five Children in Rural Dembiya, Northwest Ethiopia: A Community-Based Cross-sectional Study**

Authors: Gizaw, Zemichael, Addisu, Ayenew, and Gebrehiwot, Mulat

Source: Environmental Health Insights, 13(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/1178630219896804>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Socioeconomic Predictors of Intestinal Parasitic Infections Among Under-Five Children in Rural Dembiya, Northwest Ethiopia: A Community-Based Cross-sectional Study

Environmental Health Insights  
Volume 13: 1–6  
© The Author(s) 2019  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/1178630219896804



Zemichael Gizaw<sup>1</sup> , Ayenew Addisu<sup>2</sup> and Mulat Gebrehiwot<sup>1</sup>

<sup>1</sup>Department of Environmental and Occupational Health and Safety, Institute of Public Health, College of Medicine and Health Sciences, University of Gondar, Gondar, Ethiopia. <sup>2</sup>Department of Parasitology, School of Biomedical Science, College of Medicine and Health Sciences, University of Gondar, Gondar, Ethiopia.

## ABSTRACT

**BACKGROUND:** Soil-transmitted helminths and protozoan parasitic infections are endemic throughout the world. The problem of intestinal parasitic infection is higher among developing countries where children are the most vulnerable groups. Although health information related to parasitic infections is available globally, it is often limited in rural setups in least developed countries. This study was, therefore, conducted to assess socioeconomic predictors of intestinal parasitic infections among under-five children in rural Dembiya, Northwest Ethiopia.

**METHODS:** This cross-sectional study was conducted among 224 randomly selected households with under-five children. We used questionnaire to collect data and direct stool examination to identify intestinal parasitic infections. Adjusted odds ratio (AOR) with 95% confidence interval (CI) and  $P < .05$  was used to identify socioeconomic predictors of parasitic infections.

**RESULTS:** We found that 25.4% (95% CI = [20.2, 31.1]) under-five children had intestinal parasitic infection. *Ascaris lumbricoides* was the leading infection, which accounted 44 of 224 (19.6%). The prevalence of childhood intestinal parasitic infections was higher among households with no members whose education level is secondary and above (AOR = 3.36, 95% CI = [1.23, 9.17]). Similarly, intestinal parasitic infections were statistically associated with presence of 2 under-five children in a household (AOR = 3.56, 95% CI = [1.29, 9.82]), absence of frequent health supervision (AOR = 3.49, 95% CI = [1.72, 7.09]), larger family size (AOR = 2.30, 95% CI = [1.09, 4.85]), and poor household economic status (AOR = 2.58, 95% CI = [1.23, 5.41]).

**CONCLUSIONS:** Significant proportion of children was infected with intestinal parasitic infection in rural Dembiya. Educational status of family members, number of under-five children in a household, health supervision, family size, and wealth index were statistically associated with parasitic infections. Provision of anthelmintic drugs, health supervision, and health education targeted with transmission and prevention of infections are recommended.

**KEYWORDS:** Intestinal parasitic infections, socioeconomic predictors, under-five children, rural Dembiya

**RECEIVED:** November 24, 2019. **ACCEPTED:** November 25, 2019.

**TYPE:** Original Research

**FUNDING:** The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The study was funded by NALA foundation.

**DECLARATION OF CONFLICTING INTERESTS:** The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

**CORRESPONDING AUTHOR:** Zemichael Gizaw, Department of Environmental and Occupational Health and Safety, Institute of Public Health, College of Medicine and Health Sciences, University of Gondar, P.O.Box 196, Gondar, Ethiopia. Email: zemichael12@gmail.com

## Background

Helminth and protozoan parasites cause health problems in human. Twenty-five percent of the known human infections are caused by the helminth/protozoan group.<sup>1</sup> The burden of disease caused by infection with soil-transmitted helminths (STH) remains enormous. Helminth/protozoan infections represented greater than 40% of the burden caused by all tropical diseases.<sup>2</sup> Intestinal parasitic infections are associated with a disability-adjusted life year (DALY) loss of 5 266 000 globally.<sup>3</sup>

Intestinal parasitic infections are common in the world. However, the problem is higher in developing countries. In 2010, at least 1.3 billion people were estimated to be infected with STH.<sup>4</sup> A global-level estimate indicated that greater than 0.8 billion people had *ascariasis* in 2010,<sup>5</sup> around 0.45 billion people had hookworm,<sup>5</sup> and at least 0.23 billion people are

estimated to have *schistosomiasis*,<sup>6–8</sup> in which the most of the cases are children. In 2010, an estimated 9 million life years lost due to the major worm infections of children.<sup>9,10</sup>

The burden of intestinal parasitic infections is higher in Sub-Saharan Africa (SSA).<sup>1,5,7,8</sup> More than 90% of *schistosomiasis* cases occurred in SSA, with the largest number in Nigeria, Ethiopia, and Democratic Republic of Congo.<sup>11,12</sup> In Ethiopia, a 2005 estimate showed that 4882 children were infected with Hookworm, 1956 with *ascariasis*, 1983 with *trichiuriasis*, and 7357 with other STHs.<sup>12</sup>

The parasitic relation of worms with human has been influenced by global changes in the human sociocultural spectrum.<sup>13,14</sup> Socioeconomic factors, like income or poverty,<sup>8,15–18</sup> occupation especially farming and fishing,<sup>8,18–20</sup> number of siblings,<sup>16,20</sup> age of children,<sup>19,21,22</sup> family size,<sup>19,20</sup> households educational status,<sup>19,23</sup> and health supervision or provision of health



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without

education<sup>24,25</sup> are contributing for persistent transmission of the parasitic disease.

Although health information related to parasitic infections is available globally, health information often limited in rural setups in least developed countries including Ethiopia. This study was, therefore, done to investigate socioeconomic predictors of intestinal parasitic infections in under-five children in rural Dembiya, Northwest Ethiopia.

## Materials and Methods

### *Study design and description of study settings*

A cross-sectional survey, which is community-based, was conducted in rural Dembiya during May 2017. The study setting is described elsewhere.<sup>26</sup>

### *Sample size determination and sampling procedures*

This study is part of the baseline survey for Dembiya Water, sanitation and hygiene—neglected tropical diseases (Dembiya WASH-NTDs) project. The project was implemented to prevent intestinal parasitic infections through improved WASH. In this project, single population proportion formula was used to calculate sample size. The assumptions we used to calculate sample size were presented elsewhere.<sup>26</sup> A total of 225 children aged 6 to 59 months were selected from 5 rural kebeles (the lowest administrative units in Ethiopia). The study subjects were selected by systematic random sampling technique.

### *Data collection tools*

We used pretested and structured questionnaire to collect socio-demographic information. Direct stool examination technique was used to identify parasitic infections in children. We used standardized procedures as presented in World Health Organization's training manual on diagnosis on intestinal parasitic infections.<sup>27</sup>

Wealth index of households, one of the socioeconomic predictors of intestinal parasitic infections, was determined by principal component analysis (PCA). Asset information was gathered based on the list of assets in health and demographic surveys and other related studies to determine wealth index.<sup>28-30</sup> Variables were selected based on eigenvalues greater than 1, and variables whose components greater than 0.4 in the component matrix were considered to compute wealth index. Finally, wealth index of households was classified into poor and rich.

### *Data analysis*

Frequencies, percentages, mean or/and median, standard deviation (SD) or/and interquartile range (IQR) were used to present data. Socioeconomic predictors were selected by univariable binary logistic regression analysis on the basis of  $P < .2$  and

then analyzed by multivariable binary logistic regression for controlling the possible effect of confounders, and finally, the variables which had significant association were identified on the basis of adjusted odds ratio (AOR) with 95% confidence interval (CI) and  $P < .05$ . Hosmer and Lemeshow test was used to test the model goodness of fitness.

## Results

### *Socioeconomic information*

Two hundred twenty-five children were participated in this study. However, the data set for 1 child is incomplete for some variables and so that we considered 224 children in the analysis. Out of 224 children, 118 (52.7%) of them were women, and 166 (74.1%) of the study subjects were 24 to 59 months old. The median age was 42 months, and the IQR was 24 to 48 months. More than half 134 (59.8%) of the mothers were  $\leq 30$  years old. The minimum age of mothers participated in this study was 18 years and the maximum was 47 years. The median age was 30 years and the IQR was 25 to 35 years. One hundred seventy-nine (79.9%) of mothers did not attend formal education and 50 (22.3%) of the households had at least 1 member whose education level is secondary and above. Two hundred thirteen (95.1%) of the mothers were married at the time of the survey and 221 (98.7%) of the mothers were farmer by their occupation. One hundred twenty-six (56.3%) of the households had more than 5 family members, and 139 (57.6%) households were economically poor (Table 1).

### *Health information of rural households*

One hundred forty-one (62.9%) of the households reported that they were frequently supervised by health professionals. One hundred sixteen (51.8%) households reported as they received health messages 1 week before the time of the survey and the commonest source of information was government health workers, which accounted 112 (96.6%). One hundred thirty-eight (61.6%) households reported as they exchanged health information within the family weekly at regular basis (Table 2).

### *Prevalence of intestinal parasitic infections*

This study reported that 57 of 224 (25.4%; 95% CI = [20.2, 31.1]) of under-five children had intestinal parasitic infection. *Ascaris lumbricoides* (44 of 224), hookworm (6 of 224), *Hymenolepis nana* (3 of 224), *Enterobius vermicularis* (2 of 224), *Schistosoma mansoni* (1 of 224), and *Giardia lamblia* (1 of 224) were identified.

### *Socioeconomic factors associated with parasitic infections*

Childhood intestinal parasitic infections were statistically associated with educational status of family members, number of

**Table 1.** Sociodemographic information of households with children aged 6 to 59 months (225) in rural Dembiya, Northwest Ethiopia, May 2017.

VARIABLES	FREQUENCY	PERCENT
Sex of children		
Male	106	47.3
Female	118	52.7
Age of children		
Under 2 years	58	25.9
2 and above years	166	74.1
Mothers age		
≤30 years	134	59.8
>30 years	90	40.2
Maternal education		
No formal education	179	79.9
Have formal education	45	20.1
The household has at least 1 member whose education is secondary and above		
Yes	50	22.3
No	174	77.7
Mother's marital status		
Married	213	95.1
Not married	11	4.9
Maternal occupation		
Farmer	221	98.7
Merchant	3	1.3
Family size		
≤5	126	56.3
>5	98	43.8
Wealth index		
Poor	129	57.6
Rich	95	42.4

under-five children in a household, health supervision, family size, and wealth index. Age of children and age of mothers did not show statistically significant association with parasitic infections (Table 3). Occupational and educational status of mothers did not pass the chi-square assumption.

This study revealed that childhood parasitic infections were associated with educational status of family members. The odds of childhood parasitic infections were 3.36 times more likely to be higher among households who had no members

**Table 2.** Health information of households with children aged 6 to 59 months (n=225) in rural Dembiya, Northwest Ethiopia, May 2017.

VARIABLES	FREQUENCY	PERCENT
Health professional frequently visit the household		
Yes	141	62.9
No	83	37.1
The household received health messages last week prior to the survey		
Yes	116	51.8
No	108	48.2
Source of health messages		
Government health workers	112	96.6
School children	26	22.4
Church leaders	5	4.3
Radio	3	2.6
Community discussion	2	1.7
Households exchange health information at regular basis		
Yes	138	61.6
No	86	38.4

whose education status is secondary and above (AOR=3.36, 95% CI=[1.23, 9.17]). The presence of 2 under-five children in a household was statistically associated with intestinal parasitic infections. Childhood parasitic infections were more prevalent among households having 2 under-five children compared with their counterparts (AOR=3.56, 95% CI=[1.29, 9.82]). Intestinal parasitic infections were statistically associated with absence of health supervision. The prevalence of childhood parasitic infections was 3.49 times to be higher among households who had not been frequently supervised by health professionals (AOR=3.49, 95% CI=[1.72, 7.09]). Children who live in households whose family size is greater than 5 had more odds to have parasitic infections (AOR=2.30, 95% CI=[1.09, 4.85]). Households' economic status was also identified as a contributing factor for the occurrence of parasitic infections in children. Children from the poor families had 2.58 more chance to have parasitic infections (AOR=2.58, 95% CI=[1.23, 5.41]).

## Discussion

This study reported that 57 of 224 (25.4%; 95% CI=[20.2, 31.1]) children had intestinal parasitic infection, which was lower than the findings of studies in Wondo Genet (85.1%),<sup>31</sup> Hawassa Zuria District (51.3%),<sup>32</sup> Southern Ethiopia (41.9%),<sup>33</sup> and Chuahit (35.2%).<sup>34</sup> The prevalence reported by this study was similar to the findings in Wonji Shoa Sugar Estate (24.3%)<sup>35</sup> and Butajira town (23.3%)<sup>36</sup>; Sudan (24.9%)<sup>37</sup>;

**Table 3.** Sociodemographic factors affecting parasitic infection among children aged 6 to 59 months (n=225) in rural Dembiya, Northwest Ethiopia, May 2017.

VARIABLES	PARASITIC INFECTION		COR WITH 95% CI	AOR WITH 95% CI
	YES	NO		
The family has 1 or more members whose education level is secondary and above				
Yes	6	44	1	
No	51	123	3.04 [1.22, 7.58]	3.36 [1.23, 9.17]*
Age of children				
Under 2 years	20	38	1	
2 years and above	37	129	0.55 [0.28, 1.05]	0.62 [0.30, 1.32]
Mothers age				
≤35 years	27	107	1	
>35 years	30	60	1.98 [1.09, 3.64]	1.55 [0.73, 3.28]
Number of under-five children				
1	47	154	1	
2	10	13	2.52 [1.04, 6.12]	3.56 [1.29, 9.82]*
Health professional frequently visit households				
Yes	24	117	1	
No	33	50	3.23 [1.73, 5.99]	3.49 [1.72, 7.09]**
Family size				
≤5	22	104	1	
>5	35	63	2.63 [1.42, 4.87]	2.30 [1.09, 4.85]*
Wealth index				
Poor	43	86	2.89 [1.47, 5.68]	2.58 [1.23, 5.41]*
Rich	14	81	1	

Abbreviations: COR, crude odds ratio; CI, confidence interval; AOR, adjusted odds ratio.

Hosmer and Lemeshow test=0.376.

\*Statistically significant at  $P < .05$ .

\*\*Statistically significant at  $P < .001$ .

Bogota (26.4%)<sup>20</sup>; and Diamantina, Brazil (27.5%).<sup>18</sup> The current prevalence is also higher than the finding of a study in Nigeria (13.7%).<sup>38</sup> The prevalence of intestinal parasitic infections in children was higher in rural Dembiya. This may be due to the fact that the population in the area had poor access to sanitation. During June 2017, clean water and latrine coverage was 26.6% and 55%, respectively.<sup>10</sup> Moreover, as depicted by this study, significant proportion of the households lack WASH information. In this study, 37.1% of households reported that they were not frequently supervised by health professionals.

This study showed that educational status of family members was associated with intestinal parasitic infections in children. Childhood parasitic infections were higher among households who had no members whose education status is

secondary and above. This may be due to the fact that educated households may have awareness about the transmission and prevention methods of infectious diseases. Education encourages changes in healthy behaviors at the household level. Other similar studies also reported the relation of education with occurrence of parasitic infections.<sup>19,23,39,40</sup>

Childhood intestinal parasitic infections were associated family size and number of under-five children in a household. Childhood intestinal parasitic infections were more prevalent among households having higher family size and 2 under-five children. Other studies also identified the relationship between number of under-five children and parasitic infections.<sup>16,19,20</sup> This can be justified that children in larger families may be exposed to infections because the quality of care and attention

from parents decreases as mothers may become unable to care children and less effort is available for each individual child. Moreover, high family size affects mothers or caregivers health seeking and hygienic behaviors.<sup>40-43</sup>

This study depicted that intestinal parasitic infections were associated with health supervision or health education. The prevalence of intestinal parasitic infections was higher among households who had not been frequently supervised by health professionals. Other studies also reported the association of health supervision or health education and intestinal parasitic infections.<sup>24,25,44,45</sup> This fact can be justified that health education promotes health behaviors toward hygiene and sanitation practices. Health education increases knowledge and acceptability of interventions within the community. It also sustains integrated control of the infection.<sup>45-50</sup>

Childhood parasitic infections were significantly associated with households' economic status. Children from the poor families had more chance to have parasitic infections. The finding of this study is in line with findings of other similar studies.<sup>8,15,16,19,39</sup> This may be due to the fact that rich families may have greater opportunity to healthy measures like soap, household water treatment, toilets and other facilities, and lower income families could not afford these facilities.<sup>17,40,51</sup>

As a limitation, in this research, we did not use floatation techniques/McMaster technique to detect hookworm because the McMaster chamber was not available in the country. We used standardized wet mount preparation. We examined each specimen within 1 hour of sampling time to effectively detect hookworm. Moreover, the 95% CI for some predictor variables is wide due to small sample size.

## Conclusions

Significant proportion of children was infected with intestinal parasitic infection in rural Dembiya. Educational status of family members, number of under-five children in a household, health supervision, family size, and wealth index were statistically associated with parasitic infections. Provision of anthelmintic drugs, health supervision, and health education targeted with transmission and prevention of infections are recommended.

## Acknowledgements

The authors acknowledged NALA foundation, data collectors, field supervisors, and study participants for their contributions to this study.

## Author Contributions

All the authors actively participated during development of a research proposal, data collection, analysis and interpretation, and writing various parts of the research report. ZG prepared the manuscript. All of the authors read and approved the final manuscript.

## Availability of Data and Material

Data will be made available on requesting the primary author.

## Ethics Approval and Consent to Participate

The Institutional Review Board of the University of Gondar approved the ethical aspects of this study. There were no risks due to participation in this research. Confidentiality and privacy were maintained. Verbal informed consent was obtained from the mothers and participation was on voluntary basis. Appropriate anthelmintic drugs were given for infected children. Moreover, the researchers provided health education for mothers or caregivers.

## Consent for Publication

This manuscript does not contain any individual person's data.

## ORCID iD

Zemichael Gizaw  <https://orcid.org/0000-0002-6713-1975>

## REFERENCES

- Cleaveland S, Laurenson M, Taylor L. Diseases of humans and their domestic mammals: pathogen characteristics, host range and the risk of emergence. *Philos Trans R Soc Lond B Biol Sci.* 2001;356:991-999.
- World Health Organization (WHO). Neglected tropical diseases: PCT data-bank: schistosomiasis. [http://www.who.int/neglected\\_diseases/preventive\\_chemotherapy/sch/en/](http://www.who.int/neglected_diseases/preventive_chemotherapy/sch/en/). Updated 2015. Accessed December 20, 2017.
- World Health Organization (WHO). *Global Health Estimates (GHE)*. Geneva: WHO; 2016. [http://www.who.int/healthinfo/global\\_burden\\_disease/en/](http://www.who.int/healthinfo/global_burden_disease/en/). Accessed December 28, 2017.
- Global Atlas of Helminth Infection (GAHI). Soil-transmitted helminths. <http://www.thiswormyworld.org/maps/soil-transmitted-helminths>. Accessed December 21, 2017.
- Pullan RL, Smith JL, Jasrasaria R, Brooker SJ. Global numbers of infection and disease burden of soil transmitted helminth infections in 2010. *Parasit Vectors.* 2014;7:37.
- Colley DG, Bustinduy AL, Secor WE, King CH. Human schistosomiasis. *Lancet.* 2014;383(9936):2253-2264.
- Knowles SC, Webster BL, Garba A, et al. Epidemiological interactions between urogenital and intestinal human schistosomiasis in the context of praziquantel treatment across three West African countries. *PLoS Negl Trop Dis.* 2015;9:e0004019.
- Adenowo AF, Oyinloye BE, Ogunyinka BI, Kappo AP. Impact of human schistosomiasis in Sub-Saharan Africa. *Braz J Infect Dis.* 2015;19:196-205.
- Murray CJ, Vos T, Lozano R, et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet.* 2013;380:2197-2223.
- Hotez PJ, Alvarado M, Basanez MG, et al. The global burden of disease study 2010: interpretation and implications for the neglected tropical diseases. *PLoS Negl Trop Dis.* 2014;8:e2865.
- World Health Organization (WHO). Schistosomiasis: number of people receiving preventive chemotherapy in 2012= Schistosomiase: nombre de personnes ayant bénéficié d'une chimioprévention en 2012. *Wkly Epidemiol Rec.* 2014;89:21-28.
- Brooker S, Clements AC, Bundy DA. Global epidemiology, ecology and control of soil-transmitted helminth infections. *Adv Parasitol.* 2006;62:221-261.
- Alum A, Rubino JR, Ijaz MK. The global war against intestinal parasites—should we use a holistic approach. *Int J Infect Dis.* 2010;14:e732-e738.
- WHO/WER. Schistosomiasis and soil-transmitted helminthes infections: preliminary estimates of the number of children treated with albendazole or mebendazole. *Wkly Epidemiol Rec.* 2006;81:145-164. [www.who.int/wer/2006/wer8116.pdf](http://www.who.int/wer/2006/wer8116.pdf). Accessed December 28, 2017.
- Ngui R, Ishak S, Chuen CS, Mahmud R, Lim YA. Prevalence and risk factors of intestinal parasitism in rural and remote West Malaysia. *PLoS Negl Trop Dis.* 2011;5:e974.
- El-Masry H, Ahmed Y, Hassan A, et al. Prevalence, risk factors and impacts of schistosomal and intestinal parasitic infections among rural school children in Sohag Governorate. *Egypt J Hosp Med.* 2007;29:616-630.
- De Silva NR, Brooker S, Hotez PJ, Montresor A, Engels D, Savioli L. Soil-transmitted helminth infections: updating the global picture. *Trends Parasitol.* 2003;19:547-551.

18. Nobre LN, Silva RV, Macedo MS, Teixeira RA, Lamounier JA, Franceschini SC. Risk factors for intestinal parasitic infections in preschoolers in a low socio-economic area, Diamantina, Brazil. *Pathog Glob Health*. 2013;107:103-106.
19. Al-Mohammed HI, Amin TT, Aboulmagd E, Hablus HR, Zaza BO. Prevalence of intestinal parasitic infections and its relationship with socio-demographics and hygienic habits among male primary schoolchildren in Al-Ahsa, Saudi Arabia. *Asian Pac J Trop Med*. 2010;3:906-912.
20. Bouwmans MC, Gaona MA, Chenault MN, Zuluaga C, Pinzón-Rondon ÁM. Prevalence of intestinal parasitic infections in preschool-children from vulnerable neighborhoods in Bogotá. *Rev Univ Ind Santander Salud*. 2016;48:178-187.
21. Okpala H, Josiah S, Oranekwulu M, Ovie E. Prevalence of intestinal parasites among children in day care centres in Esan West Local Government Area, Edo State, Nigeria. *Asian J. Med. Sci*. 2014;6:34-39.
22. Jacobsen KH, Ribeiro PS, Quist BK, Rydbeck BV. Prevalence of intestinal parasites in young Quichua children in the highlands of rural Ecuador. *J Health Popul Nutr*. 2007;25:399-405.
23. Heidari A, Rokni M. Prevalence of intestinal parasites among children in day-care centers in Damghan-Iran. *Iranian J Publ Health*. 2003;32:31-34.
24. Fouamno Kamga HL, Shey Nsagha D, Suh Atanga MB, et al. The impact of health education on the prevalence of faecal-orally transmitted parasitic infections among school children in a rural community in Cameroon. *Pan Afr Med J*. 2011;8:38.
25. Kanoa B, George E, Abed Y, Al-Hindi A. Evaluation of the relationship between intestinal parasitic infection and health education among school children in Gaza city, Beit-lahia village and Jabalia refugee camp, Gaza strip, Palestine. *Islamic Univ J*. 2006;14:39-49.
26. Gizaw Z, Adane T, Azanaw J, Addisu A, Haile D. Childhood intestinal parasitic infection and sanitation predictors in rural Dembiya, Northwest Ethiopia. *Environ Health Prev Med*. 2018;23:26.
27. World Health Organization (WHO). *Training manual on diagnosis of intestinal parasites based on the WHO bench aids for the diagnosis of intestinal parasites, district laboratory practice in tropical countries* (WHO/CTD/SIP/98.2 CD-Rom). [http://usaf.phsource.us/PH/PDF/HELM/trainingmanual\\_sip98-2.pdf](http://usaf.phsource.us/PH/PDF/HELM/trainingmanual_sip98-2.pdf). Updated 2004. Accessed November 1, 2017.
28. Rutstein SO. *The DHS Wealth Index: Approaches for rural and urban areas*. Calverton, MD: Macro International. <https://dhsprogram.com/pubs/pdf/WP60/WP60.pdf>. Accessed October 23, 2017.
29. Córdova A. Methodological note: measuring relative wealth using household asset indicators (Series No. 06). *AmericasBarometer Insights*. [https://www.vanderbilt.edu/lapop/insights/I0806en\\_v2.pdf](https://www.vanderbilt.edu/lapop/insights/I0806en_v2.pdf). Updated 2009. Accessed October 21, 2017.
30. Kolenikov S, Angeles G. Socioeconomic status measurement with discrete proxy variables: is principal component analysis a reliable answer? *Rev Income Wealth*. 2009;55:128-165.
31. Nyantekyi LA, Legesse M, Belay M, et al. Intestinal parasitic infections among under-five children and maternal awareness about the infections in Shesha Kekele, Wondo Genet, Southern Ethiopia. *Ethiop J Health Dev*. 2010;24:185-190.
32. Kabeta A, Assefa S, Hailu D, Berhanu G. Intestinal parasitic infections and nutritional status of pre-school children in Hawassa Zuria District, South Ethiopia. *Afr J Microbiol Res*. 2017;11:1243-1251.
33. Unasho A. An investigation of intestinal parasitic infections among the asymptomatic children in, Southern Ethiopia. *Int J Child Health Nutr*. 2013;2:212-222.
34. Alemu A, Tegegne Y, Damte D, Melku M. Schistosoma mansoni and soil-transmitted helminths among preschool-aged children in Chuahit, Dembia district, Northwest Ethiopia: prevalence, intensity of infection and associated risk factors. *BMC Public Health*. 2016;16:422.
35. G/hiwot Y, Degarege A, Erko B. Prevalence of intestinal parasitic infections among children under five years of age with emphasis on Schistosoma mansoni in Wonji Shoa Sugar Estate, Ethiopia. *PLoS ONE*. 2014;9:e109793.
36. Shumbej T, Belay T, Mekonnen Z, Tefera T, Zemene E. Soil-transmitted helminths and associated factors among pre-school children in Butajira Town, South-Central Ethiopia: a community-based cross-sectional study. *PLoS ONE*. 2015;10:e0136342.
37. Sun C. Prevalence and associated risk factors of Intestinal Helminths infections among pre-school children (1 to 5 years old) in IDPs settlements of Khartoum state, Sudan. *J Global Health*. <https://www.gjournal.org/prevalence-and-associated-risk-factors-of-intestinal-helminths-infections-among-pre-school-children-1-to-5-years-old-in-idps-settlements-of-khartoum-state-sudan/>. Accessed November 13, 2017.
38. Achi E, Njoku O, Nnachi A, et al. Prevalence of intestinal parasitic infections among under five children in Abakaliki local government area of Ebonyi state. *Ejpmr*. 2017;4:218-222.
39. Quihui L, Valencia ME, Crompton DW, et al. Role of the employment status and education of mothers in the prevalence of intestinal parasitic infections in Mexican rural schoolchildren. *BMC Public Health*. 2006;6:225.
40. Wolde W, Bitew BD, Gizaw Z. Socioeconomic factors associated with diarrheal diseases among under-five children of the nomadic population in Northeast Ethiopia. *Trop Med Health*. 2016;44:40.
41. Lovelyn OA, Bertrand NO, Godswill N. Family and social determinants of health-seeking behaviour of caregivers of febrile children in an urban city of South-Eastern Nigeria. *Arch Med*. 2016;8:1-6.
42. Uggl C, Mace R. Parental investment in child health in sub-Saharan Africa: a cross-national study of health-seeking behaviour. *R Soc Open Sci*. 2016;3:150460.
43. Sisay S, Endalew G, Hadgu G. Assessment of mothers/care givers health care seeking behavior for childhood illness in rural Ensaro District, North Shoa Zone, Amhara Region, Ethiopia. *Global J Life Sci Biol Res*. 2015;1:20-34.
44. Anantaphruti M, Waikagul J, Maipanich W, et al. School-based health education for the control of soil-transmitted helminthiasis in Kanchanaburi province, Thailand. *Ann Trop Med Parasitol*. 2008;102:521-528.
45. Gyorkos TW, Maheu-Giroux M, Blouin B, Casapia M. Impact of health education on soil-transmitted helminth infections in schoolchildren of the Peruvian Amazon: a cluster-randomized controlled trial. *PLoS Negl Trop Dis*. 2013;7:e2397.
46. Albright JW, Basaric-Keys J. Instruction in behavior modification can significantly alter soil-transmitted helminth (STH) re-infection following therapeutic de-worming. *Southeast Asian J Trop Med Public Health*. 2006;37:48-57.
47. Asaolu S, Ofoezie I. The role of health education and sanitation in the control of helminth infections. *Acta Trop*. 2003;86:283-294.
48. Lansdown R, Ledward A, Hall A, et al. Schistosomiasis, helminth infection and health education in Tanzania: achieving behaviour change in primary schools. *Health Educ Res*. 2002;17:425-433.
49. Albonico M, Montresor A, Crompton D, Savioli L. Intervention for the control of soil-transmitted helminthiasis in the community. *Adv Parasitol*. 2006;61:311-348.
50. Bieri FA, Gray DJ, Williams GM, et al. Health-education package to prevent worm infections in Chinese schoolchildren. *N Engl J Med*. 2013;368:1603-1612.
51. Rahman A. Assessing income-wise household environmental conditions and disease profile in urban areas: study of an Indian city. *Geojournal*. 2006;65:211-227.