

Human Health Risk Assessment of Trace Metals in the Commonly Consumed Fish Species in Nakuru Town, Kenya

Authors: Esilaba, Felly, Moturi, Wilkister Nyaora, Mokua, Millicent, and Mwanyika, Terewe

Source: Environmental Health Insights, 14(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/1178630220917128>


BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Human Health Risk Assessment of Trace Metals in the Commonly Consumed Fish Species in Nakuru Town, Kenya

Felly Esilaba¹ , Wilkister Nyaora Moturi¹, Millicent Mokua¹ and Terewe Mwanyika²

¹Department of Environmental Science, Egerton University, Egerton, Kenya. ²Department of Chemistry, Egerton University, Egerton, Kenya.

Environmental Health Insights
Volume 14: 1–8
© The Author(s) 2020
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/1178630220917128



ABSTRACT: The present study was conducted to determine daily intake of cadmium (Cd), copper (Cu), and lead (Pb) and to assess non-carcinogenic human health risk caused by these trace metals in the commonly consumed fish species (*Oreochromis niloticus*, *Rastrineobola argentea*, *Lates niloticus*, and *Protopterus aethiopicus*) in Nakuru town, Kenya. Trace metal determination in the composite samples of the commonly consumed fish species was done using flame atomic absorption spectrophotometer. Cd, Cu, and Pb content in the muscle tissues of the commonly consumed fish species ranged from 0.11 ± 0.045 to 1.11 ± 0.931 mg kg⁻¹ for Cd, 0.48 ± 0.013 to 3.00 ± 0.009 mg kg⁻¹ for Cu, and 3.42 ± 0.045 to 12.78 ± 0.108 mg kg⁻¹ for Pb. Cu concentrations were within Food and Agriculture Organization (FAO) recommended limits for this trace metal in fish. In contrast, Cd and Pb had values above their respective permissible limits in fish. The assessment of human exposure to trace metals indicated that exposure doses of Cd and Cu were safe for fish consumers. Conversely, target hazard quotient (THQ) values of Pb suggested possible health risks for consumers of the commonly consumed fish species in Nakuru town, Kenya.

KEYWORDS: Trace metals, commonly consumed fish species, human health risk assessment

RECEIVED: July 29, 2019. **ACCEPTED:** March 15, 2020.

TYPE: Original Research

FUNDING: The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: *This work was supported by the National Commission for Science, Technology and Innovation (grant number NACOSTI/RCD/ST/I/7TH CALL/MSc/270).*

DECLARATION OF CONFLICTING INTEREST: The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

CORRESPONDING AUTHOR: Felly Esilaba, Department of Environmental Science, Egerton University, P.O. Box 536-20115, Egerton, Kenya. Email: fellyesilaba@gmail.com

Introduction

Fish is an important component of many meals around the world because it has high contents of proteins, essential fats, vitamins, and minerals.^{1–3} Its growing demand is evident from the vigorous growth in aquaculture and in the fact that the world's per capita fish consumption has risen from 9.9 kg in the 1960s to 14.4 kg in the 1990s to 19.7 kg in 2013.⁴ Nonetheless, increased aquatic pollution caused by industrial discharge, domestic waste, agricultural runoff, mine drainage, and accidental oil spills indicate that fish could be a serious source of human exposure to potentially hazardous substances like trace metals.^{2,5,6} Trace metals are persistent and nonbiodegradable, and once they enter aquatic ecosystems, they bioaccumulate in aquatic organisms and their concentrations are biomagnified as they move up the aquatic food chain.^{1,5,7} Trace metals in fish may be transferred and pose serious health problems to fish consumers. Cadmium, for instance, may cause cardiovascular, neurological, and reproductive disorders.^{7,8} Short-term exposure to high doses of Cd can cause nausea, vomiting, abdominal disturbances, and fatigue.⁸ In addition, it has been classified as a probable human carcinogen by the International Agency for Research on Cancer.^{9–11} Exposure to Pb has been associated with renal, muscular, and cardiovascular problems. It may also cause cognitive and development deficits in children and reproductive disorders in men and women.^{7,11–15} Unlike Cd and Pb, Cu is an essential element required for various physiological functions in the human body.¹⁶ Even so, Cu levels above the

recommended limit may lead to brain, liver, and kidney disorders.¹⁷ Short-term exposure to high doses of Cu can cause diarrhea, stomach pains, vomiting, and death due to liver and kidney failure or due to the depression of the central nervous system.¹⁷

Because of the aforementioned and many other health effects associated with dietary trace metals, there is growing interest in food safety with regard to the accumulation of these substances in food. Studies around the world show that fish and other seafood are a major source of dietary trace metals.^{18–23} Majority of these studies reveal that human exposure to trace metals from fish and seafood is within the recommended limits. Conversely, other studies indicate that consumers of fish and seafood may be exposed to hazardous levels of these toxicants and adverse health incidents may occur as a result of this exposure.^{22,24–28} Vegetables and cereals are the other major dietary sources of trace metals. Vegetables and cereals are valued components of many meals around the world. Thus, the presence of trace metals in these food groups is raising concern world over as is made evident by these studies.^{29–34}

In Kenya, various studies have assessed trace metals content in fish,^{35–39} beef,⁴⁰ honey,^{41,42} and vegetables^{43–45} to determine whether they exceeded internationally set standards for these substances in food. Nonetheless, exceedance does not always represent human health risk.⁴⁶ Thus, in the present study estimated daily intake (EDI) of Cd, Cu, and Pb via consumption



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without

further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).
Downloaded From: <https://complete.bioone.org/journals/Environmental-Health-Insights> on 18 Apr 2024
Terms of Use: <https://complete.bioone.org/terms-of-use>

of the commonly consumed fish species in Nakuru town, Kenya, were calculated and compared with the provisional tolerable daily intake (PTDI) established by the Joint FAO/WHO Expert Committee on Food Additives (JECFA).^{47–49} In addition, target hazard quotient (THQ) and hazard index (HI) were computed to assess human health risks posed by Cd, Cu, and Pb in the commonly consumed fish species in Nakuru town, Kenya.

Materials and Methods

Questionnaire survey

A questionnaire-based survey was conducted to establish the commonly consumed fish species in Nakuru town and the consumption rates of these fish species. Questionnaires were administered to 385 randomly selected participants buying fish in 5 retail outlets in Nakuru town. The questionnaire acquired basic information including gender, age, education, and income level of the respondents. It was used to acquire information on fish preferences of consumers. It was also used to determine the frequency and ingestion rates of the commonly consumed fish species in Nakuru town.

The cross-sectional survey established that *Oreochromis niloticus* (Nile tilapia), *Rastrineobola argentea* (Silver cyprinid), *Lates niloticus* (Nile perch), and *Protopterus aethiopicus* (African marbled lungfish) were the commonly consumed fish species in Nakuru town, Kenya. Moreover, the findings revealed that in a day consumers of these fish varieties consumed on average 0.027, 0.012, 0.035, and 0.023 kg of *O niloticus* (Nile tilapia), *R argentea* (Silver cyprinid), *L niloticus* (Nile perch), and *P aethiopicus* (African marbled lungfish), respectively.

Sample collection and preparation

Four commonly consumed fish varieties (*O niloticus*, *R argentea*, *L niloticus* and *P aethiopicus*) in Nakuru town were selected based on the data obtained by the cross-sectional survey. Ten pieces of *O niloticus*, 10 pieces of *L niloticus*, and 10 pieces of *P aethiopicus*, and 10 samples of *R argentea* each weighing 100 g were purchased from the first market (market 1). Ten pieces of *O niloticus* and 10 samples of *R argentea* each weighing 100 g were purchased from the second market (market 2). Ten pieces of *O niloticus* and 10 samples of *R argentea* each weighing 100 g were purchased from the third market (market 3). Ten pieces of *O niloticus* and 10 pieces of *L niloticus* were purchased from the fourth market (market 4). Ten samples of *R argentea* each weighing 100 g were purchased from the fifth market (market 5). In total, 70 fresh samples of *O niloticus*, *L niloticus*, and *P aethiopicus* and 40 sun-dried samples of *R argentea* each weighing 100 g were purchased in August 2016 from the 5 major fish retail outlets in Nakuru town, Kenya (−0.2833°, 36.06667°). The samples were put in sterile polyethylene bags and transported to the laboratory using a cool box at <5°C. After reaching the laboratory, the fish were washed using double distilled

water. The samples were carefully dissected to remove muscle tissues from the dorsal part of the fish which were washed using doubled distilled water. The muscle tissues of the same fish species from the same retail market were combined into composite samples of 10 individuals. For the small species (*R argentea*), 10 samples of this fish species each weighing 100 g from the same retail market were pooled into composite samples. Because of their small size, the entire individual fish were included in the preparation of the composite sample. The composite samples of the commonly consumed fish species were then kept frozen at −20°C until trace metal analysis.

Trace metals analysis

The composite fish samples were thawed at room temperature, oven dried for 72 hours at 105°C until constant dry weight. They were pulverized in a dry pestle-mortar then stored in sterile containers placed in desiccators awaiting trace metal analysis.

The powdered composite samples were thoroughly mixed to attain optimum homogeneity. They were then digested using Food and Agriculture Organization (FAO) procedures.^{50,51} Briefly, 1 g of homogenized composite sample was weighed into a digestion vessel. Ten milliliters digestion mixture consisting of 65% nitric acid (HNO₃) and 30% peroxide hydrogen (H₂O₂) was prepared and added to the digestion vessels, kept overnight to allow various reactions to occur. The digestion vessels were heated on a hot plate to 130°C until the solutions turned clear and the volume reduced to between 2 and 3 mL. They were allowed to cool, filtered then diluted with double distilled water to 50 mL in volumetric flasks.

Analyses of Cd, Cu, and Pb were performed using flame atomic absorption spectrophotometer (Thermo Jarrell Ash). The detection limits of Cd, Cu, and Pb were 0.0015, 0.003, and 0.01 mg/L, respectively. To ensure reliability of the results, double distilled water was used throughout the study. Glass vessels were precleaned thoroughly using detergent, soaked overnight in 20% nitric acid (HNO₃) then rinsed using double distilled water. The experimental reagents and standards used during the study were of analytical grade obtained from Sigma Aldrich. The accuracy of the instrument was ensured by running samples in triplicates. In addition, a reagent blank and a standard were run after every 3 samples to check instrumental drift. The AAS instrument was calibrated with a series of standard solutions prepared in the concentration range of 0 to 15 mg L^{−1}. A calibration curve of absorbance versus concentration was obtained and used to quantify Cd, Cu, and Pb content in the commonly consumed fish species. Trace metal concentration was expressed in milligram per kilogram dry weight.

Human health risk assessment

The risk to human health as a result of consuming the commonly consumed fish varieties sold in Nakuru town, Kenya,




Figure 1. Geographic locations of the main fish retail markets in Nakuru town (Kenya).

was evaluated by calculating EDI and THQ of trace metals. In addition, literature indicates that exposure to 2 or more trace metals from the same source may result to additive effects.⁵² Thus, THQs of Cd, Cu, and Pb were summed to generate hazard indices (HI) that represent overall health risk posed by the 3 trace metals in the commonly consumed fish species.

EDI was calculated using equation (1).^{22,53}

$$EDI = \frac{EF \times ED \times FIR \times C}{BW \times TA} \quad (1)$$

where EF is exposure frequency (365 days year⁻¹); ED is the frequency duration (70 years, equivalent to the average human lifespan); FIR is the ingestion rate of fish in kilograms person⁻¹ day⁻¹; C is the concentration of Cd, Cu, or Pb in the commonly consumed fish species in mg kg⁻¹; BW is the average body weight which is equivalent to 60.7 kg for an average African adult³⁷; and TA is the average exposure time for noncarcinogens (365 days year⁻¹ × ED). Computed EDI values were compared with PTDI values of Cd (0.001 mg kg⁻¹ day⁻¹), Cu (0.5 mg kg⁻¹ day⁻¹), and Pb (0.00357 mg kg⁻¹ day⁻¹)⁴⁷⁻⁴⁹ to determine whether the daily recommended values were exceeded or not.

Target hazard quotient a ratio between exposure to a potentially hazardous element and its reference dose⁵ was determined

using equation (2).²² Computed THQ values less than 1 indicate that the exposed population is unlikely to experience adverse effects associated with trace metals. On the other hand, computed THQ values greater than 1 suggest possible health risks for the exposed consumers.^{5,54}

$$THQ = \frac{EDI}{RfDo} \quad (2)$$

where RfDo is the oral reference dose of trace metals in mg kg⁻¹ day⁻¹ based on the safe upper level of element's oral intake for an adult. The oral RfD for Cd, Cu, and Pb is 0.001, 0.04, and 0.004 mg kg⁻¹ day⁻¹, respectively.⁵⁵

THQs of Cd, Cu, and Pb were summed to generate a hazard index using equation (3).⁵⁶ Assuming additive effects, HI is a measure of the potential risk of adverse health effects from more than 1 element.^{5,54} HI greater than 1 suggests likelihood of adverse effects on human health and the necessity for further action.^{5,54}

$$HI = \sum_{i=1}^n THQ_i \quad (3)$$

Statistical analysis

SPSS version 22 was used for statistical data analysis. Results are presented as arithmetic mean and standard deviation

Table 1. Trace metal concentration (mg kg⁻¹ dry weight) in *O niloticus*, *R argentea*, *L niloticus*, & *P aethiopicus* sold in 5 retail markets in Nakuru town, Kenya.

SAMPLING SITE	FISH SPECIES	Cd	Cu	Pb
Market 1	<i>O niloticus</i>	0.67 ± 0.035	2.26 ± 0.707	9.99 ± 0.470
Market 1	<i>L niloticus</i>	0.15 ± 0.045	0.65 ± 0.056	7.23 ± 0.149
Market 1	<i>R argentea</i>	0.22 ± 0.050	2.83 ± 0.203	4.44 ± 0.309
Market 1	<i>P aethiopicus</i>	0.75 ± 0.027	0.60 ± 0.026	6.90 ± 0.020
Market 2	<i>O niloticus</i>	0.65 ± 0.017	1.90 ± 0.115	6.30 ± 0.003
Market 2	<i>R argentea</i>	1.00 ± 0.058	2.32 ± 0.175	12.78 ± 0.108
Market 3	<i>O niloticus</i>	0.41 ± 0.009	0.48 ± 0.013	3.42 ± 0.045
Market 3	<i>R argentea</i>	0.11 ± 0.045	3.00 ± 0.009	5.33 ± 0.030
Market 4	<i>O niloticus</i>	0.17 ± 0.020	0.65 ± 0.063	7.40 ± 0.046
Market 4	<i>L niloticus</i>	0.19 ± 0.033	0.74 ± 0.017	4.00 ± 0.051
Market 5	<i>R argentea</i>	1.11 ± 0.931	2.45 ± 0.523	10.00 ± 0.850
Permissible limits of trace metals in fish		0.5	30	0.5

All values are given as means ± SD of 3 replicates of composite fish samples. Permissible limits (mg kg⁻¹) were adopted from FAO.^{19,57}

(mean ± SD). One-way analysis of variance (ANOVA) was used to assess differences among mean concentrations of Cd, Cu, and Pb in fish. It was used to examine variation in trace metal content among the commonly consumed fish species. It was also used to examine trace metal content in fish based on the markets where the fish were purchased. All statistical test were regarded significant when $P < .05$.

Results and Discussion

Trace metal concentrations in the commonly consumed fish species

The mean concentrations of Cd, Cu, and Pb in the commonly consumed fish species, ie, *O niloticus*, *R argentea*, *L niloticus* and *P aethiopicus* purchased from the 5 main retail markets in Nakuru town, Kenya, are presented in Table 1. Metal distribution in the commonly consumed fish species followed the sequence Pb > Cu > Cd, Pb having significantly higher ($P < .05$) concentrations than Cu and Cd. The results indicated that Cd, Cu, and Pb content in fish did not vary significantly ($P > .05$) among the commonly consumed fish varieties (*O niloticus*, *R argentea*, *L niloticus*, and *P aethiopicus*). The results also indicated that trace metal content in the commonly consumed fish species did not vary significantly ($P > .05$) based on the markets where the fish were purchased.

The mean concentrations of Cd in the commonly consumed fish species ranged from 0.11 ± 0.045 to 1.11 ± 0.931 mg kg⁻¹. Similar Cd concentrations in the range of 0.05 ± 0.02 to 0.95 ± 0.09 mg kg⁻¹ were reported by Oyoo-Okoth et al³⁶ in

O niloticus, *R Argentea*, and *L niloticus* from the coastal zone of Lake Victoria, Kenya. Cd concentrations in the present study also compared fairly well with 1.12 ± 1.13 and 1.66 ± 2.48 mg kg⁻¹ in cultured fish from Machakos and Kiambu counties, Kenya,³⁹ and 1.06 ± 0.004 to 1.73 ± 0.002 mg kg⁻¹ in *Cyprinus carpio* from Lake Naivasha, Kenya.³⁸ They were 2-fold higher than those reported by Oyoo-Okoth et al³⁵ in *R argentea* from Lake Victoria. According to FAO,^{19,57} permissible limit of Cd in fish is 0.5 mg kg⁻¹. Some concentrations were above this proposed limit but generally they did not significantly ($P < .05$) surpass it.

Cu content in *O niloticus*, *R argentea*, *L niloticus*, and *P aethiopicus* ranged from 0.48 ± 0.013 to 3.00 ± 0.009 mg kg⁻¹. Cu concentrations in *O niloticus*, *R Argentea*, and *L niloticus* in literature was reported in the range of 2.1 ± 1.1 to 18.8 ± 1.1 mg kg⁻¹ from the coastal zone of Lake Victoria, Kenya,³⁶ and in the range of 5.38 ± 1.27 to 6.20 ± 1.54 mg kg⁻¹ in *R argentea* from Lake Victoria, Kenya.³⁵ Similar Cu concentrations to those in the present study were reported by Mutia et al³⁸ in *C carpio* from Lake Naivasha, Kenya, but Otachi et al³⁷ reported lower Cu concentration in the muscle tissues of *Oreochromis leucostictus* from the same lake. Permissible limit of Cu in fish according to FAO is 30 mg kg⁻¹.^{19,57} The amount of Cu measured in all the samples of the commonly consumed fish species were below this permissible limit suggesting no appreciable health risk to consumers.

Pb concentrations in *O niloticus*, *R argentea*, *L niloticus*, and *P aethiopicus* varied between 3.42 ± 0.045 to 12.78 ± 0.108 mg kg⁻¹. Lower concentrations of Pb were reported in the range of

Table 2. Estimated daily intake of Cd, Cu, and Pb by consuming the commonly consumed fish species in Nakuru town, Kenya.

SAMPLING SITE	FISH SPECIES	FISH INTAKE, KG DAY ⁻¹	Cd INTAKE, MG KG ⁻¹	Cu INTAKE, MG KG ⁻¹	Pb INTAKE, MG KG ⁻¹
Market 1	<i>O niloticus</i>	0.027	0.00030	0.00099	0.00438
Market 1	<i>L niloticus</i>	0.035	0.00008	0.00037	0.00412
Market 1	<i>R argentea</i>	0.012	0.00004	0.00055	0.00086
Market 1	<i>P aethiopicus</i>	0.023	0.00028	0.00023	0.00262
Market 2	<i>O niloticus</i>	0.027	0.00028	0.00083	0.00276
Market 2	<i>R argentea</i>	0.012	0.00019	0.00045	0.00248
Market 3	<i>O niloticus</i>	0.027	0.00018	0.00021	0.00150
Market 3	<i>R argentea</i>	0.012	0.00002	0.00058	0.00104
Market 4	<i>O niloticus</i>	0.027	0.00007	0.00029	0.00325
Market 4	<i>L niloticus</i>	0.035	0.00011	0.00042	0.00228
Market 5	<i>R argentea</i>	0.012	0.00022	0.00048	0.00194
PTDI for an adult weighing 60.7 kg			0.0607	30.35	0.2168

Provisional tolerable daily intake (mg kg⁻¹ day⁻¹) established by the Joint FAO/WHO Expert Committee on Food Additives (JECFA).⁴⁷⁻⁴⁹ 60.7 kg is the average body weight of an African adult.³⁷ Abbreviation: PTDI, provisional tolerable daily intake.

0.18 ± 0.03 to 0.57 ± 0.02 mg kg⁻¹ in *R argentea* from Lake Victoria³⁵ and 0.024 ± 0.03 mg kg⁻¹ in the muscle tissues of *O leuostictus* from Lake Naivasha, Kenya.³⁷ Pb concentrations similar to those observed in the present study were reported in farmed fish from Machakos and Kiambu counties, Kenya.³⁹ FAO has set 0.5 mg kg⁻¹ as the permissible limit of Pb in fish.^{19,57} The lowest measured Pb content in fish in the present study is above this limit. Pb is not required for physiological functions in the human body; thus, short-term exposure to high doses of this trace metal may cause gastrointestinal disorders. Chronic exposure may cause cognitive and learning deficits in children. It may also cause renal, muscular, cardiovascular, and reproductive disorders.^{7,11-15}

Human health risk assessment

To assess human health risk associated with Cd, Cu, and Pb in the commonly consumed fish species, EDI, THQ, and HI were computed. Human health risks were calculated with the assumption that an adult weighing 60.7 kg³⁷ consumes 0.027, 0.012, 0.035, and 0.023 kg day⁻¹ of *O niloticus*, *R argentea*, *L niloticus*, and *P aethiopicus*, respectively. The EDI of Cd, Cu, and Pb through consumption of *O niloticus*, *R argentea*, *L niloticus*, and *P aethiopicus* from Nakuru town, Kenya, is presented in Table 2. The results show that the trends of the computed EDI values decreased in the following order Pb > Cu > Cd and ranged from 0.00086 to 0.00438, 0.00021 to 0.00099, and 0.00002 to 0.0003 mg kg⁻¹ day⁻¹, respectively.

The highest EDI value of Cd (0.0003 mg kg⁻¹ day⁻¹) was much lower than the PTDI, whose value is 0.001 mg kg⁻¹ day⁻¹ of body weight^{48,49} corresponding to 0.0607 mg kg⁻¹ for an

adult weighing 60.7 kg. Daily Cd intake through consumption of the commonly consumed fish varieties was lower than or similar to the daily intake reported for adult consumers from other developing countries—Ghana: 0.000092 mg kg⁻¹ day⁻¹,⁵⁸ Nigeria: 0.000063 mg kg⁻¹ day⁻¹,⁵⁴ Ethiopia: 0.000029 mg kg⁻¹ day⁻¹,⁵⁹ D. R. Congo: 0.00012 to 0.00094 mg kg⁻¹ day⁻¹,⁶⁰ Egypt: 0.000048 to 0.00022 mg kg⁻¹ day⁻¹,⁶¹ Iran: 0.00012 mg kg⁻¹ day⁻¹,⁶² and Bangladesh: 0.00027 to 0.00246 mg kg⁻¹ day⁻¹.⁶³ Daily Cd intake was higher than that reported for adults in Yaoundé, Cameroon, 0.000017 mg kg⁻¹ day⁻¹,⁶⁴ and roughly 26 times higher than that reported for adults from Dares salaam, Tanzania.⁶⁵

With respect to Cu, the highest daily intake (0.00099 mg kg⁻¹ day⁻¹) corresponded to consumers of *O niloticus* from Market 1. Compared with the current provisional maximum tolerable daily intake (PMTDI) whose value is 30.35 mg kg⁻¹ day⁻¹ for an adult weighing 60.7 kg,⁴⁷ the highest Cu intake was tens of thousands fold lower than its permissible daily limit. This implies that Cu in the commonly consumed fish species does not pose any threat to consumers. Lower or similar Cu EDI values are reported in literature in the range of 0.00000066 to 0.00093 mg kg⁻¹ day⁻¹.^{54,58,63-65} Higher daily intake of Cu through fish consumption is reported for consumers in Egypt, 0.0124 – 0.0250 mg kg⁻¹ day⁻¹, and Congo, 0.12586 – 0.0211428 mg kg⁻¹ day⁻¹.^{60,61,63}

As for Pb, the greatest daily intake (0.00438 mg kg⁻¹) was also observed in consumers of *O niloticus* from market 1. The PTDI of Pb according FAO/WHO⁴⁸ is 0.00357 mg kg⁻¹ day⁻¹, corresponding to 0.2168 mg kg⁻¹ day⁻¹ for an adult weighing 60.7 kg. All the EDI values of Pb were below this permissible limit. Nonetheless, they were significantly higher than those of

Table 3. Target hazard quotient and hazard index of consumers of commonly consumed fish species from 5 retail markets in Nakuru town, Kenya.

SAMPLING SITE	FISH SPECIES	FISH INTAKE, KG DAY ⁻¹	Cd THQ	Cu THQ	Pb THQ	HI
Market 1	<i>O niloticus</i>	0.027	0.295	0.025	1.168	1.488
Market 1	<i>L niloticus</i>	0.035	0.083	0.009	1.099	1.192
Market 1	<i>R argentea</i>	0.012	0.043	0.014	0.230	0.286
Market 1	<i>P aethiopicus</i>	0.023	0.284	0.006	0.699	0.989
Market 2	<i>O niloticus</i>	0.027	0.284	0.021	0.737	1.042
Market 2	<i>R argentea</i>	0.012	0.331	0.011	0.662	1.004
Market 3	<i>O niloticus</i>	0.027	0.178	0.005	0.400	0.584
Market 3	<i>R argentea</i>	0.012	0.022	0.015	0.276	0.313
Market 4	<i>O niloticus</i>	0.027	0.074	0.007	0.865	0.946
Market 4	<i>L niloticus</i>	0.035	0.106	0.011	0.609	0.725
Market 5	<i>R argentea</i>	0.012	0.216	0.012	0.518	0.745

Abbreviations: HI, hazard index; THQ, target hazard quotient.

Cd and Cu suggesting that Pb may pose serious health risk to consumers due to the likelihood of bio-accumulation over a period of time. The EDI of Pb compared very well with Pb intake of adult fish consumers elsewhere in developing countries.^{54,61,63} Lower EDI values in the range of 0.0000053 to 0.0008 mg kg⁻¹ day⁻¹ were reported for adult consumers from Cameroon, Congo, Ethiopia, Ghana, and Tanzania.^{58–60,64,65}

Target hazard quotient and hazard index

Noncarcinogenic THQs and hazard index of the 3 trace metals through consumption of the commonly consumed fish species sold in Nakuru Town, Kenya, are presented in Table 3. The trend of THQ values for consumers of the commonly consumed fish species decreased in the order of Pb > Cd > Cu. The THQ values of Cd, Cu, and Pb varied from 0.022 to 0.331, 0.005 to 0.025, and 0.230 to 1.168, respectively. Target hazard quotient is a ratio between potential exposure to a given trace metal and its oral reference dose. It is used to assess potential health risk associated with long-term exposure to dietary trace metals.^{5,54} If the computed ratio is greater than 1, the exposed population is likely to develop adverse health effects.^{55,56,66} According to the New York State Department of Health (NYSDOH), if the computed THQ value is greater than 1 but less than 5, the risk is low; if it is greater than 5 but less than 10, the risk is moderate; however, if the THQ value is greater than 10, the risk is high.⁶⁷ Computed THQ values of Cd and Cu were all below 1, suggesting that health effects associated with Cd and Cu are unlikely to occur. In contrast, THQ values of Pb for consumers of *O niloticus* and *L niloticus* bought from market 1 were above 1, indicating a likelihood—albeit low—of adverse health effects occurring.

Hazard index is the numerical sum of the computed THQ values.^{5,54} Like THQ, HI values should not exceed 1 otherwise

they would present significant health risks to consumers.^{5,54} HI values of Cd, Cu, and Pb in this study ranged from 0.286 to 1.488. THQ values of Pb were significantly higher than those of Cd and Cu and contributed between 66% and 92% to the HI values. Consumers of *O niloticus* and *L niloticus* from market 1 and *O niloticus* and *R argentea* from market 2 had HI values greater than 1, suggesting significant health risks for these consumers posed by the possible additive effects of Cd, Cu, and Pb.

Conclusion

Estimated daily intake, THQ, and hazard index were used to assess consumers' exposure to Cd, Cu, and Pb via consumption of the commonly consumed fish species (*O niloticus*, *R argentea*, *L niloticus*, and *P aethiopicus*) in Nakuru town, Kenya. Daily intake of Cd, Cu, and Pb through consumption of the commonly consumed fish species were within FAO/WHO recommended limits. THQ values of Cd and Cu suggest no health risk to consumers of the 4 commonly consumed fish species. In contrast, THQ values of Pb indicate that long-term exposure to Pb content similar to those reported in the present study may pose serious health risk to fish consumers in Nakuru town, Kenya. There is therefore a need for similar studies to be carried out regularly to monitor trends of trace metals in fish and educate consumers on safe quantities of fish to consume. Similar studies done in the future should also determine the sources of fish and trace metal contamination in aquatic ecosystems. Only then strict monitoring and mitigation measures can be implemented to reduce trace metals contents in fish and by that reduce the associated human health risks.

Author Contributions

FE, WNM, and MM conceived, designed and planned this research. WNM and MM supervised the project. FE and TW carried out the experiments. FE, WNM, MM, and TW

contributed to the analysis and the interpretation of the results. FE took the lead in writing the manuscript with significant and critical contributions from WNM, MM, and TM.

ORCID iD

Felly Esilaba  <https://orcid.org/0000-0003-1336-704X>

REFERENCES

- Fuentes-Gandara F, Herrera-Herrera C, Pinedo-Hernández J, Marrugo-Negrete J, Díez S. Assessment of human health risk associated with methylmercury in the imported fish marketed in the Caribbean. *Environ Res.* 2018;165:324-329.
- Marengo M, Durieux EDH, Ternengo S, et al. Comparison of elemental composition in two wild and cultured marine fish and potential risks to human health. *Ecotoxicol Environ Saf.* 2018;158:204-212.
- Yi Y, Yang Z, Zhang S. Ecological risk assessment of heavy metals in sediment and human health risk assessment of heavy metals in fishes in the middle and lower reaches of the Yangtze River basin. *Environ Pollut.* 2011;159:2575-2585.
- FAO. *The State of World Fisheries and Aquaculture 2016: Contributing to Food Security and Nutrition for All*. Rome, Italy: FAO; 2016.
- Javed M, Usmani N. Accumulation of heavy metals and human health risk assessment via the consumption of freshwater fish *Mastacembelus armatus* inhabiting, thermal power plant effluent loaded canal. *SpringerPlus.* 2016;5:776.
- Suami RB, Sivalingam P, Kabala CD, et al. Concentration of heavy metals in edible fishes from Atlantic Coast of Muanda, Democratic Republic of the Congo. *J Food Compos Anal.* 2018;73:1-9.
- Ullah AKMA, Maksud MA, Khan SR, Lutfa LN, Quraishi SB. Dietary intake of heavy metals from eight highly consumed species of cultured fish and possible human health risk implications in Bangladesh. *Toxicol Rep.* 2017;4:574-579.
- Faroon O, Ashizawa A, Wright S, et al. *Toxicological Profile for Cadmium*. Atlanta, GA: Agency for Toxic Substances and Disease Registry; 2012. <https://www.atsdr.cdc.gov/ToxProfiles/tp5.pdf>.
- Waalkes MP. Cadmium carcinogenesis in review. *J Inorg Biochem.* 2000;79:241-244.
- Waisberg M, Joseph P, Hale B, Beyersmann D. Molecular and cellular mechanisms of cadmium carcinogenesis. *Toxicology.* 2003;192:95-117.
- Lei L, Liang D, Yu D. Human health risk assessment of heavy metals in the irrigated area of Jinghui, Shaanxi, China, in terms of wheat flour consumption. *Environ Monit Assess.* 2015;187:647.
- Abadin H, Ashizawa A, Stevens YW, et al. *Toxicological Profile for Lead*. Atlanta, GA: Agency for Toxic Substances and Disease Registry; 2007. <https://www.atsdr.cdc.gov/toxprofiles/tp13.pdf>.
- Flora G, Gupta D, Tiwari A. Toxicity of lead : a review with recent updates. *Interdiscip Toxicol.* 2012;5:47-58.
- Navas-Acien A, Guallar E, Silbergeld EK, Rothenberg SJ. Lead exposure and cardiovascular disease—a systematic review. *Environ Health Perspect.* 2007;115:472-482.
- Mudipalli A. Lead hepatotoxicity & potential health effects. *Indian J Med Res.* 2007;126:518-527.
- Gaetke LM, Chow CK. Copper toxicity, oxidative stress, and antioxidant nutrients. *Toxicology.* 2003;189:147-163.
- Dorsey A, Ingberman L, Swarts S. *Toxicological Profile for Copper*. Atlanta, GA: Agency for Toxic Substances and Disease Registry; 2004. <https://www.atsdr.cdc.gov/toxprofiles/tp132.pdf>.
- Mok JS, Yoo HD, Kim PH, et al. Bioaccumulation of heavy metals in oysters from the Southern Coast of Korea : assessment of potential risk to human health. *Bull Environ Contam Toxicol.* 2015;94:749-755.
- Jonathan MP, Auriolos-Gamboa D, Villegas LEC, Bohórquez-Herrera J, Hernández-Camacho CJ, Ujitha SB. Metal concentrations in Demersal fish species from Santa Maria Bay, Baja California Sur, Mexico (Pacific coast). *Mar Pollut Bull.* 2015;99:356-361.
- Lepak JM, Hooten MB, Eagles-Smith CA, et al. Assessing potential health risks to fish and humans using mercury concentrations in inland fish from across western Canada and the United States. *Sci Total Environ.* 2016;571:342-354.
- Jose A, Ray JG. Toxic heavy metals in human blood in relation to certain food and environmental samples in Kerala, South India. *Environ Sci Pollut Res Int.* 2018;25:7946-7953.
- Liu Q, Liao Y, Shou L. Concentration and potential health risk of heavy metals in seafoods collected from Sanmen Bay and its adjacent areas, China. *Mar Pollut Bull.* 2018;131:356-364.
- Anandkumar A, Nagarajan R, Prabakaran K, et al. Bioaccumulation of trace metals in the coastal Borneo (Malaysia) and health risk assessment. *Mar Pollut Bull.* 2019;145:56-66.
- Abdallah MA. Bioaccumulation of heavy metals in Mollusca species and assessment of potential risks to human health. *Bull Environ Contam Toxicol.* 2013;90:552-557.
- Kaya G, Turkoglu S. Bioaccumulation of heavy metals in various tissues of some fish species and green tiger shrimp (*Penaeus semisulcatus*) from Iskenderun Bay, Turkey, and risk assessment for human health. *Biol Trace Elem Res.* 2017;180:314-326.
- Rahmani J, Fakhri Y, Shahsavani A, et al. A systematic review and meta-analysis of metal concentrations in canned tuna fish in Iran and human health risk assessment. *Food Chem Toxicol.* 2018;118:752-765.
- Wang X, Gu Y, Wang Z, Ke C, Mo M. Biological risk assessment of heavy metals in sediments and health risk assessment in bivalve mollusks from Kaozhouy-ang Bay, South China. *Mar Pollut Bull.* 2018;133:312-319.
- Atique Ullah AKM, Akter M, Musarrat M, Quraishi SB. Evaluation of possible human health risk of heavy metals from the consumption of two marine fish species *Tenualosa ilisha* and *Dorosoma cepedianum*. *Biol Trace Elem Res.* 2019;191:485-494.
- Islam MS, Ahmed MK, Habibullah-AI-Mamun M, Raknuzzaman M, Ali MM, Eaton DW. Health risk assessment due to heavy metal exposure from commonly consumed fish and vegetables. *Environ Syst Decis.* 2016;36:253-265.
- Kwon JC, Nejad ZD, Jung MC. Arsenic and heavy metals in paddy soil and polished rice contaminated by mining activities in Korea. *Catena.* 2016;148:92-100.
- Alam M, Khan M, Khan A, et al. Concentrations, dietary exposure, and human health risk assessment of heavy metals in market vegetables of Peshawar. *Environ Monit Assess.* 2018;190:505.
- Chen H, Yang X, Wang P, Wang Z, Li M, Zhao F. Dietary cadmium intake from rice and vegetables and potential health risk : a case study in Xiangtan, southern China. *Sci Total Environ.* 2018;639:271-277.
- Ji Y, Wu P, Zhang J, et al. Heavy metal accumulation, risk assessment and integrated biomarker responses of local vegetables: a case study along the Le'an river. *Chemosphere.* 2018;199:361-371.
- Orisakwe OE, Ozoani HA, Nwaogazie IL, Ezejirofor AN. Probabilistic health risk assessment of heavy metals in honey, *Manihot esculenta*, and *Vernonia amygdalina* consumed in Enugu State, Nigeria. *Environ Monit Assess.* 2019;191:424.
- Oyoo-Okoth E, Admiraal W, Osano O, Ngure V, Kraak MH, Omutange ES. Monitoring exposure to heavy metals among children in Lake Victoria, Kenya : environmental and fish matrix. *Ecotoxicol Environ Saf.* 2010;73:1797-1803.
- Oyoo-Okoth E, Admiraal W, Osano O, et al. Contribution of soil, water and food consumption to metal exposure of children from geological enriched environments in the coastal zone of Lake Victoria, Kenya. *Int J Hyg Environ Health.* 2013;216:8-16.
- Otachi EO, Körner W, Avenant-Oldewage A, Fellner-Frank C, Jirsa F. Trace elements in sediments, blue spotted tilapia *Oreochromis leucostictus* (Trewavas, 1933) and its parasite *Contracaecum multipapillatum* from Lake Naivasha, Kenya, including a comprehensive health risk analysis. *Environ Sci Pollut Res Int.* 2014;21:7339-7349.
- Mutia TM, Virani MZ, Moturi WN, Muyela B, Mavura WJ, Lalah JO. Copper, lead and cadmium concentrations in surface water, sediment and fish, *C. Carpio*, samples from Lake Naivasha: effect of recent anthropogenic activities. *Environ Earth Sci.* 2012;67:1121-1130.
- Omwenga I, Kanja L, Nguta J, Mbaria J, Irungu P. Assessment of lead and cadmium residues in farmed fish in Machakos and Kiambu counties, Kenya. *Toxicol Environ Chem.* 2014;96:58-67.
- Nyamari JM, Simiyu GM. Urban livestock and potential human health risks in Eldoret town, Kenya. *J Build L Dev.* 2007;14:100-107.
- Mbiri A, Onditi A, Oyaro N, Murago E. Determination of essential and heavy metals in Kenyan honey by atomic absorption and emission spectroscopy. *J Agric Sci Technol.* 2011;13:107-115.
- Maiyo WK, Mitei YJ, Kagwanja SM. Heavy metal contamination in raw honey, soil and flower samples obtained from Baringo and Keiyo Counties, Kenya. *Int J Emerg Sci Eng.* 2014;2:5-9.
- Onyango CM, Shibairo SI, Imungi JK, Harbinson J. The physico-chemical characteristics and some nutritional values of vegetable amaranth sold in Nairobi-Kenya. *Ecol Food Nutr.* 2008;47:382-398.
- Gallaher CM, Mwaniki D, Njenga M, Karanja NK, Winklerprins AMGA. Real or perceived: the environmental health risks of urban sack gardening in Kibera Slums of Nairobi, Kenya. *Ecobhealth.* 2013;10:9-20.
- Inoti KJ, Fanuel K, George O, Paul O. Assessment of heavy metal concentrations in urban grown vegetables in Thika Town, Kenya. *African J Food Sci.* 2012;6:41-46.
- Copat C, Bella F, Castaing M, Fallico R, Sciacca S. Heavy metals concentrations in fish from Sicily (Mediterranean Sea) and evaluation of possible health risks to consumers. *Bull Environ Contam Toxicol.* 2012;88:78-83.
- JEFCA. *Evaluation of Certain Food Additives and Contaminants: Twenty-sixth Report of the Joint FAO/WHO Expert Committee on Food Additives*. Geneva, Switzerland: World Health Organization; 1982.

48. JEFCA. *Evaluation of Certain Food Additives and Contaminants: Forty-first Report of the Joint FAO/WHO Expert Committee on Food Additives*. Geneva, Switzerland: World Health Organization; 1993.
49. JEFCA. *Evaluation of Certain Food Additives and Contaminants: Sixty-first Report of the Joint FAO/WHO Expert Committee on Food Additives*. Geneva, Switzerland: World Health Organization; 2004.
50. Dalziel J, Baker C. Analytical methods for measuring metals by atomic absorption spectrophotometry. In: Naeve H, ed. *Manual of Methods in Aquatic Environment Research: Part 9—Analyses of Metals and Organochlorines in Fish*. Rome, Italy: FAO; 1983:14-20.
51. Anandkumar A, Nagarajan R, Prabakaran K, Bing CH, Rajaram R. Human health risk assessment and bioaccumulation of trace metals in fish species collected from the Miri coast, Sarawak, Borneo. *Mar Pollut Bull*. 2018;133:655-663.
52. Chen Y, Hu W, Huang B, et al. Accumulation and health risk of heavy metals in vegetables from harmless and organic vegetable production systems of China. *Ecotoxicol Environ Saf*. 2013;98:324-330.
53. Taghizadeh SF, Davarynejad G, Asili J, et al. Health risk assessment of heavy metals via dietary intake of five pistachio (*Pistacia vera* L.) cultivars collected from different geographical sites of Iran. *Food Chem Toxicol*. 2017;107:99-107.
54. Moslen M, Miebaka CA. Concentration of heavy metals and health risk assessment of consumption of fish (*Sarotherodon melanotheron*) from an Estuarine Creek in the Niger Delta, Nigeria. *IOSR J Env Sci, Toxicol Food Technol*. 2017;11:68-73.
55. Yabanli M, Alparslan Y. Potential health hazard assessment in terms of some heavy metals determined in Demersal fishes caught in Eastern Aegean. *Bull Environ Contam Toxicol*. 2015;95:494-498.
56. Yap CK, Jusoh A. Potential human health risk assessment of heavy metals via the consumption of tilapia *Oreochromis mossambicus* collected from contaminated and uncontaminated ponds. *Environ Monit Assess*. 2015;187:584.
57. Nauen CE. *Compilation of Legal Limits for Hazardous Substances in Fish and Fishery Products* (FAO Fish Circular). Rome, Italy: FAO; 1983.
58. Kwaansa-Ansah EE, Nti SO, Opoku F. Heavy metals concentration and human health risk assessment in seven commercial fish species from Asafo Market, Ghana. *Food Sci Biotechnol*. 2018;28:569-579.
59. Dsikowitzky L, Mengesha M, Dadebo E, de Carvalho CEV, Sindern S. Assessment of heavy metals in water samples and tissues of edible fish species from Awassa and Koka Rift Valley. *Environ Monit Assess*. 2013;185:3117-3131.
60. Squadrone S, Burioli E, Monaco G, et al. Human exposure to metals due to consumption of fish from an artificial lake basin close to an active mining area in Katanga (D.R. Congo). *Sci Total Environ*. 2016;568:679-684.
61. El-Sadaawy MM, El-Said GF, Sallam NA. Bioavailability of heavy metals in fresh water *Tilapia nilotica* (*Oreochromis niloticus* Linnaeus, 1758): potential risk to fishermen and consumers. *J Environ Sci Health B*. 2013;48:402-409.
62. Miri M, Akbari E, Amrane A, et al. Health risk assessment of heavy metal intake due to fish consumption in the Sistan region, Iran. *Environ Monit Assess*. 2017;189:583.
63. Ahmed K, Baki MA, Islam S. Human health risk assessment of heavy metals in tropical fish and shellfish collected from the river Buriganga, Bangladesh. *Environ Sci Pollut Res Int*. 2015;22:15880-15890.
64. Gimou M, Pouillot R, Charrondiere UR, et al. Food additives & contaminants : part a dietary exposure and health risk assessment for 14 toxic and essential trace elements in Yaoundé: the Cameroonian total diet study. *Food Addit Contam Part A*. 2014;31:1064-1080.
65. Elisante E. Urban dietary heavy metal intake from protein foods and vegetables in Dar es Salaam. *Tanzania J Sci*. 2010;36:85-94.
66. Peters DE, Eebu C, Nkpaa KW. Potential human health risk assessment of heavy metals via consumption of root tubers from Ogoniland, Rivers State, Nigeria. *Biol Trace Elem Res*. 2018;186:568-578.
67. NYSDOH. Hopewell precision area contamination: NYSDOH procedure for evaluating potential health risks for contaminants of concern. <https://www.health.ny.gov/environmental/investigations/hopewell/appendc.htm>. Published 2007.