



Global Health Impacts of Dust Storms: A Systematic Review

Authors: Aghababaeian, Hamidreza, Ostadtaghizadeh, Abbas, Ardalan, Ali, Asgary, Ali, Akbary, Mehry, et al.

Source: Environmental Health Insights, 15(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/11786302211018390>


BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.


Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Global Health Impacts of Dust Storms: A Systematic Review

Hamidreza Aghababaeian^{1,2,3} , Abbas Ostadtaghizadeh^{1,2},
Ali Ardalan¹, Ali Asgary⁴, Mehry Akbary⁵, Mir Saeed Yekaninejad⁶
and Carolyn Stephens⁷

¹Department of Health in Emergencies and Disasters, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran. ²Center for Air Pollution Research (CAPR), Institute for Environmental Research (IER), Tehran University of Medical Sciences, Tehran, Iran. ³Department of Nursing and Emergency, Dezfoul University of Medical Sciences, Dezfoul, Iran. ⁴Disaster and Emergency Management, School of Administrative Studies, York University, Toronto, Canada. ⁵Department of Climatology, Faculty of Geographical Sciences, Kharazmi University, Tehran, Iran. ⁶Department of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran. ⁷UCL Bartlett Development Planning Unit, London School of Hygiene & Tropical Medicine, London, UK.

Environmental Health Insights
Volume 15: 1–28
© The Author(s) 2021
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/11786302211018390


ABSTRACT

BACKGROUND: Dust storms and their impacts on health are becoming a major public health issue. The current study examines the health impacts of dust storms around the world to provide an overview of this issue.

METHOD: In this systematic review, 140 relevant and authoritative English articles on the impacts of dust storms on health (up to September 2019) were identified and extracted from 28968 articles using valid keywords from various databases (PubMed, WOS, EMBASE, and Scopus) and multiple screening steps. Selected papers were then qualitatively examined and evaluated. Evaluation results were summarized using an Extraction Table.

RESULTS: The results of the study are divided into two parts: short and long-term impacts of dust storms. Short-term impacts include mortality, visitation, emergency medical dispatch, hospitalization, increased symptoms, and decreased pulmonary function. Long-term impacts include pregnancy, cognitive difficulties, and birth problems. Additionally, this study shows that dust storms have devastating impacts on health, affecting cardiovascular and respiratory health in particular.

CONCLUSION: The findings of this study show that dust storms have significant public health impacts. More attention should be paid to these natural hazards to prepare for, respond to, and mitigate these hazardous events to reduce their negative health impacts.

Registration: PROSPERO registration number CRD42018093325

KEYWORDS: Air quality, desert dust, dust storm, health, PM₁₀

RECEIVED: February 16, 2021. **ACCEPTED:** April 27, 2021.

TYPE: Review

FUNDING: The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study is a part of PhD dissertation and was funded and supported by School of public health and Center for air pollution research (CAPR), institute for environmental research (IER), Tehran University of Medical Sciences, Tehran, Iran, Grant No: 98-03-46-43717.

DECLARATION OF CONFLICTING INTERESTS: The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

CORRESPONDING AUTHOR: Abbas Ostadtaghizadeh, Department of Health in Emergencies and Disasters, School of Public Health, Tehran University of Medical Sciences, Poorsina Avenue, Tehran, Iran. Email: ostadtaghizadeh@gmail.com

Introduction

Dust storms are natural hazards and the most common sources of natural particles, including very small materials, potential allergens, and pollutants.^{1–5} Depending on the nature of the source of the dust, these materials and substances may include, quartz, silicon dioxide, oxides of magnesium, calcium, iron, and aluminum^{6,7} and sometimes a range of organic matter, anthropogenic pollutants, and salts.⁸ Dust storms carry millions of tons of soil into the air each year from thousands of kilometers away. They can last a few hours or a few days^{1–5} and distribute a large number of small particles in the air,^{9,10} increasing the amount of particles above the allowable threshold for human health.^{11,12} During a dust storm event, the concentration of PM₁₀ (particles with an aerodynamic diameter <10 μm) and

PM_{2.5} (particles with an aerodynamic diameter <2.5 μm) particles are often higher than the normal thresholds recommended by the World Health Organization (PM_{2.5}: 10 μg/m³ annual mean, 25 μg/m³ 24-hour mean. PM₁₀: 20 μg/m³ annual mean, 50 μg/m³ 24-hour mean).^{8,13} It can also exceed 6000 μg/m³ in seriously strong dust storms.¹⁴ According to the Huffman Classification of dust PM₁₀ range (μg/m³), in dusty air, light dust storm, dust storm, strong dust storm, and serious strong dust storm days, levels can be between 50 to 200, 200 to 500, 500 to 2000, 2000 to 5000, and >5000, respectively.¹⁵

Dust storms are occurring increasingly frequently in many desert areas and arid regions around the world,³ causing extensive damage and emergencies each year.^{3,16–18} Therefore, dust storms have attracted increasing attention in recent



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without

years.^{16,17,19} Researchers have demonstrated how dust storms affect various aspects of human life.¹⁹ The particles in dust storms affect weather conditions, agricultural production, human health, and the ecosystem.^{20,21} Evidence suggests that mineral aerosols affect cloud formation and precipitation and can reduce the acidity of precipitation.²² Moreover, a high density and diversity of bacteria and plant pollens have been observed during dust storms.²³ In addition to endangering the ecosystem, dust storms have direct and indirect impacts on public health and human health.^{8,20,21,24} Due to their small sizes, almost all dust storm particles, that is, airborne particles (PM) can enter the respiratory tract²⁵; larger particles are often deposited in the upper respiratory tract (nasopharyngeal region, tracheobronchial region), while smaller particles can enter deep lung tissue.^{26,27} The physical, biological, and chemical properties of these particles can cause disorders in the health of the body,^{8,24,26} and in addition to the respiratory tract, can damage other systems of the body, including the cerebral, cardiovascular, skin,^{8,24,26} blood, and immune systems.^{28,29}

Research has indicated that exposure to dust particles, which can remain in the air from hours to days,²⁴ can result in other problems like conjunctivitis, meningitis, and valley fever.^{24,26,30} In rare cases, it can even lead to death.^{26,31} Evidence further suggests that frequent exposure to dust storms can lead to increased adverse health effects^{24,32-37} in people of almost all age groups and genders.^{3,38,39} People with a history of diabetes, hypertension, cerebrovascular, or pulmonary disease are also at higher risk.⁴⁰ Many epidemiological studies have determined the health effects of dust storms by comparing outcomes during dust storm periods with outcomes during non-dust storm periods⁴¹⁻⁴³ and by assessing the correlation between dust storms or PM₁₀ exposure and health outcomes.^{32,44} Many researcher have acknowledged the existence of a significant association between dust exposure and increased morbidity or mortality, but there is no consensus in this regard to date.⁴⁵ Pérez et al. stated that increased PM during dust storms caused a significant increase in mortality rate in Barcelona.⁴⁶ Chen et al.,⁴⁷ Kashima et al.,⁴⁸ and Delangizan⁴⁹ also noted that increased PM₁₀ levels during Asian dust storms increased cardiovascular mortality. Some studies have reported that Middle Eastern dust storms can affect inflammation and coagulation markers in young adults,^{28,29} have adverse effects on pulmonary function,⁵⁰ and increase the number of asthma patients.⁵¹⁻⁵² Conversely, some studies have either ruled out the possibility of an increase in mortality or hospitalizations of patients due to dust storm exposure or do not consider the increase to be significant.^{43,53-55} For example, in studies conducted in Italy,⁵³ Greece,⁵⁴ Kuwait,⁴³ and Taipei,⁵⁵ researchers found no significant relationship between dust storms and increased risk of death. Bell,⁵⁶ Ueda,⁵⁷ and Min⁵⁸ also found that dust storms did not significantly increase hospitalizations of asthmatic patients or asthma attacks in Taipei and Japan.⁵⁶

There are mixed results and a lack of accurate and up-to-date classified data about the health impacts of dust storms on humans around the world. Moreover, the causes of dust storm-related health problems are not yet completely understood.⁵⁹ Given the importance of the impact of dust storms on human health as well as the increasing evidence of recurring and negative impacts of these storms, and because of the lack of systematic review studies, the current study conducted an extensive review of the current literature on the impacts of dust storms on human health.

Materials and Methods

This systematic review of scientific resources identified articles related to dust storms and related human health outcomes published up to 30 September 2019. PubMed, EMBASE, Scopus, and ISI WoS (Web of Science) databases were searched for articles published in relevant journals from the 28th to the 30th of October, 2019. All peer-reviewed articles from English language journals were discovered in the primary search stage. Citations and references of all relevant articles were examined and searched manually to ensure that all relevant articles were included. The primary search used the following Medical Subject Headings (MeSH terms) and keywords: Dust* OR Kosa OR Yellow sand OR Arabian Sand OR Dust Storms AND Mortality OR Disease* OR Morbidity OR Admission* OR Health* OR "Adverse affect" OR affect*.

Executive limitations: The main limitations of the current study were the lack of access to all required databases as well as the lack of access to the full text of some articles which should be obtained by correspondence with the authors of those articles. To resolve this problem, the researchers resorted to using resources from various universities inside and outside the country.

Inclusion criteria: All studies that had the full text available, that used appropriate methods and data, and that calculated the impacts of dust storms on health (eg, odds ratio, relative risk, rate ratio, regression coefficient, percentage change, excess risk, etc. in health indicators following dust storms); those in which dust storm was a major problem and those in which health indicators were analyzed were included in this study without restrictions on the publication date.

Exclusion criteria: Non-English articles, non-research letters to editors, review studies, case reports, case series, specialized articles about microorganisms, animal experiments, in vitro studies, and dust from volcanic or manmade sources like stone mines or stone and cement factories were excluded.

Data collection process: The current study followed the PRISMA guidelines (PRISMA Flow Diagram). EndNote software was used to manage the retrieved articles. After all articles were entered into the software, duplicates were identified and removed. Then, 2 researchers screened the remaining articles separately based on the inclusion and exclusion criteria by reading the titles, abstracts, and keywords. After removing unrelated papers, the full text of the remaining articles were




Figure 1. PRISMA flow diagram.

found and attached, and the quality of each paper in a standard format related to the type of study was assessed separately by the 2 researchers using JBI's critical appraisal tools. In cases of disagreement between the researchers, the third researcher helped to select the most relevant items.

Data extraction: The information required for this study was extracted using a checklist previously reviewed and prepared, which included all the characteristics of the selected articles, including type of article, publication year, first author's name, location of study, study design/methodology, health effects, PM fraction, and age/gender.

Risk of bias (quality) assessment: For quality assessment of the included papers, the Critical Appraisal Skills Program (CASP) checklist was used. The assessment was conducted by 3 independent reviewers. Discrepancies were resolved by 2 other reviewers.

Results

Search results

Out of a total of 35 712 articles searched, 140 articles met the inclusion criteria (Figure 1). The majority of them were related

to ecological, case crossover, and prospective studies; other studies included descriptive, retrospective, and Panel studies and 1 research letter (Table 1).

The current results showed that most data analyses investigated the effects of dust storms on health and used the generalized additive model (GAM) with nonlinear Poisson regression method to analyze the data in ecological and case-crossover studies.

Furthermore, most studies on the impact of dust storms on health were performed within the last decade (Chart 1).

Most health and dust storm studies included in this study were undertaken in Japan ($n=29$; 20.71%), Taiwan ($n=25$; 17.85%), Korea ($n=16$; 11.42%), China ($n=10$; 7.14%), Spain ($n=9$; 6.42%), and Iran ($n=8$; 5.71%), respectively (Figure 2).

In this review, the following adverse health effects of dust storms emerged as important:

- Non-accidental death (mortality due to respiratory, cardiovascular, or cerebrovascular disease);
- Emergency medical dispatch, hospitalization or admission, and hospital visits due to respiratory or cardiovascular diseases;

Table 1. Published studies on adverse health effects of dust storms.

REFERENCE	FIRST AUTHOR AND YEAR	STUDY LOCATION	POPULATION (AGE, GENDER)	PM FRACTION	STUDY DESIGN/ METHODOLOGY	HEALTH OUTCOMES	RESULTS
All-cause mortality							
Al et al. ⁶⁶	Al et al. (2018)	Gaziantep/Turkey	Older than 16 years	PM ₁₀	Retrospective study/ GAM	Mortality of cardiovascular diseases	Congestive cardiac failure Mortality, OR 0.95 (0.81–1.11) Acute coronary syndrome mortality, OR 0.40 (0.31–0.50)
Al-Tajer and Thalib ⁴³	Al-Tajer and Thalib (2014)	Kuwait	All ages/all gender	PM ₁₀	Ecological time series, GAM	All-causes, respiratory, cardiovascular Mortality	Respiratory mortality, RR 0.96 (0.88–1.04) Cardiovascular mortality, RR 0.98 (0.96–1.012) All-cause mortality, RR 0.99 (0.97–1.00)
Chan and Ng ³⁸	Chan and Ng (2011)	Taipei, Taiwan	All ages/all gender	PM ₁₀	Case-crossover/ conditional logistic regression models	Non-accidental, respiratory, cardiovascular, deaths	Non-accidental deaths, OR 1.019 (1.003–1.035) Above 65 years old, OR 1.025 (1.006–1.044) Cardiovascular deaths, OR 1.045 (1.0011–1.081) Respiratory deaths, OR 0.988 (1.038–0.941)
Chen et al. ³⁹	Chen et al. (2004)	Taipei, Taiwan	All ages/all gender	PM ₁₀	Case-crossover/tests of student	Daily mortality	Respiratory disease, RR 7.66% Total deaths, RR 4.92% Circulatory diseases, RR 2.59%
Crooks et al. ³	Crooks et al. (2016)	National/United States	All ages/all gender	PM ₁₀	Case-crossover/ conditional logistic regression models	Daily non-accidental mortality	Non-accidental mortality 7.4% ($p=0.011$) Lag _{3,3} 6.7% ($p=0.018$) Lag _{5,5} 2.7% ($p=0.023$)
Díaz et al. ⁶⁸	Díaz et al. (2017)	Spain: 9 region	All ages/all gender	PM ₁₀	Longitudinal ecological time series/ GAM	Daily mortality	Daily mortality values South-west, 21.20 (20.81–21.59) $p < 0.05$ South-east, 20.16 (19.88–20.45) $p < 0.05$ Canary Islands, 17.93 (17.60–18.26) $p < 0.05$
Díaz et al. ⁶⁹	Díaz et al. (2012)	Madrid (Spain)	All ages/all gender	PM ₁₀	Case-crossover design/Poisson regression model	Case-specific mortality	Respiratory death, IR 3.34% (0.36, 6.41) Circulatory causes, IR 4.19% (1.34, 7.13)
Hwang et al. ⁷⁰	Hwang et al. (2004)	Seoul, Korea	All ages/all gender	PM ₁₀	Ecological time series / GAM	Daily non accidental deaths	Non accidental deaths, 1.7% (1.6 5.3) Aged 65 years and older, 2.2% (3.5 8.3) Cardiovascular and respiratory, 4.1% (3.8 12.6)
Jimenez et al. ⁷¹	Jimenez et al. (2010)	Madrid (Spain)	Elderly	PM ₁₀ , PM _{2.5} or PM _{10-2.5}	Ecological time series/ Poisson regression models	Mortality	PM ₁₀ Total mortality, lag ₃ 1.02 (1.01–1.04) Circulatory, lag ₃ 1.04 (1.01–1.06) Respiratory, lag ₁ 1.03 (1.00–1.06)
Johnston et al. ⁷²	Johnston et al. (2011)	Sydney, Australia	All ages/all gender	PM ₁₀	Case crossover / conditional logistic regression model	Non-accidental mortality	Non-accidental mortality, lag ₃ , OR 1.16 (1.03–1.30)

(Continued)

Table 1. (Continued)

REFERENCE	FIRST AUTHOR AND YEAR	STUDY LOCATION	POPULATION (AGE, GENDER)	PM FRACTION	STUDY DESIGN/METHODOLOGY	HEALTH OUTCOMES	RESULTS
Kashima et al. ⁷³	Kashima et al. (2016)	South Korea and Japan	>65 years old/all gender	PM ₁₀	Ecological time-series analyses/specific Poisson regression models	Cause-specific mortality	All-cause mortality, lag ₀ RR 1.003 (1.001-1.005) lag ₁ , 1.001 (1.000-1.003) Cerebrovascular disease, lag ₁ RR: 1.006 (1.000-1.011)
Kashima et al. ⁴⁸	Kashima et al. (2012)	Western Japan	Aged 65 or above l	SPM	Ecological multi-city time-series analysis/Poisson regression models	Daily all-cause or cause-specific mortality	Heart disease, 0.6 (0.1-1.1) Ischemic heart disease, 0.8 (0.1-1.6) Arrhythmia, 2.1 (0.3-3.9) Pneumonia mortality, 0.5 (0.2-0.8)
Khaniabadi et al. ⁸⁷	Khaniabadi et al. (2017)	Ilam (Iran)	–	PM ₁₀	Ecological time series/air Q model	Respiratory mortality	Respiratory Mortality 7.3 (4.9-19.5)
Kim et al. ⁷⁴	Kim et al. (2012)	Seoul, Korea	General population/all gender	–	Ecological time-series/Poisson regression analyses	All-cause/cardiopulmonary mortality	The relative risk of total mortality for general population and over 75 years old increased on dusty days
Kwon et al. ⁸³	Kwon et al. (2002)	Seoul, Korea	All ages/all gender	PM ₁₀	Ecological time series/GLM with Poisson regression	Non accidental deaths	All causes, RR 1.7% (1.6, 5.3) Persons aged 65 years older, RR 2.2% (3.5, 8.3) Cardiovascular and respiratory death, RR 4.1% (3.8, 12.6)
Lee et al. ⁵⁵	Lee et al. (2014)	(Seoul, Korea; Taipei, Taiwan, Kitakyushu, Japan)	All ages/all gender	PM ₁₀	Ecological time-series using/GAM with Quasi-Poisson distribution	Mortality	Seoul: Under 65 years old (lag ₂ : 4.44%, lag ₃ : 5%, and lag ₄ : 4.39%) Kitakyushu: Respiratory mortality (lag ₂ : 18.82%) Total non-accidental mortality (lag ₀ : -2.77%, lag ₁ : -3.24%) Taipei: Over 65 years old (lag ₀ : -3.35%, lag ₁ : -3.29%) Respiratory mortality (lag ₀ : -10.62%, lag ₁ : -9.67%)
Lee et al. ⁷⁵	Lee et al. (2013)	Seven metropolitan cities of Korea	All ages/all gender	PM ₁₀	Ecological time-series/GAM with Quasi-Poisson regressions	Mortality	Lag ₀ Cardiovascular, 2.91% (0.13, 5.77) Male: 2.74% (0.74, 4.77) Lag ₂ <65 years, 2.52% (0.06, 5.04) Male 2.4% (0.43, 4.4) Lag ₃ <65 years, lag ₃ 2.49% (0.07, 4.97) Total non-accidental: 1.57% (0.11, 3.06) Male: 2.24% (0.28, 4.0) <65 years: 2.43% (0.01, 4.91) lag ₅ cardiovascular: 3.7% (0.93, 6.54)
Lee et al. ⁷⁶	Lee et al. (2007)	Seoul, Korea	All ages/all gender	PM ₁₀	Ecological time-series, GAM	Mortality	Total death, IR 0.7 (0.2, 1.3)

(Continued)

Table 1. (Continued)

REFERENCE	FIRST AUTHOR AND YEAR	STUDY LOCATION	POPULATION (AGE, GENDER)	PM FRACTION	STUDY DESIGN/METHODOLOGY	HEALTH OUTCOMES	RESULTS
Mallone et al. ⁸⁴	Mallone et al. (2011)	Rome, Italy	≥35 years/all gender	PM _{2.5} , PM _{2.5-10} , and PM ₁₀	Case-crossover/Poisson regression model	Mortality	PM _{2.5-10} Cardiac mortality, lag 0-2, IR 9.73 (4.25-15.49) Circulatory system, lag 0-2, IR 7.93 (3.20-12.88) PM ₁₀ Cardiac mortality, lag 0-2, IR 9.55 (3.81-15.61%)
Perez et al. ⁴⁶	Perez et al. (2008)	Barcelona (Spain)	All ages/all gender	PM _{2.5} and PM _{10-2.5}	Case crossover/linear regression	Daily Mortality	PM _{10-2.5} Daily mortality, Lag ₁ , OR 1.084 (1.015, 1.158)
Perez et al. ⁸⁵	Perez et al. (2012)	Barcelona (Spain)	All ages/all gender	PM ₁ , PM _{2.5} and PM ₁₀	Case-crossover/conditional logistic regression	Cause-specific mortality	PM _{10-2.5} OR Cardiovascular mortality, (lag ₁) 1.085 (1.01-1.15) <i>p</i> < 0.05 Respiratory mortality, (lag 2) 1.109 (0.978, 1.257) <i>p</i> < 0.1 PM _{2.5-1} OR Cardiovascular mortality, (lag ₁) 1.074 (0.998, 1.156) <i>p</i> < 0.1
Renzi et al. ⁷⁷	Renzi et al. (2018)	Sicily, Italy	All ages/all gender	PM ₁₀	Ecological time-series/Poisson conditional regression model	Mortality	Non-accidental mortality, (lag ₀₋₅) IR 3.8% (3.2, 4.4) Cardiovascular, IR 4.5% (3.8, 5.3) Respiratory IR 6.3% (5.4, 7.2)
Pirsaheb et al. ⁵⁰	Pirsaheb et al. (2016)	Kermanshah, Iran	All ages/all gender	PM ₁₀	Descriptive studies/spearman test	Death from cardiovascular and respiratory disease	Increased dust concentrations increase the risk of cardiovascular mortality
Schwartz et al. ⁸⁸	Schwartz et al. (1999)	Six United States, cities	All ages/all gender	PM ₁₀	Ecological/GAM with Poisson regression	Mortality	Mortality, RR 0.99 (0.81-1.22)
Sajani et al. ⁵³	Sajani et al. (2011)	Emilia-Romagna (Italy)	All ages/all gender	PM ₁₀	Case crossover/conditional logistic regression	Mortality	Respiratory mortality, OR 22.0 (4.0-43.1) Natural, OR 1.04 (0.99-1.09) Cardiovascular mortality, OR 1.04 (0.96-1.12)
Stafoggia et al. ⁷⁶	Stafoggia et al. (2016)	Southern European cities-Spain, France, Italy, Greece	All ages/all gender	PM ₁₀	Case-crossover/Poisson regression models	Mortality	Natural mortality lag ₀₋₁ , IR 0.65% (0.24-1.06)
Shahsavani et al. ⁷⁹	Shahsavani et al. (2019)	Tehran and Ahvaz, IRAN	All ages/all gender	PM ₁₀	Case crossover/conditional Poisson regression models	Mortality	Daily mortality 3.28 (2.42-4.15)
Tobias et al. ⁸⁰	Tobias et al. (2011)	Madrid (Spain)	All ages/all gender	PM _{2.5} and PM _{10-2.5}	Case-crossover/conditional logistic regression models	Mortality	PM _{10-2.5} Each increase of 10 µg/m ³ of PM _{10-2.5} increased Total mortality, 2.8% (<i>P</i> = 0.01)

(Continued)

Table 1. (Continued)

REFERENCE	FIRST AUTHOR AND YEAR	STUDY LOCATION	POPULATION (AGE, GENDER)	PM FRACTION	STUDY DESIGN/METHODOLOGY	HEALTH OUTCOMES	RESULTS
Wang and Lin ⁸¹	Wang and Lin (2015)	Metropolitan Taipei	All ages/all gender	PM ₁₀	Ecological time series/distributed lag non-linear model	Mortality	All-cause mortality lag ₀₋₅ , RR 1.10 (1.04–1.17) Elders 1.10 (1.02–1.18) Elderly circulatory Mortality lag ₀₋₅ , RR 1.21 (1.02–1.44)
Samoli et al. ⁵⁴	Samoli et al. (2011)	Athens, Greece	All ages/all gender	PM ₁₀	Ecological time series/Poisson regression models	Mortality	Mortality 0.71% (0.40-0.99)
Neophytou et al. ⁸²	Neophytou et al. (2013)	Nicosia, Cyprus	All ages/all gender	PM ₁₀	Ecological time-series/GAM	Mortality	Total non accidental, IR 0.13% (1.03, 1.30) Cardiovascular mortality, IR 2.43 (0.53–4.37) Respiratory mortality, IR 0.79 (4.69, 3.28)
Goto et al. ⁶⁰	Goto et al. (2010)	Western Japan	All ages/all gender	–	Ecological time-series/Spearman's rank correlation	Bronchial asthma mortality	Asthma mortality ($r=0.268$, $n=8$, $P>0.05$)
Achilleos et al. ⁴¹	Achilleos et al. (2019)	Kuwait	All ages/all gender	Poor visibility (AOD > 0.4)	Ecological time-series/generalized additive model (GAM)/Poisson regression models	Mortality	Rate ratio: 1.02, (1.00–1.04)
Emergency dispatch or air medical retrieval service							
Holyoak et al. ⁹⁰	Holyoak et al. (2011)	Queensland, Australia	–	–	Ecological retrospective review/simple t-test	Air medical retrieval service for respiratory and injury cases	Respiratory cases 62.5% increased Injury cases 13.3% increased
Aghababaeian et al. ⁴²	Aghababaeian et al. (2019)	Iran/dezful	All ages/all gender	PM ₁₀	Ecological time-series/GAM	Emergency dispatch of cardiovascular, respiratory and traffic accident missions	RR of Emergency dispatch Lag ₂ 1.008 (1.001–1.016)/ female/18–60 years/>60 years Lag ₃ 1.008 (1.00 1.01) Lag ₄ 1.008 (1.00–1.01) Lag ₅ 1.008 (1.00–1.01) Lag ₆ 1.007 (1.00–1.01) Lag ₇ 1.006 (1.000–1.01) Lag ₀₋₇ 1.06 (1.01–1.12) Lag ₀₋₁₄ 1.09 (1.01–1.17) >60 years 1.28 (1.08–1.52) Cardiovascular Problems Lag ₀₋₁₄ 1.33 (1.17–1.50) Respiratory problems Lag ₀₋₁₄ 1.13 (0.93–1.38) Traffic Accident Trauma Lag ₀₋₁₄ 1.03 (0.94–1.13)

(Continued)

Table 1. (Continued)

REFERENCE	FIRST AUTHOR AND YEAR	STUDY LOCATION	POPULATION (AGE, GENDER)	PM FRACTION	STUDY DESIGN/METHODOLOGY	HEALTH OUTCOMES	RESULTS
Kashima et al. ⁸⁹	Kashima et al. (2014)	Okayama, Japan	Elderly people	SPM	Ecological time-series/Poisson regression with GAM	Emergency ambulance calls	All causes, Lag 0 1.009 (1.002–1.017) Cardiovascular, lag ₀₋₃ 1.02 (1.00–1.03) Cardiovascular, Lag ₀ 1.016 (1.001–1.032) Cerebrovascular, Lag ₀ 1.028 (1.007–1.049) Pulmonary, Lag ₀ 1.005 (0.986–1.025)
Ueda et al. ⁶¹	Ueda et al. (2012)	Nagasaki, Japan	All ages/all gender	SPM	Case-crossover/conditional logistic regression	Emergency ambulance dispatches	All causes lag ₀₋₃ 12.1% (2.3–22.9) Cardiovascular diseases 20.8% (3.5–40.9)
Visits							
Akpinar-Elici et al. ¹³⁷	Akpinar-Elici et al. (2015)	Grenada, Caribbean	All ages/all gender	–	Ecological/regression analysis	Asthma visits	Asthma (R ² =0.036, <i>p</i> <0.001)
Cadelis et al. ¹³⁸	Cadelis et al. (2014)	Guadeloupe (Caribbean)	Children/all gender	PM ₁₀ , PM _{2.5-10}	Case-crossover/t-test and Mann-Whitney	Visits of children due to asthmatic conditions	PM ₁₀ Lag ₀ IR 9.1% (7.1–11.1) Lag ₀₋₁ IR 5.1% (1.8–7.7) PM _{2.5-10} Lag ₀ IR 4.5% (3.3–5) Lag ₀₋₁ IR: 4.7% (2.5–6.5)
Carlsen et al. ¹⁴²	Carlsen et al. (2015)	Reykjavik, Iceland	All ages/all gender	PM ₁₀	Ecological time-series study/generalized additive regression model	Emergency hospital visits	Emergency hospital visits 5.8% (<i>p</i> =0.02)
Chan et al. ¹⁴³	Chan et al. (2008)	Taipei, Taiwan	All ages/all gender	PM ₁₀	Ecological time-series/Poisson regression model and paired t-test	Emergency visits	Cardiovascular visits 1.5 (0.3–2.6) Ischemic heart diseases visits 0.7 (0.1–1.4) Cerebrovascular visits 0.7 (0.1–1.3) Chronic obstructive pulmonary disease (COPD) visits 0.9 (0.1–1.7)
Chien et al. ¹⁴⁴	Chien et al. (2014)	Taipei, Taiwan	Children	PM ₁₀	Ecological studies/structural additive regression modeling	Conjunctivitis clinic visits	Conjunctivitis visits Preschool children 1.48% (0.79, 2.17) Schoolchildren. 9.48% (9.03, 9.93)
Chien et al. ¹⁴⁶	Chien et al. (2012)	Taipei, Taiwan	Children	PM ₁₀	Ecological/STAR model and autoregressive correlation	Respiratory diseases visits	Respiratory visits Preschool children 2.54% (2.43, 2.66) Schoolchildren 5.03% (4.87, 5.20)
Hefflin et al. ¹⁴⁷	Hefflin et al. (1994)	Washington, United States	All ages/all gender	PM ₁₀	Ecological/multivariable analysis using generalized estimating equations	Emergency room visits for respiratory disorders	Daily number of emergency visits for bronchitis, IR 3.5% Daily Number of emergency room visits, IR 4.5%
Lin et al. ¹⁴⁸	Lin et al. (2016)	Taipei, Taiwan	All ages/all gender	PM ₁₀	Ecological time series/DLNM	Emergency room visits	All causes visits, RR 1.10 (1.07, 1.13) Respiratory visits, RR 1.14 (1.08, 1.21)

(Continued)

Table 1. (Continued)

REFERENCE	FIRST AUTHOR AND YEAR	STUDY LOCATION	POPULATION (AGE, GENDER)	PM FRACTION	STUDY DESIGN/METHODOLOGY	HEALTH OUTCOMES	RESULTS
Liu and Liao ¹⁴⁹	Liu and Liao (2017)	Taiwan	All ages/all gender	PM _{2.5}	Case-crossover/conditional logistic regression	Emergency visits	Cardiovascular, OR 2.92 (1.22–5.08) Respiratory, OR 1.86 (1.30–2.91)
Merrifield et al. ¹⁴¹	Merrifield et al. (2013)	Sydney, Australia	All ages/all gender	PM ₁₀	Ecological time-series/distributed-lag Poisson generalized models	Emergency visits	Asthma visits, RR 1.23, ($p < 0.01$) All visits, R 1.04, ($p < 0.01$) Respiratory visits, RR 1.20, ($p < 0.01$) Cardiovascular visits, RR 0.91, ($p = 0.09$)
Nakamura et al. ¹³⁹	Nakamura et al. (2016)	Nagasaki, Japan	children aged 0–15 years/all gender	SPM	Case-crossover/conditional logistic models	Pediatric emergency visits for respiratory diseases	School children Bronchial asthma visits, lag ₃ OR 1.83 (1.212–2.786) Lag ₄ 1.829 (CI, 1.179–2.806) Preschool children Respiratory visit, lag ₀ , OR 1.244 (1.128–1.373) Lag day 1, OR 1.314 (1.189–1.452) Lag day 2, OR 1.273 (1.152–1.408)
Park et al. ⁶²	Park et al. (2015)	Chuncheon, Gangwon-do, Korea	All ages/all gender	PM ₁₀	Ecological retrospective study/Poisson regression model	Hospital visits for airway diseases	Asthma visits, RR 1.10 ($P < 0.05$) COPD visits, RR 1.29 ($P < 0.05$)
Wang et al. ⁶³	Wang et al. (2016)	Minqin, China	All ages/all gender	–	Ecological time series/generated regression model	Pulmonary tuberculosis (PTB) visits	PTB visits, R ₂ = 0.685
Park et al. ¹⁴⁰	Park et al. (2016)	Seoul and Incheon, Korea	11–20, 51–70 and 490 years/all gender	PM ₁₀	Case-crossover/T-tests and Poisson regression model	Asthma exacerbation	Asthma related visits Lag ₀ , RR 0.96 (0.95–0.98) Lag ₁ , RR 1.27 (1.25–1.29) Lag ₂ , RR 1.12 (1.10–1.14) Lag ₃ , RR 1.25 (1.23–1.26) Lag ₄ , RR 1.13 (1.12–1.15) Lag ₅ , RR 1.06 (1.04–1.07) Lag ₆ , RR 0.82 (0.81–0.81)
Yu et al. ¹²	Yu et al. (2012)	Taipei (Taiwan)	Children	PM ₁₀	Ecological studies/STAR model/generalized additive mode	Children's respiratory health risks	All children Lag ₀ –3.66 Lag ₁ –2.05 Lag ₂ 1.78 Lag ₃ 2.40 Lag ₄ 0.66 Lag ₅ 1.74 Lag ₆ –1.01 Lag ₇ 2.26

(Continued)

Table 1. (Continued)

REFERENCE	FIRST AUTHOR AND YEAR	STUDY LOCATION	POPULATION (AGE, GENDER)	PM FRACTION	STUDY DESIGN/METHODOLOGY	HEALTH OUTCOMES	RESULTS
Yang ¹⁴⁵	Yang (2006)	Taipei, Taiwan	All ages/all gender	PM ₁₀	Case-crossover/Poisson regression model	Conjunctivitis visit	Lag ₀ RR 1.02 (0.88–7.99) Lag ₁ RR 0.99 (0.86–7.46) Lag ₂ RR 0.95 (0.83–6.93) Lag ₃ RR 0.97 (0.85–7.11) Lag ₄ RR 1.11 (0.97–9.41) Lag ₅ RR 0.95 (0.84–6.86)
Lorentzou et al. ¹²²	Lorentzou et al. (2019)	Heraklion in Crete Island, Greece	All ages/all gender	PM ₁₀	Ecological retrospective analysis/one-way ANOVA and Pearson Correlation	Emergency department visits	Correlation All cases 0.313 $p=0.128$ Allergy cases 0.929 $p=0.000$ Dyspnea cases 0.464 $p=0.041$
Trianti et al. ⁵²	Trianti et al. (2017)	Athens, Greece	Aged 18 years and Upper/all gender	PM ₁₀	Ecological study/mixed Poisson model	Respiratory morbidity/emergency room visits	Respiratory visits, IR 1.95% (0.02, 3.91) Asthma visits, IR 38% ($p < 0.001$) COPD visits, IR 57% ($p < 0.001$) Respiratory infections visits, IR 60% ($p < 0.001$)
Yang et al. ¹⁵⁰	Yang et al. (2015)	Wuwei, China	All ages/all gender	PM _{2.5}	Ecological time-series/GAM	Respiratory and cardiovascular outpatient visits	Respiratory outpatient Male, RR 1.217 (1.08, 1.606) Female, RR 1.175 (1.025, 1.347) Cardiovascular outpatient Male, RR 1.146 (1.056, 1.243) Female, RR 1.105 (1.017, 1.201)
Long-term health effects							
Altindag et al. ³²	Altindag et al. (2017)	Korea	Infant	PM ₁₀	Cohort/linear regression models	Birth weight, a binary indicator of low birthweight, gestation, premature birth, and fetal growth	Birth Weight, -0.232 ($P=0.10$) Low birth weight, 0.0001 ($P=0.000$) Gestation -0.001 ($P=0.001$) Prematurity, 0.0001 ($P=0.000$) Growth, -0.005 ($P=0.003$)
Dadvand et al. ¹⁶⁹	Dadvand et al. (2011)	Barcelona/Spain	Pregnant woman	PM ₁₀	Cohort/linear regression models-logistic regression model	Pregnancy complications	Birth weight -2.1 (-5.8, 1.7) Gestation 0.5 (0.4, 0.6) Preeclampsia 0.98 (0.91, 1.07)
Li et al. ³³	Li et al. (2018)	Between northern and southern China.	Aged 10–15 years, all gender	–	Cohort/fixed-effect model	Children's cognitive function	Reduction in word scores, 0.20 (0.06, 0.35) Reduction in mathematics scores 0.18 (0.10, 0.25)
Viel et al. ³⁴	Viel et al. (2019)	Guadeloupe (French West Indies)	909 pregnant women	PM ₁₀	Cohort/multivariate logistic regression models	Preterm births	OR 1.40, (1.08–1.81)

(Continued)

Table 1. (Continued)

REFERENCE	FIRST AUTHOR AND YEAR	STUDY LOCATION	POPULATION (AGE, GENDER)	PM FRACTION	STUDY DESIGN/METHODOLOGY	HEALTH OUTCOMES	RESULTS
Tong et al. ³⁶	Tong et al. (2017)	Southwestern United States	All ages/all gender	PM ₁₀	Research letter/correlation coefficient	Valley fever	Correlation coefficient Maricopa, 0.51 Pima, 0.36–0.41
Ma et al. ⁴⁴	Ma et al. (2017)	Western China	All ages/all gender	TSP, PM ₁₀	Ecological time series/Pearson correlation coefficient	Measles incidence	The correlation coefficient for TSP Entire Lanzhou city, 0.291 Downtown Lanzhou, 0.346 The correlation coefficient for PM ₁₀ Entire Lanzhou city, 0.260 Downtown Lanzhou, 0.342 Dust events, Excess measles Zhangye, 39.1 (17.3–87.6) Lanzhou, 149.9 (7.1–413.4) Jiuquan, 31.3 (20.6–63.5)
Hospitalization or admission							
Aili and Oanh ⁹¹	Aili and Oanh (2015)	China/Taklimakan Desert	All ages/all gender	TSP	Ecological time series/GAM	Daily number of outpatients Daily number of inpatients	Respiratory outpatients, RR 1.01 (1.00–1.02) Respiratory inpatients, RR 0.99 (0.99–1.00) Digestion outpatients, RR 1.005 (0.99–1.01) Digestion inpatients, RR 1.001 (0.999–1.002) Circulatory outpatients, RR 1.010 (1.003–1.016) Circulatory inpatients, RR 1.001 (0.999–1.002) Gynecology outpatients, RR 1.008 (1.002–1.014) Gynecology inpatients, RR 0.999 (0.997–1.001) Pediatrics outpatients, RR 1.010 (1.002–1.018) Pediatrics inpatients, RR 1.001 (0.999–1.002) ENT outpatients, RR, 1.007 (1.002–1.012) ENT inpatients, RR, 1.002 (0.998–1.004)
Al et al. ⁸⁶	Al et al. (2018)	Gaziantep/Turkey	Older than 16 years	PM ₁₀	Retrospective study/GAM	Morbidity of cardiovascular diseases admitted to emergency department	Congestive cardiac failure admission, OR 1.003 (0.972–1.036) Hospitalization, OR 2.209 (2.069–2.359) Acute coronary syndrome admission, OR 1.150 (1.135–1.166) Hospitalization, OR 1.304 (1.273–1.336)
Alangari et al. ¹²⁶	Alangari et al. (2015)	Riyadh, Saudi Arabia	Children 2–12 years	PM ₁₀	Ecological/correlation coefficient	Patient presented to the emergency department (ED) with acute asthma	Acute asthma, $r = -0.14$, ($P = 0.45$) Admission rate, $r = -0.08$, ($P = 0.65$)

(Continued)

Table 1. (Continued)

REFERENCE	FIRST AUTHOR AND YEAR	STUDY LOCATION	POPULATION (AGE, GENDER)	PM FRACTION	STUDY DESIGN/METHODOLOGY	HEALTH OUTCOMES	RESULTS
Alessandrini et al. ⁹²	Alessandrini et al. (2013)	Rome, Italy	Less than 14 years or 35 years or more	PM _{2.5} , PM _{2.5-10} and PM ₁₀	Ecological time-series/GAM	Respiratory, cardiac and cerebrovascular hospitalizations	PM _{2.5} Cardiac diseases, lag ₀₋₁ -2.41 (-0.21, 5.09) Cerebrovascular diseases, lag ₀ -2.14 (-4.73, 0.53) Respiratory diseases, lag ₀₋₅ -0.52 (-5.33, 4.53) Respiratory diseases ₀₋₁₄ -2.14 (-9.09, 5.35) PM _{2.5-10} (IF) Cardiac diseases, lag ₀₋₁ 3.93 (1.58, 6.34) Cerebrovascular diseases, lag ₀ 1.68 (-0.70, 4.11) Respiratory Diseases, lag ₀₋₅ 4.77 (-0.57, 10.40) Respiratory diseases lag ₀₋₁₄ -1.20 (-8.52, 6.71) PM ₁₀ Cardiac diseases, lag ₀₋₁ 3.37 (1.11, 5.68) Cerebrovascular diseases, lag ₀ 2.64 (0.06, 5.29) Respiratory Diseases, lag ₀₋₅ 3.59 (0.18, 7.12) Respiratory diseases, lag ₀₋₁₄ -0.04 (-4.64, 4.78)
Al-Hemoud et al. ⁹³	Al-Hemoud et al. (2018)	Kuwait	All ages/all gender	PM ₁₀	Ecological time series/GAM	Daily morbidity	Bronchial asthma, $r=0.292$ Respiratory infection Lower, $r=0.737$ upper, $r=0.839$
Al-Taiar ⁵¹	Al-Taiar (2012)	Kuwait	All ages/all gender	PM ₁₀	Ecological time series/generalized/GAM	Daily emergency admissions due to asthma and respiratory causes	Asthma admission, RR 1.07 (1.02-1.12) Respiratory admission, RR 1.06 (1.04-1.08)
Barnett ¹²⁷	Barnett (2012)	Brisbane, Australia	All ages/all gender	PM ₁₀	Ecological time series/Poisson regression model	Emergency admissions to hospital	Emergency admissions 39% (5, 81%)
Bell et al. ⁵⁶	Bell et al. (2008)	Taipei, Taiwan	All ages/all gender	PM ₁₀	Ecological time-series/Poisson time-series model	Cause-specific hospital admissions	Ischemic heart disease, Lag ₂ 16.17 (1.17, 33.39)
Chan et al. ¹³⁵	Chan et al. (2018)	Nationwide/Taiwan	All ages/all gender	Total atmospheric PM	Ecological time-series/autoregressive model-ARMAX regression	Diabetes hospitalization	Diabetes lag1 27.41 ($p=0.04$)
Chen and Yang ⁹⁴	Chen and Yang (2005)	Taipei, Taiwan	All ages/all gender	PM ₁₀	Case-crossover/tests of student	Daily hospital admissions for cardiovascular disease (CVD)	CVD, lag _{RR} (3.65%) $P > 0.05$

(Continued)

Table 1. (Continued)

REFERENCE	FIRST AUTHOR AND YEAR	STUDY LOCATION	POPULATION (AGE, GENDER)	PM FRACTION	STUDY DESIGN/METHODOLOGY	HEALTH OUTCOMES	RESULTS
Cheng et al. ¹¹⁹	Cheng et al. (2008)	Taipei, Taiwan	All ages/all gender	PM ₁₀	Case-crossover/Poisson regression models	Daily pneumonia hospital admissions	Pneumonia admissions lag ₀ RR 1.03 (0.98–1.08) lag ₁ RR 1.04 (1.00–1.09) lag ₂ RR 1.04 (0.99–1.09) lag ₃ RR 1.03 (0.99–1.08)
Chiu et al. ¹²¹	Chiu et al. (2008)	Taipei, Taiwan	All ages/all gender	PM ₁₀	Case-crossover/Poisson regression models	COPD admissions	COPD, Lag ₃ , RR 1.057; (0.982–1.138)
Dong et al. ⁵⁹	Dong et al. (2007)	large cities of Korea	All ages/all gender	PM ₁₀	Ecological/correlation coefficients	Hospitalization	Seoul 0.652 Busan 0.377 Daegu 0.681 Incheon 0.736 Kwangju 0.481 Daejeon 0.652 Ulsan 0.702 Jeju-do 0.129
Ebenstein et al. ¹⁰⁷	Ebenstein et al. (2015)	Israel, Jerusalem and Tel Aviv	All ages/all gender	PM ₁₀	Ecological/IV methodology/Poisson regression approach	Respiratory hospital admissions	Respiratory admissions IR 0.8% COPD 0.01 (0.003) Asthma 0.008 (0.003) Respiratory abnormalities 0.006 (0.002)
Ebrahimi et al. ⁹²	Ebrahimi et al. (2014)	Sanandaj, Iran	All ages/all gender	PM ₁₀	Ecological/Pearson's correlation coefficient, linear regression model	Emergency admissions for cardiovascular and respiratory diseases	Cardiovascular 0.48 ($P < 0.05$) Respiratory patients 0.19 ($P > 0.05$)
Ebrahimi et al. ⁶⁴	Geravandi et al. (2017)	Ahvaz/Iran	All ages/all gender	PM ₁₀	Ecological/non-parametric Mann-Whitney U test/correlation coefficients	Hospital admissions for Respiratory diseases	Respiratory diseases ($r = 0.53$)
Grineski et al. ¹¹	Grineski et al. (2011)	El Paso, Texas, United States	All ages/all gender	PM _{2.5}	Case-crossover/conditional logistic regression	Hospital admissions for Asthma and Acute bronchitis	Asthma 1.11 (0.96–1.28) All ages 1.23 (0.99–1.55)
Kamouchi et al. ¹³¹	Kamouchi et al. (2012)	Fukuoka, Japan	20 years and older/all gender	–	Case-crossover/conditional logistic regression	Ischemic stroke	Overall////Atherothrombotic Z D7 lag ₀₋₁ , OR 1.07 0.93–1.23//1.44 1.08–1.91 lag ₀₋₂ , OR 1.04 0.97–1.18//1.48 1.14–1.93 lag ₀₋₃ , OR 1.02 0.90–1.15//1.37 1.06–1.76 lag ₀₋₄ , OR 1.02 0.90–1.14//1.35 1.06–1.73 lag ₀₋₅ , OR 1.02 0.91–1.15//1.35 1.06–1.72
Kanatani et al. ¹¹⁵	Kanatani et al. (2010)	Toyama, Japan	Children	–	Case-crossover/generalized estimating equations logistic and Conditional logistic regression	Asthma hospitalization	OR 1.88 (1.04–3.41; $P = 0.037$)

(Continued)

Table 1. (Continued)

REFERENCE	FIRST AUTHOR AND YEAR	STUDY LOCATION	POPULATION (AGE, GENDER)	PM FRACTION	STUDY DESIGN/METHODOLOGY	HEALTH OUTCOMES	RESULTS
Kang et al. ¹²⁰	Kang et al. (2012)	Taipei, Taiwan	All ages/all gender	PM ₁₀	Ecological time series/ Kruskal–Wallis test/ auto-regressive integrated moving average (ARIMA) method	Pneumonia hospitalization	Pneumonia admissions ($P=0.001$)
Kang et al. ¹³²	Kang et al. (2013)	Taiwan	All ages/all gender	PM	Ecological time series/ ARIMA method (auto-regressive integrated moving average)	Stroke hospitalization	Stroke admissions (239.6), post-DS days (249.2) ($p<0.001$)
Kashima et al. ⁴⁰	Kashima et al. (2017)	Okayama, Japan	Elderly	SPM	Case-crossover/ conditional logistic regression analyses	Susceptibility of the elderly to disease	Respiratory OR: 1.09 (1.00, 1.19) Cardiovascular OR: 0.99 (0.97, 1.01) Cerebrovascular OR: 1.15 (1.01, 1.31)
Khaniabadi et al. ⁸⁷	Khaniabadi et al. (2017)	Khorrarnabad (Iran)	All ages/all gender	PM ₁₀	Ecological time series/ AirQ model	Hospitalizations for chronic obstructive pulmonary disease (COPD)	COPD, ER, 7.3% (4.9, 19.5)
Khaniabadi et al. ⁹⁵	Khaniabadi et al. (2017)	Ilam, Iran	All ages/all gender	PM ₁₀	Ecological time series/ AirQ model	Cardiovascular and respiratory admissions	Respiratory diseases 4.7% (3.2–6.7%) Cardiovascular diseases, 4.2% (2.6–5.8%)
Ko et al. ¹³⁶	Ko et al. (2016)	Fukuok- western Japan	Men, Women ratio 30,15 Age, 49.6 ± 22.7	–	Cohort design/t-test	Acute conjunctivitis	Conjunctivitis scores $P<0.05$
Kojima et al. ⁹⁸	Kojima et al. (2017)	Kumamoto, Japan	20 years of age or older/all gender	PM _{2.5}	Case-crossover/ conditional logistic regression model	Acute myocardial infarction (AMI)	AMI OR, 1.46 (1.09–1.95) Non ST-segment OR 2.03 (1.30–3.15)
Lai and Cheng ¹⁰⁹	Lai and Cheng (2008)	Taipei, Taiwan	All ages/all gender	PM ₁₀	Case-control/Z test	Respiratory admissions	Elderly RR 3.44; (0.03–380.1) All age RR 1.04; (0.30–3.16) Pre-school RR, 1.01 (0.26–3.89)
Lee and Lee ¹⁷	Lee and Lee (2014)	Seoul, Korea	All ages/all gender	PM ₁₀	Ecological time series patterns/paired t-test	Daily asthma patients	Lag ₀ , 3.79 $p=0.4$ Lag ₁ , 4.85 $p=0.3$ Lag ₂ , 11.02 $p=0.1$ Lag ₃ , 15.46 $p=0.06$ Lag ₄ , 18.05 $p=0.03$ Lag ₅ , 17.76 $p=0.02$ Lag ₆ , 18.18 $p=0.01$
Lorentzou et al. ¹²²	Lorentzou et al. (2019)	Heraklion in Crete Island, Greece	All ages/all gender	PM ₁₀	Ecological/one-way ANOVA and Pearson correlation	COPD morbidity	COPD exacerbations, 3.0 (0.8–5.2) Dyspnea admissions, 0.71 ($p=0.001$) COPD admissions, 0.813 ($p=0.000$)

(Continued)

Table 1. (Continued)

REFERENCE	FIRST AUTHOR AND YEAR	STUDY LOCATION	POPULATION (AGE, GENDER)	PM FRACTION	STUDY DESIGN/METHODOLOGY	HEALTH OUTCOMES	RESULTS
Matsukawa et al. ⁹⁹	Matsukawa et al. (2014)	Fukuoka, Japan	Patients aged ≥20 years/all gender	SPM	Case-crossover/conditional logistic regression model	Incidence of acute myocardial infarction	AMI Lag ₄ OR 1.33 (1.05–1.69) Lag ₀₋₄ OR 1.20 (1.02–1.40)
Menendez et al. ¹²⁸	Menendez et al. (2017)	Gran Canaria, Spain	Adults (age 14–80 years) and >80/all gender	PM ₁₀	Epidemiological survey/(ANOVA) and Spearman correlation coefficients (ρ)	Health condition of the allergic population	ρ (p-values) Pneumony 0.2 (0.5) Asthma 0.8 (0.0) COPD 0.0 (1.0)
Meng and Lu ⁹⁶	Meng and Lu (2007)	Minqin, China	All ages/all gender	–	Ecological time-series/GAM	Daily hospitalization for respiratory and cardiovascular diseases	Respiratory hospitalization, lag ₃ RR Male 1.14 (1.01–1.29) Female 1.18 (1.00–1.41) Respiratory infection, Male, RR 1.28 (1.04–1.59) Pneumonia, Lag ₆ Males, RR 1.17 (1.00–1.38) Hypertension, Lag ₃ Males, RR 1.30 (1.03, 1.64)
Middleton et al. ⁹⁷	Middleton et al. (2008)	Nicosia, Cyprus	All ages/all gender	PM ₁₀	Ecological time-series/GAM	Respiratory and cardiovascular morbidity	All-cause 4.8% (0.7, 9.0) Cardiovascular 10.4% (–4.7, 27.9)
Nakamura et al. ¹⁰³	Nakamura et al. (2015)	All-Japan	All ages/all gender	SPM	Case-crossover/conditional logistic models	Out-of-hospital cardiac arrests	Cardiac arrests, lag, OR Model 1 1.00 (0.97–1.19) Model 2 1.08 (0.97–1.20)
Nastos, et al. ¹⁰⁴	Nastos, et al. (2011)	Crete Island, Greece	All ages/all gender	–	Ecological time series-HYSPLIT 4 model of air resources laboratory of NOAA	Cardiovascular and respiratory syndromes	Respiratory five-fold increased Cardiovascular didn't increased significant
Pirsaheb et al. ⁵⁰	Pirsaheb et al. (2016)	Kermanshah, Iran	All ages/all gender	PM ₁₀	Ecological/regression	Respiratory disease	Respiratory infection P ≤ 0.05 Chronic pulmonary disease P ≤ 0.05 COPD P > 0.05 Angina P > 0.05 Asthma P > 0.05
Prospero et al. ¹²⁹	Prospero et al. (2008)	Caribbean	Aged 18 years and under/all gender	–	Ecological time series/Mann-Whitney rank-sum test, two-tailed	Pediatric asthma	Pediatric asthma, P > 0.05
Radmanesh et al. ¹³³	Radmanesh et al. (2019)	Abadan, Iran	All ages/all gender	PM ₁₀	Ecological studies/Pearson coefficient	Hospital admission for cerebral ischemic attack, epilepsy and headaches	Cerebral ischemic attack, r: 0.113 p=0.3 Epilepsy, r: 0.492 p=0.03 Headaches, r: 0.009 p=0.9
Reyes et al. ¹¹⁰	Reyes et al. (2014)	Madrid (Spain)	All ages/all gender	PM _{10-2.5}	Ecological time series/conditional logistic regression model	Hospital admissions	Respiratory admissions, Lag, RR 1.031 (1.002 1.060)

(Continued)

Table 1. (Continued)

REFERENCE	FIRST AUTHOR AND YEAR	STUDY LOCATION	POPULATION (AGE, GENDER)	PM FRACTION	STUDY DESIGN/METHODOLOGY	HEALTH OUTCOMES	RESULTS
Rutherford, et al. ¹¹⁸	Rutherford, et al. (1999)	Brisbane, Australia	All ages/all gender	TSP	Cross sectional/paired two-tailed t-tests	Impact on asthma severity	Asthma severity, $P \leq 0.05$ In General $P > 0.05$
Stafoggia et al. ⁷⁶	Stafoggia et al. (2016)	Southern European cities-Spain, France, Italy, Greece	All ages/all gender	PM ₁₀	Case-crossover/Poisson regression models	Hospital admissions	Admissions, IR Cardiovascular, age ≥ 15 0.32 (-0.24, 0.89) Respiratory, age ≥ 15 0.70 (-0.45, 1.87) Respiratory, age 0–14 2.47 (0.22, 4.77)
Tam et al. ¹⁰¹	Tam et al. (2012)	Hong Kong	All ages/all gender	PM ₁₀ -2.5	Case-crossover/t-test/Poisson regression model	Daily emergency admissions for cardiovascular diseases	PM ₁₀ -2.5 Ischemic heart disease, RR 1.04 (1.00, 1.08)
Tao et al. ¹¹¹	Tao et al. (2012)	Lanzhou, China	All ages/all gender	PM ₁₀	Ecological/Poisson regression model into GAM model	Respiratory diseases admissions	Respiratory hospitalizations, RR Male, 1.148 $P > 0.05$ Female 1.144 $P > 0.05$
Teng et al. ¹⁰⁰	Teng et al. (2016)	Taipei, Taiwan	All ages/all gender	PM ₁₀	Ecological time series/autoregressive with exogenous variables model	Daily acute myocardial infarction hospital admissions	AMI hospitalizations, 3.2 more
Thalib and Al-Taiar ⁵¹	Thalib and Al-Taiar (2012)	Kuwait	All ages/all gender	PM ₁₀	Ecological time series study/GAM	Asthma admissions	Asthma, RR 1.07 (1.02–1.12) Respiratory admission, RR 1.06 (1.04–1.08)
Ueda et al. ⁵⁷	Ueda et al. (2010)	Fukuoka, Japan	children under 12 years of age/all gender	SPM	Case-crossover/conditional logistic regression	Hospitalization for asthma	Asthma hospitalization, lag _{2,3} OR 1.041 (1.013–1.070)
Vodonos et al. ¹²³	Vodonos et al. (2014)	Be'er Sheva, Israel	All ages/all gender	PM ₁₀	Ecological time series/GAM	Hospitalizations due to exacerbation of COPD	COPD exacerbation: IR 1.16 ($p < 0.001$)
Vodonos et al. ²	Vodonos et al. (2015)	Be'er Sheva, Israel	Above 18 years old/all gender	PM ₁₀	Case crossover/GAM	Cardiovascular Morbidity	Acute coronary syndrome (lag1); OR = 1.007 (1.002–1.012).
Wang et al. ¹¹³	Wang et al. (2014)	Taiwan,	All ages/all gender	PM ₁₀	Ecological time series/ARIMAX regression model	Asthma admissions	Asthma, Lag _{1,3} average of 17–20 ($p < 0.05$) more hospitalized
Wang et al. ¹²⁵	Wang et al. (2015)	Minqin County, China	Above 40 years old/all gender	–	Case-control/comparison/Student's t test	Human respiratory system	Chronic rhinitis, OR 3.14 (1.77–5.55) Chronic bronchitis, OR 2.46 (1.42–4.28) Chronic cough, OR 1.78 (1.24–2.56)
Watanabe et al. ¹¹⁴	Watanabe et al. (2014)	Western Japan	Aged ≥ 18 years old/all gender	SPM	Descriptive/telephone survey/t-test. Multiple regression analysis	Worsening asthma	Worsening asthma 11–22% Pulmonary function of asthma patients –0.367 $p = 0.003$

(Continued)

Table 1. (Continued)

REFERENCE	FIRST AUTHOR AND YEAR	STUDY LOCATION	POPULATION (AGE, GENDER)	PM FRACTION	STUDY DESIGN/METHODOLOGY	HEALTH OUTCOMES	RESULTS
Yang et al. ¹³⁴	Yang et al. (2005)	Taipei, Taiwan	All ages/all gender	PM ₁₀	Case-crossover/Poisson regression model	Stroke admissions	Hemorrhagic stroke, Lag ₃ RR 1.15 (1.01–10.10)
Yang et al. ¹⁰²	Yang et al. (2009)	Taipei, Taiwan	All ages/all gender	PM ₁₀	Case-crossover/Poisson regression model	Hospital admissions for congestive heart failure	CHF, Lag ₁ RR 1.114 (0.993–1.250)
Yang et al. ¹¹⁶	Yang et al. (2005)	Taipei, Taiwan	All ages/all gender	PM ₁₀	Case-crossover studies/Poisson regression model	Daily admissions for asthma	Asthma lag ₂ 8% ($p > 0.05$)
Al et al. ⁸⁶	Al et al. (2018)	Gaziantep, Turkey	All ages/all gender	PM ₁₀	Retrospective study/GAM	Cardiovascular diseases admitted to ED	Cardiac failure, OR Admission 1.003 (0.972–1.036) $P = 0.833$ Hospitalization 2.209 (2.069–2.359) $P = 0.001$
Gyan et al. ¹¹²	Gyan et al. (2005)	Caribbean island of Trinidad	Patients aged 15 years and under	–	Ecological/Poisson regression model	Pediatric asthma accident and emergency admissions	Admission rate increased 7.8–9.25
Bennett et al. ¹⁰⁵	Bennett et al. (2006)	Lower Fraser Valley, British Columbia, Canada	All ages/all gender	PM ₁₀	Ecological time-series/Chi-squared	Hospital admissions	hospitalizations Respiratory 0.89, $\chi^2 = 0.71$ Cardiac 0.91, $\chi^2 = 0.54$
Cheng et al. ⁶⁵	Cheng et al. (2008)	Taipei, Taiwan	All ages/all gender	PM ₁₀	Case-crossover/Poisson regression model	Daily pneumonia hospital admissions	Pneumonia admissions, RR 1.032 (0.980–1.086) Lag ₁ 1.049 (1.002–1.098) Lag ₂ 1.044 (0.999–1.092) Lag ₃ 1.037 (0.993–1.084)
Wilson et al. ¹²⁴	Wilson et al. (2012)	Hong Kong	All ages/all gender	PM ₁₀	Case-crossover/Poisson regression model	Daily emergency admissions for respiratory diseases	COPD, lag ₂ RR 1.05 (1.01–1.09)
Wiggs et al. ¹³⁰	Wiggs et al. (2003)	Karakalpakstan, Uzbekistan	Children/all gender	PM ₁₀	Ecological	Respiratory health	Decreased the rate of respiratory health problems
Pulmonary function							
Hong et al. ¹⁶²	Hong et al. (2010)	Seoul, Korea	Children/all gender	PM _{2.5} and PM ₁₀	Prospective/linear mixed-effects mode	Pulmonary function of school children	PM _{2.5} ($P > 0.05$) PM ₁₀ ($P > 0.05$)
Kurai et al. ¹⁵⁷	Kurai et al. (2017)	Yonago, Tottori, western Japan	School children/adults	PM _{2.5}	Descriptive/longitudinal/Linear mixed models	Respiratory function	Lag ₀ , -1.76 (-3.30, -0.21) Lag ₀₋₁ , -1.54 (-2.84, -0.25) Lag ₀₋₂ , -1.05 (-2.21, 0.11) Lag ₀₋₃ , -1.09 (-2.18, -0.01)

(Continued)

Table 1. (Continued)

REFERENCE	FIRST AUTHOR AND YEAR	STUDY LOCATION	POPULATION (AGE, GENDER)	PM FRACTION	STUDY DESIGN/METHODOLOGY	HEALTH OUTCOMES	RESULTS
Watanabe et al. ¹⁶¹	Watanabe et al. (2016)	western Japan	Schoolchildren	SPM	A panel study/linear mixed models	Pulmonary function	Peak expiratory flow (PEF) -3.62 (-4.66 , -2.59)
Watanabe et al. ⁶⁶	Watanabe et al. (2015)	western Japan	Schoolchildren	SPM	Longitudinal follow-up study/linear mixed models	Pulmonary function	PEF 2012 -8.17 (-11.40 , -4.93) 2013 -1.17 (-4.07 , 1.74)
Yoo et al. ¹⁶⁰	Yoo et al. (2008)	Seoul, Korea	Children	PM ₁₀	Prospective/Pearson correlation tests/paired t-test	Respiratory symptoms and peak expiratory flow	PEF decreased ($p < 0.05$)
Watanabe et al. ¹⁵⁸	Watanabe et al. (2016)	Western Japan	Aged 18 years	SPM	Panel study/linear mixed models	Pulmonary function	PEF, in allergic patients with Asthma $_{16.3}$ ($_{-32.9}$, 0.4) $P=0.06$ Rhinitis $_{7.0}$ ($_{-19.5}$, 5.5) $P=0.27$ Conjunctivitis $_{-3.9}$ ($_{-38.8}$, 30.9) $P=0.83$ Dermatitis $_{5.6}$ ($_{-21.3}$, 10.2) $P=0.49$ Food allergy 0.4 ($_{-23.3}$, 23.9) $P=0.98$
Watanabe et al. ¹⁵⁹	Watanabe et al. (2015)	Western Japan	Aged >18 years	SPM	Panel study/linear regression analysis	Pulmonary function in adult with asthma	PEF 0.01 (-0.62 , 0.11)
Park et al. ¹⁶³	Park et al. (2005)	Incheon, Korea	Ages of 16 and 75 years/ all gender	PM ₁₀	Cohort/t-test/GAM with Poisson log-linear regression	Peak expiratory flow rates and respiratory symptoms of asthmatics	PEF 1.05 (0.89 – 1.24)
O'Hara et al. ¹⁶⁶	O'Hara et al. (2001)	Karakalpakstan, Uzbekistan	Children aged 7 to 11	PM ₁₀	Cross-sectional survey/multivariate regression model	Lung function	There was an inverse relationship between dust event and Lung function
Other impacts							
Lee et al. ¹⁶⁸	Lee et al. (2019)	Korean national	All ages/all gender	PM ₁₀	Case-crossover/conditional logistic regression	Risk of suicide	Suicide risk, 13.1% (4.5 – 22.4) $P=0.002$
Soy et al. ¹⁰⁶	Soy et al. (2016)	Mardin, Turkey	All gender/18 to 65 years	PM ₁₀	Prospective study/pairs t-test	Quality of life(QoL) in patients with or without asthma	GoL, AR 2.5-fold higher SF-36, AR 1.9-fold higher
Islam et al. ¹⁶⁷	Islam et al. (2019)	Saudi Arabia	All ages/all gender	–	Ecological/panel regression models	Road traffic accidents	$P \leq 0.05$
Mu et al. ¹¹⁷	Mu et al. (2010)	Choyr City, Mongolia	44.2 ± 17.3 /all gender	–	Cross-sectional/student's t-test/multiple regression analysis	Health-related Quality of Life	Decreased HRQL $P < 0.05$

(Continued)

Table 1. (Continued)

REFERENCE	FIRST AUTHOR AND YEAR	STUDY LOCATION	POPULATION (AGE, GENDER)	PM FRACTION	STUDY DESIGN/METHODOLOGY	HEALTH OUTCOMES	RESULTS
Sing and symptom							
Higashi et al. ¹⁵¹	Higashi et al. (2014)	Japan	Aged 23–84 years all gender	PM _{2.5}	Panel study/logistic regression with a generalized estimating equation	Daily cough occurrence in patients with chronic cough	Grade 1, 1.111 (0.995, 1.239) Grade 2, 1.171 (1.006, 1.363) Grade 3, 1.357 (1.029, 1.788) Grade 4, 1.414 (0.983, 2.036)
Higashi et al. ¹⁵²	Higashi et al. (2014)	Kanazawa, Japan	Between 23 and 84	TSP	Cohort study, McNamara's test	Cough and allergic symptoms in adult with chronic cough	Cough $p=0.02$
Watanabe et al. ⁶⁷	Watanabe et al. (2012)	Japan	Age 63.4 ± 15.2/ all gender	SPM	Descriptive telephone survey/multivariate logistic regression analysis	Lower respiratory tract symptoms in asthma patients	Exacerbation 4% Unaffected 48%
Otani et al. ¹⁵³	Otani et al. (2011)	Yonago, Japan	all gender/mean age of 36.2 ± 12.5 years	SPM	Ecological Time-series/t test/Pearson's correlation coefficient	Daily symptoms	All symptoms ($p=0.020$) Skin symptom ($p<0.001$)
Onishi et al. ¹⁵⁴	Onishi et al. (2012)	Yonago, Japan	All gender/mean age-SD: 36.2–12.5 years	SPM	Prospective/Wilcoxon's rank test	Symptom nasal/ocular/respiratory/throat/skin symptoms	All symptom increased
Mu et al. ³⁵	Mu et al. (2011)	Mongolia	35–44/all gender	–	Descriptive studies/cross-sectional study/multiple logistic regression analysis	Eye and respiratory system symptoms	Itchy eye $P=0.3$ Bloodshot eye $P=0.02$ Lacrimation $P=0.001$ Respiratory system $P>0.05$
Majbaudain et al. ¹⁵⁵	Majbaudain et al. (2016)	Yonago, Japan	Mean age of 33.57 ± 1/all gender	SPM	Prospective web-based survey/student's t-test	Daily symptoms	Ocular, $r=0.47$ ($P<0.01$) Nasal, $r=0.61$ ($P<0.001$) Skin, $r=0.445$ ($P<0.05$)
Kanatani et al. ¹⁶⁴	Kanatani et al. (2016)	Kyoto, Tottori, Toyama, Japan	Pregnant women	SPM	Observational study/Cohort/conditional logistic regression analysis	Allergic symptoms	Allergic symptoms, OR 1.10 (1.04–1.18)
Yoo et al. ¹⁶⁰	Yoo et al. (2008)	Seoul, Korea	Children	PM ₁₀	Prospective/Pearson correlation tests/paired t-test	Respiratory symptoms in children with mild asthma	Cough 42.9 ± 20.8 ($p<0.05$) Runny/stuffed nose 53.8 ± 19.2 ($p<0.05$) Sore throat 24.2 ± 13.5 ($p<0.05$) Eye irritation 24.5 ± 18.1 ($p<0.05$) Limited physical activity 16.2 ± 12.5 Nocturnal awakening 15.7 ± 14.1 Shortness of breath 20.1 ± 13.8 ($p<0.05$) Wheeze 16.7 ± 7.1 ($p<0.05$)

(Continued)

Table 1. (Continued)

REFERENCE	FIRST AUTHOR AND YEAR	STUDY LOCATION	POPULATION (AGE, GENDER)	PM FRACTION	STUDY DESIGN/METHODOLOGY	HEALTH OUTCOMES	RESULTS
Watanabe et al. ¹⁵⁹	Watanabe et al. (2015)	Western Japan	Aged >18 years	SPM	Panel study/linear regression analysis	Respiratory symptoms in adult patients with asthma	All symptom 0.04 (0.03, 0.05)
Park et al. ¹⁶³	Park et al. (2005)	Incheon, Korea	Ages of 16 and 75 years/all gender	PM ₁₀	Prospective study/t-test/GAM with Poisson log-linear regression	Respiratory symptoms of asthmatics	Nighttime symptoms RR 1.05 (0.99–1.17)
O'Hara et al. ¹⁶⁶	O'Hara et al. (2001)	Karakalpakstan, Uzbekistan	Children aged 7 to 11	PM ₁₀	Descriptive studies/cross-sectional survey/multivariate regression model	Respiratory symptoms and lung function	There is an apparent inverse relationship between total dust exposure and respiratory health
Watanabe et al. ¹⁶⁵	Watanabe et al. (2011)	Western Japan	At least 18 years old	SPM	Cross-sectional telephone survey/multivariate logistic regression analysis	Worsening asthma	Aggravated lower respiratory tract symptoms in asthma patients
Meo et al. ¹⁵⁶	Meo et al. (2013)	Riyadh, Saudi Arabia	Age 28.6 ± 3.14 years/all gender	–	Descriptive studies/Chi square test	General health complaints	OR Wheeze 4.18 (2.36–7.41) Cough 4.13 (2.28–7.46) Acute asthmatic attack 6.7 (4.09–10.99) Psychological disturbances 3.72 (2.48–5.57) Eye irritation/redness 7.89 (4.4–14.16) Headache 4.17 (2.8–6.2) Body ache 1.24 (0.82–1.88) Sleep disturbance 4.16 (2.77–6.22) Runny nose 31.9 (14.33–70.96)

Abbreviations: p, Spearman correlation coefficients; AOD, aerosol optical depth; AMI, acute myocardial infarction; ACS, acute coronary syndrome; CHF, congestive heart failure; COPD, chronic obstructive pulmonary disease; GAM, generalized additive model; IHD, ischemic heart diseases; IR, increase risk; OR, odds ratio; PM, particulate matter; PM₁₀, particles less than 10 µm in aerodynamic diameter; PM_{2.5-10}, particles less than 2.5 µm in aerodynamic diameter; PM_{2.5-10}, particles with an aerodynamic diameter >2.5 µm and <10 µm; PTB, pulmonary tuberculosis; GoL, quality of life; RR, relative risk; SPM, suspended particulate matter; TSP, total suspended particulate.




Chart 1. Number of studies of the impact of dust storms on health in different years.




Figure 2. Locations of dust storms and health impact research, 1994–2019.

- Daily symptoms such as nasopharyngeal, skin, or ocular symptoms, and decreased pulmonary function (Table 1).

The current analysis indicated that the effects of dust storms on health can be divided into 2 general sections: short- and long-term effects. **Short-term effects** have been defined herein as human health problems that occurred during or immediately after a dust storm, and **long-term effects** are defined as human health problems that occurred after a long exposure to several periods of dust storms.

Short-Term Health Effects

The short-term effects included all-cause mortality, emergency dispatch or air medical retrieval service, hospitalization or admission, healthcare visits, daily symptoms, decreased pulmonary function, and other problems.

Mortality

Thirty-three articles from almost all regions discussed mortality due to dust storms by means of different health problems,

such as increased total non-accidental deaths,^{3,38,39,41,46,53-55,68-82} cardiovascular deaths,^{3,38,39,48,50,53,70,74,77,82-85} mortality due to acute coronary syndrome (ACS),^{3,70,81,86} and respiratory mortality.^{48,53,55,77,87} Some studies reported, however, that the number of cases was not increased significantly for all-causes,^{43,88} respiratory,^{38,43} cardiovascular,⁴³ or cerebrovascular mortality.⁶⁹ Neophytou et al.⁸² in Nicosia reported that associations for respiratory mortality was -0.79 ($-4.69, 3.28$) on dust storm days. Lee et al.⁵⁵ in Taipei found that dust storms have a protective effect on non-accidental deaths, respiratory deaths, and death in people >65 years of age.

Emergency dispatch or air medical retrieval service

Four articles discussed the emergency medical services required due to dust storm, focusing on different health problems. This review observed an increased relative risk of all medical emergency dispatches and a significant increase in cardiovascular dispatches,⁴² increased daily ambulance calls due to respiratory, cardiovascular, and all causes,⁸⁹ and an increase in emergency

dispatches due to cardiovascular, respiratory, injury and all causes.⁹⁰

Hospitalization or admission

Sixty-two articles from almost all regions discussed hospitalization or admission due to dust storms by means of different health problems or diseases. The results indicated that in many studies, dust storms were associated with an increased risk of hospital admission due to cardiovascular, cerebrovascular, and respiratory diseases, among others.

Cardiovascular disease (CVD) hospitalizations or admissions. In relation to cardiovascular diseases and the effect of dust storms, 17 studies stated that dust storms can increase: (1) the risk of circulatory outpatients and inpatients⁹¹; (2) odds ratio of admission and hospitalization due to congestive cardiac failure⁸⁶ and acute coronary syndrome^{2,86}; (3) effects on cardiac diseases⁹²; (4) risk of CVD hospitalization or admission^{40,78,93-97}; (5) emergency admissions for CVD⁹²; (6) the impacts on acute myocardial infarction (AMI)⁹⁸⁻¹⁰⁰; (7) emergency hospital admissions for ischemic heart diseases (IHD)¹⁰¹; (8) hospital admissions for congestive heart failure (CHF)¹⁰²; and (9) inpatient hospitalization due to cardiac failure.⁸⁶ However, some studies reported non-significant results, such as no association between dust storms and out-of-hospital cardiac arrests¹⁰³ and no significant changes in admissions concerning cardiovascular syndromes.¹⁰⁴ Also, some reported no significant association between increased dust particles and angina.⁵⁰ Bennett et al.¹⁰⁵ reported that the dust storms were not associated with an excess of CVD hospitalizations.

Respiratory disease hospitalizations or admissions. Regarding respiratory diseases related to dust storms, 35 studies stated that dust storms can increase the risk of respiratory outpatients,⁹¹ respiratory disease hospitalizations or admissions,^{11,40,43,51,57,78,92,93,96,104,106-114} cases of bronchial asthma,⁹³ asthma-related hospitalizations or admissions,^{51,57,112-116} cases of aggravated asthma disease,^{117,118} daily pneumonia admissions,^{119,120} hospital admissions for chronic obstructive pulmonary disease (COPD),^{50,87,121-123} emergency hospital admissions for COPD,¹²⁴ emergency admissions for respiratory diseases,⁹² admitted patients suffering from respiratory infection,⁵⁰ and the prevalence of chronic bronchitis, cough, and rhinitis.¹²⁵

Surprisingly, several studies did not find any link between dust storms and negative health outcomes, such as no significant effect on asthma exacerbations in Riyadh,¹²⁶ no significant change in the risk of emergency admission in dust events,¹²⁷ and no association between sandstorms and risk of hospital admission for asthma or pneumonia patients.⁵⁶ Moreover, some studies reported no statistically significant relationship between increased dust levels and pulmonary function, allergic disease, emergency admission, or drug use¹²⁸; no significant relationship between increased risk of chronic obstructive pulmonary

disease, asthma, and angina and increased concentration of dust storms^{50,129}; And no excess risk of respiratory hospitalizations.¹⁰⁵ Only two studies found a decrease in respiratory problems after dust storms, like a decreased risk of respiratory inpatients in Taklimakan Desert,⁹¹ and a lower rate of respiration problems among children in areas with higher levels of dust deposition as reported by Wiggs et al.¹³⁰

Cerebrovascular diseases hospitalizations or admissions. Regarding the correlation between cerebrovascular diseases and dust storms, 6 studies stated that dust storms can increase the risk of cerebrovascular diseases,^{40,92} the incidence of athero-thrombotic brain infarction,¹³¹ stroke admission rates,¹³² hospital admissions for epilepsy problems, cerebral ischemic attacks, and various types of headaches,¹³³ and daily intracerebral hemorrhagic (ICH) stroke admissions.¹³⁴ Bell et al.,⁵⁶ however, reported that sandstorms have no significant relationship with the risk of admission to cerebrovascular patients. Moreover, Yang et al.¹³⁴ stated that there is no significant association between the risk of ischemic stroke and dust storms.

Other diseases hospitalizations or admissions. Aili et al.⁹¹ reported that the risk of digestion outpatients and inpatients, gynecology outpatients, pediatrics outpatients and inpatients, and ENT outpatients and inpatients was increased during dust storms. Chan et al.¹³⁵ also stated that dust storms were significantly associated with diabetes admissions for females. Furthermore, Ko et al.¹³⁷ stated that dust storms can increase the risk of conjunctivitis.

Healthcare visits

Nineteen articles studied the daily number of healthcare visits due to dust storms for different health problems. Except for 1 article, all others reported that dust storms are associated with an increased daily number of healthcare visits due to asthma-related health problems¹³⁷⁻¹⁴¹ cardiac, respiratory, and stroke diagnoses,¹⁴² emergency healthcare visits for IHD, CVD, and COPD,¹⁴³ conjunctivitis clinic visits,^{144,145} children clinic visits for respiratory problems,^{139,146} healthcare visits for respiratory diseases,^{52,139,146,147} healthcare visits for all causes, circulatory, and respiratory diseases,¹⁴⁸ and for cardiovascular and respiratory problems.^{149,150} Lorentzou et al.¹²² also reported a large increase in emergency visits related to dyspnea during dust storms; however, no clinically significant increase was observed in the total number of emergency visits.

Daily symptoms

Twenty articles studied the daily symptoms resulting from dust storms. In 2 studies, Higashi et al.^{151,152} showed the effects of Kosa on cough. Otani et al.¹⁵³ found that the scores for symptoms (nasopharyngeal, ocular, respiratory, and skin) were significantly higher when related to dust storms. Onishi et al.¹⁵⁴

reported that all symptoms (nasal, ocular, respiratory, throat, and skin) increased after exposure to dust storms. Mu et al.³⁵ also reported that an increased risk of eye lacrimation occurrence is associated with dust events. Majbauddin et al.¹⁵⁵ reported a positive correlation between the increased concentration of dust storms and ocular, nasal, and skin symptoms. Similarly Meo et al.¹⁵⁶ observed that sandstorms can increase complaints of sleep and psychological disturbances as well as other problems like eye irritation, cough, wheeze, headache, and runny nose.

Pulmonary function

Nine articles discussed pulmonary function in relation to dust storms, and the evidence is conflicting. Kurai et al.,¹⁵⁷ Watanabe et al.,^{158,159} Yoo et al.¹⁶⁰ and Watanabe et al.¹⁶¹ all found that dust storms have a significant, negative effect on pulmonary function. Other studies, including Hong et al.,¹⁶² Watanabe et al.¹⁵⁹ and Park et al.¹⁶³ found no significant relationship between pulmonary function and dust storms. Kanatani et al. found that dust storms can increase the risk of allergic symptoms in pregnant women.¹⁶⁴ Yoo et al.,¹⁶⁰ reported a significant increase in respiratory symptoms during dust storms, and Watanabe et al.¹⁵⁹ reported that sand and dust storms are significantly associated with respiratory symptoms. Moreover, Park et al.¹⁶³ reported a relationship between nighttime symptoms and particulate matter levels during dust storms. Watanabe et al.¹⁶⁵ also stated that dust storms worsen respiratory symptoms in asthmatic patients, but some studies like O'Hara et al.¹⁶⁶ reported that pulmonary function was better in children who were more exposed to dust storms than in those with low exposure to dust.

Other impacts

Some articles explored the relationship of dust storms with road traffic accidents, risk of suicide, placental abruption, and health-related quality of life. Islam et al.¹⁶⁷ found that sandstorms and the number of vehicles were significantly responsible for road traffic accidents. Soy et al.¹⁰⁶ reported that dust storms can have adverse effects on the quality of life of patients with asthma and allergies. Mu et al.¹¹⁷ reported that dust storms can decrease health-related quality of life in everyone exposed to them. Lee et al.¹⁶⁸ reported that exposure to dust storms was associated with an increased risk of suicide (13.1%; $p = 0.002$).

Long-Term Health Effects

Six articles discussed the long-term adverse health effects caused by dust storms by means of different outcomes, like reduced birth weight, baby's birth weight <2.5 Kg, gestation/gestational age >37 weeks and premature birth,³² and decreased cognitive function in children.³³ Preterm births³⁴ were correlated with Valley fever incidences³⁶ and increased spring

measles incidence.⁴⁴ Only one article was observed to indicate no significant effect of desert dust storms on pregnancy consequences.¹⁶⁹

Discussion

In this study, the majority of valid scientific databases were searched to find articles and studies related to the health effects of dust storms. Other similar studies have used fewer scientific databases in their search. The final number of articles included in this study is higher than that in all previous studies.^{24,26} The current results showed that the model most used to evaluate the health effects of dust storms was the GAM method. In this regards, Ramsay 2003 stated, "*Such methods eliminate the need to specify a parametric form for secular trends and allow a greater degree of robustness against model misspecification.*"¹⁷⁰ The results of the current study also showed that most dust storm studies have been carried out in Japan, Taiwan, and South Korea, which may be due to the large number of dust storms occurring in Northeast Asia. This area is exposed to yellow dust storms caused by strong winds on the Loos Plateau and the Gobi and Talkmanistan Deserts, and as yellow dust storms became so prevalent in that area within the last two decades, researchers in the area have studied their health effects.^{152,171}

The review results showed that most studies around the world confirm the adverse effects of dust storms on health. The relevant health problems were categorized into long-term and short-term impacts. Few studies were found that focused on the long-term impacts of dust storms on human and public health; however, those studies found showed that dust storms may increase the risks for problems in pregnancy and childbirth, children's cognitive problems, and infectious diseases. In line with the risks of birth as well as cognitive problems in children, animal studies have shown that the fetal brain is easily exposed to air pollutants, because in the human fetus, the blood-brain barrier has not yet developed; therefore, the fetal brain is exposed to pollutants and is sensitive to blood changes caused by them.¹⁻³ Furthermore, new research on humans has shown that environmental pollutants can possibly create inflammation, oxidative stress, and vascular damage to the fetal brain after passing through the placenta.⁴⁻⁷ Researchers have studied the effects of PM from dust storms on maternal health during pregnancy and birth problems, and they refer to variations in maternal host-defense mechanisms, maternal-placental exchanges, oxidative pathways, and endocrine dysfunction as possible causes of these problems.⁸ Ultimately, the evidence from infectious diseases shows that pathogenic microorganisms are abundant in dust storms,⁹ and dust storms can spread these microorganisms over a large area. Therefore, it can be argued that microorganisms that are suspended or attached to dust particles can be transferred from one part to another and may induce infectious diseases at various destinations by dust storms.^{10,11} More studies have been conducted on the short-term impacts of dust storms. The majority of these studies indicate the effects of dust storms on important body systems,

including the cardiovascular, respiratory and cerebral systems, which lead to the increased incidence of clinical symptoms and severity of symptoms; increased emergency visits, ambulance dispatches, and hospitalizations or admissions; decreased lung capacity; and eventually death.

Most studies show that dust storms increase the risk of cardiovascular problems, the number of cardiovascular emergency medical dispatches, cardiovascular visits, the number of cardiovascular symptoms among patients referring to the hospital, cardiovascular admissions or hospitalizations, and deaths due to cardiovascular disease. Although the exact mechanism for the effects of dust storms on heart problems has not yet been determined,¹² studies show that fine particles in dust storms can enter lung tissue and the bloodstream through chemical interactions,¹³ causing a thrombolytic and inflammatory process and the secretion of cytokines in the body.^{14,15} Moreover, the toxicity of some of these substances in the body reduces the contractibility of the heart, increases vasoconstriction, and increases blood pressure.^{14,18-20} Therefore, the above cases may confirm the effects of dust storms on cardiovascular health.

The results of the current study showed that according to most articles, the risk of death following respiratory problems; the risk of admission and hospitalization due to respiratory disorders like pneumonia, asthma, and chronic obstructive pulmonary disease and other respiratory problems; respiratory symptoms; and healthcare visits associated with dust storms have increased. Other results showed that dust storms reduce lung capacity and function.

The results of studies have shown that 1 mechanism of dust storms is that small particulates in dust storms are likely to trigger an innate immune response by T-lymphocytes in the body and respiratory system, which can cause chronic inflammation and advanced COPD.²²⁻²⁵ PM can also play a significant role in respiratory oxidative stress, increase pulmonary inflammation, increase atopic responses and Immunoglobulin E production in respiratory problems (especially asthma), and exacerbate symptoms.²⁶ Another mechanism that may cause respiratory illnesses following a dust storm is the presence of pathogens such as microorganisms and fungi³⁷ as well as some minerals such as silica in some of these storms. These particles enter the airway after dust storms and exacerbate the disease or cause respiratory problems in people at risk.²² For example, neutrophilic pulmonary inflammation may be caused by bacterial and fungal debris in dust particles to which individuals are exposed. Some of this debris includes lipopolysaccharide (LPS), a cell wall glycolipid of gram-negative bacteria, and β -glucan, which is the most important constituent of the fungal wall. Both of them are clearly observed in dust storms along with dust particles.^{22,38,39} Although the precise mechanisms for pneumonia are yet to be found, some studies have suggested that high amounts of particles in dust storms can cause oxidative stress, induce inflammation, increase blood clotting, disrupt defense cells, and cause immune system fluctuations,

ultimately inducing alveolar inflammation and exacerbating lung disease.^{3,40,41}

In 2009, Calderon Garosia stated that pollutants in dust storms can cause problems such as cardiovascular, respiratory, liver, and skin toxicity through systemic inflammation⁴² and may induce a pre-inflammatory systemic response in cytokines, which may disrupt the HPA axis and ultimately cause mood swings and psychological problems, including suicidal thoughts.⁴²⁻⁴⁴ In addition, chemical components found in dust storms can enter the brain through the mucosa and olfactory system.⁴² After entering the nervous system, they may accumulate in the anterior cortex of the brain and cause problems in emotional regulation and impulse control.⁴⁵ Some researchers also suggest that some mechanisms are associated with placental abruption due to dust storms, such as microbiological and chemical substances in dust storms that induce an inflammatory response in the body.^{46,47} Inflammation and ischemia increase the risk of decidual bleeding, followed by hematoma formation and placental abruption.^{48,49} There is also some speculation that as lipopolysaccharide has been found in Asian dust storms, the activity of this endotoxin in the body may lead to premature birth due to chorioamnionitis, which is also associated with placental abruption.^{50,51}

The current review shows that some studies have also linked dust storms with some other health problems, such as increased road accidents, increased suicide risk, increased premature placental abruption, ocular problems, and reduced quality of life. These issues could be further studied in areas prone to dust storms. Islam¹¹ stated that the reduced field of vision, the lack of dust storm warning systems, and traffic due to dust and sand storms can be considered as reasons for the recent increase in number of road accidents. Dust particulates in these storms can also cause acute ocular problems such as tears and conjunctivitis in people due to their inflammatory effects.⁵² In terms of the quality of life, Mu⁵³ stated that an increase in health problems and clinical symptoms that are associated with allergens and ocular problems such as conjunctivitis dust storms reduce the quality of life.

Despite all the significant effects of dust storms on health, this review found some studies that presented no significant association between dust storms and health problems including all-cause and respiratory mortality,^{43,88} cardiovascular,¹⁰³⁻¹⁰⁵ cerebral,¹³⁴ and respiratory problems.¹²⁷⁻¹²⁹ Moreover, some studies reported that dust storms may have a protective effect against non-accidental and respiratory death⁵⁵ and other pulmonary problems.^{91,130,166}

However, O'Hara stated that although the lack of matching of exposed and non-exposed groups in nutritional, economic, and social problems may play a role in the insignificance of the effects of dust storms on health, the chemical and physical nature of the particles in dust storms are of more importance than their total amounts.^{55,166} Differences in the chemical and physical nature of particulate matters may cause different health outcomes in varying regions.⁵⁵ Another reason for the

difference may be the use of rapid early warning systems in some countries. Lee justified the protective effects of dust storms on death, stating that in Taipei, a complex rapid early warning system is used for dust storms, and the use of this system may produce protective effects of dust storms on mortality.⁵⁵ Finally, almost all of the reviewed articles reported on a group of diseases or deaths that were studied, while dust storms may not affect all diseases and deaths.²² This may be another reason for these differing results.

Conclusion

This systematic review presents an accurate and comprehensive study of all aspects of human health in relation to dust storms. For the first time in the world, this in-depth and unique study was conducted to summarize the short-term and long-term effects of dust storms. To date, this amount of reliable data on this issue has never been investigated. As the results showed, despite the short-term effects dust storms have on human health (including adverse effects on the respiratory, skin, ocular, cardiovascular, and cerebral systems as well as increased mortality and morbidity) that may occur immediately after each dust storm, the frequency of dust storms in an area is also an important factor. In addition to exacerbating short-term health effects, they may also cause long-term health effects, which may include health problems for pregnant mothers, fetuses and infants, in the cognitive function of children, and increases in some infectious diseases. Therefore, as climate change and drought have caused this phenomenon to endanger the lives of many people around the world, and as the health and well-being of people is a main priority in any country, it is recommended that more studies be conducted in countries exposed to dust storms to examine the health effects of these storms in order to better understand the mechanisms through which dust storms impact human and public health and to develop a better strategy for preparing for, preventing, and mitigating the destructive effects of these storms.

Acknowledgements

Many thanks to Institute for School of public health and Environmental Research (IER) of Tehran University of Medical Sciences (TUMS), for supported current study.

Author Contributions

“HA and AOT designed the study; HA collected the data; HA and AOT analyzed and interpreted the data. HA, AOT, A Ardalan, MA, MY, CS and A Asgary prepared the manuscript. All authors contributed to the drafting and final review of the manuscript. The author (s) read and approved the final manuscript.”

Ethical Approval

Current study was approved by the Ethics Committee of Tehran University of Medical Sciences (TUMS) Ethics Code: IR.TUMS.SPH.REC.1399.004, and also all methods were

performed in accordance with the relevant guidelines and regulations.

ORCID iD

Hamidreza Aghababaeian  <https://orcid.org/0000-0003-3339-5507>

REFERENCES

- Shao Y, Wyrwoll K-H, Chappell A, et al. Dust cycle: an emerging core theme in Earth system science. *Aeolian Res.* 2011;2(4):181-204.
- Vodonos A, Friger M, Katra I, et al. Individual effect modifiers of dust exposure effect on cardiovascular morbidity. *PLoS One.* 2015;10(9):e0137714.
- Crooks JL, Cascio WE, Percy MS, Reyes J, Neas LM, Hilborn ED. The association between dust storms and daily non-accidental mortality in the United States, 1993–2005. *Environ Health Perspect.* 2016;124(11):1735-1743.
- Almeida-Silva M, Almeida SM, Freitas M, Pio C, Nunes T, Cardoso J. Impact of Sahara dust transport on Cape Verde atmospheric element particles. *J Toxicol Environ Health Part A.* 2013;76(4-5):240-251.
- Ardon-Dryer K, Mock C, Reyes J, Lahav G. The effect of dust storm particles on single human lung cancer cells. *Environ Res.* 2020;181:108891.
- Middleton N, Tozer P, Tozer B. Sand and dust storms: underrated natural hazards. *Disasters.* 2019;43(2):390-409.
- Krueger BJ, Grassian VH, Cowin JP, Laskin A. Heterogeneous chemistry of individual mineral dust particles from different dust source regions: the importance of particle mineralogy. *Atmos Environ.* 2004;38(36):6253-6261.
- Goudie AS, Middleton NJ. *Desert Dust in the Global System.* Springer Science & Business Media; 2006.
- Schweitzer MD, Calzadilla AS, Salamo O, et al. Lung health in era of climate change and dust storms. *Environ Res.* 2018;163:36-42.
- Fairlie TD, Jacob DJ, Park RJ. The impact of transpacific transport of mineral dust in the United States. *Atmos Environ.* 2007;41(6):1251-1266.
- Grineski SE, Staniswalis JG, Bulathsinhala P, Peng Y, Gill TE. Hospital admissions for asthma and acute bronchitis in El Paso, Texas: do age, sex, and insurance status modify the effects of dust and low wind events? *Environ Res.* 2011;111(8):1148-1155.
- Johnston F, Hanigan I, Henderson S, Morgan G, Bowman D. Extreme air pollution events from bushfires and dust storms and their association with mortality in Sydney, Australia 1994–2007. *Environ Res.* 2011;111(6):811-816.
- WHO. Air quality guidelines for particulate matter, ozone, nitrogen dioxide and sulphur dioxide. Global update 2005. Summary of Risk Assessment. 2006.
- Song Z, Wang J, Wang S. Quantitative classification of northeast Asian dust events. *J Geophys Res Atmos.* 2007;112(D4):D007048.
- Hoffmann C, Funk R, Wieland R, Li Y, Sommer M. Effects of grazing and topography on dust flux and deposition in the Xilingele grassland, Inner Mongolia. *J Arid Environ.* 2008;72(5):792-807.
- Hamidi M, Kavianpour MR, Shao Y. Synoptic analysis of dust storms in the Middle East. *Asia-Pac J Atmos Sci.* 2013;49(3):279-286.
- Shepherd G, Terradellas E, Baklanov A, et al. Global assessment of sand and dust storms; 2016. <https://repositorio.aemet.es/handle/20.500.11765/4495>
- UNCCD. Sand and dust storms: United Nations Convention to Combat Desertification; 2020. <https://www.unccd.int/actions/sand-and-dust-storms>
- Modarres R. Regional maximum wind speed frequency analysis for the arid and semi-arid regions of Iran. *J Arid Environ.* 2008;72(7):1329-1342.
- Akhlaq M, Sheltami TR, Mouftah HT. A review of techniques and technologies for sand and dust storm detection. *Rev Environ Sci Bio/Technol.* 2012;11(3):305-322.
- Hahnenberger M, Nicoll K. Meteorological characteristics of dust storm events in the eastern Great Basin of Utah, USA. *Atmos Environ.* 2012;60:601-612.
- Duce RA. Sources, distributions, and fluxes of mineral aerosols and their relationship to climate. *Aerosol forcing of clim.* 1995;6:43-72.
- Jiang D, Liu Z, Cao C, Kou Z, Wang R. *Desertification and Ecological Restoration of Keerqin Sandy Land.* China Environmental Science Press; 2003.
- Goudie AS. Desert dust and human health disorders. *Environ Int.* 2014;63:101-113.
- Liu R-T, Huza MA. Filtration and indoor air quality: a practical approach. *ASHRAE J.* 1995;37(2):24-31.
- Zhang X, Zhao L, Tong DQ, Wu G, Dan M, Teng B. A systematic review of global desert dust and associated human health effects. *Atmosphere.* 2016;7(12):158.

27. Hsieh N-H, Liao C-M. Assessing exposure risk for dust storm events-associated lung function decrement in asthmatics and implications for control. *Atmos Environ*. 2013;68:256-264.
28. Jaafari J, Naddafi K, Yunesian M, et al. The acute effects of short term exposure to particulate matter from natural and anthropogenic sources on inflammation and coagulation markers in healthy young adults. *Sci Total Environ*. 2020;735:139417.
29. Jaafari J, Naddafi K, Yunesian M, et al. Associations between short term exposure to ambient particulate matter from dust storm and anthropogenic sources and inflammatory biomarkers in healthy young adults. *Sci Total Environ*. 2020;761:144503.
30. Agier L, Deroubaix A, Martiny N, Yaka P, Djibo A, Broutin H. Seasonality of meningitis in Africa and climate forcing: aerosols stand out. *JR Soc Interface*. 2013;10(79):20120814.
31. Panikkath R, Jumper CA, Mulkey Z. Multilobar lung infiltrates after exposure to dust storm: the Haboob Lung Syndrome. *Am J Med*. 2013;126(2):e5-e7.
32. Altindag DT, Baek D, Mocan N. Chinese yellow dust and Korean infant health. *Soc Sci Med*. 2017;186:78-86.
33. Li Z, Chen L, Li M, Cohen J. Prenatal exposure to sand and dust storms and children's cognitive function in China: a quasi-experimental study. *Lancet Planet Health*. 2018;2(5):e214-e222.
34. Viel JF, Mallet Y, Raghoulmandan C, et al. Impact of Saharan dust episodes on preterm births in Guadeloupe (French West Indies). *Occup Environ Med*. 2019;76(5):336-340.
35. Mu H, Battsetseg B, Ito TY, Otani S, Onishi K, Kurozawa Y. Health effects of dust storms: subjective eye and respiratory system symptoms in inhabitants in Mongolia. *J Environ Health*. 2011;73(8):18-20.
36. Tong DQ, Wang JXL, Gill TE, Lei H, Wang B. Intensified dust storm activity and valley fever infection in the southwestern United States. *Geophys Res Lett*. 2017;44(9):4304-4312.
37. Yang Z, Meng Z, Pan J, Yan X. Epidemiological survey on the diseases of the respiratory and circulatory system among the people in a region where dust events occurred frequently. *J Environ Occup Med*. 2008;1(4):96.
38. Chan CC, Ng HC. A case-crossover analysis of Asian dust storms and mortality in the downwind areas using 14-year data in Taipei. *Sci Total Environ*. 2011;410-411:47-52.
39. Chen YS, Sheen PC, Chen ER, Liu YK, Wu TN, Yang CY. Effects of Asian dust storm events on daily mortality in Taipei, Taiwan. *Environ Res*. 2004;95(2):151-155.
40. Kashima S, Yorifuji T, Suzuki E. Are people with a history of disease more susceptible to a short-term exposure to Asian dust? A case-crossover study among the elderly in Japan. *Epidemiology*. 2017;28(Suppl 1):S60-S66.
41. Achilleos S, Al-Ozairi E, Alahmad B, et al. Acute effects of air pollution on mortality: a 17-year analysis in Kuwait. *Environ Int*. 2019;126:476-483.
42. Aghababaeian H, Dastoorpoor M, Ghasemi A, Kiarsi M, Khanjani N, Ahvazi LA. Cardiovascular and respiratory emergency dispatch due to short-term exposure to ambient PM10 in Dezful, Iran. *J Cardiovasc Thorac Res*. 2019;11(4):264-271.
43. Al-Taiar A, Thalib L. Short-term effect of dust storms on the risk of mortality due to respiratory, cardiovascular and all-causes in Kuwait. *Int J Biometeorol*. 2014;58(1):69-77.
44. Ma Y, Zhou J, Yang S, Zhao Y, Zheng X. Assessment for the impact of dust events on measles incidence in western China. *Atmos Environ*. 2017;157:1-9.
45. Giannadaki D, Pozzer A, Lelieveld J. Modeled global effects of airborne desert dust on air quality and premature mortality. *Atmos Chem Phys*. 2014;14(2):957.
46. Perez L, Tobias A, Querol X, et al. Coarse particles from Saharan dust and daily mortality. *Epidemiology*. 2008;19(6):800-807.
47. Chen Y-S, Yang C-Y. Effects of Asian dust storm events on daily hospital admissions for cardiovascular disease in Taipei, Taiwan. *J Toxicol Environ Health Part A*. 2005;68(17-18):1457-1464.
48. Kashima S, Yorifuji T, Tsuda T, Eboshida A. Asian dust and daily all-cause or cause-specific mortality in western Japan. *Occup Environ Med*. 2012;69(12):908-915.
49. Delangizan S, Jafari Motlagh Z. Dust phenomenon affects on cardiovascular and respiratory hospitalizations and mortality a case study in Kermanshah, during march-september 2010-2011. *Iran J Health Environ*. 2013;6(1):65-76.
50. Ghozikali MG, Ansarin K, Naddafi K, et al. Short-term effects of particle size fractions on lung function of late adolescents. *Environ Sci Pollut Res*. 2018;25(22):21822-21832.
51. Thalib L, Al-Taiar A. Dust storms and the risk of asthma admissions to hospitals in Kuwait. *Sci Total Environ*. 2012;433:347-351.
52. Trianti SM, Samoli E, Rodopoulou S, Katsouyanni K, Papiris SA, Karakatsani A. Desert dust outbreaks and respiratory morbidity in Athens, Greece. *Environ Health*. 2017;16(1):72.
53. Sajani SZ, Miglio R, Bonasoni P, et al. Saharan dust and daily mortality in Emilia-Romagna (Italy). *Occup Environ Med*. 2011;68(6):446-451.
54. Samoli E, Kougea E, Kassomenos P, Analitis A, Katsouyanni K. Does the presence of desert dust modify the effect of PM10 on mortality in Athens, Greece? *Sci Total Environ*. 2011;409(11):2049-2054.
55. Lee H, Honda Y, Lim YH, Guo YL, Hashizume M, Kim H. Effect of Asian dust storms on mortality in three Asian cities. *Atmos Environ*. 2014;89:309-317.
56. Bell ML, Levy JK, Lin Z. The effect of sandstorms and air pollution on cause-specific hospital admissions in Taipei, Taiwan. *Occup Environ Med*. 2008;65(2):104-111.
57. Ueda K, Nitta H, Odajima H. The effects of weather, air pollutants, and Asian dust on hospitalization for asthma in Fukuoka. *Environ Health Prev Med*. 2010;15(6):350-357.
58. Min PK, Kim CW, Yun YJ, et al. Effect of yellow sand on respiratory symptoms and diurnal variation of peak expiratory flow in patients with bronchial asthma. *J Asthma Allergy Clin Immunol*. 2001;21(6):1179-1186.
59. Dong T, Duong T, Lee BK, Huh YS, Cho SW, editors. An analysis on the effects of Asian dust event on human health in large cities of Korea based on the number of patients. 2007 International Forum on Strategic Technology, IFOST; 2007.
60. Goto K, Nmor JC, Kurahashi R, et al. Relationship between influx of yellow dust and bronchial asthma mortality using satellite data. *Sci Res Essays*. 2010;5(24):4044-4052.
61. Ueda K, Shimizu A, Nitta H, Inoue K. Long-range transported Asian dust and emergency ambulance dispatches. *Inhal Toxicol*. 2012;24(12):858-867.
62. Park J, Lim MN, Hong Y, Kim WJ. The influence of Asian dust, haze, mist, and fog on hospital visits for airway diseases. *Tuberc Res Dis*. 2015;78(4):326-335.
63. Wang Y, Wang R, Ming J, et al. Effects of dust storm events on weekly clinic visits related to pulmonary tuberculosis disease in Minqin, China. *Atmos Environ*. 2016;127:205-212.
64. Yu HL, Chien LC, Yang CH. Asian dust storm elevates children's respiratory health risks: a spatiotemporal analysis of children's clinic visits across Taipei (Taiwan). *PLoS One*. 2012;7(7):e41317.
65. Watanabe M, Noma H, Kurai J, et al. Decreased pulmonary function in school children in Western Japan after exposures to Asian desert dusts and its association with interleukin-8. *Biomed Res Int*. 2015;2015:583293.
66. Watanabe M, Kurai J, Igishi T, et al. Influence of Asian desert dust on lower respiratory tract symptoms in patients with asthma over 4 years. *Yonago Acta Med*. 2012;55(2):41-48.
67. Ramsay TO, Burnett RT, Krewski D. The effect of concurrency in generalized additive models linking mortality to ambient particulate matter. *Epidemiology*. 2003;14(1):18-23.
68. Diaz J, Linares C, Carmona R, et al. Saharan dust intrusions in Spain: health impacts and associated synoptic conditions. *Environ Res*. 2017;156:455-467.
69. Diaz J, Tobias A, Linares C. Saharan dust and association between particulate matter and case-specific mortality: a case-crossover analysis in Madrid (Spain). *Environ Health*. 2012;11:11.
70. Hwang SS, Cho SH, Kwon HJ. Effects of the Asian dust events on mortality in the spring of 2002, Seoul, Korea. *Epidemiology*. 2004;15(4):S126.
71. Jimenez E, Linares C, Martinez D, Diaz J. Role of Saharan dust in the relationship between particulate matter and short-term daily mortality among the elderly in Madrid (Spain). *Sci Total Environ*. 2010;408(23):5729-5736.
72. Kashima S, Yorifuji T, Bae S, Honda Y, Lim YH, Hong YC. Asian dust effect on cause-specific mortality in five cities across South Korea and Japan. *Atmos Environ*. 2016;128:20-27.
73. Kim HS, Kim DS, Kim H, Yi SM. Relationship between mortality and fine particles during Asian dust, smog-Asian dust, and smog days in Korea. *Int J Environ Health Res*. 2012;22(6):518-530.
74. Lee H, Kim H, Honda Y, Lim YH, Yi S. Effect of Asian dust storms on daily mortality in seven metropolitan cities of Korea. *Atmos Environ*. 2013;79:510-517.
75. Lee JT, Son JY, Cho YS. A comparison of mortality related to urban air particles between periods with Asian dust days and without Asian dust days in Seoul, Korea, 2000-2004. *Environ Res*. 2007;105(3):409-413.
76. Renzi M, Forastiere F, Calzolari R, et al. Short-term effects of desert and non-desert PM10 on mortality in Sicily, Italy. *Environ Int*. 2018;120:472-479.
77. Stafoggia M, Zauli-Sajani S, Pey J, et al. Desert dust outbreaks in southern Europe: Contribution to daily PM10(0) concentrations and short-term associations with mortality and hospital admissions. *Environ Health Perspect*. 2016;124(4):413-419.
78. Shahsavani A, Tobias A, Querol X, et al. Short-term effects of particulate matter during desert and non-desert dust days on mortality in Iran. *Environ Int*. 2019;134:105299.
79. Tobias A, Perez L, Diaz J, et al. Short-term effects of particulate matter on total mortality during Saharan dust outbreaks: a case-crossover analysis in Madrid (Spain). *Sci Total Environ*. 2011;412-413:386-389.
80. Wang YC, Lin YK. Mortality associated with particulate concentration and Asian dust storms in Metropolitan Taipei. *Atmos Environ*. 2015;117:32-40.

81. Neophytou AM, Yiallourou P, Coull BA, et al. Particulate matter concentrations during desert dust outbreaks and daily mortality in Nicosia, Cyprus. *J Expo Sci Environ Epidemiol*. 2013;23(3):275-280.
82. Kwon HJ, Cho SH, Chun Y, Lagarde F, Pershagen G. Effects of the Asian dust events on daily mortality in Seoul, Korea. *Environ Res*. 2002;90(1):1-5.
83. Mallone S, Stafoggia M, Faustini A, Gobbi GP, Marconi A, Forastiere F. Saharan dust and associations between particulate matter and daily mortality in Rome, Italy. *Environ Health Perspect*. 2011;119(10):1409-1414.
84. Perez L, Tobías A, Querol X, et al. Saharan dust, particulate matter and cause-specific mortality: a case-crossover study in Barcelona (Spain). *Environ Int*. 2012;48:150-155.
85. Pirsahab M, Bakhshi S, Almasi A, et al. Evaluating the effect of dust phenomenon on respiratory disease and death from cardiovascular and respiratory disease in Kermanshah (2008-2013). *Int J Pharm Technol* 2016;8(2):13908-13918.
86. Al B, Bogan M, Zengin S, et al. Effects of dust storms and climatological factors on mortality and morbidity of cardiovascular diseases admitted to ED. *Emerg Med Int*. 2018;2018:3758506.
87. Khaniabadi YO, Daryanoosh SM, Amrane A, et al. Impact of middle eastern dust storms on human health. *Atmos Pollut Res*. 2017;8(4):606-613.
88. Schwartz J, Norris G, Larson T, Sheppard L, Claiborn C, Koenig J. Episodes of high coarse particle concentrations are not associated with increased mortality. *Environ Health Perspect*. 1999;107(5):339-342.
89. Kashima S, Yorifuji T, Suzuki E. Asian dust and daily emergency ambulance calls among elderly people in Japan. *J Occup Environ Med*. 2014;56(12):1277-1283.
90. Holyoak AL, Aitken PJ, Elcock MS. Australian dust storm: impact on a state-wide air medical retrieval service. *Air Med J*. 2011;30(6):322-327.
91. Aili A, Oanh NTK. Effects of dust storm on public health in desert fringe area: case study of northeast edge of Taklimakan Desert, China. *Atmos Pollut Res*. 2015;6(5):805-814.
92. Alessandrini ER, Stafoggia M, Faustini A, Gobbi GP, Forastiere F. Saharan dust and the association between particulate matter and daily hospitalisations in Rome, Italy. *Occup Environ Med*. 2013;70(6):432-434.
93. Al-Hemoud A, Al-Dousari A, Al-Shatt A, Al-Khayat A, Behbehani W, Malak M. Health impact assessment associated with exposure to PM10 and dust storms in Kuwait. *Atmosphere*. 2018;9(1):6.
94. Khaniabadi YO, Fanelli R, De Marco A, et al. Hospital admissions in Iran for cardiovascular and respiratory diseases attributed to the Middle Eastern Dust storms. *Environ Sci Pollut Res Int*. 2017;24(20):16860-16868.
95. Meng Z, Lu B. Dust events as a risk factor for daily hospitalization for respiratory and cardiovascular diseases in Minqin, China. *Atmos Environ*. 2007;41(33):7048-7058.
96. Middleton N, Yiallourou P, Kleanthous S, et al. A 10-year time-series analysis of respiratory and cardiovascular morbidity in Nicosia, Cyprus: the effect of short-term changes in air pollution and dust storms. *Environ Health*. 2008;7:39.
97. Ebrahimi SJ, Ebrahimzadeh L, Eslami A, Bidarpoor F. Effects of dust storm events on emergency admissions for cardiovascular and respiratory diseases in Sanandaj, Iran. *J Environ Health Sci Eng*. 2014;12:110.
98. Kojima S, Michikawa T, Ueda K, et al. Asian dust exposure triggers acute myocardial infarction. *Eur Heart J*. 2017;38(43):3202-3208.
99. Matsukawa R, Michikawa T, Ueda K, et al. Desert dust is a risk factor for the incidence of acute myocardial infarction in Western Japan. *Circ Cardiovasc Qual Outcomes*. 2014;7(5):743-748.
100. Teng JCY, Chan YS, Peng YI, Liu TC. Influence of Asian dust storms on daily acute myocardial infarction hospital admissions. *Public Health Nurs*. 2016;33(2):118-128.
101. Tam WWS, Wong TW, Wong AHS. Effect of dust storm events on daily emergency admissions for cardiovascular diseases. *Circ J*. 2012;76(3):655-660.
102. Yang CY, Cheng MH, Chen CC. Effects of Asian dust storm events on hospital admissions for congestive heart failure in Taipei, Taiwan. *J Toxicol Environ Health A*. 2009;72(5):324-328.
103. Nakamura T, Hashizume M, Ueda K, et al. The relationship between Asian dust events and out-of-hospital cardiac arrests in Japan. *J Epidemiol*. 2015;25(4):289-296.
104. Nastos PT, Kampanis NA, Giaouzaki KN, Matzarakis A. Environmental impacts on human health during a Saharan dust episode at Crete Island, Greece. *Meteorol Z*. 2011;20(5):517-529.
105. Bennett CM, McKendry I, Kelly S, Denike K, Koch T. Impact of the 1998 Gobi dust event on hospital admissions in the Lower Fraser Valley, British Columbia. *Sci Total Environ*. 2006;366(2-3):918-925.
106. Ebenstein A, Frank E, Reingewertz Y. Particulate matter concentrations, sandstorms and respiratory hospital admissions in Israel. *Isr Med Assoc J*. 2015;17(10):628-632.
107. Geravandi S, Sicard P, Khaniabadi YO, et al. A comparative study of hospital admissions for respiratory diseases during normal and dusty days in Iran. *Environ Sci Pollut Res*. 2017;24(22):18152-18159.
108. Lai LW, Cheng WL. The impact of air quality on respiratory admissions during Asian dust storm periods. *Int J Environ Health Res*. 2008;18(6):429-450.
109. Reyes M, Diaz J, Tobias A, Montero JC, Linares C. Impact of Saharan dust particles on hospital admissions in Madrid (Spain). *Int J Environ Health Res*. 2014;24(1):63-72.
110. Tao Y, An XQ, Sun ZB, Hou Q, Wang Y. Association between dust weather and number of admissions for patients with respiratory diseases in spring in Lanzhou. *Sci Total Environ*. 2012;423:8-11.
111. Gyan K, Henry W, Lacaille S, et al. African dust clouds are associated with increased paediatric asthma accident and emergency admissions on the Caribbean island of Trinidad. *Int J Biometeorol*. 2005;49(6):371-376.
112. Wang CH, Chen CS, Lin CL. The threat of Asian dust storms on asthma patients: a population-based study in Taiwan. *Glob Public Health*. 2014;9(9):1040-1052.
113. Watanabe M, Kurai J, Shimizu E. Influence of Asian dust storm on asthma in western Japan. *Genes Environ*. 2014;36(3):137-144.
114. Kanatani KT, Ito I, Al-Delaimy WK, Adachi Y, Mathews WC, Ramsdell JW. Desert dust exposure is associated with increased risk of asthma hospitalization in children. *Am J Respir Crit Care Med*. 2010;182(12):1475-1481.
115. Yang CY, Tsai SS, Chang CC, Ho SC. Effects of Asian dust storm events on daily admissions for asthma in Taipei, Taiwan. *Inhal Toxicol*. 2005;17(14):817-821.
116. Lee JW, Lee KK. Effects of Asian dust events on daily asthma patients in Seoul, Korea. *Meteorol Appl*. 2014;21(2):202-209.
117. Rutherford S, Clark E, McTainsh G, Simpson R, Mitchell C. Characteristics of rural dust events shown to impact on asthma severity in Brisbane, Australia. *Int J Biometeorol*. 1999;42(4):217-225.
118. Cheng MF, Ho SC, Chiu HF, Wu TN, Chen PS, Yang CY. Consequences of exposure to Asian dust storm events on daily pneumonia hospital admissions in Taipei, Taiwan. *J Toxicol Environ Health A*. 2008;71(19):1295-1299.
119. Kang JH, Keller JJ, Chen CS, Lin HC. Asian dust storm events are associated with an acute increase in pneumonia hospitalization. *Ann Epidemiol*. 2012;22(4):257-263.
120. Chiu HF, Tiao MM, Ho SC, Kuo HW, Wu TN, Yang CY. Effects of Asian dust storm events on hospital admissions for chronic obstructive pulmonary disease in Taipei, Taiwan. *Inhal Toxicol*. 2008;20(9):777-781.
121. Lorentzou C, Kouvarakis G, Kozyrakis GV, et al. Extreme desert dust storms and COPD morbidity on the island of Crete. *Int J Chron Obstruct Pulmon Dis*. 2019;14:1763-1768.
122. Vodonos A, Friger M, Katra I, et al. The impact of desert dust exposures on hospitalizations due to exacerbation of chronic obstructive pulmonary disease. *Air Q Atmos Health*. 2014;7(4):433-439.
123. Tam WW, Wong TW, Wong AH, Hui DS. Effect of dust storm events on daily emergency admissions for respiratory diseases. *Respirology*. 2012;17(1):143-148.
124. Wang JY, Li S, Wang SG, Shang KZ. Effects of long-term dust exposure on human respiratory system health in Minqin County, China. *Arch Environ Occup Health*. 2015;70(4):225-231.
125. Alangari AA, Riaz M, Mahjoub MO, Malhis N, Al-Tamimi S, Al-Modaihsh A. The effect of sand storms on acute asthma in Riyadh, Saudi Arabia. *Ann Thorac Med*. 2015;10(1):29-33.
126. Barnett AG, Fraser JF, Munck L. The effects of the 2009 dust storm on emergency admissions to a hospital in Brisbane, Australia. *Int J Biometeorol*. 2012;56(4):719-726.
127. Menendez I, Derbyshire E, Carrillo T, et al. Saharan dust and the impact on adult and elderly allergic patients: the effect of threshold values in the northern sector of Gran Canaria, Spain. *Int J Environ Health Res*. 2017;27(2):144-160.
128. Prospero JM, Blades E, Naidu R, Mathison G, Thani H, Lavoie MC. Relationship between African dust carried in the Atlantic trade winds and surges in pediatric asthma attendances in the Caribbean. *Int J Biometeorol*. 2008;52(8):823-832.
129. Wiggs GF, O'hara SL, Wegerdt J, Van Der Meer J, Small I, Hubbard R. The dynamics and characteristics of aeolian dust in dryland Central Asia: possible impacts on human exposure and respiratory health in the Aral Sea basin. *Geogr J*. 2003;169(2):142-157.
130. Kamouchi M, Ueda K, Ago T, Nitta H, Kitazono T. Relationship between Asian dust and ischemic stroke: a time-stratified case-crossover study. *Stroke*. 2012;43(11):3085-3087.
131. Kang JH, Liu TC, Keller J, Lin HC. Asian dust storm events are associated with an acute increase in stroke hospitalisation. *J Epidemiol Community Health*. 2013;67(2):125-131.
132. Radmanesh E, Maleki H, Goudarzi G, et al. Cerebral ischemic attack, epilepsy and hospital admitted patients with types of headaches attributed to PM10 mass concentration in Abadan, Iran. *Aeolian Res*. 2019;41:100541.
133. Yang CY, Chen YS, Chiu HF, Goggins WB. Effects of Asian dust storm events on daily stroke admissions in Taipei, Taiwan. *Environ Res*. 2005;99(1):79-84.
134. Chan YS, Teng JCY, Liu TC, Peng YI. Asian dust storms and diabetes hospitalization: a nationwide population-based study. *Air Q Atmos Health*. 2018;11(10):1243-1250.

135. Ko R, Hayashi M, Hayashi H, et al. Correlation between acute conjunctivitis and Asian dust on ocular surfaces. *J Toxicol Environ Health A*. 2016;79(8):367-375.
136. Akpinar-Elci M, Martin FE, Behr JG, Diaz R. Saharan dust, climate variability, and asthma in Grenada, the Caribbean. *Int J Biometeorol*. 2015;59(11):1667-1671.
137. Cadet G, Tourres R, Molinie J. Short-term effects of the particulate pollutants contained in Saharan dust on the visits of children to the emergency department due to asthmatic conditions in Guadeloupe (French Archipelago of the Caribbean). *PLoS One*. 2014;9(3):e91136.
138. Nakamura T, Hashizume M, Ueda K, et al. Asian dust and pediatric emergency department visits due to bronchial asthma and respiratory diseases in Nagasaki, Japan. *J Epidemiol*. 2016;26(11):593-601.
139. Park YS, Kim JH, Jang HJ, Tae YH, Lim DH. The effect of Asian dust on asthma by socioeconomic status using national health insurance claims data in Korea. *Inhal Toxicol*. 2016;28(1):1-6.
140. Merrifield A, Schindler S, Jalaludin B, Smith W. Health effects of the September 2009 dust storm in Sydney, Australia: did emergency department visits and hospital admissions increase? *Environ Health*. 2013;12:32.
141. Carlsen HK, Gislason T, Forsberg B, et al. Emergency hospital visits in association with volcanic ash, dust storms and other sources of ambient particles: a time-series study in Reykjavik, Iceland. *Int J Environ Res Public Health*. 2015;12(4):4047-4059.
142. Chan CC, Chuang KJ, Chen WJ, Chang WT, Lee CT, Peng CM. Increasing cardiopulmonary emergency visits by long-range transported Asian dust storms in Taiwan. *Environ Res*. 2008;106(3):393-400.
143. Chien LC, Lien YJ, Yang CH, Yu HL. Acute increase of children's conjunctivitis clinic visits by Asian dust storms exposure - a spatiotemporal study in Taipei, Taiwan. *PLoS One*. 2014;9(10):e109175.
144. Yang CY. Effects of Asian dust storm events on daily clinical visits for conjunctivitis in Taipei, Taiwan. *J Toxicol Environ Health A*. 2006;69(18):1673-1680.
145. Chien LC, Yang CH, Yu HL. Estimated effects of Asian dust storms on spatiotemporal distributions of clinic visits for respiratory diseases in Taipei children (Taiwan). *Environ Health Perspect*. 2012;120(8):1215-1220.
146. Hefflin BJ, Jalaludin B, McClure E, et al. Surveillance for dust storms and respiratory diseases in Washington state, 1991. *Arch Environ Health*. 1994;49(3):170-174.
147. Lin YK, Chen CF, Yeh HC, Wang YC. Emergency room visits associated with particulate concentration and Asian dust storms in metropolitan Taipei. *J Expo Sci Environ Epidemiol*. 2016;26(2):189-196.
148. Liu ST, Liao CY, Kuo CY, Kuo HW. The effects of PM2.5 from Asian dust storms on emergency room visits for cardiovascular and respiratory diseases. *Int J Environ Res Public Health*. 2017;14(4):428.
149. Yang ZH, Zhang YX, Zhang QX, Zhang J, Lu B, Meng ZQ. Association of PM10 derived from dust events with daily outpatient number for cardiovascular diseases. *Zhongguo Huanjing Kexue/China Environ Sci*. 2015;35(1):277-284.
150. Higashi T, Kambayashi Y, Ohkura N, et al. Effects of Asian dust on daily cough occurrence in patients with chronic cough: a panel study. *Atmos Environ*. 2014;92:506-513.
151. Higashi T, Kambayashi Y, Ohkura N, et al. Exacerbation of daily cough and allergic symptoms in adult patients with chronic cough by Asian dust: a hospital-based study in Kanazawa. *Atmos Environ*. 2014;97:537-543.
152. Otani S, Onishi K, Mu H, Kurozawa Y. The effect of Asian dust events on the daily symptoms in Yonago, Japan: a pilot study on healthy subjects. *Arch Environ Occup Health*. 2011;66(1):43-46.
153. Onishi K, Kurosaki Y, Otani S, Yoshida A, Sugimoto N, Kurozawa Y. Atmospheric transport route determines components of Asian dust and health effects in Japan. *Atmos Environ*. 2012;49:94-102.
154. Majbaudhin A, Onishi K, Otani S, Kurosaki Y, Kurozawa Y. Association between Asian dust-borne air pollutants and daily symptoms on healthy subjects: a web-based pilot study in Yonago, Japan. *J Environ Public Health*. 2016;2016:8280423.
155. Meo SA, Al-Kherajji MF, Alfaraj ZF, Alwehaibi NA, Alderehim AA. Respiratory and general health complaints in subjects exposed to sandstorm at Riyadh, Saudi Arabia. *Pak J Med Sci*. 2013;29(2):642-646.
156. Kurai J, Watanabe M, Noma H, et al. Estimation of the effects of heavy Asian dust on respiratory function by definition type. *Genes Environ*. 2017;39:25.
157. Watanabe M, Noma H, Kurai J, et al. Differences in the effects of Asian dust on pulmonary function between adult patients with asthma and those with asthma-chronic obstructive pulmonary disease overlap syndrome. *Int J COPD*. 2016;11:183-190.
158. Watanabe M, Noma H, Kurai J, et al. Association of sand dust particles with pulmonary function and respiratory symptoms in adult patients with asthma in western Japan using light detection and ranging: a panel study. *Int J Environ Res Public Health*. 2015;12(10):13038-13052.
159. Yoo Y, Choung JT, Yu J, Kim DK, Koh YY. Acute effects of Asian dust events on respiratory symptoms and peak expiratory flow in children with mild asthma. *J Korean Med Sci*. 2008;23(1):66-71.
160. Watanabe M, Noma H, Kurai J, et al. Association between pulmonary function and daily levels of sand dust particles assessed by light detection and ranging in schoolchildren in western Japan: a panel study. *Allergol Int*. 2016;65(1):56-61.
161. Hong Y-C, Pan X-C, Kim S-Y, et al. Asian dust storm and pulmonary function of school children in Seoul. *Sci Total Environ*. 2010;408(4):754-759.
162. Park JW, Lim YH, Kyung SY, et al. Effects of ambient particulate matter on peak expiratory flow rates and respiratory symptoms of asthmatics during Asian dust periods in Korea. *Respirology*. 2005;10(4):470-476.
163. Kanatani KT, Hamazaki K, Inadera H, et al. Effect of desert dust exposure on allergic symptoms: a natural experiment in Japan. *Ann Allergy Asthma Immunol*. 2016;116(5):425.
164. Watanabe M, Yamasaki A, Burioka N, et al. Correlation between Asian dust storms and worsening asthma in Western Japan. *Allergol Int*. 2011;60(3):267-275.
165. O'Hara SL, Wiggs GFS, Wegerdt J, et al. Dust exposure and respiratory health amongst children in the environmental disaster zone of Karakalpakstan, Central Asia: preliminary findings of the ASARD project. *Environ Health Risk*. 2001;5:71-82.
166. Islam MM, Alharthi M, Alam MM. The impacts of climate change on road traffic accidents in Saudi Arabia. *Climate*. 2019;7(9):103.
167. Soy FK, Yazici H, Kulduk E, et al. The effects of dust storms on quality of life of allergic patients with or without asthma. *Kulak Burun Bogaz Ihtis Derg*. 2016;26(1):19-27.
168. Mu H, Battsetseg B, Ito TY, Otani S, Onishi K, Kurozawa Y. Effects of Asian dust storm on health-related quality of life: a survey immediately after an Asian dust storm event in Mongolia. *Int J Health Res*. 2010;3(2):87-92.
169. Lee H, Jung J, Myung W, et al. Association between dust storm occurrence and risk of suicide: case-crossover analysis of the Korean national death database. *Environ Int*. 2019;133(Pt A):105146.
170. Dadvand P, Basagana X, Figueras F, et al. Saharan dust episodes and pregnancy. *J Environ Monit*. 2011;13(11):3222-3228.
171. Hashizume M, Ueda K, Nishiwaki Y, Michikawa T, Onozuka D. Health effects of Asian dust events: a review of the literature. *Jpn J Hyg*. 2010;65(3):413-421.