

## **Impacts of High Concentration, Medium Duration Coal Mine Fire Related PM<sub>2.5</sub> on Cancer Incidence: 5-Year Follow-Up of the Hazelwood Health Study**

Authors: Yu, Pei, Guo, Yuming, Gao, Caroline X, Dimitriadis, Christina, Ikin, Jillian F, et al.

Source: Environmental Health Insights, 15(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/11786302211059722>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Impacts of High Concentration, Medium Duration Coal Mine Fire Related PM<sub>2.5</sub> on Cancer Incidence: 5-Year Follow-Up of the Hazelwood Health Study

Environmental Health Insights  
Volume 15: 1–4  
© The Author(s) 2021  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/11786302211059722



Pei Yu<sup>1</sup>, Yuming Guo, Caroline X Gao<sup>1</sup>, Christina Dimitriadis, Jillian F Ikin<sup>1</sup>, Anthony Del Monaco<sup>1</sup>, David Brown, Malcolm R Sim and Michael J Abramson

School of Public Health and Preventive Medicine, Monash University, Melbourne, VIC, Australia.

**ABSTRACT:** No studies have investigated the cancer outcomes from high level medium duration coal mine fire fine particulate matter  $\leq 2.5\mu\text{m}$  in diameter (PM<sub>2.5</sub>) exposure. We included 2208 Morwell residents (exposed) and 646 Sale residents (unexposed) who participated in the Hazelwood Health Study Adult Survey. Competing risk regression models were used to evaluate relationships between coal mine fire exposure and cancer incidence, adjusting for known confounders. There were 137 cancers in the exposed and 27 in the unexposed over 14 849 person-years of follow-up. A higher risk of cancer incidence was observed for Morwell participants (HR = 1.67 [95% CI 1.05–2.67]), but no evidence to suggest associations between PM<sub>2.5</sub> exposure and incidence of all cancers (HR = 1.02 [95% CI 0.91–1.13]), or site-specific cancers. There is no strong evidence that exposure to high concentrations of mine fire-related PM<sub>2.5</sub> over a prolonged period could explain the higher risk in exposed population in this study.

**KEYWORDS:** Coal mine fire smoke, fine particulate matter, cancer incidence, data linkage cohort, Hazelwood Health Study

**RECEIVED:** October 4, 2021. **ACCEPTED:** October 20, 2021.

**TYPE:** Short Report

**FUNDING:** The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: PY was supported by the China Scholarship Council (201906210065). YG was supported by a Career Development Fellowship of the Australian National Health and Medical Research Council (APP1163693). The Hazelwood Health Study is funded by the Victorian Department of Health, however this paper presents the views of the authors and does not represent the views of the Department.

**DECLARATION OF CONFLICTING INTERESTS:** The author(s) declared the following potential conflicts of interest with respect to the research, authorship, and/or publication of this article: MJA holds investigator initiated grants from Pfizer and Boehringer-Ingelheim for unrelated research. He has undertaken an unrelated consultancy for and received assistance with conference attendance from Sanofi. He has also received a speaker's fee from GSK. All other authors declare no competing interests.

**CORRESPONDING AUTHOR:** Yuming Guo, School of Public Health and Preventive Medicine, Monash University, 553 St Kilda Rd, Melbourne, VIC 3004, Australia. Email: yuming.guo@monash.edu

## Introduction

Coal mine fires cause substantial environmental and economic harms, and have been increasingly shown to be associated with adverse effects on human health.<sup>1</sup> However, there are no studies exploring the association between coal mine fire exposure with malignant cancers.<sup>2</sup> Up to 62 compounds with potentially harmful health impacts are found in coal mine fire smoke, including fine particulate matter  $\leq 2.5\mu\text{m}$  in diameter (PM<sub>2.5</sub>)<sup>1</sup> which has been classified as a Group 1 carcinogen to humans.<sup>3</sup> Associations between long-term (years) ambient total PM<sub>2.5</sub> exposure and cancer risk have been demonstrated in many studies,<sup>4</sup> however the effect of medium-term (weeks to months) exposure at much higher concentration is unknown. The composition of PM<sub>2.5</sub> from coal mine fires may differ from other sources and have unique health impacts.<sup>5</sup> Five years after the Hazelwood mine fire, which burned for 6 weeks in south-eastern Australia, we investigated the association between this high concentration, medium duration exposure event and cancer incidence.

## Methods

### Population and study design

The Hazelwood mine fire burned from 9 February 2014 to approximately 31 March 2014 in the Latrobe Valley, a rural area in the State of Victoria, Australia. The Hazelwood Health Study (HHS) ([www.hazelwood-healthstudy.org.au](http://www.hazelwood-healthstudy.org.au)) was commissioned by the Victoria

Department of Health and Human Services to investigate the long-term effects of the Hazelwood mine fire.<sup>6</sup> The Adult Survey is one of the streams of research in HHS and the cohort comprised adult residents from the towns of Morwell (located adjacent to the mine and highly exposed) and Sale (60 km away and unexposed).<sup>6</sup> Participants of this study were members of the Adult Survey cohort who consented to linkage with the Victorian Cancer Registry (VCR).

VCR data were obtained for the period 1st January 2008 and 31st December 2019 which comprised demographics, cancer site, International Classification of Disease (ICD) codes, count of tumours and vital status. Death data were obtained from the VCR supplemented with reports by next of kin and Ryerson newspaper index (<https://ryersonindex.org>) searching. Participants were followed up until 31 December 2019 or date of death, whichever was earlier.

Monash University Human Research Ethics Committee (MUHREC) approved the Hazelwood Adult Survey & Health Record Linkage Study.

### Mine-fire related PM<sub>2.5</sub> exposure

The Commonwealth Scientific and Industrial Research Organisation (CSIRO) modelled temporal and spatial PM<sub>2.5</sub> concentrations during the mine fire using high-resolution meteorological and dispersion models incorporating local wind data and an appropriate plume rise mechanism.<sup>7</sup> Participants'



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without

**Table 1.** Participant characteristics at time of interview.


CHARACTERISTIC	MORWELL (N=2208)	SALE (N=646)	P-VALUE
Mean daily mine-fire related PM <sub>2.5</sub> exposure (µg/m <sup>3</sup> ), Mean (SD)	15.3 (11.7)	0.0 (0.3)	
Age at the survey, mean (SD) years	58.1 (16.4)	58.0 (17.4)	.914
0-59	1079 (48.9%)	320 (49.5%)	.765
≥60	1129 (51.1%)	326 (50.5%)	
Sex			.112
Male	1018 (46.1%)	275 (42.6%)	
Female	1190 (53.9%)	371 (57.4%)	
Marital status			.062
Married	1346 (61.0%)	420 (65.0%)	
Not married	862 (39.0%)	226 (35.0%)	
Education			<.001
Secondary up to year 10	704 (32.1%)	147 (22.9%)	
Secondary year 11-12	435 (19.8%)	106 (16.5%)	
Certificate (trade/apprenticeship/technicians)	759 (34.6%)	266 (41.4%)	
University or other tertiary institute degree	296 (13.5%)	124 (19.3%)	
Missing <sup>a</sup>	14	3	
Employment			<.001
Paid employment (FT, PT, self-employed)	909 (41.5%)	292 (45.3%)	
Other (student/volunteer/home-duties/retired)	1007 (45.9%)	310 (48.1%)	
Unemployed	101 (4.6%)	12 (1.9%)	
Not working due to ill-health	176 (8.0%)	30 (4.7%)	
Missing <sup>a</sup>	15	2	
Occupational exposure <sup>b</sup>			<.001
Not exposed	1244 (56.3%)	407 (63.0%)	
Exposed in coal mine or power station	375 (17.0%)	21 (3.3%)	
Exposed, but not coal mine/station	589 (26.7%)	218 (33.7%)	
Smoking			.033
Never <sup>c</sup>	1070 (48.7%)	345 (53.9%)	
Current smoker	376 (17.1%)	87 (13.6%)	
Former smoker	750 (34.2%)	208 (32.5%)	
Missing <sup>a</sup>	14	6	
Alcohol drinking <sup>d</sup>			.011
Non-drinker	546 (25.4%)	150 (23.5%)	
Low risk	790 (36.8%)	205 (32.2%)	
High risk	810 (37.7%)	282 (44.3%)	
Missing <sup>a</sup>	62	9	

<sup>a</sup>Participants did not answer this question in the questionnaire.

<sup>b</sup>Participants' self-reported job held for at least 6 months which involved breathing a lot of dust, fumes, gas, vapour or mist.

<sup>c</sup>Participants self-reported never smoked or smoked occasionally.

<sup>d</sup>Participants with alcohol use disorders identification test consumption score of 0 indicated a non-drinker, 1 to 3 for men, 1 to 2 for women indicated a low risk drinker and ≥4 for men, ≥3 for women indicated a high-risk drinker.



**Figure 1.** Forest plots showing adjusted Hazard Ratios (HRs with 95% CI) for incidence of all cancers, and site-specific cancers, per 10-µg/m<sup>3</sup> increase in mine-fire related PM<sub>2.5</sub> exposure for all participants (A) and incidence for participants living in Morwell compared with Sale (B). Hazard ratios were estimated from competing risk regression model adjusted for sex, age, marital status, education, employment status, smoking and drinking behaviour and occupational exposure. <sup>a</sup>Analyses for breast cancer were conducted in females only, while analyses for cancers of male reproductive organs, for example, prostate were conducted in males only. <sup>b</sup>Crude cancer incidence per 1000 (‰) population.

mine-fire related PM<sub>2.5</sub> exposures were estimated by mapping their detailed time location diaries, collected as a part of Adult Survey, onto the modelled PM<sub>2.5</sub> data.<sup>6</sup>

*Statistical analysis*

Subjects diagnosed with a primary malignant cancer (included ICD10: C00-C97, D45-D46, D47.1, D47.3-D47.5) from 6 months after the fire (9 August 2014) were defined as incident cases. Competing risk regression models were developed to quantify the association between region and PM<sub>2.5</sub> exposures on cancer incidence by calculating hazard ratios (HRs) and 95% confidence intervals (CIs) and controlling for known confounders including age, sex, marital status, education, employment, occupational exposure, smoking and alcohol drinking. Statistical significance was defined as a 2-side P-value <.05. R software (V3.4.3, www.R-project.org) was used to perform all data analyses.

**Results**

There were 164 cancers (137 in the exposed group, 27 in the unexposed group) over 14849 person-years of follow-up. Risk factors listed in Table 1 were compared between Morwell and Sale. Morwell residents had lower levels of education, employment and risky drinking. For the period 5 years after the fire

(9 August 2014–31 December 2019), a higher risk of overall cancer incidence was observed in Morwell compared with Sale (HR = 1.67 [95% CI 1.05–2.67]), after adjustment for known confounders. However increments in mine fire-related PM<sub>2.5</sub> exposure were not found to be associated with overall cancer incidence (HR = 1.02 [95% CI 0.91–1.13]) or any site-specific cancers during this 5 years period. All urinary tract cancers in the study period were from Morwell and thus not able to estimate an HR (Figure 1).

**Discussion**

Previous analysis showed that, prior to the mine fire, overall cancer incidence in the Morwell and Sale regions were similar to rural and regional Victoria.<sup>8</sup> We observed an increased incidence of cancer in 5 years of follow-up after the fire in the exposed versus unexposed town, however there was no strong evidence that these cancers were associated with mine fire related PM<sub>2.5</sub> exposure.

It’s possible that homogeneity in PM<sub>2.5</sub> levels within the exposed participants contributed to the lack of any observed association between increments in PM<sub>2.5</sub> and cancer incidence. Although there was no effect related to exposure found, the limited number of cases and the long-latency period for many types of cancers likely limited our ability to identify associations. Another possibility for the difference

between the 2 townships was related to risk factors that were not measured in the study, for example, diet. For urinary cancers, all cases were from the exposed town, which warrants further investigation using larger population level data to better understand regional differences.

Although no statistically significant association was found with PM<sub>2.5</sub> 5 years after the exposure, this is to our knowledge the first study to investigate an association between coal mine fire related PM<sub>2.5</sub> and cancer incidence. Further follow-up is needed to provide more certainty around any effects of the Hazelwood mine fire exposure on cancer incidence.

### Acknowledgements

We thank participants in the HHS Adult Survey, Victorian Cancer Registry.

### Author Contributions

Conception and design: PY, MS, MJA. Development of methodology: PY, YG, MJA. Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): CD, DB, JI, ADM. Analysis and interpretation of data: PY, CG, ADM. Writing, review and/or revision of the manuscript: PY, CG, YG, MS, MJA, JI, ADM. Administrative, technical or material support (ie, reporting or organizing data,


constructing databases): CG, JI, DB. Study supervision: JI, YG, MS, MJA.

### ORCID iDs

Pei Yu  <https://orcid.org/0000-0002-7485-1751>

Caroline X Gao  <https://orcid.org/0000-0002-0987-2759>

Jillian F Ikin  <https://orcid.org/0000-0003-0794-0129>

Anthony Del Monaco  <https://orcid.org/0000-0002-9352-0552>

### REFERENCES

- Melody SM, Johnston FH. Coal mine fires and human health: what do we know? *Int J Coal Geol.* 2015;152:1-14.
- Roy D, Singh G, Seo YC. Coal mine fire effects on carcinogenicity and non-carcinogenicity human health risks. *Environ Pollut.* 2019;254:113091.
- Loomis D, Grosse Y, Lauby-Secretan B, et al. The carcinogenicity of outdoor air pollution. *Lancet Oncol.* 2013;14:1262-1263.
- Yu P, Guo S, Xu R, et al. Cohort studies of long-term exposure to outdoor particulate matter and risks of cancer: a systematic review and meta-analysis. *Innovation.* 2021;2:100143.
- Wyzga RE, Rohr AC. Long-term particulate matter exposure: attributing health effects to individual PM components. *J Air Waste Manag Assoc.* 2015;65:523-543.
- Ikin J, Carroll MTC, Walker J, et al. Cohort profile: the Hazelwood health study adult cohort. *Int J Epidemiol.* 2021;49:1777-1778.
- Luhar AK, Emmerson KM, Reisen F, Williamson GJ, Cope ME. Modelling smoke distribution in the vicinity of a large and prolonged fire from an open-cut coal mine. *Atmos Environ.* 2020;229:117471.
- Sim M, Dimitriadis C, Gao C, Del Monaco A. *Hazelinks – cancer incidence analysis (First data extraction)*. Version 1.0. Hazelwood Health Study; 2017.