

Environmental Predictors of Undernutrition Among Under-5 Children at Dabat District Health Facilities, Northwest Ethiopia, 2023

Authors: Melese, Mihret, Esubalew, Dereje, Malede, Asmamaw, Birhan, Tsegaye Adane, and Azanaw, Jember

Source: Environmental Health Insights, 18(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/11786302241252735>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.



Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Environmental Predictors of Undernutrition Among Under-5 Children at Dabat District Health Facilities, Northwest Ethiopia, 2023

Environmental Health Insights
Volume 18: 1–9
© The Author(s) 2024
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/11786302241252735



Mihret Melese¹, Dereje Esubalew², Asmamaw Malede³ ,
Tsegaye Adane Birhan⁴  and Jember Azanaw⁴

¹Department of Human Physiology, College of Medicine and Health Sciences, University of Gondar, Gondar, Ethiopia. ²Department of Human Physiology, College of Medicine and Health Science, Ambo University, Ambo, Ethiopia. ³Department of Environmental Health, College of Medicine and Health Sciences, Wollo University, Dessie, Ethiopia. ⁴Department of Environmental and Occupational Health and Safety, Institute of Public Health, College of Medicine and Health Sciences, University of Gondar, Gondar, Ethiopia.

ABSTRACT

BACKGROUND: In Ethiopia, undernutrition remains a significant public health concern throughout the year due to persistent household food insecurity. The intensifying conflict in the Dabat district, involving the Ethiopian government and the Tigray Liberation Front, has increased the threat of undernutrition among under-5 children. The cessation of humanitarian aid has worsened food insecurity, increasing the vulnerability of the children in the region. However, there was no data showing the prevalence and environmental predictors of undernutrition in the Dabat district, northwestern Ethiopia. Therefore, the objective of this study was to determine the prevalence of undernutrition and identify environmental predictors in this study setting.

METHODS: This institutional-based cross-sectional study was conducted from January to March 2023. A total of 400 under-5 children were included in this study using systematic random sampling techniques with a 100% response rate. The study included all under-5 children who visited healthcare facilities during the data collection period. However, it excluded children under-5 who had physical deformities, were critically ill, or had congenital abnormalities or known chronic diseases such as HIV/AIDS, tuberculosis, or chronic heart diseases. Binary logistic regression was used to determine environmental predictors of undernutrition, with statistical significance at a *P*-value of $\leq .05$.

RESULTS: The overall prevalence of undernutrition was found to be 12% (95% CI: 8.72–21.5). Among those affected, 32.8% (95% CI: 21.5–39.8) were stunted, 37.9% (95% CI: 28.8–47.50) were underweight, and 29.3% (95% CI: 21.3–29.7) were wasted. The environmental predictors of undernutrition were latrine use, recent diarrhea episodes, mothers' occupation and place of residence, water treatment before consumption, water sources and storage, water extraction methods, and mothers' handwashing habits.

CONCLUSIONS AND RECOMMENDATIONS: The study found a higher prevalence of undernutrition among under-5 children compared to the 2019 Ethiopia Mini Demographic and Health Survey. This study recommended raising awareness about establishing private toilets, promoting water treatment, proper latrine use, and handwashing practices. Training on personal hygiene and economic support for households should also be provided. Healthcare providers should offer quality health services for under-5 children. Further research is needed to explore specific nutrient deficiencies using laboratory methods.

PLAIN LANGUAGE SUMMARY

What is Already Known on This Topic? Undernutrition is the main cause of child mortality, physical, and mental retardation. What This Study Adds? This research aimed to determine the current prevalence and factors that contribute to malnutrition in Dabat, Northwest Ethiopia, 2023.

KEYWORDS: Prevalence, stunting, underweight, wasting, Ethiopia

RECEIVED: November 30, 2023. **ACCEPTED:** April 18, 2024.

TYPE: Research Article

FUNDING: The author(s) received no financial support for the research, authorship, and/or publication of this article.

DECLARATION OF CONFLICTING INTERESTS: The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

CORRESPONDING AUTHOR: Mihret Melese, Department of Human Physiology, College of Medicine and Health Sciences, University of Gondar, Gondar, Ethiopia.
Email: mihhret86@gmail.com

Introduction

Undernutrition is a medical condition resulting from inadequate food intake, insufficient intake of the right combination of nutrients, or the body's inability to properly use the food consumed. It manifests itself in 3 forms: stunting, wasting, and underweight.¹ In 2020, global statistics revealed that approximately 149 million under-5 children were stunted, while approximately 45.4 million were wasted. Moreover,

approximately 38.9 million under-5 children were identified as overweight.^{1–3} Undernutrition significantly increases children's susceptibility to disease and mortality. Globally, nearly 45.4 million deaths among under-5 children are associated with undernutrition, with a substantial portion occurring in developing countries.¹ According to 2015 statistics in Ethiopia, 10% of children under-5 years of age are underweight, 29% are overweight, and 44% are stunted.⁴



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without




Figure 1. The location of the Dabat district zone in the Amhara region, Ethiopia, 2023.

Undernutrition in under-5 children can lead to stunted growth, weakened immune systems, cognitive impairment, and higher mortality from illnesses like diarrhea and pneumonia. It may also result in chronic health issues later in life and perpetuate a cycle of poverty.

Addressing undernutrition is crucial to ensure healthy development and break the cycle of negative outcome.⁵⁻⁸

Several factors contribute to the occurrence of undernutrition among under-5 children. These include sociodemographic variables such as age, employment status, and educational background of parents, as well as factors such as contaminated water and sanitation, food insecurity, limited access to healthcare services, environmental pollution, inadequate housing, issues related to climate change, and a lack of education and awareness.⁹⁻¹²

People living in areas affected by conflict often struggle with issues such as food insecurity, disruptions in medical care, increased susceptibility to infectious diseases, and restricted access to humanitarian aid. These conditions increase the risk of undernutrition among under-5 children.¹¹⁻¹⁴ In Ethiopia, undernutrition remains a significant public health concern throughout the year due to persistent household food insecurity. The intensifying conflict in the Dabat district, involving the Ethiopian government and the Tigray Liberation Front, has increased the threat of undernutrition among under-5 children. The cessation of humanitarian aid has worsened food insecurity, increasing the vulnerability of the children in the region. However, there was no data showing the prevalence and environmental predictors of undernutrition in the Dabat district, northwestern Ethiopia. Therefore, the objective of this study was to determine the prevalence of undernutrition and identify environmental predictors in this study setting.

Methods

Study design and area

We conducted an institutional-based cross-sectional study in the Dabat district from January to March 2023. The district is

located approximately 775 km from Addis Ababa and 70 km from Gondar city. In the Dabat district, there are 39 rural kebeles and 4 urban kebeles, with an estimated 51 444 households and a total population of 203 050 residents. It also has 1 primary hospital, 6 health centers, and 35 health posts (Figure 1).

Study population

The source population for this study consisted of all under-5 children who visited healthcare facilities in the Dabat district. The study population included all under-5 children accessing health facilities during the data collection period.

Eligibility criteria and inclusion criteria

Inclusion criteria. All under-5 children who visited selected health facilities in the Dabat district during the data collection period.

Exclusion criteria. Under-5 children with physical deformities, those who were critically ill, and those with congenital abnormalities or known chronic diseases such as HIV/AIDS, tuberculosis, or chronic heart disease were excluded from the study.

Sample size and sampling technique

The sample size was determined using the single population proportion from the Ethiopia Demographic and Health Survey (EDHS) 2016 report on the Amhara Region, which reported a prevalence of 46%.¹³

$$n_i = \frac{(Z_{\alpha/2})^2 * p(1-p)}{d^2} = \frac{(1.96)^2 * 0.46(1-0.46)}{(0.05)^2} = 381,$$

With a 5% contingency rate, the final sample size was $N_f = 19 + 381 = 400$, where n_i = the initial sample size $z_{\alpha/2} = 1.96$ (the critical value for a normal distribution at a 95% confidence level). $P = .46$ (the proportion of stunting). $d = 0.05$ (the level of precision or acceptable error). N_f = final sample size

variables; $P = .46$ (the proportion of stunting); $d = 0.05$ (the level of precision or acceptable error); $N_f =$ final sample size.

Variables

Dependent variable. Undernutrition (Yes/No).

The independent variables included sociodemographic factors (age of the child, sex, number of children under 5 in the household, mother's educational status, residence, income, occupation) and environmental factors (water treatment practices, source of drinking water, ownership of latrine facilities, handwashing practices after toilet use, occurrence of diarrheal disease).

Operational definition

Malnutrition. It refers to an imbalance between nutrient requirements and intake, which can adversely impact growth, development, and other relevant outcomes.¹⁴

Undernutrition. It was defined as under-5 children with either a z score for weight-for-height (WHZ) $\leq -2SD$ or weight-for-age (WAZ) $\leq -2SD$, or a height-for-age (HAZ) $\leq -2SD$.¹⁵

Stunting. The HAZ score was considered $\leq (SD)$.¹⁵

Wasting. It was defined as a WHZ score of $\leq -2SD$.

Underweight was defined as a study participant with a weight-for-age z score (WAZ) ≤ -2 standard deviations (SDs).¹⁵

Mid-upper arm circumference (MUAC). We used the MUAC cut-off points of any child with a MUAC of 126 mm who was declared undernourished to evaluate whether a child was under- or overnourished.¹⁵

Data collection tools. Primarily, the questionnaire was organized in the English language by reviewing related literature and translated into the Amharic language to facilitate the communication between the data collector and study participants. The tool included data on sociodemographic characteristics and clinical and environmental factors. All measurements were conducted using a standardized protocol with calibrated equipment adhering to international standards for anthropometric measurements. Weight measurements for children aged 2 years and younger were taken using a baby scale, with the children wearing light clothing. Moreover, their height was measured horizontally using a wooden UNICEF height sheet. For children older than 2 years, weight was assessed while they stood on the center of a scale, wearing lightweight clothing, and without shoes. To measure a child's height, they were required to stand upright on a wooden height measuring board facing forward with their head held high. Subsequently, the anthropometric data were entered into the WHO Antero Plus software version 3.2.2 for analysis.¹⁶ After receiving the necessary training, 2 female clinical nurses and 2 female midwifery professionals collected all of this data. Prior to data collection, the

tool's validity and reliability were assessed, and the Cronbach's alpha was .757.

Statistical analysis

The data were entered and cleaned using EpiInfo version 7, while analysis was conducted using SPSS V.26 software. Descriptive analysis was used to generate frequencies, percentages, mean, and standard deviations for the study variables. The chi-squared test was employed to assess associations between dependent and independent variables. Next, bivariable and multivariable binary logistic regression analyses were performed. In the bivariable analysis, predictor variables with a P -value $< .2$ were selected for multivariable analyses. Subsequently, in multivariable logistic regression, variables with $P \leq .05$ were identified as environmental predictor variables of undernutrition. The Hosmer Lemeshow test was used to assess the fitness of the final model.

Ethical considerations

Ethical approval was obtained from the Institutional Review Board of the University of Gondar (IRB/346/2023). According to the Helsinki Declaration, under-5 children are incapable of providing informed consent on their own.¹⁷ Thus, their mothers or other caregivers provided their informed consent.

Results

Background characteristics of the study subjects

In this study, a total of 400 under-5 children were included, 227 of whom (56.8%) were female. The mean age of the under-5 children was 30.68 (± 3.49 SD) months. Approximately 37% of mothers had a diploma or above educational background. Of the participants included in this study, 62% came from rural areas (Table 1).

Environmental health characteristics

The findings indicate that the majority of study participants (83.7%) reported consuming treated water and having continuous access to it. Among the 400 households, 92 (23.0%) engaged in open defecation. Additionally, one-fifth of the study participants reported never washing their hands after using the toilet (refer to Table 2 for details).

Prevalence of undernutrition

The study found that the overall prevalence of undernutrition was 12% (95% CI: 8.72-21.5). Among the undernourished patients, 32.8% (95% CI: 21.5-39.8) were stunted based on their height-for-age z -score (HAZ). Additionally, 37.9% (95% CI: 28.8-47.5) were classified as underweight according to their weight-for-age z -score (WAZ), and 29.3% (95% CI:

Table 1. Background characteristics of the study participants in the Dabat district, Northwest Ethiopia, 2023.

VARIABLES	CATEGORIES	FREQUENCY	(%)
Age of the child in month	<30	173	43.3
	31-45	132	33.0
	≥46	95	23.8
Mean age = 30.68 ± 3.491(Std. Dev)			
Sex	Male	173	43.3
	Female	227	56.8
No. of the under-5 y of age	1	350	87.5
	≥2	50	12.5
Residence	Rural	152	38.0
	Urban	248	62.0
Mothers educational status	Not read and write	115	28.7
	Read and write	46	1.5
	Primary education	14	3.5
	Secondary education	77	19.3
	Diploma and above	146	6.5
Vaccination status	Partial vaccinated	37	9.2
	Fully vaccinated	363	90.8
Mothers occupation	Housewife	144	36.0
	Government employee	204	51.0
	Private	52	13.0


Table 2. Environmental characteristics of the study participants in the Dabat district, Northwest Ethiopia, 2023.

VARIABLES	CATEGORIES	FREQUENCY	PERCENT
Latrine ownership	Not at all	92	23.0
	Commune latrine	53	13.3
	Private latrine	255	63.7
Major water source	Pipe	245	61.3
	Protected well or spring	81	20.3
	Unprotected well or spring	63	15.8
	River	11	2.8
Does the water available at all time	Yes	75	18.8
	No	322	80.5

(Continued)

Table 2. (Continued)

VARIABLES	CATEGORIES	FREQUENCY	PERCENT
Time to fetch water	<30 min	131	32.8
	≥30 min	269	67.2
Household water treatment	Yes	65	16.3
	No	335	83.7
Do you cover drinking water storage always?	Yes	332	83.0
	No	68	17.0
How do you take water from the drinking water storage container?	Pouring	303	75.8
	Dipping	97	24.2
Do you wash fruits or vegetables before consumption?	Rarely	77	19.3
	Sometimes	49	12.3
	Always	274	68.5
Do you trim a child's nails when grown	Rarely	49	12.2
	Sometimes	65	16.3
	Always	286	71.5
Mothers/ caregivers hand washing after toilet	Never	81	20.3
	Sometimes	70	17.4
	Always	249	62.3

**Figure 2.** Undernutrition status of 5-children in Dabat district, Northwest Ethiopia, 2023.

21.3–29.7) were wasted based on their weight-for-height z-score (WHZ) (refer to Figure 2).

Factors associated with undernutrition

Multivariable logistic regression analysis showed that undernutrition was significantly associated with latrine utilization, diarrhea within 2 weeks, mothers' occupation, residence, drinking untreated water, latrine ownership, water sources, water storage status, ways of taking water from storage, and mothers' hand washing after toilet. The odds of stunting were

higher among under-5 children whose parents reported improper latrine use compared to those whose families reported proper latrine use (AOR = 9.76; 95% CI: 3.29–15.71). The odds of becoming stunted, wasted, and underweight were higher among the children affected by diarrhea (AOR = 11.67; 95% CI: 4.86–28.07), (AOR = 9.16; 95% CI: 3.92–21.39), and (AOR = 8.57, 95% CI: 4.09–17.93), respectively. Compared to mothers who were government employees, housewife mothers had a higher likelihood of their children experiencing wasting (AOR = 5.53, 95% CI: 42.03–15.05) and being underweight (AOR = 3.58, 95% CI: 1.27–10.06). Under-5 children in rural areas had higher odds of stunting (AOR = 8.51; 95% CI: 3.35–22.16), wasting (AOR = 2.98; 95% CI: 1.44–3.76), and being underweight (AOR = 6.07; 95% CI: 2.88–12.80). Under-5 children from households with improper drinking water storage had higher odds of stunting (AOR = 2.83; 95% CI: 3.99–14.57) and wasting (AOR = 3.81; 95% CI: 3.99–14.57) compared to those from households that stored their drinking water properly. Under-5 children whose mothers never or occasionally washed their hands after using the toilet had significantly higher odds of wasting compared to those whose mothers always washed their hands (AOR = 2.20, 95% CI: 1.91–5.53). Similarly, children whose mothers never or sometimes washed their hands after using the toilet had higher odds of being underweight compared to those whose mothers always washed their hands, with adjusted odds ratios of 2.51 (95% CI: 2.21–8.83) and 1.51 (95% CI: 1.12–4.34), respectively (Table 3).

Discussion

The study findings showed that undernutrition was more prevalent among children under-5 (refer to Figure 1). This aligns with a study conducted in Debre Birhan, Ethiopia, 15.8%.¹⁸ However, our reported results is higher than those reported in a study conducted in South Africa, 3.8%¹⁹ and lower than in a study conducted in Dera district, northwest Ethiopia, 69.8%.²⁰

The differences observed in the results could be attributed to variations in sample size, study design, and a range of cultural, environmental, and social factors, including dietary habits.²¹ The prevalence of stunting was higher among children under-5 (as shown in Figure 1), which is consistent with findings from Dessie town, northeast Ethiopia, where the prevalence was 36.8%,²² in Lesotho, 33.3%,²³ in Botswana, 38.7%,²⁴ in India, 35.5%,²⁵ and in Nepal, 36%.²⁶ However, this result was higher than that reported in China, 4.4%.²⁷ This could be attributed to China being one of the most developed countries globally, leading to better nutritional behaviors compared to Ethiopia.²⁸ Another possible explanation for this difference may be due to economic disparities. Households with limited income often struggle with inadequate financial resources, leading to inconsistent access to sufficient food to maintain an active and healthy lifestyle. This situation often leads to inconsistent eating patterns and a dependence on inexpensive,

calorie-dense foods that lack vital nutrients. As a result, this can cause undernutrition in the long run.²⁹ In this study, the prevalence of underweight among under-5 children was significantly high (see Figure 1), which is similar to the prevalence of underweight reported in studies in Tanzania 46.0%,³⁰ India, 36%,³¹ and Bangladesh, 32.8%.³²

However, the present outcome exceeds that of a study conducted in China (3.9%)³³ and earlier studies in Ethiopia (27.6%).²² The possible explanation for the variation in the prevalence of underweight could be differences in economic development and child feeding status.²⁸

In this study, the prevalence of wasting was higher among under-5 children, as shown in Figure 1. This finding is consistent with previous research conducted in Ethiopia, where the prevalence rates were reported to be 7.3% and 11.5%^{22,34}; India, 10.5%³⁵; Tanzania, 24.7%³⁰; and China, 4.0%.³³ However, the current prevalence of wasting was lower than the 28.5% reported in other studies conducted in India.³¹ The discrepancy in the prevalence of underweight, stunting, and wasting across studies could be due to feeding-related cultural practices rendered to children in complementary and deterring nourishing practices.^{36,37} Moreover, in the last 2 years, Ethiopia has undergone significant political liberalization alongside a notable increase in violent conflicts. This surge in violence has resulted in the displacement of the Amhara community, leading to increased poverty and undernutrition among the Amhara people, particularly affecting children.³⁸

In the final model, the analysis showed a significant association between undernutrition and several factors, such as latrine use, recent episodes of diarrhea, mothers' occupation and place of residence, water treatment practices before consumption, latrine ownership, sources of water, water storage conditions, methods of extracting water from storage, and mothers' handwashing habits after toilet use (see Table 3). Children of mothers with no formal education were more likely to be underweight, wasted, and stunted than children of educated mothers were. This result is supported by earlier research conducted globally.^{30,39–42}

The observed difference may result from educated mothers having the skills to care for their children effectively and higher incomes to support their families. Overall, education equips mothers with the knowledge, skills, and resources necessary to ensure their children receive proper nutrition and healthcare, reducing the risk of undernutrition.^{43–45}

The prevalence of underweight and wasting among under-children was higher among housewife mothers compared to those employed by the government. This result is consistent with earlier research.^{46,47} One possible reason for this difference is that mothers who are economically empowered in their careers can use their income to provide their children with nutritious food and access quality medical care to prevent illness.^{48,49} Another factor linked to underweight, wasting, and stunting in under-5 children was the source of water.

Table 3. Bivariable and multivariable logistic regression analysis of undernutrition with predictor variables among under-5 children in the Dabat district, Northwest Ethiopia, 2023.

PREDICTOR VARIABLES	OUTCOME VARIABLES					
	STUNTING		WASTING		UNDERWEIGHT	
	COR (95% CI)	AOR (95% CI)	COR (95% CI)	AOR (95% CI)	COR (95% CI)	AOR (95% CI)
Proper latrine utilization						
No	13.04 (7.664-22.22)	9.76 (3.29-15.71)*	13.04 (7.63-22.21)	0.80 (0.24-2.66)	2.93 (1.85-4.65)	0.78 (0.32-1.99)
Yes	1	1	1	1	1	1
Age of under-5 in a month						
<24	17.83 (0.98-26.76)				2.19 (1.27-3.79)	3.10 (1.60-5.98)
24-36	35 (0.98-47.98)				0.63 (0.86-1.58)	0.89 (0.49-1.83)
≥37	1	1	1	1	1	1
Mothers educational status						
Illiterate	47.67 (0.96-66.47)		22.48 (12.20-47.67)	0.59 (0.12-2.71)	3.10 (1.84-5.25)	0.62 (0.22-1.78)
Read and write	37.44 (0.45-53.19)		10.49 (2.25-38.85)	0.90 (0.22-3.70)	1.20 (0.57-2.53)	1.28 (0.32-5.02)
Primary education	23.77 (0.67-44.69)		10.12 (4.96-21.95)	1.65 (0.16-16.94)	1.70 (0.53-5.39)	1.04 (0.48-2.26)
Secondary education	18.22 (0.67-37.44)		7.84 (3.97-33.57)	0.94 (0.26-3.08)	1.00 (0.53-1.99)	0.47 (0.13-1.70)
Diploma and above (R)	1	1	1	1	1	1
Mother's occupation						
Housewife	20.74 (10.40-41.36)	4.62 (1.78-16.66)	15.73 (10.39-21.35)	5.53 (2.03-15.05)**	4.65 (2.11-10.23)	3.58 (1.27-10.06)*
Private work	3.55 (1.76-7.19)	1.67 (0.46-5.59)	4.55 (1.76-7.19)	1.88 (0.64-5.49)	1.64 (0.76-3.58)	3.42 (1.22-9.53)*
Government employee (R)	1	1	1	1	1	1
Residence						
Rural	9.84 (5.80-16.67)	2.98 (1.44-3.76)*	9.84 (5.80-16.67)	8.51 (3.35-22.16)*	3.26 (2.11-5.04)	6.07 (2.88-12.80)*
Urban (R)	1	1	1	1	1	1
Drinking water treatment before consumption						
No	3.67 (2.93-14.44)	4.65 (1.18-18.32)**	7.02 (3.97-12.42)	3.70 (1.03-13.49)*	2.85 (1.68-4.90)	1.03 (0.47-2.30)
Yes (R)	1	1	1	1	1	1

(Continued)

Table 3. (Continued)

PREDICTOR VARIABLES	OUTCOME VARIABLES					
	STUNTING		WASTING		UNDERWEIGHT	
	COR (95% CI)	AOR (95% CI)	COR (95% CI)	AOR (95% CI)	COR (95% CI)	AOR (95% CI)
Latrine ownership						
No latrine at all	5.42 (1.02-29.35)	4.37 (2.12-6.57)*	6.75 (4.95-12.98)	6.05 (1.19-30.78)**	4.06 (2.47-6.69)	0.90 (0.31-2.56)
Common latrine	1.91 (0.47-7.74)	2.28 (0.66-7.91)	3.39 (2.70-4.03)*	1.94 (0.52-7.29)	1.18 (0.68-2.28)	0.70 (0.31-1.54)
Private latrine (R)	1	1	1	1	1	1
Water sources						
Pipe (R)	1	1	1	1	1	1
Protected well or spring	7.05 (3.25-19.33)	6.40 (2.43-17.53)**	0.29 (0.15-0.57)	0.47 (0.13-1.73)	2.20 (1.29-3.77)	1.94 (1.07-3.75)*
Unprotected well or spring	9.32 (5.18-47.70)	9.22 (5.00-12.27)**	0.05 (0.02-0.07)	1.01 (0.22-4.50)	3.95 (2.22-7.05)	1.67 (0.53-5.31)
River	6.07 (5.46-15.59)	12.52 (5.48-17.08)**	1.27 (0.52-6.026)	1.04 (0.54-3.74)	17.75 (4.23-49.38)	5.06 (1.57-25.50)*
Proper water storage						
No	23.55 (11.95-46.39)	2.83 (1.07-3.12)*	23.55 (11.95-46.39)	3.81 (3.99-14.57)*	3.44 (2.00-5.97)	0.915 (0.33-2.50)
Yes (R)	1	1	1	1	1	1
Ways of taking water from storage						
Dipping	17.04 (9.67-30.02)	0.44 (0.15-1.25)	17.04 (9.67-30.01)	2.49 (1.92-6.71)*	4.02 (2.46-6.56)	1.85 (0.88-3.93)
Pouring	1	1	1	1	1	1
Trimming child fingernails						
Rarely	14.08 (6.94-228.56)	0.66 (0.19-2.29)	14.08 (6.94-28.56)	0.69 (0.20-2.34)	3.05 (1.64-5.66)	1.18 (0.49-2.84)
Sometimes	6.704 (2.92-15.36)	1.80 (0.62-5.18)	6.70 (2.92-15.36)	1.69 (0.60-4.71)	0.81 (0.44-1.48)	0.58 (0.28-1.23)
Always	1	1	1	1	1	1
Mothers hand washing after toilet						
Never	1.59 (0.81-3.12)	2.65 (0.76-7.83)	3.80 (1.73-8.32)	2.20 (1.91-5.53)*	2.92 (1.73-4.92)	2.51 (2.21-8.83)*
Sometimes	0.60 (0.30-1.18)	0.30 (0.22-3.45)	10.24 (5.77-18.18)	3.41 (1.53-9.34)*	2.02 (1.06-3.87)	1.51 (1.12-4.34)*
Always	1	1	1	1	1	1

Abbreviation: R, Reference.
 Hosmer and Lemeshow test for stunting = 0.743.
 Hosmer and Lemeshow test for wasting = 0.632.
 Hosmer and Lemeshow test for wasting = 0.347.
 *P < .05. **P < .01.

Participants who used water from a river, open well, or spring had higher prevalence of stunting, underweight, and wasting in comparison to those who used tap water. This finding is consistent with previous studies.^{50,51} Similarly, improper water storage or consumption of untreated water increased the risk of stunting, wasting, and underweight among under-5 children. This outcome aligns with findings from earlier studies conducted in Chad and Pakistan.⁵²⁻⁵⁵ This is likely due to the link between undernutrition and frequent diarrhea and intestinal infections, often caused by poor hygiene practices and limited access to clean water.^{56,57} Under-5 children whose mothers rarely or never washed their hands after using the toilet had higher rates of wasting and underweight compared to those whose mothers consistently washed their hands. This finding is consistent with previous research.⁵⁸ Frequent hand-washing reduces the spread of microorganisms in food and water, lowering the risk of children contracting infections and improving the overall health of under-5 children.⁵⁹

Those who used shared latrines or practiced open defecation had a higher risk of stunting, underweight, and wasting compared to individuals with private latrines, aligning with previous research findings.^{60,61} The inconsistency may stem from open defecation's impact on soil and water, harming health, especially among under-5 children in developing countries, through various pathways.⁴²

Another key predictor of undernutrition was recent diarrhea. Under-5 children who had diarrhea were more likely to experience stunting, wasting, and underweight compared to those without diarrhea. This aligns with previous research.^{62,63} This could be due to diarrhea, which can lead to nutrient deficiencies and electrolyte loss, causing stunting, wasting, and underweight in under-5 children.⁶⁴ Children in rural areas were at a higher risk of stunting, wasting, and underweight compared to those in urban areas.⁶⁵ One possible reason for this difference is that children in urban areas often come from educated households and have better access to medical care and resources that support a healthy lifestyle.⁶⁶

Limitations of the Study

Demonstrating a cause-and-effect relationship was not feasible due to the cross-sectional study design. Furthermore, an additional limitation of this research was the impact of postwar events on the data collection process.

Conclusion and Recommendation

The study found a higher prevalence of undernutrition among under-5 children compared to the 2019 Ethiopia Mini Demographic and Health Survey. This study recommended raising awareness about establishing private toilets, promoting water treatment, proper latrine use, and handwashing practices. Training on personal hygiene and economic support for households should also be provided. Healthcare providers should offer quality health services for under-5 children.

Further research is needed to explore specific nutrient deficiencies using laboratory methods.



Author Contributions

All the authors contributed equally.

Data Availability

The data available from the primary authors.

ORCID iDs

Asmamaw Maledo  <https://orcid.org/0000-0003-3491-3839>
Tsegaye Adane Birhan  <https://orcid.org/0000-0003-3006-6169>

REFERENCES

- World Health Organization. *Levels and Trends in Child Malnutrition: UNICEF*. World Health Organization; 2021.
- Maniragaba VN, Atuhaire LK, Rutayisire PC. Undernutrition among the children below five years of age in Uganda: a spatial analysis approach. *BMC Public Health*. 2023;23:390.
- Yiga P. *Child Growth and Determinant Factors: A Case Study of Burundi*. Unpublished dissertation Universiteit Gent. 2016.
- Asfaw M, Wondaferash M, Taha M, Dube L. Prevalence of undernutrition and associated factors among children aged between six to fifty nine months in Bule Hora district, South Ethiopia. *BMC Public Health*. 2015;15:41.
- Clark H, Coll-Seck AM, Banerjee A, et al. A future for the world's children? A WHO-UNICEF-Lancet commission. *Lancet*. 2020;395:605-658.
- Kinyoki DK, Berkley JA, Moloney GM, et al. Environmental predictors of stunting among children under-five in Somalia: cross-sectional studies from 2007 to 2010. *BMC Public Health*. 2016;16:654-659.
- John-Joy Owolade A, Abdullateef RO, Adesola RO, Olaloye ED. Malnutrition: an underlying health condition faced in sub Saharan Africa: challenges and recommendations. *Ann Med Surg*. 2022;82:1-3.
- Gudu E, Obonyo M, Omballa V, et al. Factors associated with malnutrition in children < 5 years in western kenya: a hospital-based unmatched case control study. *BMC Nutr*. 2020;6:33-37.
- Darsene H, Geleto A, Gebeyehu A, Meseret S. Magnitude and predictors of undernutrition among children aged six to fifty nine months in Ethiopia: a cross sectional study. *Arch Public Health*. 2017;75:29-11.
- Kassie GW, Workie DL. Determinants of under-nutrition among children under five years of age in Ethiopia. *BMC Public Health*. 2020;20:399-411.
- Sahiledengle B, Mwanri L, Petrucka P, et al. Determinants of undernutrition among young children in Ethiopia. *Sci Rep*. 2022;12:20945-21018.
- Forsido SF, Tsegaye NK, Tamiru D, Belachew T, Hensel O. Undernutrition and associated factors among children under 2 years of age in Jimma Zone, southwest Ethiopia. *J Public Health*. 2022;30:2911-2919.
- Mengesha DK, Merkeb Y. Prevalence of undernutrition and potential risk factors among children under 5 years of age in Amhara region, Ethiopia: evidence from 2016 Ethiopian demographic and health survey. *J Nutr Sci*. 2021;10:10-e22.
- Dipasquale V, Cucinotta U, Romano C. Acute malnutrition in children: pathophysiology, clinical effects and treatment. *Nutrients*. 2020;12:2413.
- Van den Broeck J, Willie D, Younger N. *The World Health Organization Child Growth Standards: Expected Implications for Clinical and Epidemiological Research*. Springer; 2009:247-251.
- Aladag Eyibilen A, Güven M, Atış, Erkokmaz. Role of oxidative stress in hearing impairment in patients with type two diabetes mellitus. *J Laryngol Otol*. 2009;123:957-963.
- Goodyear MD, Krleza-Jeric K, Lemmens T. The declaration of Helsinki. *BMJ*. 2007;335:624-625.
- Menalu MM, Bayleyegn AD, Tizazu MA, Amare NS. Assessment of prevalence and factors associated with malnutrition among under-five children in Debre Berhan town, Ethiopia. *Int J Gen Med*. 2021;14:1683-1697.
- Ekhloenetale M, Tudeme G, Onikan A, Ekhloenetale CE. Socioeconomic inequalities in hidden hunger, undernutrition, and overweight among under-five children in 35 sub-Saharan Africa countries. *J Egypt Public Health Assoc*. 2020;95:9.
- Habtamu Z. Prevalence of under-nutrition and its associated factors among 6-23 months old children of employed and unemployed mothers in urban settings of

- Dera District Northwest Ethiopia 2022: A Comparative cross-sectional study. 2022;10:1-10.
21. Bitew ZW, Alemu A, Ayele EG, Worku T. Dietary diversity and practice of pregnant and lactating women in Ethiopia: a systematic review and meta-analysis. *Food Sci Nutr*. 2021;9:2686-2702.
 22. Sewenet T, W/Selassie M, Zenebe Y, Yimam W, Woretaw L. Undernutrition and associated factors among children aged 6-23 months in Dessie Town, north-eastern Ethiopia, 2021: a community based cross-sectional study. *Front Pediatr*. 2022;10:1-11.
 23. Pitsoyame K. *Determinants of Under-Five Mortality in Lesotho*. North-West University; 2022.
 24. Mahgoub SE, Nnyepi M, Bandeke T. Factors affecting prevalence of malnutrition among children under three years of age in Botswana. *Afr J Food Agric Nutr Dev*. 2006;6:1-15.
 25. Kim R, Rajpal S, Joe W, et al. Assessing associational strength of 23 correlates of child anthropometric failure: an econometric analysis of the 2015-2016 National family health survey, India. *Soc Sci Med*. 2019;238:1-10.
 26. Adhikari RP, Shrestha ML, Acharya A, Upadhaya N. Determinants of stunting among children aged 0-59 months in Nepal: findings from Nepal demographic and health survey, 2006, 2011, and 2016. *BMC Nutr*. 2019;5:37-10.
 27. Wondimu H, Dejene K. Determinants of under-five malnutrition, significant changes, and policy implications in the Ethiopian demographic health survey, 2019. *Discov Sustain*. 2022;3:16.
 28. Zhang L, Gao J. Exploring the effects of international tourism on China's economic growth, energy consumption and environmental pollution: evidence from a regional panel analysis. *Renew Sustain Energy Rev*. 2016;53:225-234.
 29. Seligman HK, Laraia BA, Kushel MB. Food insecurity is associated with chronic disease among low-income NHANES participants. *J Nutr*. 2010;140:304-310.
 30. Mgongo M, Chotta NAS, Hashim TH, et al. Underweight, stunting and wasting among children in Kilimanjaro region, Tanzania; a population-based cross-sectional study. *Int J Environ Res Public Health*. 2017;14:509.
 31. Sukla P, Borkar A. Nutritional status of pre-school children [1-5 years] in rural area of Chhattisgarh state. *Int J Community Med Public Health*. 2018;5:2099-2103.
 32. Rahman SMJ, Ahmed NAMF, Abedin MM, et al. Investigate the risk factors of stunting, wasting, and underweight among under-five Bangladeshi children and its prediction based on machine learning approach. *PLoS One*. 2021;16:1-11.
 33. Li H, Yuan S, Fang H, et al. Prevalence and associated factors for stunting, underweight and wasting among children under 6 years of age in rural Hunan Province, China: a community-based cross-sectional study. *BMC Public Health*. 2022;22:483-512.
 34. Gelu A, Edris M, Derso T, Abebe Z. Undernutrition and associated factors among children aged 6-59 months living in slum areas of Gondar city, northwest Ethiopia: a cross-sectional study. *Pediatric Health Med Ther*. 2018;9:81-88.
 35. Murarkar S, Gothankar J, Doke P, et al. Prevalence and determinants of undernutrition among under-five children residing in urban slums and rural area, Maharashtra, India: a community-based cross-sectional study. *BMC Public Health*. 2020;20:1559-9.
 36. Fledderjohann J, Vellakkal S, Stuckler D. Breastfeeding, pregnant, and non-breastfeeding nor pregnant women's food consumption: A matched within-household analysis in India. *Sex Reprod Healthc*. 2016;7:70-77.
 37. Okidi L, Ongeng D, Muliro PS, Matofari JW. Disparity in prevalence and predictors of undernutrition in children under five among agricultural, pastoral, and agro-pastoral ecological zones of Karamoja sub-region, Uganda: a cross sectional study. *BMC Pediatr*. 2022;22:316-16.
 38. Yusuf S. Drivers of ethnic conflict in contemporary Ethiopia. *Institute for security studies monographs*. 2019;2019:v-46.
 39. Chirande L, Charwe D, Mbwana H, et al. Determinants of stunting and severe stunting among under-fives in Tanzania: evidence from the 2010 cross-sectional household survey. *BMC Pediatr*. 2015;15:165-213.
 40. Mohammad K, Kassab M, Gamble J, Creedy DK, Foster J. Factors associated with birth weight inequalities in Jordan. *Int Nurs Rev*. 2014;61:435-440.
 41. Keino S, Plasqui G, Ettyang G, van den Borne B. Determinants of stunting and overweight among young children and adolescents in sub-Saharan Africa. *Food Nutr Bull*. 2014;35:167-178.
 42. Rahman MHU, Malik MA, Chauhan S, et al. Examining the linkage between open defecation and child malnutrition in India. *Child Youth Serv Rev*. 2020;117:1-10.
 43. Black RE, Victora CG, Walker SP, et al. Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet*. 2013;382:427-451.
 44. Frost MB, Forste R, Haas DW. Maternal education and child nutritional status in Bolivia: finding the links. *Soc Sci Med*. 2005;60:395-407.
 45. Mirowsky J, Ross CE. *Education, Social Status, and Health*. Aldine Transaction; 2003.
 46. Fakir AM, Khan MW. Determinants of malnutrition among urban slum children in Bangladesh. *Health Econ Rev*. 2015;5:59-11.
 47. S Y, K I. Prevalence and determinants of undernutrition among school age slum children in Dhaka City, Bangladesh. *J Nutr Health Sci*. 2016;3:1.
 48. Ndukwu CI, Egbuonu I, Ulasi TO, Ebenebe JC. Determinants of undernutrition among primary school children residing in slum areas of a Nigerian city. *Niger J Clin Pract*. 2013;16:178-183.
 49. Negash C, Whiting SJ, Henry CJ, Belachew T, Hailemariam TG. Association between maternal and child nutritional status in Hula, rural southern Ethiopia: a cross sectional study. *PLoS One*. 2015;10:e0142301.
 50. Gebregyorgis T, Tadesse T, Atenafu A. Prevalence of thinness and stunting and associated factors among adolescent school girls in Adwa town, North Ethiopia. *Int J Food Sci*. 2016;2016:1-8.
 51. Altare C, Delbiso TD, Mutwiri GM, Koplow R, Guha-Sapir D. Factors associated with stunting among pre-school children in southern highlands of Tanzania. *J Trop Pediatr*. 2016;62:390-408.
 52. Morrison AL, Lewthwaite H, Houghton LA, et al. Child undernutrition in households with microbiologically safer drinking water and 'improved water' in Tanna, Vanuatu. *J Water Health*. 2020;18:416-429.
 53. Doocy S, Tappis H, Villemainot N, et al. Point-of-use water treatment improves recovery rates among children with severe acute malnutrition in Pakistan: results from a site-randomized trial. *Public Health Nutr*. 2018;21:3080-3090.
 54. Srivastava A, Mahmood SE, Srivastava PM, Shrotriya VP, Kumar B. Nutritional status of school-age children - a scenario of urban slums in India. *Arch Public Health*. 2012;70:8-8.
 55. Panigrahi A, Das SC. Undernutrition and its correlates among children of 3-9 years of age residing in slum areas of Bhubaneswar, India. *Sci World J*. 2014;2014:1-9.
 56. Harrison LH, Naidu TG, Drew JS, de Alencar JE, Pearson RD. Reciprocal relationships between undernutrition and the parasitic disease visceral leishmaniasis. *Rev Infect Dis*. 1986;8:447-453.
 57. Ulijaszek SJ. Relationships between undernutrition, infection, and growth and development. *Hum Evol*. 1996;11:233-248.
 58. Wie GT, Tsegaye D. Determinants of acute malnutrition among children aged 6-59 months visiting public health facilities in Gambella town, southwest Ethiopia: unmatched case-control study. *Nutr Diet Suppl*. 2020;12:147-156.
 59. Cumming O, Cairncross S. Can water, sanitation and hygiene help eliminate stunting? Current evidence and policy implications. *Matern Child Nutr*. 2016;12:91-105.
 60. Av S, Sharma R. Understanding the impact of open defaecation on child diarrhoea and nutrition indicators. *J Health Manag*. 2019;21:487-496.
 61. Singh SK, Srivastava S, Chauhan S. Inequality in child undernutrition among urban population in India: a decomposition analysis. *BMC Public Health*. 2020;20:1852-1915.
 62. Lima AA, Guerrant RL. Persistent diarrhea in children: epidemiology, risk factors, pathophysiology, nutritional impact, and management. *Epidemiol Rev*. 1992;14:222-242.
 63. Siddiqui FJ, Belayneh G, Bhutta ZA. Nutrition and diarrheal disease and enteric pathogens. In: Humphries DL, Vermund SH, eds *Nutrition and Infectious Diseases: Shifting the Clinical Paradigm*. Springer; 2021:219-241.
 64. Dodos J, Mattern B, Lapegue J, Altmann M, Aissa MA. Relationship between water, sanitation, hygiene, and nutrition: what do link NCA nutrition causal analyses say? *Waterlines*. 2017;36:284-304.
 65. Mbogori T, Kimmel K, Zhang M, Kandiah J, Wang Y. Nutrition transition and double burden of malnutrition in Africa: a case study of four selected countries with different social economic development. *AIMS Public Health*. 2020;7:425-439.
 66. De P, Chattopadhyay N. Effects of malnutrition on child development: evidence from a backward district of India. *Clin Epidemiol Glob Health*. 2019;7:439-445.