

## **Recanalisation of a congenital extrahepatic portosystemic shunt previously attenuated with cellophane banding in a cat**

Source: Journal of Feline Medicine and Surgery Open Reports, 4(2)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/2055116918804023>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



## **Recanalisation of a congenital extrahepatic portosystemic shunt previously attenuated with cellophane banding in a cat**

James Crowley, Timothy Foo, Lara Boland and Laurencie Brunel

*JFMS Open Rep* 2018; 4; 1–5. DOI: 10.1177/2055116918795717

The authors acknowledge that the article contains the incorrect details on page 2. The correct sentence is ‘The cat received 740 mgI/kg of an iso-osmolar iodinated intravenous contrast medium (Omnipaque; GE Healthcare) using a concentration of 370 mgI/ml at 2 ml/kg administered over 20 s via the cephalic vein by means of an automated injector (Empower CTA; EZ-EM).