



Histoplasma capsulatum, Toxoplasma gondii, Bartonella henselae and Bartonella clarridgeiae coinfections in an indoor-only Siamese cat

Authors: Fuller, Alexandra, Černá, Petra, Jost, Haley, Williams, Maggie, Shropshire, Sarah, et al.

Source: Journal of Feline Medicine and Surgery Open Reports, 8(2)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/20551169221118553>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



***Histoplasma capsulatum*, *Toxoplasma gondii*, *Bartonella henselae* and *Bartonella clarridgeiae* coinfections in an indoor-only Siamese cat**

Journal of Feline Medicine and Surgery Open Reports
1–6

© The Author(s) 2022

Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/20551169221118553

journals.sagepub.com/home/jfmsopenreports

This paper was handled and processed by the American Editorial Office (AAFP) for publication in *JFMS Open Reports*



Alexandra Fuller, Petra Černá , Haley Jost, Maggie Williams, Sarah Shropshire and Michael R Lappin

Abstract

Case summary A 6-year-old male castrated Siamese cat was referred for acute-onset blindness and mydriasis. Physical examination revealed serous retinal detachment with panuveitis and systemic hypertension. Abdominal ultrasound showed suspected dilation of the cisterna chyli and abdominal lymphadenopathy. Aspirates of mesenteric lymph nodes revealed intrahistiocytic yeast organisms with mild-to-moderate pyogranulomatous inflammation. Fungal culture and *ITS1* sequencing of the lymph node aspirates confirmed infection with *Histoplasma capsulatum*. PCR performed on whole blood was positive for *Bartonella henselae* and *Bartonella clarridgeiae*, and *Toxoplasma gondii* IgG and IgM antibodies were detected in serum. The cat was prescribed prednisolone (0.5 mg/kg PO q24h), itraconazole (10 mg/kg PO q24h), clindamycin (13 mg/kg PO q12h), amlodipine (0.625 mg PO q24h), prednisolone acetate 1% drops (q6h) and ophthalmic lubricant for both eyes (q6h). The cat was doing well at home after 2 weeks of prednisolone, itraconazole and clindamycin administration, and no abnormalities were detected on repeat abdominal ultrasound 3 months later. The retinal detachment showed mild-to-moderate improvement at that time, but vision was not regained.

Relevance and novel information This is the first report of coinfection of *H capsulatum*, *T gondii*, *B henselae* and *B clarridgeiae* in an indoor-only cat living in a *H capsulatum* non-enzootic area with no known travel history.

Keywords: Infectious; histoplasmosis; toxoplasmosis; bartonellosis

Accepted: 22 July 2022

Case description

A 6-year-old male castrated Siamese cat was referred for acute-onset blindness with suspicion of bilateral retinal detachment. The patient was previously healthy and had no other clinical signs apart from dental disease, with multiple extractions performed 7 weeks prior to presentation. The patient was adopted approximately 4 years previously from a rescue center in Colorado, had no known travel history and was a predominantly indoor-only cat with supervised access to a patio. There was another cat and two dogs in the household, which were all reported to be healthy.

On physical examination, the patient's vital signs were normal and an approximately 1 × 1 cm ulcerative gingival lesion where the left first mandibular molar contacted

a previous extraction site was noted. The remainder of the physical examination was unremarkable and the patient's weight was 5.4 kg. The cat was hypertensive (185 mmHg; Doppler) and ophthalmic examination showed bilateral serous retinal detachment with retinal and vitreal hemorrhage (Figure 1a), bilateral mydriasis (Figure 1b) and absent menace and pupillary light reflexes. The

Department of Clinical Sciences, Colorado State University, Fort Collins, CO, USA

Corresponding author:

Petra Černá MRCVS, AFHEA, CertAvdFB, Department of Clinical Sciences, Colorado State University, 300 W Drake Road, Fort Collins, CO 80523-1601, USA
Email: petra.cerna@colostate.edu



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons

Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

Downloaded From: <https://complete.bioone.org/journals/Journal-of-Feline-Medicine-and-Surgery-Open-Reports> on 13 Sep 2024

Terms of Use: <https://complete.bioone.org/terms-of-use>




Figure 1 (a) Retinal detachment with retinal and vitreal hemorrhage. (b) Bilateral mydriasis in a Siamese cat. (c) Panuveitis with hyphemic clot, mydriasis and retinal detachment in the left eye. (d) Cytology of the mesenteric lymph nodes showing intrahistiocytic yeast with mild-to-moderate pyogranulomatous inflammation with *Histoplasma capsulatum* (red arrow)

patient also had aqueous flare with keratic precipitates, with mild hyphema in the left eye (OS; Figure 1c). Tonometry (TonoVet) revealed intraocular pressures of 6 mmHg in the right eye (OD) and 18 mmHg OS.

A complete blood cell count (CBC) showed mild neutrophilia ($16.6 \times 10^3/\mu\text{l}$; reference interval [RI] 2.0–12) and basophilia ($400 \times 10^3/\mu\text{l}$ [RI 0–100]).¹ Serum biochemical panel showed hypocholesterolemia (82 mg/dl [RI 95–270]), hypophosphatemia (2.6 mg/dl [RI 3.0–6.0]), hypocalcemia (8.6 mg/dl [RI 9.2–11.1]), hypomagnesemia (1.8 mg/dl [RI 2.0–2.7]), hypoalbuminemia (2.5 g/dl

[RI 3.1–4.4]), low albumin:globulin ratio (0.64 [RI 0.8–1.6]) and hypokalemia (3.67 mEq/L [RI 3.7–5.4]). Total thyroxine was within normal limits, urinalysis was unremarkable, and the cat was seronegative for feline leukemia virus antigen and feline immunodeficiency virus antibodies (SNAP FeLV/FIV; IDEXX Laboratories).

Thoracic radiographs were unremarkable, but abdominal ultrasound showed suspected dilation of the cisterna chyli with mesenteric and iliac lymphadenopathy. Aspirates of the mesenteric lymph nodes revealed low-to-moderate numbers of intrahistiocytic yeast organisms

(round-to-ovoid, measuring approximately 2–4 μm), with a thin clear halo, containing a basophilic (often lateralized) internal structure and rarely displaying budding, with mild-to-moderate pyogranulomatous inflammation (low-to-moderate numbers of macrophages, and a slightly increased number of neutrophils and plasma cells; Figure 1d). The lymphoid population was mixed, with small-sized lymphocytes predominating and low numbers of intermediate-to-large lymphocytes. Fungal culture of the lymph node aspirates confirmed infection with *Histoplasma capsulatum*. The aspirates were also evaluated for DNA from *Bartonella* species and *Toxoplasma gondii* using conventional PCR assays (positive or negative results provided) available commercially at an American Association of Veterinary Laboratory Diagnosticians certified laboratory (Veterinary Diagnostic Laboratory, Colorado State University), which were negative. These tests are based on previously reported assays.^{2,3} However, DNA from *B henselae* and *Bartonella clarridgeiae* was amplified from whole blood. While *Bartonella* species IgG was not detected in serum, the *T gondii* titers for IgG and IgM were 1:256 and 1:64, respectively (Veterinary Diagnostic Laboratory, Colorado State University). Urine was submitted for quantitative detection of *Histoplasma* and *Blastomyces* species antigen (MiraVista Diagnostics), which were positive at 0.55 ng/ml (RI 0.20–20.00) and positive but below the limit of quantification (RI 0.2–14.7), respectively.

The cat was administered amlodipine (0.625 mg PO q24h [Amlodipine Besylate; Ascend]) to control hypertension, clindamycin (13 mg/kg PO q12h [Clindamycin Hydrochloride; Zoetis]) to treat the *T gondii* infection, generic itraconazole capsule (10 mg/kg PO q24h [Itraconazole; Amneal]) to treat the *H capsulatum* infection, prednisolone (0.5 mg/kg PO q24h [PrednisTab; LLOYD]) to reduce systemic inflammation and prednisolone acetate ophthalmic solution 1% (one drop OU q6h; SANDOZ) to reduce uveal inflammation and the likelihood of anterior lens luxation. An ophthalmic lubricant (one drop OU q6h [I-Drop Vet Plus; I-MED Animal Health]) was also used. Systolic blood pressure was normal the following day (140 mmHg; Doppler) and the cat was discharged to the owners.

Three weeks later, a serum biochemistry panel was performed to monitor liver enzyme activity due to itraconazole administration and was found to be unremarkable. There was a mild improvement in the retinal detachment, but uveitis persisted and there was no return of vision. Hyphema of the left eye resolved approximately 8 weeks following initial presentation (Figure 2). Systemic prednisolone was discontinued after 3 months, but the patient continued to be administered itraconazole, amlodipine and prednisolone acetate. There were no new physical examination findings, but the cat became



Figure 2 Resolution of the hyphema of the left eye approximately 8 weeks following initial presentation. Mild plaque of an old hemorrhage and fibrin on the anterior lens capsule is visible. There is also the presence of early fibrosis of the lens capsule in the left eye

severely hypertensive (215 mmHg; Doppler) despite therapy, and the amlodipine dose was increased (to 1.25 mg PO q24h). One week later, the physical examination was still normal and blood pressure was normal (125 mmHg; Doppler). No abnormalities were noted on the CBC and serum biochemistry panel. Repeat abdominal ultrasound showed resolution of previously suspected dilation of the cisterna chyli and abdominal lymphadenopathy. *Histoplasma* species antigen was not detected in the urine and PCR on whole blood for *Bartonella* species was negative. *T gondii* IgG (1:256) and IgM (1:128) were still detected in serum, and the *Bartonella* species IgG assay was still negative at that time. At the time of manuscript submission (10 months after diagnosis), the cat was still blind with absent menace OU and bilateral retinal detachment but otherwise systemically well.

Discussion

Retinal detachment in cats is commonly associated with systemic hypertension, hyperthyroidism and kidney disease, as well as infections such as mycobacterial infection, cryptococcosis and histoplasmosis.^{4,5} Chorioretinitis in cats has been commonly reported with toxoplasmosis, cryptococcosis and histoplasmosis.⁴ This case report describes serous retinal detachment and panuveitis in a predominantly indoor cat with systemic hypertension and evidence of coinfections with *H capsulatum*, *T gondii*, *B henselae* and *B clarridgeiae*. Owing to the detection of *H*

capsulatum by cytology of the mesenteric nodes, culture and urine antigen assay, the infection was confirmed, and this organism was most likely associated with some of the clinical manifestations in this case.⁶ However, whether the fungus was directly involved with the ocular findings is unclear as chorioretinitis was not apparent, and aqueous or vitreous paracentesis was not performed. Based on the blood pressure findings over time without detection of other common causes of hypertension, it is possible some of the ocular changes (retinal detachment) were related to essential hypertension that occurred concurrently. The role of *T gondii* and the two *Bartonella* species in the clinical findings is less clear as both these genera commonly infect cats without causing clinical signs of disease.

The most unique finding in this case is the documentation of *H capsulatum* in a primarily indoor cat in a state generally considered to be non-enzootic. *H capsulatum* is a soil-borne fungal pathogen that is endemic to the Mississippi and Ohio River valleys, and the route of transmission is primarily by inhalation of spores from soil, particularly those contaminated by bird and bat droppings.⁷ Histoplasmosis in indoor-only cats is less common than in cats with outdoor access; however, transmission to indoor-only cats through potted plants or unfurnished basements has been described.⁸ Nevertheless, the agent has been detected in Montana, another Rocky Mountain state,⁹ and so may be more common in this region of the USA than previously believed. It is also possible that the organism was present in potting soil or other materials such as wood shavings that originated in enzootic areas; this cat had supervised access to a patio and could have been exposed to the organism there. Another possibility is that the cat could have been infected prior to adoption, but since this was 4 years ago, it seems less likely. The weakly positive result of the *Blastomyces* species quantitative antigen was attributed to cross-reactivity with *Histoplasma* species antigen. As multiple drugs were used in the management of this case, it is hard to determine which was related to the clinical improvement. However, as the urine *H capsulatum* test became negative and the mesenteric lymphadenopathy resolved, it is likely that itraconazole had the expected treatment effect.

T gondii infection in cats in the Rocky Mountain states is common and can infect cats housed indoors if the owners feed undercooked meat or transport hosts enter the home and are ingested by the cats.¹⁰ Most cats that are infected with *T gondii* have no clinical signs of disease, but ocular manifestations with uveitis, iritis or chorioretinitis are common when illness occurs.^{11,12} As *T gondii* is not thought to be eliminated from the body after primary infection, the detection of IgG and IgM antibodies confirms the agent infected the cat described here, but it is impossible to determine the

role the agent played in the clinical findings. *T gondii* DNA was not amplified from the aspirates from the mesenteric nodes, suggesting it was not involved with that manifestation of disease. Amplification of *T gondii* DNA in aqueous humor collected by aqueous paracentesis could have been performed to further document the role of the agent in the ocular manifestations of disease but was not carried out in this case.¹¹ The *T gondii* IgG titer could have been present for years in this cat, but IgM generally is recognized during acute or activated disease, suggesting a potential role in the clinical findings. After treatment, IgM titers can increase transiently, likely from naive B-cell activation by *T gondii* antigens released from tissue cysts. Clindamycin is an appropriate drug for the treatment of clinical feline toxoplasmosis.^{13–15} However, it is unknown whether the positive treatment response in the cat described here related to the administration of this drug or the others that were prescribed. As clindamycin is a static drug, *T gondii* antibody titers do not necessarily decrease with successful therapy as the agent persists in tissues and recurrence is possible.¹⁴

Bartonella species are vector-transmitted, blood-borne, intracellular, Gram-negative bacteria that can cause disease in mammals, with cats being recognized as the main reservoirs for some species.^{16,17} While fleas are the most common vectors for both *Bartonella* species detected in this cat, current infestations were not detected.¹⁸ However, infection could have occurred previously, and low-grade bacteremia could have persisted for months. Some cats, dogs and people with *Bartonella* species infection have antibodies below the detectable limit of the assay, particularly if concurrent immunodeficiencies occur. *Bartonella* species DNA was not amplified from the aspirates from the mesenteric lymph nodes, suggesting that the agent was not involved with that manifestation of disease. As discussed for *T gondii*, aqueous paracentesis for *Bartonella* species PCR assay could have been used to further evaluate the role of *Bartonella* species in the ocular manifestations of disease.¹¹ While clindamycin is not generally used for treatment of *Bartonella* species infection in cats,¹⁹ this drug may have an effect against some strains of the bacterium. As culture and antibiotic susceptibility testing was not performed, whether this antibiotic had a treatment effect in this cat is unknown. However, *Bartonella* species DNA was not amplified from blood from a post-treatment sample, suggesting a possible clindamycin treatment effect. While doxycycline or quinolones may have been more effective for *Bartonella* species infection, we chose to start with clindamycin alone initially because of *T gondii* infection.²⁰

H capsulatum, *T gondii* and *Bartonella* species can infect otherwise healthy cats and coinfections with *T gondii* and *Bartonella* species are common in outdoor cats that hunt and have fleas.²¹ However, detection of all three

agents in one indoor cat seems unusual and could possibly suggest immune deficits may have been present. Persistence of *T gondii* IgM has been common in cats with immunodeficiency.¹⁴ These findings could suggest an inadequate B lymphocyte response in this cat. A primary immunodeficiency could be considered, although this is rare in cats and immune profiling is currently not commercially available for cats.²² CBC did not reveal any changes to the white blood cell counts in this patient; however, CBC does not provide information about white blood cell function. Functional assays can be used to further assess neutrophil and lymphocyte function, but these tests are only available in certain research laboratories.²² Primary immunodeficiencies, although rare, exist in cats. A recent paper reported an autosomal recessive leukocyte adhesion (CD18) deficiency in a cat.²³ Whether immune function deficits existed cannot be determined with the available data. However, the rapid response to therapy suggests that any immune deficits that may have occurred were minimal.

Conclusions

The patient was successfully treated with an antihypertensive agent, anti-inflammatory agents and antimicrobial agents, and at the time of writing, was being treated with itraconazole with no systemic signs of coinfections with *H capsulatum*, *T gondii* and *Bartonella* species. However, the patient failed to regain vision in either eye. This case report reinforces previous reports suggesting that even cases believed to be indoor-only without a pertinent history of exposure should be screened for infectious agents if deemed appropriate by the clinical presentation. In addition, more specific diagnostic tests to assess immunocompetency in cats are needed.

Conflict of interest The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding The authors received no financial support for the research, authorship, and/or publication of this article.

Ethical approval The work described in this manuscript involved the use of non-experimental (owned or unowned) animals. Established internationally recognized high standards ('best practice') of veterinary clinical care for the individual patient were always followed and/or this work involved the use of cadavers. Ethical approval from a committee was therefore not specifically required for publication in *JFMS Open Reports*. Although not required, where ethical approval was still obtained it is stated in the manuscript.

Informed consent Informed consent (verbal or written) was obtained from the owner or legal custodian of all animal(s) described in this work (experimental or non-experimental animals, including cadavers) for all procedure(s) undertaken (prospective or retrospective studies). No animals or people are

identifiable within this publication, and therefore additional informed consent for publication was not required.

ORCID iD Petra Černá  <https://orcid.org/0000-0002-4300-8534>

References

- 1 Acierno MJ, Brown S, Coleman AE, et al. **ACVIM consensus statement: guidelines for the identification, evaluation, and management of systemic hypertension in dogs and cats.** *J Vet Intern Med* 2018; 32: 1803–1822.
- 2 Jensen WA, Fall MZ, Rooney J, et al. **Rapid identification and differentiation of *Bartonella* species using a single-step PCR assay.** *J Clin Microbiol* 2000; 38: 1717–1722.
- 3 Burney DP, Lappin MR, Spilker M, et al. **Detection of *Toxoplasma gondii* parasitemia in experimentally inoculated cats.** *J Parasitol* 1999; 85: 947–951.
- 4 La Croix NC. **Ocular manifestations of systemic disease in cats.** *Clin Tech Small Anim Pract* 2005; 20: 121–128.
- 5 Meekins JM. **Acute blindness.** *Top Companion Anim Med* 2015; 30: 118–125.
- 6 Hanzlicek A, Meinkoth J, Renschler J, et al. **Antigen concentrations as an indicator of clinical remission and disease relapse in cats with histoplasmosis.** *J Vet Intern Med* 2016; 30: 1065–1073.
- 7 Kauffman CA. **Histoplasmosis: a clinical and laboratory update.** *Clin Microbiol Rev* 2007; 20: 115–132.
- 8 Reinhart JM, KuKanich KS, Jackson T, et al. **Feline histoplasmosis: fluconazole therapy and identification of potential sources of *Histoplasma* species exposure.** *J Feline Med Surg* 2012; 14: 841–848.
- 9 Skillman D, Riek L, Davis B, et al. **Histoplasmosis in a state where it is not known to be endemic – Montana, 2012–2013.** *MMWR Morb Mortal Wkly Rep* 2013; 62: 834–837.
- 10 Vollaire MR, Radecki SV and Lappin MR. **Seroprevalence of *Toxoplasma gondii* antibodies in clinically ill cats in the United States.** *Am J Vet Res* 2005; 66: 874–877.
- 11 Powell CC, McInnis CL, Fontenelle JP, et al. ***Bartonella* species, feline herpesvirus-1, and *Toxoplasma gondii* PCR assay results from blood and aqueous humor samples from 104 cats with naturally occurring endogenous uveitis.** *J Feline Med Surg* 2010; 12: 923–928.
- 12 Lappin M, Roberts S, Davidson M, et al. **Enzyme-linked immunosorbent assays for the detection of *Toxoplasma gondii*-specific antibodies and antigens in the aqueous humor of cats.** *J Am Vet Med Assoc* 1992; 201: 1010–1016.
- 13 Dubey J, Lindsay DS and Lappin MR. **Toxoplasmosis and other intestinal coccidial infections in cats and dogs.** *Vet Clin Small Anim Pract* 2009; 39: 1009–1034.
- 14 Lappin MR. **Update on the diagnosis and management of *Toxoplasma gondii* infection in cats.** *Top Companion Anim Med* 2010; 25: 136–141.
- 15 Lappin MR, Greene CE, Winston S, et al. **Clinical feline toxoplasmosis: serologic diagnosis and therapeutic management of 15 cases.** *J Vet Intern Med* 1989; 3: 139–143.
- 16 Álvarez-Fernández A, Breitschwerdt EB and Solano-Gallego L. ***Bartonella* infections in cats and dogs including zoonotic aspects.** *Parasites Vectors* 2018; 11. DOI:10.1186/s13071-018-3152-6.

- 17 Breitschwerdt EB. **Feline bartonellosis and cat scratch disease.** *Vet Immunol Immunopathol* 2008; 123: 167–171.
- 18 Guptill L. **Bartonellosis.** *Vet Microbiol* 2010; 140: 347–359.
- 19 Pendle S, Ginn A and Iredell J. **Antimicrobial susceptibility of *Bartonella henselae* using Etest methodology.** *J Antimicrob Chemother* 2006; 57: 761–763.
- 20 Brunt J, Guptill L, Kordick DL, et al. **American Association of Feline Practitioners 2006 panel report on diagnosis, treatment, and prevention of *Bartonella* spp. infections.** *J Feline Med Surg* 2006; 8: 213–226.
- 21 Chalkowski K, Wilson AE, Lepczyk CA, et al. **Who let the cats out? A global meta-analysis on risk of parasitic infection in indoor versus outdoor domestic cats (*Felis catus*).** *Biol Lett* 2019; 15. DOI: 10.1098/rsbl.2018.0840.
- 22 DeBey MC. **Primary immunodeficiencies of dogs and cats.** *Vet Clin Small Anim Pract* 2010; 40: 425–438.
- 23 Bauer TR Jr, Pratt SM, Palena CM, et al. **Feline leukocyte adhesion (CD 18) deficiency caused by a deletion in the integrin β 2 (ITGB 2) gene.** *Vet Clin Pathol* 2017; 46: 391–400.