

## **Small-Scale Trials Suggest Increasing Applications of Natular™ XRT and Natular™ T30 Larvicide Tablets May Not Improve Mosquito Reduction in Some Catch Basins**

Authors: Harbison, Justin E., Henry, Marlon, Corcoran, Peter C., Zazra, Dave, and Xamplas, Christopher

Source: Environmental Health Insights, 10(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/EHI.S36722>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Small-Scale Trials Suggest Increasing Applications of Natular™ XRT and Natular™ T30 Larvicide Tablets May Not Improve Mosquito Reduction in Some Catch Basins

Justin E. Harbison<sup>1,2</sup>, Marlon Henry<sup>2</sup>, Peter C. Corcoran<sup>2</sup>, Dave Zazra<sup>2</sup> and Christopher Xamplas<sup>2</sup>

<sup>1</sup>Department of Public Health Sciences, Loyola University Medical Center, Loyola University Chicago, Maywood, IL, USA. <sup>2</sup>North Shore Mosquito Abatement District, Northfield, IL, USA.

**ABSTRACT:** Stormwater catch basins are commonly treated with larvicides by mosquito control agencies to reduce local populations of mosquito species capable of transmitting West Nile virus. Recent evidence suggests that extended-release larvicides formulated to last up to 180 days in catch basins may not be effective in some basins due to chronic flushing, rapid dissolution, or burying of treatment in sump debris. To investigate if increasing the number of applications could improve effectiveness, a small study was performed over 13 weeks in 2015 to evaluate two extended-release larvicides (Natular™ XRT 180-day tablets and Natular™ T30 30-day tablets) and a larvicide oil (CocoBear™). Over the course of 13 weeks, three groups of eight basins were monitored for mosquitoes, each group receiving Natular™ XRT, Natular™ T30, or CocoBear™ larvicides. All basins received a single application at the beginning of the study period. Once mosquitoes in a basin surpassed the treatment threshold during weekly monitoring, an additional application of the associated larvicide was given to that basin. The number of applications during the study period ranged from 1 to 10 for CocoBear™ basins, 1 to 7 for T30 basins, and 2 to 8 for XRT basins. Overall, the average number of applications and the cost of larvicide per basin were 4.4 applications at \$0.66 per CocoBear™ basin, 4.4 applications at \$6.26 per T30 basin, and 4 applications at \$16.56 per XRT basin. Basins treated with XRT and T30 needed reapplications more often than expected, yet were no more effective than CocoBear™, suggesting that increasing the frequency of application of these larvicide formulations may not provide increased mosquito reduction in some basins.

**KEYWORDS:** larvicide, catch basins, mosquitoes, cost

**CITATION:** Harbison et al. Small-Scale Trials Suggest Increasing Applications of Natular™ XRT and Natular™ T30 Larvicide Tablets May Not Improve Mosquito Reduction in Some Catch Basins. *Environmental Health Insights* 2016;10:31–34 doi: 10.4137/EHI.S36722.

**TYPE:** Short Report

**RECEIVED:** October 20, 2015. **RESUBMITTED:** November 18, 2015. **ACCEPTED FOR PUBLICATION:** November 20, 2015.

**ACADEMIC EDITOR:** Timothy Kelley, Editor in Chief

**PEER REVIEW:** Five peer reviewers contributed to the peer review report. Reviewers' reports totaled 1009 words, excluding any confidential comments to the academic editor.

**FUNDING:** Authors disclose no external funding sources.

**COMPETING INTERESTS:** Authors disclose no potential conflicts of interest.

**CORRESPONDENCE:** jharbison@luc.edu

**COPYRIGHT:** © the authors, publisher and licensee Libertas Academica Limited. This is an open-access article distributed under the terms of the Creative Commons CC-BY-NC 3.0 License.

Paper subject to independent expert blind peer review. All editorial decisions made by independent academic editor. Upon submission manuscript was subject to anti-plagiarism scanning. Prior to publication all authors have given signed confirmation of agreement to article publication and compliance with all applicable ethical and legal requirements, including the accuracy of author and contributor information, disclosure of competing interests and funding sources, compliance with ethical requirements relating to human and animal study participants, and compliance with any copyright requirements of third parties. This journal is a member of the Committee on Publication Ethics (COPE).

Published by Libertas Academica. Learn more about this journal.

## Introduction

Stormwater catch basins are common sources of permanent or semipermanent standing water that are routinely targeted for mosquito larvicide applications by local mosquito control programs to reduce local mosquito populations and incidence of mosquito-borne diseases.<sup>1–16</sup> A common practice is to treat the stagnant water in all or most of the tens of thousands or sometimes hundreds of thousands of basins that can exist in an agency's operational area with at least one application of an extended-release larvicide formulated to last up to 150–180 days. Since the mid-1990s, extended-release larvicides (eg, Altosid® XR, FourStar®, and Natular™ XRT) have been used by the North Shore Mosquito Abatement District (NSMAD), a publicly funded mosquito control agency serving the northeast Chicago metropolitan area, to reduce mosquitoes in ~50,000 catch basins each season. Extended-release larvicide formulations are appealing logistically as a single treatment could theoretically reduce mosquito larvae and pupae for an entire season (ie, May through September). However, there is growing evidence that these products do not provide control for as long as expected.<sup>17–23</sup> The cause of the

control failures is not well understood but is likely related to the catch basins' structure, hydrology, and characteristics of the sediments and debris that they collect. Recently, analyses from three years of small-scale larvicide effectiveness trials conducted by NSMAD<sup>20</sup> and a year of district-wide catch basin monitoring<sup>21</sup> revealed that extended-release larvicide applications fail to control mosquitoes in a significant portion of basins. Because catch basins capture and retain runoff and debris, extended-release larvicides applied to these structures may be susceptible to being flushed out of basins, dissolving more rapidly than designed, or becoming buried in debris captured in the sump. All of these phenomena can influence the effectiveness of larvicides.

The objective of this project was to determine if additional applications of extended-release larvicides could improve the effectiveness of these products in basins in which flushing, rapid dissolution, or burying may occur. To do this, three mosquito larvicides, Natular™ XRT 180-day tablet (Clarke Mosquito Control Products, Inc.), Natular™ T30 30-day tablet (Clarke Mosquito Control Products, Inc.), and CocoBear™ larvicide oil (Clarke Mosquito Control Products, Inc.), were evaluated



during the 2015 season in catch basins located in the same small residential area used for previous larvicide trials.<sup>20</sup>

## Methods

All trials were conducted between June and September 2015 in catch basins located within a 0.7 km<sup>2</sup> residential area near the center of the NSMAD operational area. As the study area has been used in previous NSMAD catch basin studies, there were several years of data on the present study's catch basins to inform the present trials. Prior analyses found that extended-release larvicides failed to exhibit any control of mosquitoes over three consecutive years for 12 individual catch basins in the study area. Therefore, each of the three larvicides, XRT, T30, and CocoBear™, was randomly assigned to eight basins that included four of these *failure* basins and four in which larvicides appeared to be effective. XRT is formulated with 6.25% spinosad, T30 is formulated with 8.33% spinosad, and CocoBear™ is formulated with 10% mineral oil. CocoBear™ is not considered as an extended-release larvicide but was included for control purposes to compare the extended-release larvicides (one and three applications expected, respectively, for XRT and T30) with a larvicide in which weekly reapplications would be expected. The use of untreated basins as a more stringent control was not incorporated in this study over concerns of that these untreated basins could pose a risk to residents within and near the study area.

Beginning the first week of June and ending the first week of September, each of the 24 study basins was monitored for the presence of mosquito larvae and pupae by removing the circular grate of each structure with a manhole hook and taking two dip samples using a standard 350-mL dipper. Mosquitoes collected in a dip sample were counted, with the number of pupae and fourth instar larvae noted. All 24 basins received their respective larvicide treatments in the second week of monitoring. Each subsequent week, a basin was found to have reached or surpassed an average of 12 mosquito larvae and/or pupae per dip, and that basin received an additional treatment of its respective larvicide. The 12-mosquito threshold was found by dip samples taken from untreated basins in the study area from 2011 to 2014 that showed that the average number of mosquitoes from a total of 1,752 dip samples in untreated basins was 13.7 (95% confidence interval [CI] = 12.5–14.9). The total number (sum of two dips) of mosquitoes in basins across weeks was compared among the treatments using a repeated measures analysis of variance. The number and the total cost of larvicide applications were also recorded each week. In 2015 a single XRT tablet cost NSMAD \$4.14, \$1.43 for a T30 tablet, and estimated \$0.15 per single basin for CocoBear™. Precipitation data were downloaded from a nearby weather station of the National Oceanic and Atmospheric Administration's National Weather Service Forecast Office located at the Chicago O'Hare Airport (<http://www.nws.noaa.gov/climate/index.php?wfo=lot>).

## Results

All basins except two (one treated with CocoBear™ [CBa] and one with T30 [T30h]) surpassed the threshold for retreatment at least once (Table 1). The number of applications to a single basin during the study period ranged from 1 to 10 for CocoBear™ basins, 1 to 7 for T30 basins, and 2 to 8 for XRT basins (Table 1). Generally, there did not appear to be a large difference among the effectiveness of larvicide treatments each week (Table 2). The mean total mosquitoes (sum of two dips) and 95% CI were 17.13 (11.58–22.67) for CocoBear™ basins, 21.38 (15.05–27.70) for T30 basins, and 21.70 (14.97–28.43) for XRT basins over 12 weeks with active treatment. There was no statistical significance found between treatments:  $F(2, 21) = 0.32$ ,  $P = 0.73$ . The percentages of the fourth instars and pupae observed from the total mosquitoes found among the treatments each week ranged from 6.9% to 87.9% for CocoBear™, 9.0% to 78.2% for T30, and 9.2% to 75.3% for XRT. The total number of applications to the eight basins was 35 for CocoBear™ and T30 basins and 32 for XRT basins. Overall, the average number of applications and the cost of larvicide per basin were 4.4 applications at \$0.66 per CocoBear™ basin, 4.4 applications at \$6.26 per T30 basin, and 4 applications at \$16.56 per XRT basin.

## Discussion

Over the monitoring period, basins appeared to hold similar numbers of mosquitoes across the three larvicide treatments, suggesting, at least in the study basins, approximately equal effectiveness for the amount and type of larvicides used. Certainly, it appears that the use of the extended-release tablets (both XRT and T30) may not be the most cost-effective way to reduce mosquitoes in the study basins. Most extended-release basins needed more than the expected applications throughout the monitoring period, yet their effectiveness was no better than that of the oil. The use of the active ingredient (spinosad) of two extended-release pesticides for mosquito control, including the effect on nontarget species, has been described at length elsewhere.<sup>24–30</sup> Although larvicide oils are not formulated to be effective for longer durations, it did appear to be more cost effective than tablets in the study basins. The cost of labor for larvicide applications appears to be about the same for the three products as the total applications were similar. However, the cost of product is significantly lower for CocoBear™ per application, making it the most economical treatment.

Larvicide oils are designed to leave a thin film on the surface of treated water potentially killing larvae and pupae via contact or suffocation. However, because this film tends to break down quickly when the water surface is disturbed (eg, from an influx of rain and runoff), this type of larvicide is considered to be effective for a much shorter duration than the extended-release larvicides. Alternatively, the XRT and T30 tablets are formulated to slowly release their active ingredients over time, killing mosquito larvae and



**Table 1.** A comparison of the number of larvicide applications made to eight CocoBear™ oil-treated, eight Natular™ T30-treated, and eight Natular™ XRT-treated catch basins from June to September 2015. T indicates that the threshold for treatment ( $\geq 12$  mosquitoes per dip) was surpassed and an application of the associated larvicide to the basin was made.

BASIN	WEEK NUMBER													
	23	24	25	26	27	28	29	30	31	32	33	34	35	TOTAL
CocoBear Oil														
CBa	T													1
CBb	T		T				T		T			T	T	6
CBc	T							T						2
CBd	T											T	T	3
CBe	T		T		T	T	T	T		T	T	T	T	10
CBf	T							T						2
CBg	T			T				T				T	T	5
CBh	T						T	T		T		T	T	6
T30 30-day Tablets*														
T30a	T			T	T			T		T		T	T	7
T30b	T							T				T		3
T30c	T		T							T				3
T30d	T			T	T	T		T						5
T30e	T							T		T		T		4
T30f	T			T				T		T		T	T	6
T30g	T			T	T					T		T	T	6
T30h	T													1
XRT 180-day Tablets														
XRTa	T			T										2
XRTb	T						T					T		3
XRTc	T		T				T	T				T		5
XRTd	T						T			T		T		4
XRTe	T			T	T					T		T		5
XRTf	T		T	T	T	T		T				T	T	8
XRTg	T					T								2
XRTh	T							T				T		3

**Note:** \*The duration of T30 tablets is formulated to last four weeks, and three treatments would be expected for 13 weeks on weeks 23, 28, and 33.

pupae through contact or ingestion. Since the NSMAD was established in 1927 and until the mid-1990s, the larvicide oils were predominantly applied to the district's catch basins. Oils are somewhat easier to apply than the extended-release larvicides because often the applicator does not need to leave his or her vehicle to treat basins using a long nozzle to spray the larvicide into basins. As such, all NSMAD basins could usually receive three to four oil treatments in a season. However, our results suggest that three to four applications of oil, as well as T30 and XRT, may not be enough for some basins. Currently, it takes approximately two months or more for

NSMAD technicians to treat all basins with extended-release larvicides, and often most basins do not receive an additional treatment before the seasonal technicians become unavailable. Looking for longer lasting alternatives to extended-release tablets and more clearly identifying areas with a significant number of basins in which larvicides fail will be important next steps.

### Acknowledgment

The authors thank Roger S. Nasci for review and insightful comments on the initial draft of this article.



**Table 2.** A comparison of average mosquitoes per two dips  $\pm$  SE taken weekly from eight CocoBear™ oil-treated, eight Natular™ T30-treated, and eight Natular™ XRT-treated catch basins from June to September 2015 and weekly rainfall. All basins received a single application of their associated larvicide at week 23.

WEEK	COCOBEAR™	T30	XRT	RAINFALL (cm)
24	0.0 $\pm$ 0.0	0.06 $\pm$ 0.06	0.0 $\pm$ 0.0	8.33
25	4.88 $\pm$ 1.95	5.56 $\pm$ 1.81	4.5 $\pm$ 1.99	3.18
26	7.25 $\pm$ 4.17	20.31 $\pm$ 5.54	12.06 $\pm$ 3.93	2.01
27	3.06 $\pm$ 1.64	7.87 $\pm$ 2.63	9.94 $\pm$ 3.78	2.18
28	5.37 $\pm$ 2.27	7.31 $\pm$ 1.46	11.19 $\pm$ 3.18	5.05
29	6.13 $\pm$ 2.25	1.12 $\pm$ 0.45	11.5 $\pm$ 3.45	0.00
30	17.88 $\pm$ 4.96	18.88 $\pm$ 4.59	12.69 $\pm$ 3.03	1.35
31	4.44 $\pm$ 1.61	2.69 $\pm$ 0.96	1.19 $\pm$ 0.67	0.00
32	11.44 $\pm$ 3.93	18.63 $\pm$ 3.89	21.06 $\pm$ 7.47	0.94
33	3.63 $\pm$ 2.03	1.13 $\pm$ 0.64	1.31 $\pm$ 0.52	1.32
34	24.5 $\pm$ 6.07	27.81 $\pm$ 6.98	31.0 $\pm$ 5.1	1.88
35	14.19 $\pm$ 3.67	16.88 $\pm$ 4.6	13.75 $\pm$ 5.88	0.00

## Author Contributions

Conceived, designed, and performed the experiments: JEH, MH. Analyzed the data: JEH, MH. Wrote the first draft of the article: JEH, DZ, CX, PCC, MH. All authors reviewed and approved the final article.

## REFERENCES

- Lauret TH. Problems in the treatment of residential catch basins. *Calif Mosq Control Assoc.* 1953;22:73–4.
- Munstermann LE, Craig GB Jr. *Culex* mosquito populations in the catch basins of northern St. Joseph County, Indiana. *Proc Indiana Acad Sci.* 1977;86:246–52.
- Knepper RG, Leclair AD, Strickler JD, et al. Evaluation of methoprene (Altosid XR) sustained-release briquets for control of *Culex* mosquitoes in urban catch basins. *J Am Mosq Control Assoc.* 1992;8:228–30.
- McCarry MJ. Efficacy and persistence of Altosid pellets against *Culex* species in catch basins in Michigan. *J Am Mosq Control Assoc.* 1996;12:144–6.
- Raval-Nelson P, Soin K, Tolerud S. Analysis of *Bacillus sphaericus* in controlling mosquito populations in urban catch basins. *J Environ Health.* 2005;67:28–31.
- Rey JR, O'Meara GF, O'Connell SM, et al. Factors affecting mosquito production from stormwater drains and catch basins in two Florida cities. *J Vector Ecol.* 2006;31(2):334–43.
- Kobayashi M, Kasai S, Sawabe K, et al. Distribution and ecology of potential vector mosquitoes of West Nile Fever in Japan. *Global Environ Res.* 2008;12:27–33.
- Tedesco C, Ruiz M, McLafferty S. Mosquito politics: local vector control policies and the spread of West Nile Virus in the Chicago region. *Health Place.* 2010;16:1188–95.
- Anderson JF, Ferrandino FJ, Dingman DW, et al. Control of mosquitoes in catch basins in Connecticut with *Bacillus thuringiensis israelensis*, and spinosad. *J Am Mosq Control Assoc.* 2011;27:45–55.
- Poletti P, Messeri G, Ajelli M, et al. Transmission potential of Chikungunya virus and control measures: the case of Italy. *PLoS One.* 2011;6:e18860.
- Guidi V, Lüthy P, Tonolla M. Comparison between diflubenzuron and a *Bacillus thuringiensis israelensis*- and *Lysinibacillus sphaericus*-based formulation for the control of mosquito larvae in urban catch basins in Switzerland. *J Am Mosq Control Assoc.* 2013;29:138–45.
- Jackson MJ, Gow JL, Evelyn MJ, et al. Modelling factors that affect the presence of larval mosquitoes (Diptera: Culicidae) in stormwater drainage systems to improve the efficacy of control programmes. *Can Entomol.* 2013;145:674–85.
- Manrique-Saïde P, Arisqueta-Chablé C, Geded-Moreno E, et al. An assessment of the importance of subsurface catch basins for *Aedes aegypti* adult production during the dry season in a neighborhood of Merida, Mexico. *J Am Mosq Control Assoc.* 2013;29:164–7.
- Montarsi F, Martini S, Dal Pont M, et al. Distribution and habitat characterization of the recently introduced invasive mosquito *Aedes koreicus* [Hulecoetomyia koreica], a new potential vector and pest in north-eastern Italy. *Parasit Vectors.* 2013;6:292.
- Arana-Guardia R, Baak-Baak CM, Loroño-Pino MA, et al. Stormwater drains and catch basins as sources for production of *Aedes aegypti* and *Culex quinquefasciatus*. *Acta Trop.* 2014;134:33–42.
- Ocampo CB, Mina NJ, Carabali M, et al. Reduction in dengue cases observed during mass control of *Aedes* (Stegomyia) in street catch basins in an endemic urban area in Colombia. *Acta Trop.* 2014;132C:15–22.
- Stockwell PJ, Wessell N, Reed DR, et al. A field evaluation of four larval mosquito control methods in urban catch basins. *J Am Mosq Control Assoc.* 2006;22:666–71.
- Harbison JE, Henry M, Xamplas C, et al. A comparison of FourStar™ briquets and Natular™ XRT tablets in a North Shore suburb of Chicago, IL. *J Am Mosq Control Assoc.* 2014;30:68–70.
- Harbison JE, Henry M, Xamplas C, et al. Evaluation of *Culex pipiens* populations in a residential area with a high density of catch basins in a suburb of Chicago, Illinois. *J Am Mosq Control Assoc.* 2014;30:228–30.
- Harbison JE, Sinacore JM, Henry M, et al. Identification of larvicide-resistant catch basins from 3 years of larvicide trials in a suburb of Chicago, IL. *Environ Health Insights.* 2015;8(S2):1–7.
- Harbison JE, Layden JE, Xamplas C, et al. Observed loss and ineffectiveness of mosquito larvicides applied to catch basins in the northern suburbs of Chicago IL, 2014. *Environ Health Insights.* 2015;9:1–5.
- Harbison JE, Zazra D, Henry M, et al. Assessment of reactive catch basin larvicide treatments toward improved water quality using FourStar® briquets and CocoBear™ larvicide oil. *J Am Mosq Control Assoc.* 2015;31:283–5.
- Pawelek KA, Niehaus P, Salmeron C, et al. Modeling dynamics of *Culex pipiens* complex populations and assessing abatement strategies for West Nile Virus. *PLoS One.* 2014;9:e108452.
- Su T, Cheng ML. Cross resistances in spinosad-resistant *Culex quinquefasciatus* (Diptera: Culicidae). *J Med Entomol.* 2014;51:428–35.
- Su T, Cheng ML. Laboratory selection of resistance to spinosad in *Culex quinquefasciatus* (Diptera: Culicidae). *J Med Entomol.* 2014;51:421–7.
- Marina CF, Bond JG, Muñoz J, et al. Efficacy and non-target impact of spinosad, Bti and temephos larvicides for control of *Anopheles* spp. in an endemic malaria region of southern Mexico. *Parasit Vectors.* 2014;7:55.
- Jones OM, Ottea J. The effects of spinosad on *Culex quinquefasciatus* and three nontarget insect species. *J Am Mosq Control Assoc.* 2013;29:346–51.
- Lawler SP, Dritz DA. Efficacy of spinosad in control of larval *Culex tarsalis* and chironomid midges, and its nontarget effects. *J Am Mosq Control Assoc.* 2013;29:352–7.
- Su T, Cheng ML. Resistance development in *Culex quinquefasciatus* to spinosad: a preliminary report. *J Am Mosq Control Assoc.* 2012;28:263–7.
- Marina CF, Bond JG, Muñoz J, et al. Spinosad: a biorational mosquito larvicide for use in car tires in southern Mexico. *Parasit Vectors.* 2012;5:95.