

Estimation of Fitness of Normal and Stylopized Paddy Pest, White Leafhopper Cofana spectra (Distant) (Hemiptera: Cicadellidae), in West Bengal, India through Correlation of Life History Traits

Authors: Mitra, Sangita, Harsha, Rupa, Hazra, Niladri, and Mazumdar, Abhijit

Source: International Journal of Insect Science, 6(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/IJIS.S13029>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Estimation of Fitness of Normal and Stylopedized Paddy Pest, White Leafhopper *Cofana spectra* (Distant) (Hemiptera: Cicadellidae), in West Bengal, India through Correlation of Life History Traits

Sangita Mitra, Rupa Harsha, Niladri Hazra and Abhijit Mazumdar

Entomology Research Unit, Department of Zoology, The University of Burdwan, Burdwan, West Bengal, India.

ABSTRACT: The assessment of the morphological and reproductive features of white rice leafhopper *Cofana spectra* (Distant) was carried out using selected characters that bear importance in determining the fitness at the individual and population levels. Morphometric measurements of the individuals reared in the laboratory as normal and stylopedized with the strepsipteran parasitoid, *Halictophagus australensis* Perkins, were recorded and analyzed. A *t*-test was performed to justify whether parasitization by *H. australensis* affected the traits. Correlations and regression analyses were carried out to deduce the difference in relative importance of the morphological features in the life history of *C. spectra* and their variation because of stylopedization by *H. australensis*. A principal component analysis (PCA) was applied on the morphometric data to further substantiate the difference observed in the traits. In case of stylopedized white leafhopper (WLH), fecundity was inhibited almost completely irrespective of seasons. The effect of stylopedization on the life history traits of *C. spectra* has been noted that supports its possible use in biocontrol.

KEYWORDS: *Cofana spectra*, leafhopper, rice pest, parasitoid, Strepsiptera, *Halictophagus australensis*

CITATION: Mitra et al. Estimation of Fitness of Normal and Stylopedized Paddy Pest, White Leafhopper *Cofana spectra* (Distant) (Hemiptera: Cicadellidae), in West Bengal, India through Correlation of Life History Traits. *International Journal of Insect Science* 2014;6 33–42 doi:10.4137/IJIS.S13029.

RECEIVED: August 18, 2013. **RESUBMITTED:** January 28, 2014. **ACCEPTED FOR PUBLICATION:** January 29, 2014.

ACADEMIC EDITOR: Helen Hull-Sanders, Editor in Chief

TYPE: Original Research

FUNDING: Authors disclose no funding sources.

COMPETING INTERESTS: Authors disclose no potential conflicts of interest.

COPYRIGHT: © the authors, publisher and licensee Libertas Academica Limited. This is an open-access article distributed under the terms of the Creative Commons CC-BY-NC 3.0 License.

CORRESPONDENCE: abhijitau@rediffmail.com

Introduction

The white leafhopper (WLH) *Cofana spectra* (Distant) is a minor rice pest that rarely occurs at huge populations to cause yield loss.¹ WLH is widely distributed in the tropics from Africa; south and southeast Asian countries such as India, Indonesia, Malaysia, Philippines, Sri Lanka, and Taiwan; Pacific; and Australia.² It is very common in West Bengal and other regions of India where paddy cultivation is practiced.^{3,4} Both nymphs and adults are xylem feeders and suck sap from leaves.⁵ The paddy plants experience damage as a result of sap sucking and oviposition by *C. spectra* followed by secondary fungal and bacterial infection on the affected part. *C. spectra* is also a vector of pathogenic virus like rice yellow mottle virus (RYMV) in West Africa.⁶ The RYMV pathogen is included under the Sobemovirus group and is transmitted mechanically through rice plant injuries and by insect vectors.⁷ Infestation

of WLHs results in tillering diminution; moderate infestation causes loss of ear production and leaves turn brown and curl; and severe infestation results in the death of paddy plants.⁴ In Philippines, *C. spectra* co-occur with pest insects such as *Nephotettix virescens* (Distant), *Nephotettix nigropictus* (Stål), *Recilia dorsalis* (Motschulsky), *Sogatella furcifera* Horváth, and *Nilaparvata lugens* Stål, and the predatory *Cyrtorhinus* sp., coccinellids and spiders.⁸ Similar observations are made in Sri Lanka where among paddy pests, Cicadellidae (81.3%) dominates over Delphacidae (18.3%), and among Cicadellidae, *C. spectra* accounts for 14.2% in Yala—minor cropping season (April–August). A minor variation occurs in the Maha—major cropping season (October–February) with a higher percentage of *C. spectra*.⁹ The host of *Halictophagus australensis* Perkins has been confirmed as *C. spectra*. This endoparasitoid (Plates 1 and 2) is distributed in south-east Asia, Japan, Australia, and




Plate 1. Adult female of *Halictophagus australensis* (neotenic).



Plate 2. Adult male of *Halictophagus australensis*.

India.^{10,11} The percentage of stylopization of hoppers varies between 1 and 63% depending on the seasons.^{12,13} Stylopization of *C. spectra* by *Halictophagus spectrus* Yang in Malaysia was first reported by Hirashima and Kifune¹⁴ and Barrion and Litsinger.¹⁵ They listed several natural enemies attacking the hopper showing 11–50% variation of stylopization depending on different biological and environmental factors such as the host population, weather conditions, rice cultivars, age of the paddy plants, and time of occurrence.

Several morphological and reproductive features are viewed as life history traits and are useful parameters to gage the fitness of the individual insect.^{16,17} Life history traits are those that are associated with the survival and reproduction,¹⁸ which include growth rate; size at birth; age; and size at maturity, fecundity, and longevity; and frequency of reproduction. Variations in life history traits may be because of genotypes and their interactions with the environmental factors.¹⁹ Even if the genotypic variations are absent or minimal, the life history traits may be different quantitatively because of environmental impacts that are substantiated through observations in the insects of the orders Lepidoptera,²⁰ Hemiptera,²¹ and Diptera.²² Subsequently, life history traits differ quantitatively in a population bearing similar genotype but under different environmental regulations. Adjustment in the life history traits of an individual insect is obvious in response to varied abiotic factors for its survival. Alternations in environmental factors such as food, temperature (abiotic), and biotic factors such as crowding, predators, and parasitoids can provide useful insights to understand the importance of the life history traits in a pest species. For instance, the parasitoid *Elenchus tenuicornis* Kirby causes an effect on the morphology, anatomy, ethology, and mortality of its host *Javesella pellucida* (Fabricius).²³ Similarly, external genitalia are lost in the delphacid hosts, *S. furcifera* and *N. lugens*, because of stylopization by Elenchids and Drynids, respectively.²

In insects, fecundity is a positive function of the body size and longevity.²⁴ Longevity, on the other hand, depends on the feeding pattern and calorie content in the food.²² Variations in the life history traits are adaptive providing maximal fitness

to the species that can be explained in terms of the trade-offs in the life history traits.²⁵ The trade-offs are significant determinants for individual survival and maintenance of the population. Agricultural pests and their vectors face a continuous threat from environmental stochastic factors and the synthetic chemicals. Similarly, a parasitoid may induce a significant variation in these important life history traits in the host species.^{26,27} This is evident from the effects of parasitoid on the internal and external reproductive structures, as a manifestation of parasitoid infection in delphacid *J. pellucida*,²³ cicadellid *Ulopa reticulata* (Fabricius),^{28,29} and various other plants and leafhoppers.^{30,31} According to Kathirithamby,³² stylopization lengthens the life span of the host (macrynobionts), sex characters of stylopized male delphacid are lost, and the male resembles superficially a normal female but all other homopteran host families appear normal from outside but internal genitalia may be affected.³³

In course of development, value of one life history trait is compensated by the other, or the cost of one phenotypic expression is mitigated through the benefits of the other in terms of fitness value. In the present study, an attempt has been made to quantify the life history traits such as longevity, body size, and fecundity of the host, *C. spectra*. The effects of the strepsipteran parasitoid, *H. australensis*, on these life history traits have been compared. Such variations may predict the effect of the parasitoid on the host *C. spectra* parasitized by Strepsiptera and may be used to determine the potentiality for regulation of population of *C. spectra*.

Materials and Methods

Paddy seedlings of *Oryza sativa* Linnaeus were transplanted in earthen pots containing dried loamy soil and water up to 3 cm from the soil level with holes plugged with cotton to prevent drainage of water. The pots were placed on plastic plates containing water, and 5% 1,2,3,4,5,6-hexachlorocyclohexane (HCH) insecticide was sprinkled as dust around the plates to repel the predators such as ants and spiders. NPK was used as a fertilizer for luxuriant foliage growth of the paddy plants. As the plants grew about 30 cm, those were covered with mylar




Plate 3. White leafhopper *Cofana spectra* stylized by *Halictophagus australensis* (Hyperparasitism indicated by arrow).




Plate 4. White leafhopper *Cofana spectra* stylized by *Halictophagus australensis* (Indicated by arrow).

cages of PVC sheet, for protection from other insects. These cages were cylindrical and covered at one end with a nylon net of mesh size 0.47 mm^2 for aeration. Five to six seedlings were planted in the center of the tub, which was used for stock culture. Single plant was used for mating, oviposition, and emergence of nymph of *C. spectra*. WLHs of both sexes collected from the field were reared and multiplied successfully in the cages kept in the net house of the Department of Zoology, University of Burdwan. The nymphs hatched out from the stock culture were collected and released again in another plant. The nymph was transferred to different cages singly. In this way, WLH was raised in the net house and maintained during the experiment. Population of WLH became evident when air temperature was $26\text{--}32^\circ\text{C}$ and humidity $55\text{--}75\%$.

The *C. spectra* infected by endoparasitoid *H. australensis* (Plates 3 and 4) were released in a cage for getting triungulins. Sometimes, infected *C. spectra* kept singly in a glass cylinder with a bulbous region before the wider region ends. Both ends of the cylinders were covered with nylon net of mesh size 0.47 mm^2 for aeration. A small hole was made in the center of the net at the narrow end of each cylinder for insertion of rooted seedling of height 20 cm keeping the root outside, and the hole was then plugged with non-absorbent cotton. The experimental cylinders were then placed in an inclined position on a rack in water-filled tray in such a way as roots of seedlings were kept immersed in water. The seedling remained alive for four to five days, and was then replaced by a fresh one. The hoppers harboring strepsipteran female ready for emerging of triungulins (Plate 5) from eggs were released to the experimental cage for inoculation of the first and second instar nymphs of the captive host. Inoculation was found more in live hopper for its mobility. Morbid hoppers harboring female parasitoids kept in moist condition were noticed to release triungulins. As a result, a large number of triungulins came out for consecutive three days from live or morbid hoppers. As soon as free living and host-seeking triungulin comes in contact with a host, it normally enters the body via the abdominal cuticle, though

entry via the host's tarsi has recently been recorded.³⁴ Entry is gained through a combination of enzymatic (eg chitinase) and physical activity. Fluid is secreted from the mouth and appears to partially dissolve the host's cuticle. In this way, inoculation was carried out to obtain stylized hoppers.

Both normal and stylized adult hoppers after death were kept in 70% ethanol. The hoppers were processed and mounted on microslides following the method of DasGupta and Wirth.³⁵ Measurements of different parts of WLH were taken using microscope and micrometer, and are expressed in centimeter (cm).

The assessment of the morphological features of *C. spectra* was made using selected characters that bear importance in



Plate 5. Triungulin (First instar larva) of Strepsiptera *Halictophagus australensis*.

determining the fitness at the individual and population levels. Morphometric measurements of the individuals reared in the laboratory as normal and infected with the strepsipteran parasitoid, *H. australensis*, were recorded and analyzed. Correlations and regression analyses were performed to infer the difference in relative importance of the morphological traits in the life history of WLHs and their variation attributable to stylopization. A principal component analysis (PCA)³⁶ was carried out on the morphometric data to further validate the difference observed in the traits. The morphological features considered were as follows:

- For both male and female: wing length (WL), body length (BL), width of seventh sternum (WS), femur length (FL), leg length (LL)
- For male: aedeagus length (AL)
- For female: ovipositor length (OL).

An assessment of degree of dimorphism was made following Sharmila Bharathi et al³⁷ to deduce about the difference in value of the morphological feature toward a particular sex. A comparison was made between normal and infected individuals for each of these traits. A *t*-test³⁸ was applied to validate whether parasitization by *H. australensis* affected the traits. The null hypothesis was that the values for the normal and infected individuals for any trait are not significantly different from zero (H0: value for a trait expressed as a ratio of normal/infected = 1). As morphological traits are highly correlated,^{37,39} the data on all the morphological features were applied for PCA following Gotelli and Ellison.⁴⁰ The basis for using PCA is to reduce the number of initial observed variables and redundancy in the relationship. Because PCA is a variable reduction procedure, it is useful when the data are obtained on a number of variables (life history traits or morphological parameters) and there is some redundancy in these variables. In this case, redundancy means that some of the variables are correlated with one another, possibly because they are measuring the same construct (like the relationship between FL and WL might be similar to BL and WL, expressed in terms of

correlation coefficient as indicator). Because of this redundancy, it should be possible to reduce the observed variables into a smaller number of principal components (artificial variables) that will account for most of the variance in the observed variables. Thus, the method was applied to infer about the relationship between life history traits. The initial observed data are transformed into eigenvalues of the components/factors that indicate how much of the variance is explained by the components/factors using XLSTAT 2009 version.⁴¹ The relative importance of the variables toward their representation as a component is explained through the factor loadings. As the life history traits are highly correlated and add to fitness of organisms as a whole, an analysis to reduce the redundancy of the variables is essential to infer about the relative importance of these traits to the life history of the organisms.^{33,39,42}

Results

The morphometric features of both sexes of normal and infected *C. spectra* are presented in Table 1. The degree of dimorphism was prominent in the following traits: WL, BL, FL, LL, and WS, having a trend toward females. The observed values for each trait were higher in females than males, and thus the females are larger in size (Fig. 1). These differences were statistically significant at $P < 0.05$ level (Table 2). Significant difference in these morphological traits was observed between normal and infected individuals of *C. spectra* (Fig. 2, Table 3). In all instances, the ratio between normal and infected (*N/I*) traits was less than 1, indicating that the infected individuals had a higher measurement for these traits. These suggest that the parasitization of *C. spectra* favored positively all traits except LL in males and OL in females. The PCA was appropriate as revealed by the measure of sampling adequacy, which was >0.5 , and Bartlett's sphericity test for the justification of the PCA was significant in all cases (Tables 4–7). The results are presented in a tabular form in sequence of correlation matrix, eigenvalues, and factor loadings. For the normal individuals, a single factor was accountable for the observed variances in the data, whereas for the infected individuals, two factors

Table 1. The values (range, mean \pm SE) of different morphological traits of *C. spectra* for normal males: $n = 21$ and normal females: $n = 31$, and for infected males: $n = 12$ and infected females: $n = 21$.

WL	BL	WS	FL	LL	AL
Normal male					
6–6.6, 5.62 \pm 0.49	6.4–8, 7.29 \pm 0.42	1.04–1.28, 1.11 \pm 0.06	1.4–1.8, 1.49 \pm 0.11	5.8–7.86, 6.65 \pm 0.58	0.24–0.32, 0.28 \pm 0.03
Infected male					
5.4–7, 6.06 \pm 0.45	7.2–9.2, 7.91 \pm 0.56	1.04–1.6, 1.29 \pm 0.2	1.44–2, 1.67 \pm 0.15	5.8–7.4, 6.52 \pm 0.42	0.28–0.44, 0.35 \pm 0.06
WL	BL	WS	FL	LL	OL
Normal female					
5.4–7.8, 6.61 \pm 0.58	6.8–9, 7.93 \pm 0.5	1.28–1.84, 1.55 \pm 0.13	1.4–1.92, 1.68 \pm 0.14	6.08–7.52, 6.93 \pm 0.35	2.24–3.04, 2.64 \pm 0.21
Infected female					
5.6–7.8, 7.01 \pm 0.49	7.4–10, 8.68 \pm 0.7	1.44–2.12, 1.71 \pm 0.18	1.6–2.2, 1.86 \pm 0.16	6.6–7.92, 7.36 \pm 0.38	2.2–3, 2.57 \pm 0.22




Figure 1. The box plot representations of the degree of dimorphisms in the morphological traits of *C. spectra*.

Notes: When the values are greater than 1, it signifies that the trait is biased toward female. The blue marks are outliers, the red cross is the mean value, and the box represents the lower (25%) and upper (75%) quartiles with the mean values marked as line.

Abbreviations: WLDD, WL degree of dimorphism; BWDD, body weight degree of dimorphism; WSDD, WS degree of dimorphism; FLDD, FL degree of dimorphism; LLDD, LL degree of dimorphism.

explained the observed variation. It might be because of the low sample size for the infected males and females or the effect of parasitization that yielded two components. The ordination of variables along the component values is shown in Figures 3 and 4. For the normal individuals, a single factor was accountable for the observed variances in the data, whereas for the infected individuals, two factors explained the observed variation. It might possibly be because of the low sample size of the infected males and females or the effect of parasitization that yielded two components. From the PCA tables and the ordination of the variables, it can be deduced that the life history features are altered for both infected males and females of *C. spectra*. The factor loadings for the traits were different

Table 2. The results of *t*-test indicating that the degree of dimorphism is inclined toward females, because all values have a mean of >1 .

MORPHOLOGICAL TRAIT	WLDD	BLDD	WSDD	FLDD	LLDD
Normal <i>C. spectra</i>	5.059	5.378	12.673	5.935	2.989
Infected <i>C. spectra</i>	3.958	2.579	7.366	2.769	3.266

All *t*-values are significant at $P < 0.05$.

Abbreviations: WLDD, WL degree of dimorphism; BLDD, BL degree of dimorphism; WSDD, WS degree of dimorphism; FLDD, FL degree of dimorphism; LLDD, LL degree of dimorphism.




Figure 2. The difference between the morphological traits of *C. spectra* expressed as a ratio of the normal to infected individuals (normal males: $n = 21$; normal females: $n = 31$; infected males: $n = 12$; infected females: $n = 21$).

from normal and suggestive of the effect of parasitization by Strepsiptera. The relationship between the selected morphological traits is represented as regression equations (Table 8). The relationship was similar to that observed in other insects like mosquito *Culex pipiens quinquefasciatus*.¹⁶ It was noted that the OL and WL relationship, and LL and WL relationship were asymmetrical in normal compared to infected *C. spectra*. This is further suggestive of the effects of parasitization by *H. australensis*. The observation of the differences in these life history traits is indicative of differential fitness and thus reproductive success of *C. spectra*. As life history traits are indicators of the reproductive success and fitness, it is pertinent to evaluate the reproductive features of the pest *C. spectra* to infer about the relative contribution of these traits toward the event of reproduction and fecundity.

Table 3. Results of *t*-test on the deviation of the morphological traits of infected individuals from the mean values of normal individuals for respective sexes of *C. spectra*.

FEMALE TRAITS	WL	BL	WS	FL	LL	OL
<i>t</i> -value	2.623	4.377	3.299	3.872	4.121	1.442
MALE TRAITS	WL	BW	WS	FL	LL	AL
<i>t</i> -value	1.314	2.387	2.892	2.304	0.048	3.658

Values that are significant at $P < 0.05$ level are marked as bold.

Table values: for infected males, $t_{0.05(1),12} = 1.782$; for infected females, $t_{0.05(1),21} = 1.714$.

Table 4. The results of PCA of the morphological traits of normal male *C. spectra*. **A.** Bartlett sphericity test $\chi^2 = 84.246$, $df = 15$, $P < 0.001$. **B.** Eigenvalues and factor loadings for the morphometric variables. The highest factor loadings are marked in bold.

A					
VARIABLES	WL	BL	WS	FL	LL
BL	0.824				
WS	0.898	0.701			
FL	0.752	0.658	0.547		
LL	0.554	0.545	0.417	0.660	
AL	0.532	0.410	0.490	0.234	0.163
B					
	FACTOR 1		FACTOR 2		
WL	0.964		0.097		
BL	0.878		-0.017		
WS	0.860		0.235		
FL	0.816		-0.375		
LL	0.692		-0.528		
AL	0.553		0.705		
Eigenvalue	3.891		0.981		
Variability (%)	64.847		16.349		
Cumulative (%)	64.847		81.196		

Discussion

Analysis of morphological attributes indicates existence of degree of dimorphism in *C. spectra* having a tendency toward females as shown in Figure 1. This means that the females are larger in size and live for longer period of time than the males (Table 1). This will have important consequence as pest, where the females can be expected to inflict more damage than the males. Sexual size dimorphism is also an indicator of genetic variability.^{37,43} The difference in the morphological traits was also prominent between normal and infected individuals of *C. spectra*. For all these traits, the parasitized individuals had a higher value than the normal ones (Fig. 2). Morphometric studies on the external features of pest insects allow interpretation of the importance of morphological features as life history traits. Many of the morphological

Table 5. Results of PCA of the morphological traits of normal female *C. spectra*. Bartlett sphericity test $\chi^2 = 94.019$, $df = 15$, $P < 0.001$.

A. Correlation matrix (values in bold indicate significance at $P < 0.05$). **B.** Eigenvalues and factor loadings for the morphometric variables. The highest factor loadings are marked in bold.

A					
VARIABLES	WL	BL	WS	FL	LL
BL	0.485				
WS	0.665	0.349			
FL	0.823	0.631	0.549		
LL	0.534	0.156	0.103	0.529	
OL	0.495	0.501	0.222	0.592	0.164
B					
	FACTOR 1		FACTOR 2		
WL	0.912		0.163		
BW	0.707		-0.408		
WS	0.660		-0.160		
FL	0.939		0.063		
LL	0.534		0.793		
OL	0.670		-0.354		
Eigenvalues	3.382		0.977		
Variability (%)	56.374		16.289		
Cumulative (%)	56.374		72.664		

Table 6. Results of PCA of the morphological traits of infected male *C. spectra*. Bartlett sphericity test $\chi^2 = 42.09$, $df = 15$, $P < 0.001$.

A. Correlation matrix (values in bold indicate significance at $P < 0.05$). **B.** Eigenvalues and factor loadings for the morphometric variables. The highest factor loadings are marked in bold.

A					
VARIABLES	WL	BL	WS	FL	LL
BL	0.407				
WS	0.392	0.417			
FL	0.178	0.545	-0.086		
LL	0.158	0.714	0.219	0.892	
LA	-0.043	0.414	-0.334	0.628	0.658
B					
	FACTOR 1		FACTOR 2		
WL	0.337		0.664		
BL	0.824		0.341		
WS	0.182		0.872		
FL	0.884		-0.245		
LL	0.956		-0.059		
LA	0.714		-0.547		
Eigenvalue	3.030		1.680		
Variability (%)	50.494		27.997		
Cumulative (%)	50.494		78.491		



Table 7. Results of PCA of the morphological traits of infected female *C. spectra*. Bartlett sphericity test $\chi^2 = 60.21$, $df = 15$, $P < 0.001$.

A. Correlation matrix (values in bold indicate significance at $P < 0.05$. **B.** Eigenvalues and factor loadings for the morphometric variables. The highest factor loadings are marked in bold.

A					
VARIABLES	WL	BL	WS	FL	LL
BL	0.839				
WS	0.608	0.657			
FL	0.506	0.398	0.364		
LL	0.049	0.075	0.113	0.368	
OL	0.624	0.638	0.566	0.389	−0.061
B					
	FACTOR 1		FACTOR 2		
WL	0.896		−0.103		
BL	0.891		−0.140		
WS	0.797		−0.070		
FL	0.639		0.510		
LL	0.169		0.913		
OL	0.791		−0.261		
Eigenvalue	3.296		1.197		
Variability (%)	54.936		19.955		
Cumulative (%)	54.936		74.891		

features are considered as life history traits because of their contribution to the reproductive success of the species.^{16,22} Such morphometric studies are important from the perspective of assessing the importance of the pest and its damage potential.⁴⁴ Using multivariate analyses, the significance of morphological features has been further evaluated in Heteroptera,³³ Hymenoptera,⁴⁵ and Orthoptera.⁴⁶ In the context of the present study, the consequences of parasitization can be deduced from the deviation in the morphological features and life history traits of the hosts. In case of *C. spectra* parasitized by *H. australensis*, the changes in the morphological features were prominent. The WL, BL, and LL of stylopized individuals were increased compared to the normal ones. The OL in females and AL in males were also affected suggesting that parasitization affects the traits, which contribute to the reproductive success of WLH. The multivariate analyses revealed that the morphological traits are highly correlated and that the changes in one or the other character have correlated changes in rest of the features. For instance, regression analysis supports that a change in the OL in female *C. spectra* infected by *H. australensis* leads to a change in the relationship with the corresponding WL. Such changes can be deduced as an effect of induced response because of the parasitoid and can be viewed as phenotypically plastic trait.^{17,19,47,48} A trade-off in such characters²³ is prominent because the features like WL and LL are higher in infected *C. spectra* compared to OL. Thus, the morphological features that can be considered as life history traits are affected by endoparasitoid.

In the present study, the morphometric traits of WLHs were significantly affected by stylopization. The OL is reduced but the AL (Fig. 2) is incremented in stylopized *C. spectra*. In addition, the styles are also significantly different between normal and parasitized males. The morphological characters of nymphs and adults remain unaffected in case of stylopized *Nilaparvata* sp., but the sexual characters are affected because of parasitization.²⁶ The PCA analysis substantiates the differences in the normal and parasitized *C. spectra* at the level of morphological traits. It is reflected in the correlation coefficient and ordination along the component axis. This possibly suggests that the morphological variations occurred as a whole though individual differences remain obscured. Alternately, this difference may have been pronounced if larger samples were considered. The population level is expected to vary under field conditions as the seasonal variation in the fecundity of *C. spectra* is prominent. Although fecundity exhibited a positive relationship with the oviposition events and period, it varied with seasons, suggesting that the population level varied between the months.^{49,50} Thus, this observation supports that seasonal variation occurs in the relative abundance of *C. spectra*. Such variation in abundance of *C. spectra* is also observed in other parts of the world.^{8,51}

Life history theory depicts the pattern of longevity, reproduction, and development in the light of natural selection. It is also a theory of life cycle and fitness as shaped by natural selection.²⁰ In this view, the study of life history and related traits of pest insects is important to describe the population characteristics and possible trade-off between traits to infer about the possible damage potential.⁴⁴ In the present study, the effect of parasitization by *H. australensis* on the life history features of *C. spectra* has been noted that supports its use in biological control. The stylopized females rarely mate and oviposit. If they lay eggs, the clutch size is smaller than the normal one but the size and morphology of the eggs do not vary significantly. The average length of eggs of normal and stylopized females measures 1.08 mm ($n = 35$) and 1.12 mm ($n = 12$), and the hatchability is less than normal, which is in accordance with Cronin and Strong.⁵² Naturally, fecundity varies with environmental factors.⁵³ This is evident for fecundity of *C. spectra*, which remains high in February to March and October to November, and low in April to June. The number of offspring produced can be used as a surrogate to measure the fitness of an ovipositing female and the population as a whole.^{18,25,41,44} In case of stylopized *C. spectra*, fecundity is almost repressed irrespective of seasons demonstrating that the reproductive success is reduced by the strepsipteran parasitoid.

Stylopization reduces the pest status of the host.³¹ The biological control of leafhoppers using predators and parasitoids can be a viable alternative.⁵² For instance, Strepsiptera damage the sex organ, cause sterilization (reproductive death), and suppress the population of *C. spectra* in West Africa⁴⁹ and thus can serve as an agent to suppress pest population, along with predators like spiders.⁸ Thus, Strepsiptera can act as a biocontrol




Figure 3. A biplot showing the ordination of morphological traits of male *C. spectra*.

agent for the hopper acting as paddy pest. As reflected in the findings of the present study, *H. australensis* can affect the life history of *C. spectra* detrimentally and impart regulatory effect on the population growth. However, to prove the viability of *H. australensis* as potential biological control agents, further studies are required on the environmental factors and population interactions of *H. australensis* and *C. spectra*. Nonetheless, keeping in view the effects of synthetic pesticides on non-target species and costs associated with the use of pesticides, the Strepsiptera *H. australensis* is a better alternative because it is host specific and does not interfere with the functioning of the non-target organisms. Further, the effect of parasitoid can be retrospective because reproduction of pest is affected, indicating that it has a long-term effect on the population dynamics of the pest.

Acknowledgments

We are indebted to Professor P. K. Chaudhuri for rendering valuable suggestions and to the Head of the Department of Zoology, The University of Burdwan for providing DST-FIST-sponsored laboratory facility. We are also thankful to Dr. G. Aditya for the statistical work.

Author Contributions

Conceived and designed the experiments: AM, SM. Analyzed the data: SM. Wrote the first draft of the manuscript: NH. Contributed to the writing of the manuscript: RH. Agree with manuscript results and conclusions: SM, RH, NH, AM. Jointly developed the structure and arguments for the paper: AM, RH. Made critical revisions and approved final version: NH. All authors reviewed and approved of the final manuscript.




Figure 4. A biplot showing the ordination of morphological traits of female *C. spectra*.

Table 8. The regression equations using WL as explanatory variable for LL and AL of males, and LL and OL of females of normal and infected *C. spectra*.

NORMAL	INFECTED
Male	
LL(y) = 3.795 + 0.484WL(x) $r^2 = 0.307$; df = 1, 19; $F = 8.401$; $P < 0.009$	LL(y) = 5.424 + 0.202WL(x) $r^2 = 0.025$; df = 1, 10; $F = 0.254$; not significant
AL(y) = 0.125 + 0.03WL(x) $r^2 = 0.283$; df = 1, 19; $F = 7.494$; $P < 0.013$	AL(y) = 0.380 - 0.005WL(x) $r^2 = 0.002$; df = 1, 10; $F = 0.02$; not significant
Female	
LL(y) = 4.773 + 0.327WL(x) $r^2 = 0.285$; df = 1, 29; $F = 11.588$; $P < 0.001$	LL(y) = 7.09 + 0.04WL(x) $r^2 = 0.002$; df = 1, 22; $F = 0.052$; not significant.
OL(y) = 1.458 + 0.179WL(x) $r^2 = 0.245$; df = 1, 29; $F = 9.404$; $P < 0.005$	OL(y) = 0.598 + 0.0282WL(x) $r^2 = 0.389$; df = 1, 22; $F = 14.06$; $P < 0.001$

DISCLOSURES AND ETHICS

As a requirement of publication the authors have provided signed confirmation of their compliance with ethical and legal obligations including but not limited to compliance with ICMJE authorship and competing interests guidelines, that the article is neither under consideration for publication nor published elsewhere, of their compliance with legal and ethical guidelines concerning human and animal research participants (if applicable), and that permission has been obtained for reproduction of any copyrighted material. This article was subject to blind, independent, expert peer review. The reviewers reported no competing interests.

REFERENCES

1. Dale D. Insect pests of the rice plant—their biology and ecology. In: Heinrichs EA, ed. *Biology and Management of Rice Insects*, Manila, Philippines: Wiley Eastern Ltd., India & IRRI; 1994:363–487.
2. Heinrichs EA, Barrion AT. Rice-feeding insects and selected natural enemies in west Africa—biology, ecology, identification. Los Baños, Philippines: IRRI; Abidjan, Côte d'Ivoire: WARDA—The Africa Rice Center; 2004:243.
3. Pathak MD, Khan ZR. *Insect Pests of Rice*. Manila, Philippines: International Rice Research Institute; 1994.

4. Sam MD, Chelliah S. Biology on the white leafhopper on rice. *Int Rice Res Newsl.* 1984;9(1):22.
5. Wilson MR, Claridge MF. *Handbook for the Identification of Leafhoppers and Planthoppers of Rice.* Wallingford, Oxon, United Kingdom: CAB International for International Institute of Entomology in Association with Natural Resources Institute; 1991:x+142.
6. Nwilele FE, Traore AK, Asidi A, Sere NY, Onasanya A, Abo ME. New records of insect vectors of rice yellow mottle virus (RYMV) in Cote d'Ivoire, west Africa. *J Ent.* 2009;6:198–206.
7. Abo ME, Alegbejo MD, Sy AA. The insect vectors of rice yellow mottle virus: their mode of transmission and feeding effect on rice. *ESN Occas Publ.* 2000;32:83–90.
8. Litsinger JA, Canapi BL, Bandong JP, Lumaban MD, Raymundo FD, Barrion AT. Insect pests of rainfed wetland rice in the Philippines: population densities, yield loss, and insecticide management. *Int J Pest Manage.* 2009;55(3): 221–242.
9. Bambaradeniya CNB, Edirisinghe JP. Composition, structure and dynamics of arthropod communities in a rice agro-ecosystem. *Cey J Sci (Bio Sci).* 2008; 37(1):23–48.
10. Kathirithamby J. Description and biological notes of Halictophagidae (Strepsiptera) from Australia with a checklist of the world genera and species. *Invert Taxon.* 1992;6(1):159–196.
11. Mazumdar A, Chaudhuri PK. A catalogue of Strepsipteran species (Insecta) of India. *Proc Zool Soc.* 2004;57(1):43–46.
12. Miura T, Hirashima Y, Wongsiri T. Egg and nymphal parasites of rice leafhoppers and planthopper. A result of field studies in Thailand in 1977. *Esakia.* 1978;13:21–44.
13. Pena N, Shepard M. Seasonal incidence of parasitism of brown planthoppers, *Nilaparvata lugens* (Homoptera: Delphacidae), green leafhoppers *Nephotettix* spp. and whitebacked planthoppers, *Sogatella furcifera* (Homoptera: Cicadellidae) in Laguna Provinces, Philippines. *Environ Entomol.* 1986;15:263–267.
14. Hirashima Y, Kifune T. Strepsipterous parasites of Homoptera injurious to the rice plant in Sarawak, Borneo with description of a new species. *Esakia.* 1978;11:53–58.
15. Barrion AT, Litsinger JA. Parasites of the rice white leaf hopper *Cofana spectra* (Dist.) in the Philippines. *Int Rice Res Newsl.* 1983;8(6):20.
16. Agnew P, Haussy C, Michalakakis Y. Effects of density and larval competition on selected life history traits of *Culex pipiens quinquefasciatus* (Diptera: Culicidae). *J Med Entomol.* 2000;37(5):732–735.
17. Via S, Gomulkiewicz R, De Jong G, Scheiner SM, Schlichting CD, Vantienderen PH. Adaptive phenotypic plasticity: consensus and controversy. *Trends Ecol Evol.* 1995;10(5):212–217.
18. Stearns SC, Hoekstra RF. *Evolution an Introduction.* 2nd ed. Oxford University Press USA; 2005.
19. Gotthard K, Nylin S. Adaptive plasticity and plasticity as an adaptation: a selective review of plasticity in animal morphology and life history. *Oikos.* 1995;74(1):3–17.
20. Nylin S, Gotthard K. Plasticity in life-history traits. *Ann Rev Entomol.* 1998;43: 63–83.
21. Pfenning B, Poethke HJ. Variability in the life history of the water strider *Gerris lacustris* (Heteroptera: Gerridae) across small spatial scales. *Ecol Entomol.* 2006;31:123–130.
22. Agnew P, Mallorie H, Sidobre C, Michalakakis Y. A minimalist approach to the effects of density—dependent competition on insect life-history traits. *Ecol Entomol.* 2002;27:396–402.
23. Raatikainen M. Bionomics, enemies and population dynamics of *Javesella pel-lucida* (F) (Hom, Delphacidae). *Ann Agr Fenn.* 1967;6(2):1–147.
24. Ambruster P, Hutchinson RA. Pupal mass and wing length as indicators of fecundity in *Aedes albopictus* and *Aedes geniculatus* (Diptera: Culicidae). *J Med Entomol.* 2002;39(4):699–704.
25. Stearns SC. Trade-offs in life-history evolution. *Funct Ecol.* 1989;3:259–268.
26. Kathirithamby J. *Elenchus* sp. (Strepsiptera: Elenchidae), a parasitoid of *Nilaparvata lugens* (Stål) (Homoptera: Delphacidae) in Peninsular Malaysia. In: Heong KL, Lee BS, Lim TH, Teoh CH, Ibrahim Y, eds. *Proc Int Conf Plant Prot Tropics.* 1982; Malaysian Plant Protection Society, Kuala Lumpur, Malaysia: 349–361.
27. Jervis MA, Ellers J, Harvey JA. Resource acquisition, allocation and utilization of parasitoid reproductive strategies. *Annu Rev Entomol.* 2008;53:361–385.
28. Waloff N. The life history and description of *Halictophagus silwoodensis*. sp.n. (Strepsiptera) and its host *Ulopa reticulata* (Cicadellidae) in Britain. *Syst Ent.* 1981;6:103–113.
29. Waloff N. *Halictophagus silwoodensis* Waloff (Strepsiptera, Halictophagidae) and its host *Ulopa reticulata* (F.) (Auchenorrhyncha, Cicadellidae). *Acta Entomol Fenn.* 1981;38:52–53.
30. Kathirithamby J. Review of the order Strepsiptera. *Syst Ent.* 1989;14(1):41–92.
31. Kathirithamby J. Host parasitoid associations of Strepsiptera: anatomical and developmental consequences. *Int J Insect Morphol Embryol.* 1998;27(1):39–51.
32. Kathirithamby J. Host-parasitoid associations in Strepsiptera. *Ann Rev Entomol.* 2009;54:227–249.
33. David JR, Gibert P, Mignon-Grasteau S, et al. Genetic variability of sexual size dimorphism in a natural population of *Drosophila melanogaster*: an isofemale-line approach. *J Gen.* 2003;82:79–88.
34. Kathirithamby J. Stand tall and they still get you in your Achilles foot-pad. *Proc Biol Sci.* 2001;268:2287–2289.
35. DasGupta SK, Wirth WW. Revision of oriental species of *Stilobezzia* Kieffer (Diptera, Ceratopogonidae). *Bull U S Nat Mus.* 1968;283:1–164.
36. Manly BFF. *Multivariate Statistical Methods: A Primer.* 2nd ed. London: Chapman and Hall; 1994.
37. Sharmila Bharathi N, Prasad NG, Mallikarjun S, Joshi A. Correlates of sexual dimorphism for dry weight and development time in five species of *Drosophila*. *J Zool.* 2004;264:87–95.
38. Zar JH. *Biostatistical Analysis.* 4th ed. India: Pearson Education; 1999.
39. Daly HV. Insect morphometrics. *Ann Rev Entomol.* 1985;30:415–438.
40. Gotelli NJ, Ellison AM. *A Primer of Ecological Statistics.* Sunderland, MA, USA: Sinauer Associates; 2004:479.
41. Addinsoft. *XLSTAT, Version 2009.* France: Addinsoft; 2009.
42. Rubin-de-Celis VE, Gassen DN, Callegari-Jacques SM, Valente VLS, Oliveira AK. Morphometric observations on three populations of *Schizaphis graminum* (Rondani), a main wheat aphid pest in Brazil. *Ana Soc Entomol Brasil.* 1997;26(3):417–428.
43. Iglesias MS, Gaspe MS, Valverde CA. A longitudinal study of two species of *Belostoma* Latreille (Heteroptera: Belostomatidae): allometry and ontogeny. *Neotrop Entomol.* 2008;37(6):662–667.
44. Nylin S. Life history perspectives on pest insects: What's the use? *Austral Ecol.* 2001;26:507–517.
45. Chaizawong T, Deowanish S, Wongsiri S, Sylvester HA, Rinderer TH, DeGuzman L. Multivariate morphometric study of *Apis florea* in Thailand. *J Apic Res.* 2004;43(3):123–127.
46. Nespolo RF, Castañeda LE, Roff DA. Dissecting the variance-covariance structure in insect physiology: the multivariate association between metabolism and morphology in the nymphs of the sand cricket (*Gryllus firmus*). *J Insect Physiol.* 2005;51:913–921.
47. Stearns SC. The evolutionary significance of phenotypic plasticity. *BioScience.* 1989;39(7):436–445.
48. Dewitt TJ, Sih A, Wilson DS. Costs and limits of phenotypic plasticity. *Trends Ecol Evol.* 1998;13(2):77–81.
49. Hafez M. Seasonal fluctuations of population density of the cabbage aphid, *Brevicoryne brassicae* (L.), in the Netherlands, and the role of its parasite, *Aphidius (Diaeretiella) rapae* (Curtis). *Tijdschr Plantenziekten Ned.* 1961;67:445–548.
50. Satpathi CR, Katti G, Prasad YG. Effect of seasonal variation on life table of brown plant hopper *Nilaparvata lugens* Stål on rice plant in eastern India. *Middle-East J Sci Res.* 2011;10(3):370–373.
51. Oyediran O, Ndongidila A, Heinrich EA. Strepsipteran parasitism of white leafhoppers, *Cofana* spp. (Hemiptera: Cicadellidae) in lowland rice in Côte d'Ivoire. *Int J Pest Manage.* 2000;46(2):141–147.
52. Cronin JT, Strong DR. Parasitoid interactions and their contribution to the stabilization of Auchenorrhyncha populations. In: Denno RF, Perfect TJ, eds. *Planthoppers: Their Ecology and Management.* New York: Chapman and Hall; 1994:400–428.
53. Baldrige RS, Blocker HD. Parasites of leafhoppers (Homoptera: Cicadellidae) from Kansas grassland. *J Kansas Entomol Soc.* 1980;53(2):441–447.