

Phytochrome Signaling Mechanisms

Authors: Li, Jigang, Li, Gang, Wang, Haiyang, and Wang Deng, Xing

Source: The Arabidopsis Book, 2011(9)

Published By: The American Society of Plant Biologists

URL: <https://doi.org/10.1199/tab.0148>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

First published on August 29, 2011: e0148. doi: 10.1199/tab.0148

This chapter is an updated version of a chapter originally published on July 6, 2004, e0074.1. doi: 10.1199/tab.0074.1

Phytochrome Signaling Mechanisms

Jigang Li^{a,b}, Gang Li^b, Haiyang Wang^b, and Xing Wang Deng^{a,b,1}

^aPeking-Yale Joint Center for Plant Molecular Genetics and Agro-biotechnology, State Key Laboratory of Protein and Plant Gene Research, College of Life Sciences, Peking University, Beijing 100871, China

^bDepartment of Molecular, Cellular, and Developmental Biology, Yale University, New Haven, Connecticut, 06520-8104

¹Address correspondence to xingwang.deng@yale.edu

Phytochromes are red (R)/far-red (FR) light photoreceptors that play fundamental roles in photoperception of the light environment and the subsequent adaptation of plant growth and development. There are five distinct phytochromes in *Arabidopsis thaliana*, designated phytochrome A (phyA) to phyE. phyA is light-labile and is the primary photoreceptor responsible for mediating photomorphogenic responses in FR light, whereas phyB-phyE are light stable, and phyB is the predominant phytochrome regulating de-etiolation responses in R light. Phytochromes are synthesized in the cytosol in their inactive Pr form. Upon light irradiation, phytochromes are converted to the biologically active Pfr form, and translocate into the nucleus. phyB can enter the nucleus by itself in response to R light, whereas phyA nuclear import depends on two small plant-specific proteins FAR-RED ELONGATED HYPOCOTYL 1 (FHY1) and FHY1-LIKE (FHL). Phytochromes may function as light-regulated serine/threonine kinases, and can phosphorylate several substrates, including themselves *in vitro*. Phytochromes are phosphoproteins, and can be dephosphorylated by a few protein phosphatases. Photoactivated phytochromes rapidly change the expression of light-responsive genes by repressing the activity of CONSTITUTIVE PHOTOMORPHOGENIC 1 (COP1), an E3 ubiquitin ligase targeting several photomorphogenesis-promoting transcription factors for degradation, and by inducing rapid phosphorylation and degradation of Phytochrome-Interacting Factors (PIFs), a group of bHLH transcription factors repressing photomorphogenesis. Phytochromes are targeted by COP1 for degradation via the ubiquitin/26S proteasome pathway.

INTRODUCTION

As sessile organisms, plants have acquired a high degree of developmental plasticity to optimize their growth and reproduction in response to their ambient environment, such as light, temperature, humidity, and salinity. Plants utilize a wide range of sensory systems to perceive and transduce specific incoming environmental signals. Light is one of the key environmental signals that influences plant growth and development. In addition to being the primary energy source for plants, light also controls multiple developmental processes in the plant life cycle, including seed germination, seedling de-etiolation, leaf expansion, stem elongation, phototropism, stomata and chloroplast movement, shade avoidance, circadian rhythms, and flowering time (Deng and Quail, 1999; Wang and Deng, 2003; Jiao et al., 2007).

Plants can monitor almost all facets of light, such as direction, duration, quantity, and wavelength by using at least four major classes of photoreceptors: phytochromes (phys) primarily responsible for absorbing the red (R) and far-red (FR) wavelengths (600-750 nm), and three types of photoreceptors perceiving the blue (B)/ultraviolet-A (UV-A) region of the spectrum (320-500 nm): cryptochromes (crys), phototropins (phot), and three newly recognized LOV/F-box/Kelch-repeat proteins ZEITLUPE (ZTL), FLAVIN-BINDING KELCH REPEAT F-BOX (FKF), and LOV KELCH REPEAT PROTEIN 2 (LKP2). In addition, UV RESISTANCE LOCUS 8 (UVR8) was recently shown to be a UV-B (282-320

nm) photoreceptor (Rizzini et al., 2011). These photoreceptors perceive, interpret, and transduce light signals, via distinct intracellular signaling pathways, to modulate photoresponsive nuclear gene expression, and ultimately leading to adaptive changes at the cell and whole organism levels.

The past two decades have seen dramatic progress in molecular characterization and understanding of the photobiology and photochemistry of the phytochrome photoreceptors in higher plants. This chapter aims to highlight some of the most recent progress in elucidating the molecular, cellular and biochemical mechanisms of phytochrome signaling in *Arabidopsis*. Interested readers are encouraged to read the accompanying reviews on other related subjects, such as photomorphogenesis (Nemhauser and Chory, 2002), cryptochromes (Yu et al., 2010), phototropins (Pedmale et al., 2010), and the circadian clock (McClung et al., 2002).

PLANT PHYTOCHROMES

The Discovery and Action Modes of Phytochromes

The term phytochrome, meaning “plant color”, was originally coined to describe the proteinous pigment that controls photoperiod detection and floral induction of certain short-day plants (such as cocklebur and soybean) (Garner and Allard, 1920), and

the reversible seed germination of lettuce (c.v. Grand Rapids) by R and FR light (Borthwick et al., 1952). R light promotes seed germination, whereas subsequent FR light treatment abolishes R light induction of seed germination. The germination response of lettuce seeds repeatedly treated with R/FR cycles is determined by the last light treatment. Thus R/FR photoreversibility is a characteristic feature of this response. In addition, the law of reciprocity applies to this response, i.e. the response is dependent on the total amount of photons received irrespective of the duration of light treatment.

Over the years, three action modes for phytochromes have been defined, i.e. low-fluence responses (LFRs), very-low-fluence responses (VLFRs) and high-irradiance responses (HIRs) (Table 1). The above-mentioned F/FR reversible response is characteristic of LFRs. LFRs also induce other transient responses, such as changes in ion flux, leaf movement, chloroplast rotation, and changes in gene expression (Haupt and Hader, 1994; Roux, 1994; Vince-Prue, 1994). VLFRs are activated by extremely low light intensities of different wavelengths (FR, R and B); examples include light-induced expression of the light-harvesting chlorophyll *a/b*-binding protein (*LHCB*) gene and light induction of seed germination. HIRs depend on prolonged exposure to relatively high light intensities, and are primarily responsible for the control of seedling de-etiolation (e.g. inhibition of hypocotyl elongation and promotion of cotyledon expansion) under all light qualities (Mustilli and Bowler, 1997; Casal et al., 1998; Neff et al., 2000; Table 1).

Chromophores and Two Reversible Forms of Phytochromes

Photoreversibility occurs because phytochromes exist as two distinct but photoreversible forms *in vivo*: the R light-absorbing form (Pr) and the FR light-absorbing form (Pfr). The Pr form absorbs maximally at 660 nm, whereas the Pfr form absorbs maximally at 730 nm (Quail, 1997a; Figure 1). The Pfr forms of phytochromes are generally considered to be the biologically active forms. It should be noted that in addition to their maximal absorptions of R and FR wavelengths, phytochromes also weakly absorb B light (Furuya and Song, 1994; Figure 1).

Phytochromes are soluble proteins and exist as homodimers. The molecular mass of the apoprotein monomer is approximately 125 kDa. Phytochrome apoproteins are synthesized in the cytosol, where they assemble autocatalytically with a linear tetrapyrrole chromophore, phytochromobilin (PΦB). The synthesis of PΦB is accomplished by a series of enzymatic reactions in the plastid that begins with 5-aminolevulinic acid (Figure 2A). The early steps in the PΦB pathway are shared with chlorophyll and heme biosynthesis. The committed step is the oxidative cleavage of heme by a ferredoxin-dependent heme oxygenase (HO) to form biliverdin IX (BV). BV is subsequently reduced to 3Z-PΦB by the enzyme PΦB synthase. Both 3Z-PΦB and its isomerized form 3E-PΦB can serve as functional precursors of the phytochrome chromophore. PΦB is then exported to the cytosol, where it binds to the newly synthesized apo-PHYs to form holo-PHYs (Terry, 1997; Figure 2A). The chromophore is attached via a thioether linkage to an invariant cysteine in a well-conserved domain among all phytochromes (see below).

The intrinsic photochemical activity of the chromophore prosthetic group allows phytochromes to convert between the two




Figure 1. Absorption spectra of phytochromes. Absorption spectra of the two forms (Pr and Pfr) of phytochromes. The Pr form absorbs maximally at 660 nm, while the Pfr form absorbs maximally at 730 nm. The visible light range of the human eye is approximately 380-700 nm. The light spectrum was adapted from Kami et al. (2010). Reprinted with permission from Elsevier.

Table 1. Diagnostic Features of Different Phytochrome Action Modes

Action Mode	Fluence Requirements	Photoreversibility	Reciprocity
VLFR	0.1 $\mu\text{mol}/\text{m}^2$ - 1 $\mu\text{mol}/\text{m}^2$	No	Yes
LFR	1 - 1000 $\mu\text{mol}/\text{m}^2$	Yes	Yes
HIR	> 1000 $\mu\text{mol}/\text{m}^2$	No	No

VLFR: very-low-fluence response;
LFR: low-fluence response;
HIR: high-irradiance response.

forms. Phytochromes are synthesized in the Pr form in dark-grown seedlings. It has been widely accepted that absorption of R light triggers a “Z” to “E” isomerization in the C15-C16 double bond between the C and D rings of the linear tetrapyrrole, resulting in the FR-absorbing Pfr form (Andel et al., 1996; Figure 2B). However, a recent NMR analysis showed that the A pyrrole ring around C4-C5 double bond rotates during photoconversion (Ulijasz et al., 2010). This discrepancy should be resolved in future studies. In addition, the Pr-to-Pfr transition is associated with rearrangement of the protein backbone (Figure 2B). The active Pfr form can be converted back to the inactive Pr form, either by a slow non-photoinduced reaction (dark reversion) or much faster upon absorption of FR light (Mancinelli, 1994; Quail, 1997a; Fankhauser, 2001; Figure 2B). This property allows phytochrome to function as a R/FR-dependent developmental switch.





Figure 2. Arabidopsis phytochrome chromophore.

(A) The biosynthesis pathway of Arabidopsis phytochrome chromophore. Image adapted from Kohchi et al. (2001).

(B) Red (R) light triggers a "Z" to "E" isomerization in the C15-C16 double bond between the C and D rings of the linear tetrapyrrole (upper panel), which is accompanied by rearrangement of the apoprotein backbone (lower panel; adapted from Bae and Choi, 2008). This results in the photoconversion of phytochromes from the Pr form to the Pfr form. Please note that the chromophore ring A rather than D is rotated during photoconversion according to a recent NMR analysis (Uliasz et al., 2010). The discrepancy needs to be resolved in future studies. Far-red (FR) light converts the Pfr form back to the Pr form.

Upper panel image reprinted from Bae and Choi (2008) with permission, from the Annual Review of Plant Biology, Volume 59 © 2008 by Annual Reviews (www.annualreviews.org).




Figure 3. The phylogenetic tree of the five phytochrome species from *Arabidopsis thaliana*.

PHYB and PHYD share ~80% amino acid sequence identity, and constitute a branch of the gene family. PHYE itself, PHYA and PHYC form two other branches of the evolutionary family tree.

Image adapted from Clack et al. (1994). Reprinted with permission from Springer.

It is generally assumed that all phytochromes have the same chromophore. Arabidopsis mutants defective in the PΦB-synthetic pathway have been isolated. These mutants (*hy1* and *hy2*) have dramatically reduced levels of PΦB and consequently of functional phytochromes, and thus exhibit severely impaired photomorphogenesis (Parks and Quail, 1991). The Arabidopsis *HY1* locus encodes a heme oxygenase (AtHO1) responsible for much of PΦB synthesis in Arabidopsis (Davis et al., 1999a; Muramoto et al., 1999). Three additional *HO* genes were found in the Arabidopsis genome, designated *AtHO2* to *AtHO4* (Davis et al., 2001). The Arabidopsis *HY2* locus, likely a unique gene in the Arabidopsis genome, encodes the phytochromobilin synthase (Kohchi et al., 2001).

It should be pointed out that in addition to PΦB, phycocyanobilin (PCB), the chromophore of the light-harvesting pigment phycocyanin, can also bind phytochrome resulting in Pr and Pfr spectra that are slightly blue shifted compared with the PΦB adducts (Lagarias and Rapoport, 1980). This finding allowed the reconstitution of photoreversible phytochromes by expressing recombinant phytochrome proteins in yeast and assembling them *in vitro*.

The Phytochrome Gene Family

In *Arabidopsis thaliana*, there are five phytochromes, designated phytochrome A (phyA) to phyE. They are encoded by five distinct members of the phytochrome gene family and are classified into two groups according to their stability in light (Sharrock and Quail, 1989). phyA is a type I (light labile) phytochrome, and phyB to phyE are all type II (light stable) phytochromes. phyA is most abundant in dark-grown seedlings, whereas its level drops rapidly upon exposure to R or white (W) light. In light-grown plants, phyB is the most abundant phytochrome, whereas phyC-phyE are less abundant (Clack et al., 1994; Hirschfeld et al., 1998; Sharrock and Clack, 2002).

Sequence analysis suggests that these phytochromes can be clustered into three subfamilies: phyA/phyC, phyB/phyD, and phyE (Figure 3). Analysis of reconstituted recombinant phyA, phyB, phyC and phyE proteins revealed that they have similar but not identical spectral properties (Kunkel et al., 1996; Remberg et al., 1998; Eichenberg et al., 2000). Orthologs of Arabidopsis *PHY* genes are present in most, if not all, higher plants (Clack et al., 1994; Sharrock and Quail, 1989; Mathews and Sharrock, 1997).




Figure 4. The domain structure of Arabidopsis phyA and phyB molecules.

H, hinge; NTE, N-terminal extension; PAS, Per (period circadian protein), Arnt (Ah receptor nuclear translocator protein), and Sim (single-minded protein); GAF, cGMP-stimulated phosphodiesterase, *Anabaena* adenylate cyclases and *Escherichia coli* FhlA; PHY, phytochrome; PRD, PAS-related domain; HKRD, histidine kinase-related domain. The chromophore is attached to a conserved cysteine residue in the GAF domain. The numbers indicate the positions of each domain.

Image adapted from Bae and Choi (2008). Reprinted with permission, from the Annual Review of Plant Biology, Volume 59 © 2008 by Annual Reviews (www.annualreviews.org).

General Structures of Phytochromes

The phytochrome molecule consists of an N-terminal domain (~ 70 kDa) and a C-terminal domain (~ 55 kDa), connected by a flexible hinge region (Figure 4). The N-terminal domain can be further divided into four consecutive subdomains: N-terminal extension (NTE), Per-Arnt-Sim (PAS), GAF, and PHY, while the C-terminal domain can also be divided into two subdomains: the PAS-related domain (PRD) containing two PAS repeats, and the histidine kinase-related domain (HKRD) (Figure 4).

Among the N-terminal subdomains, the NTE domain is uniquely present in plant phytochromes, whereas the PAS, GAF and PHY domains are also found in phytochrome-like proteins of various organisms (see below). Among the C-terminal subdomains, the PRD domain is unique to plant phytochromes, whereas the HKRD domain is also found in phytochrome-like proteins (Rockwell et al., 2006; Bae and Choi, 2008; Nagatani, 2010). The chromophore is attached to a conserved cysteine residue in the GAF domain of plant phytochromes (Figure 4). The PAS domains can be used either as platforms for protein-protein interactions, or as response modules to small ligands or changes in light conditions, oxygen levels, and redox potentials (Quail, 1997a; Neff et al., 2000). The HKRD domain lacks a critical histidine residue, and thus may be an evolutionary remnant rather than an active histidine kinase (Boylan and Quail, 1996). The putative dimerization motifs of phytochromes are also localized in the C-terminal half of the phytochrome molecules (Quail, 1997a).

A number of point mutations in the C-terminal domains of both phyA and phyB do not affect photoreversibility but eliminate the biological activity (Quail et al., 1995; Quail, 1997a), suggesting that the C-terminal domain is essential for proper downstream signaling. Consistent with this idea, a domain-swapping and deletion analysis suggested that the N terminus of phytochrome is essential for its specific photosensory properties, while the C termini of phyA and phyB are interchangeable and function as

the output domains (Wagner et al., 1996). However, this notion was later challenged by the finding that the N-terminal domain of phyB, when dimerized and localized in the nucleus, confers much higher photosensitivity than the full-length phyB (Matsushita et al., 2003). These results suggest that the N-terminal domain of phyB transduces the light signal to downstream targets, whereas the C-terminal domain attenuates the activity of phyB (Matsushita et al., 2003). Similarly, the N-terminal domain of phyA also showed a partial physiological activity when dimerized and localized in the nucleus (Mateos et al., 2006). A further study demonstrated that dimers of the N-terminal 450-aa fragment of phyB (lacking the PHY domain) can still transduce the light signal upon nuclear localization (Oka et al., 2004). Therefore, these reports suggest that the N-terminal 450-aa fragment (encompassing the NTE, PAS and GAF domains) constitutes the core signaling domain of phytochrome.

PHYTOCHROME FUNCTIONS

In most instances, the roles of individual phytochromes are studied in the context of specific responses and/or developmental stages. Loss-of-function studies of monogenic phytochrome mutant and higher order mutants of various combinations combined with gain-of-function analyses are revealing the roles of individual phytochromes in regulating different aspects of plant development. It is clear now that individual phytochromes play both unique and overlapping roles throughout the life cycle of plants, regulating a range of developmental processes from seed germination to the timing of reproductive development (Table 2). Phytochrome functions in Arabidopsis development were recently reviewed by Franklin and Quail (2010), and this chapter will mainly discuss the roles of phytochromes in seed germination, seedling de-etiolation, and shade avoidance.

Table 2. Different Roles of Phytochrome Family Members in Seedling and Early Vegetative Development

Phytochrome Members	Primary Photosensory Activities	Primary Physiological Roles
phyA	VLFRs FR-HIRs	Seed germination under a broad spectrum of light conditions (UV, visible, FR); Seedling de-etiolation under FRc; promoting flowering under LD.
phyB	LFRs R-HIRs EOD-FR (R/FR ratio)	Seed germination under Rc; Seedling de-etiolation under Rc; Shade avoidance response (petiole and internode elongation, flowering).
phyC	R-HIRs	Seedling de-etiolation under Rc.
phyD	EOD-FR (R/FR ratio)	Shade avoidance response (petiole and internode elongation, flowering).
phyE	LFRs EOD-FR (R/FR ratio)	Seed germination; Shade avoidance response (petiole and internode elongation, flowering).
VLFRs: very-low-fluence responses; LFRs: low-fluence responses; HIRs: high-irradiance responses; FR: far-red light; R: red light;		FRc: continuous far-red light; Rc: continuous red light; LD: long day light condition; EOD-FR: end-of-day far-red light; R/FR ratio: red/far-red light ratio.

Seed Germination

As mentioned above, the involvement of a R/FR-reversible photoreceptor in mediating seed germination was first demonstrated by Harry Borthwick and colleagues in 1952 by studying the reversible germination of Grand Rapids lettuce seeds by R and FR treatments (Borthwick et al., 1952). The involvement of individual phytochromes in mediating Arabidopsis seed germination has been documented in many mutant studies. At least three phytochromes, i.e. phyA, phyB and phyE are involved in the control of Arabidopsis seed germination. phyA is responsible for the irreversible VLFR responses triggered by a wide variety of irradiations (ultraviolet, visible and FR light), while phyB controls the R/FR photoreversible LFRs (Reed et al., 1994; Botto et al., 1996; Shinomura et al., 1996). Seed germination can be promoted by both VLFRs and LFRs (Table 2). In addition, phyA promotes germination in continuous FR light in the HIR mode (Johnson et al., 1994; Reed et al., 1994; Hennig et al., 2002). However, phyE was also found to play a role in controlling seed germination in continuous FR light. This could be either because phyE is directly involved in the photoperception of FR light for this response, or because phyA requires phyE to mediate seed germination (Hennig et al., 2002). It is interesting that ambient temperature modulates the light-regulation of Arabidopsis seed germination, and different phytochromes display altered functional hierarchies at different temperatures (Heschel et al., 2007).

Seedling De-etiolation

Dark-grown seedlings undergo skotomorphogenesis (etiolation) and are characterized by long hypocotyls, closed cotyledons and apical hooks, and development of the proplastids into etioplasts. Light-grown seedlings undergo photomorphogenesis (de-etiolation) and are characterized by short hypocotyls, open and expanded cotyledons, and development of the proplastids into

green mature chloroplasts (McNellis and Deng, 1995). Phytochromes perform a variety of overlapping functions in regulating seedling de-etiolation.

Mutants deficient in phyA display a wild-type photomorphogenic phenotype in W and R light. However, when grown in continuous FR light, *phyA* mutants display a skotomorphogenic phenotype (Figure 5), confirming that phyA is the primary photoreceptor responsible for perceiving and mediating various responses to FR light (Dehesh et al, 1993; Nagatani et al., 1993; Parks and Quail, 1993; Whitelam et al., 1993; Reed et al., 1994). Comparative transcriptional profiling of etiolated wild-type and *phyA* mutants subjected to FR light treatments revealed more than 800 *phyA*-regulated genes, providing the first insight into the *phyA* transcriptional network (Tepperman et al., 2001). It should be noted that *phyA* mutants also display elongated hypocotyls in continuous B light (Whitelam et al., 1993; Neff and Chory, 1998), suggesting that phyA also plays a pivotal role in perceiving and transducing B light.

phyB is the predominant phytochrome regulating de-etiolation in W and R light (Figure 5). However, transcriptional profiles of etiolated *phyB* mutants subjected to R light treatments did not differ dramatically from the wild-type controls (Tepperman et al., 2004). Subsequent studies showed that phyA plays a dominant role in regulating rapid gene expression responses to R light treatments (Tepperman et al., 2006). Moreover, the long hypocotyl and reduced cotyledon expansion phenotypes were enhanced in *phyA phyB* double mutants relative to *phyB* monogenic mutants in R light (Figure 5), revealing a role for phyA in responding to R light which is normally masked in the presence of phyB (Neff and Van Volkenburgh, 1994; Reed et al., 1994; Casal and Mazzella, 1998; Neff and Chory, 1998).

Mutants deficient in phyC exhibit a partial loss of sensitivity to R light, with longer hypocotyls and smaller cotyledons than the wild-type controls, indicating that phyC functions in regulating seedling de-etiolation in R light (Franklin et al., 2003a; Monte et




Figure 5. Phenotypes of 4-d-old wild-type (WT), *phyA*, *phyB* and *phyA phyB* mutant plants grown in darkness (D) or under continuous white (W), far-red (FR), red (R) and blue (B) light conditions.

al., 2003). However, no additive phenotype was observed in *phyB phyC* double mutants compared with their monogenic mutants, suggesting that *phyC* function is dependent on *phyB* (Monte et al., 2003). This might be explained by a recent finding that *phyC* does not homodimerize but rather forms heterodimers with *phyB* *in vivo* (Clack et al., 2009). Moreover, despite showing more sequence similarity to *PHYA* rather than to *PHYB*, *PHYD* and *PHYE* (Figure 3), *phyC* seems not to play a role in mediating seedling de-etiolation in FR light (Franklin et al., 2003a; Monte et al., 2003). However, the hypocotyls of *phyA phyC* double mutants are significantly longer than those of *phyC* monogenic mutants in R light (Franklin et al., 2003a; Monte et al., 2003), again confirming the contribution of *phyA* to seedling establishment under R light.

Although *PHYD* shows high sequence similarity to *PHYB* (Figure 3), the role of *phyD* in seedling de-etiolation in R seems minor, as it was reported that the Wassilewskija (Ws) ecotype of *Arabidopsis* contains a natural *phyD* deletion but its seedlings display only marginally longer hypocotyls in R than plants containing an introgressed *PHYD* gene (Aukerman et al., 1997). However, a synergistic relationship was observed between *phyB* and *phyD* in R, as the hypocotyls of *phyB phyD* double mutants are more than additively longer than those of each monogenic mutant (Aukerman et al., 1997). The contribution of *phyE* to seedling de-etiolation seems negligible, as it was shown that monogenic *phyE* mutants were indistinguishable from wild-type control plants in a variety of light conditions (Devlin et al., 1998).

As mentioned above, phytochromes also weakly absorb B light (Figure 1). In addition, phytochromes were shown to modulate cryptochrome-mediated seedling de-etiolation and phototropin-mediated phototropic curvature of *Arabidopsis* hypocotyls in B light (Parks et al., 1996; Ahmad and Cashmore, 1997; Hamazato et al., 1997; Janoudi et al., 1997; Casal and Mazzella, 1998; Neff and Chory, 1998). Direct physical interactions between phytochromes and cryptochromes were also reported (Ahmad et al., 1998; Mas et al., 2000). Therefore, phytochromes co-act with B/UV-A light photoreceptors to regulate seedling de-etiolation in B light.

Shade Avoidance

Plant development is regulated not only by the difference between light and darkness, but also by light quality, in particular the change of light quality due to shading by other plants. Light passed through or reflected from living vegetation is depleted in R and B wavebands, which are absorbed by chlorophyll and carotenoid pigments used for photosynthesis, leading to a reduction in the ratio of R to FR wavelengths (R:FR). This allows plants to initiate a suite of developmental responses called shade avoidance syndrome (SAS), which elevates leaves towards unfiltered daylight and enables plants to overtop competitors (reviewed in Smith and Whitelam, 1997). These responses include elongation of stems and petioles, accelerated flowering time, and increased apical dominance. The ability of plants to monitor their light environments and change their architecture provides them with a competitive strategy to survive and complete their life cycle in dense stands.

Reductions in R:FR ratio favor the conversion of phytochrome molecules to their inactive Pr form. Therefore the shade avoidance syndrome must be suppressed under high R:FR ratio conditions.




Figure 6. Phenotypes of 3-week-old wild-type (WT), *phyA*, *phyB*, *phyA phyB*, *phyB phyD phyE*, *phyA phyB phyD phyE* plants grown under white light conditions (16-h light/8-h dark).

In this sense, shade avoidance is due to the relief of suppression rather than the induction of physiological responses. *phyB* is the predominant suppressor of shade avoidance responses in high R:FR, as *phyB*-deficient plants display a constitutive shade avoidance phenotype (elongated petiole and early flowering) (Nagatani et al., 1991; Somers et al., 1991; Figure 6). The shade avoidance responses enabled by low R:FR ratios can be effectively phenocopied by end-of-day far-red (EOD-FR) treatments. This led to the discovery of the roles of *phyD* and *phyE* in shade avoidance. Although the monogenic *phyD* mutant plants have no obviously abnormal phenotype to EOD-FR, plants impaired in both *phyB* and *phyD* display significantly longer hypocotyls under either R or W light, and flower earlier than the *phyB* monogenic mutants, suggesting that *phyB* and *phyD* function redundantly in suppressing shade avoidance (Aukerman et al., 1997; Devlin et al., 1999). As with *phyD*, monogenic *phyE* mutants show no phenotypic alterations unless in the *phyB* mutant background, and the *phyB phyE* double mutants flower much earlier than the *phyB* monogenic mutants (Devlin et al., 1998; Franklin et al., 2003b).

As *phyA* is the primary photoreceptor sensing FR wavelengths, enrichment of FR in transmitted/reflected light can lead to enhanced *phyA* signaling in the HIR mode. Therefore, the action of *phyA* can substitute for the loss of *phyB*, *phyD* and *phyE* activity due to their conversion to the inactive Pr forms. Indeed, Arabidopsis *phyA* mutant seedlings display enhanced shade avoidance responses relative to wild-type control plants when grown in low R:FR conditions (Johnson et al., 1994; Smith et al., 1997; Salter et al., 2003). Despite the relatively close phylogenetic relationship between *phyA* and *phyC* (Figure 3), no role for *phyC* in mediating shade avoidance responses has been reported. This conclusion is further confirmed by the observations that *phyA phyB phyD phyE* quadruple mutants display insensitivity to reductions in R:FR ratio and EOD-FR treatments (Franklin et al., 2003b).

LIGHT-REGULATED SUBCELLULAR LOCALIZATION OF PHYTOCHROMES

As mentioned above, phytochromes are synthesized in the cytosol in their inactive Pr forms. It was widely accepted before the mid-1990s that phytochromes were cytoplasmic photoreceptors based on early biochemical and immunocytochemical studies (Nagy and Schafer, 2002). However, extensive studies conducted in the last decade have established the notion that phytochromes must enter the nucleus to trigger most light responses (Nagatani, 2004; Kevei et al., 2007; Fankhauser and Chen, 2008). Thus, light-regulated translocation of the photoreceptors from the cytoplasm into the nucleus is a key event in the phytochrome signaling cascade. Recent publications are beginning to shed light on the molecular mechanisms underlying this central control step.

Regulation of *phyB* Nuclear Localization

phyB nuclear accumulation is efficiently initiated by continuous R light, and to a lesser extent by continuous B light, but completely ineffective by FR light. Single pulses of R, FR and B light cannot induce *phyB* nuclear accumulation (Gil et al., 2000). Moreover, *phyB* nuclear transport by R light is reversible by FR light, a typical characteristic of LFR (Kircher et al., 1999). A similar regulation of subcellular localization was also reported for *phyC*, *phyD* and *phyE* (Kircher et al., 2002). It should be noted that there is a weak, but detectable level of *phyB*-*phyE* present in the nucleus of dark-grown plants (Kircher et al., 2002).

A structure-function analysis first demonstrated that the C-terminal half of *phyB* (amino acids 594-1172) is sufficient to localize GUS to the nucleus, suggesting that the C-terminal domain of *phyB* harbors a putative nuclear localization signal (NLS) (Sakamoto and Nagatani, 1996). This result was confirmed by a later study, which showed that GFP fused to the N-terminal half of *phyB* (amino acids 1-651) localizes to the cytoplasm, whereas GFP fused to the C-terminal half of *phyB* (amino acids 625-1172)

localizes to the nucleus (Matsushita et al., 2003). Subsequent further analyses of various truncations of the phyB C-terminal domain revealed that the PRD domain of phyB (amino acids 594-917) is both necessary and sufficient for nuclear localization, indicating that a putative NLS resides in this domain (Chen et al., 2005). Interestingly, this study also showed that the N-terminal photosensory GAF-PHY domains interact with the PRD domain in a light-dependent manner, thus providing a mechanistic link between light-dependent Pr/Pfr conformational alterations of phyB and the unmasking of the NLS that regulates phyB nuclear accumulation (Chen et al., 2005).

Regulation of phyA Nuclear Localization

The nuclear accumulation pattern of phyA is quite distinct from that of phyB. Firstly, all light illuminations (FR, R and B) are effective in inducing phyA nuclear translocation. A single, brief pulse of FR, R or B light induces phyA nuclear import as well (Hisada et al., 2000; Kim et al., 2000; Kircher et al., 2002). Therefore, phyA nuclear import is mediated by VLFR and HIR. Secondly, phyA nuclear translocation is very rapid (within minutes), whereas phyB nuclear import is relatively slow that takes hours (Kircher et al., 1999; Kim et al., 2000; Kircher et al., 2002). Finally, in contrast to phyB-phyE, phyA is exclusively localized in the cytosol in etiolated seedlings (Kircher et al., 2002).

FAR-RED ELONGATED HYPOCOTYL 1 (FHY1) and FHY1-LIKE (FHL). The rapid nuclear translocation of phyA, and the fact that phyA itself does not contain a typical NLS suggest the existence of an efficient transport machinery responsible for phyA nuclear import. Indeed, two small plant-specific proteins, FHY1 and FHL, have been shown to play an essential role in facilitating phyA nuclear translocation. The history of FHY1 essentially parallels the history of the molecular genetic analysis of phyA signaling pathway in Arabidopsis. The *fhy1* mutant was firstly reported in 1993, together with two other mutants, i.e. *fhy2* and *fhy3*, and all these *fhy* mutants develop elongated hypocotyls in FR light (Whitelam et al., 1993). This pioneering report showed that *FHY2* locus corresponds to *PHYA*, while the *FHY1* and *FHY3* genes were cloned as separate loci in 2001 and 2002, respectively (Desnos et al., 2001; Wang and Deng, 2002). Subsequent studies identified a FHY1-like protein, named FHL, based on its sequence homology to FHY1 (Zhou et al., 2005).

FHY1 and FHL are two small proteins (202 and 201 amino acids, respectively) in Arabidopsis that were found to have homologs in both monocot and dicot plant species (Genoud et al., 2008; Li et al., 2010). Each protein contains an NLS and a nuclear exclusion signal (NES) at their N-termini and a septin-related domain (SRD) at their C termini (Desnos et al., 2001; Zhou et al., 2005). *In vitro* binding assays showed that both proteins are capable of homo- and hetero-dimerization through their C-terminal domains (Zhou et al., 2005). The NLS and SRD motifs are functionally important, because removal of either motif disrupts the function of FHY1 (Zeidler et al., 2004).

Earlier studies involving FHY1 suggested that FHY1 is responsible for mediating a branch of phyA signaling (Barnes et al., 1996). However, microarray analysis showed that all genes affected by *phyA* mutation are also affected by *fhy1* mutation, although to a lesser degree (Wang et al., 2002). The roles of

FHY1 in phyA signaling remained obscure until it was reported that FHY1/FHL physically interact with the Pfr form of phyA *in vitro* and in yeast cells through their SRD motifs, and that FHY1/FHL are required for nuclear accumulation of phyA since phyA is localized only in the cytosol of *fhy1 fhl* double mutants (Hiltbrunner et al., 2005, 2006). This conclusion was extended by a later report that the major function of FHY1/FHL is to act as adaptor proteins to chaperone photoactivated phyA into the nucleus (Genoud et al., 2008). Evidence supporting this proposed mode of action of FHY1 includes, first, sequence alignments show that the N-terminal NLS and the C-terminal phyA-interacting motifs are the only conserved motifs among all FHY1 homologs. Consistently, an artificial FHY1 consisting of a virus NLS motif and the C-terminal phyA-interacting motif of Arabidopsis FHY1 could rescue *fhy1* mutant phenotypes and colocalizes with phyA in the nucleus. Second, FHY1 becomes functionally dispensable in transgenic seedlings expressing a constitutively nuclear phyA, i.e. if phyA could enter the nucleus by itself (Genoud et al., 2008).

FHY1/FHL specifically control the subcellular localization of phyA but not phyB, because *fhy1/fhl* mutants only show long-hypocotyl phenotype in FR but not R light, and phyB nuclear import is not affected in the *fhy1* mutant (Hiltbrunner et al. 2005). As each of FHY1 and FHL has a functional monopartite NLS and NES that are indeed involved in the nuclear localization and exclusion of FHY1 (Zeidler et al., 2004), it is possible that phyA utilizes the NLS of FHY1/FHL for its nuclear transport. The fact that FHY1 homologs are widely distributed in angiosperms suggests that the mechanism uncovered in Arabidopsis may be conserved in higher plants (Genoud et al., 2008). However, interaction of phyA with FHY1/FHL alone appears to be insufficient for phyA nuclear translocation, because phyA-402, containing a missense mutation in the HKRD domain of phyA, is still capable of interacting with FHY1/FHL but does not translocate into the nucleus in R light (Muller et al., 2009).

Based on the evidence discussed above, one would assume that FHY1/FHL only function for phyA nuclear transport. However, this assumption was challenged by a recent report, which showed that FHY1/FHL might transmit phyA signals to downstream transcription factors (Yang et al., 2009). FHY1/FHL physically interact with two well-characterized transcription factors in phyA signaling network, LONG HYPOCOTYL IN FAR-RED 1 (HFR1) and LONG AFTER FAR-RED LIGHT 1 (LAF1), both *in vitro* and *in vivo*. Analysis of double and triple mutants showed that HFR1 and LAF1 independently transmit phyA signals downstream of FHY1 and FHL. Intriguingly, FHY1 was shown to mediate the assembly of a PHYA/FHY1/HFR1 signaling complex *in vitro*, suggesting that such kind of phyA signaling complexes may be assembled *in vivo* (Yang et al., 2009).

FHY3 and FAR-RED IMPAIRED RESPONSE1 (FAR1). The distinct role of FHY1/FHL in phyA nuclear accumulation suggests that any factor regulating *FHY1/FHL* transcriptionally or post-transcriptionally may indirectly affect phyA nuclear import. This notion is true, as demonstrated by the functional studies on FHY3 and FAR1. When the *FHY3* gene was cloned in 2002, sequence alignments revealed that FHY3 shares high homology with FAR1, a previously identified phyA signaling component (Hudson et al., 1999; Wang and Deng, 2002). The function of FHY3 and FAR1 was not clear at that time, but some reports showed that FHY3 positively regulates the transcript levels of *FHY1*, and that FHY3/

FAR1 proteins share substantial similarity to *Mutator*-like element (MULE) transposases and may work as transcriptional regulators (Desnos et al., 2001; Hudson et al., 2003).

The breakthrough was made in 2007, when a study unequivocally demonstrated that FHY3 and FAR1 are transposase-derived transcription factors directly binding to the *FHY1/FHL* promoters via a specific *cis*-element called the FHY3/FAR1 binding site (FBS), and activating *FHY1/FHL* gene expression (Lin et al., 2007). The N-terminal C2H2 zinc finger domains of FHY3/FAR1 are essential for DNA binding, whereas the entire C-terminal regions are required for their transcriptional activation activity (Lin et al., 2007, 2008). Thus, FHY3 and FAR1 define a new type of transposase-derived transcription factors. Moreover, phyA nuclear accumulation is abolished in the *thy3 far1* double mutant, indicating that as the key transcriptional activators of *FHY1/FHL* expression, FHY3/FAR1 indirectly control phyA nuclear accumulation (Lin et al., 2007). This conclusion was further supported by another report that constitutively nuclear phyA could rescue the *thy3* mutant phenotypes (Genoud et al., 2008).

The discovery of FHY3/FAR1's role in phyA signaling invites the further question of how *FHY1/FHL* expression is down-regulated by phyA signaling. A negative feedback mechanism(s) controlling *FHY1/FHL* expression should exist, as previous studies showed that *FHY1/FHL* transcript levels were rapidly down-regulated when dark-grown plants were exposed to FR light (Desnos et al., 2001; Lin et al., 2007). A recent study showed that ELONGATED HYPOCOTYL 5 (HY5), a well-characterized bZIP transcription factor involved in promoting photomorphogenesis under various light conditions, plays a major role in this process (Li et al., 2010). HY5 achieves its goal by two distinct mechanisms. The first mechanism involves steric hindrance as HY5 directly binds ACGT-containing elements (ACEs) less than 10 bp away from the FHY3/FAR1 binding sites in the *FHY1/FHL* promoters. Thus, HY5's occupation of the ACEs consequently decreases the accessibility of the *FHY1/FHL* promoters to FHY3/FAR1. The second mechanism is called "sequestration" through the physical interactions between HY5 and FHY3/FAR1, a mechanism also used in the regulation of some plant bHLH transcription factors (de Lucas et al., 2008; Feng et al., 2008; Hornitschek et al., 2009). Therefore, HY5 acts as a repressor of *FHY1/FHL* expression by modulating the transcriptional activities of FHY3 and FAR1 (Li et al., 2010; Figure 7).

Nuclear Bodies (NBs)

Upon import into the nucleus, both phyA and phyB localize to discrete subnuclear foci, called nuclear bodies or speckles. Although most studies on NBs were conducted in cells overexpressing fluorescent-protein-tagged phytochromes, native phyA and phyB have each been shown to localize to NBs by immunocytochemical studies, suggesting that the formation of NBs is not an artifact because of the overexpression of phytochromes (Hisada et al., 2000; Kircher et al., 2002). The pattern of NBs is highly dynamic and directly regulated by light quality, quantity, and periodicity, and closely correlates to phytochrome-mediated responses (Kircher et al., 2002; Chen et al., 2003). However, the precise nature of NBs is still unknown. But several phytochrome signaling components colocalize to NBs, suggesting that NBs play important roles in phytochrome signaling (Chen, 2008).




Figure 7. Control of *FHY1/FHL* expression and phyA nuclear accumulation.

FHY1 and FHL are required for phyA nuclear accumulation (Hiltbrunner et al., 2005, 2006; Genoud et al., 2008). FHY3 and FAR1 are two transposase-derived transcription factors that directly activate *FHY1/FHL* transcription, and thus indirectly regulate phyA nuclear accumulation and subsequent responses (Lin et al., 2007). phyA is localized exclusively in the cytosol in darkness in its inactive Pr form. Upon light exposure, the Pfr form of phyA is imported into the nucleus by FHY1/FHL, and thus triggers phyA signaling leading to multiple light responses, including the reduction of COP1 in the nucleus and accumulation of HY5 (Osterlund and Deng, 1998; Osterlund et al., 2000), and feedback regulation of *FHY3* and *FAR1* transcript levels (Lin et al., 2007). HY5 plays dual roles in phyA signaling: promoting photomorphogenesis, and down-regulating *FHY1/FHL* transcript levels by modulating the activities of the transcriptional activators FHY3 and FAR1 (Li et al., 2010). FHY3 and FHY1 (indicated by larger letters) are the more predominant players in the phyA signaling process compared to their respective homologs FAR1 and FHL. Pr: R-absorbing form of phyA (inactive); Pfr: FR-absorbing form of phyA (active). Arrow, positive regulation; bar, negative regulation. Image adapted from Li et al. (2010).

Based on the kinetics of phyB-GFP localization during the dark-to-light transition, two types of NBs have been defined for phyB. Within minutes of light exposure, small and transient phyB-GFP NBs appear (Bauer et al., 2004). Interestingly, phyB colocalizes with PIF3 in the early transient NBs, and its localization to the early NBs is PIF3-dependent *in vivo* (Bauer et al., 2004). These early transient NBs disappear after 10-15 min in the light (Kevei et al., 2007), and interestingly, the disappearance of these early NBs correlates with the light-induced and phytochrome-dependent degradation of PIF3 (Al-Sady et al., 2006), thus implying that these early phyB NBs are the sites for phyB-PIF3 interaction, and their disappearance is due to PIF3 degradation (Chen, 2008).

After the disappearance of these early phyB NBs, longer R light treatment (2-3 h) leads to the appearance of larger and more stable NBs (Nagatani, 2004; Kevei et al., 2007; Chen, 2008). Interestingly, the size and number of these NBs depend on the fluence rate of R light (Chen et al., 2003). As increasing fluence rates of R not only induced a change in the pattern of phyB NBs but also enhanced inhibition of hypocotyl elongation, it was pro-

posed that the formation of phyB NBs plays a role in the regulation of phyB-mediated signal transduction (Chen et al., 2003). The observations that PIF3 colocalizes with phyB only in the early but not late NBs indicate that the components of the early and late NBs may be different (Bauer et al., 2004).

phyA rapidly enters the nucleus in response to FR light and also forms early and late NBs (Bauer et al., 2004). PIF3 colocalizes with phyA in the early transient NBs but not late stable NBs (Bauer et al., 2004), consistent with the report that light-activated phyA also interacts with PIF3 and contributes to its degradation (Al-Sady et al., 2006). FHY1 and FHL, two phyA-interacting proteins required for phyA nuclear accumulation (see above), colocalize with phyA in the early transient NBs (Hiltbrunner et al., 2005, 2006). However, although a short R treatment also induces translocation of phyA into the nucleus, NB formation, and colocalization with PIF3, extended R light results in the complete loss of the phyA-GFP fluorescence (Bauer et al., 2004), possibly due to the photolabile nature and rapid degradation of phyA (see below).

Because both phyA and PIF3 are localized to NBs before their degradation, it has been proposed that NBs of phytochromes are sites for protein degradation (Bauer et al., 2004; Seo et al., 2004; Al-Sady et al., 2006). Recently, Chen et al. (2010) used a confocal microscopy-based screen to identify a gene, *HEMERA* (*HMR*), required for the localization of phyB-GFP to large NBs in high fluence rate of R light. Characterization of *hmr* mutants, localization of HMR protein within cells, and analysis of its biochemical function indicate that HMR is a specific and early phytochrome signaling component required for light-dependent proteolysis of phyA, PIF1, and PIF3 (Chen et al., 2010). Moreover, HMR is predicted to be structurally similar to the multiubiquitin-binding protein, RAD23, and can partially rescue yeast *rad23* mutants, thus suggesting that phytochrome nuclear bodies may serve as sites of proteolysis (Chen et al., 2010).

PHYTOCHROMES AS LIGHT-REGULATED KINASES

How do phytochromes initiate their signal transduction upon photo-activation? A long-standing but much disputed hypothesis is that phytochromes act as light-regulated kinases (Wong et al., 1986; Kim et al., 1989). This hypothesis was initially supported

by the observation that purified preparations of phyA catalyzed phosphorylation of serine residues on the photoreceptor itself, i.e. autophosphorylation activity (Wong et al., 1986; Yeh and Lagarias, 1998). The discovery of phytochrome-like photoreceptors in bacteria, collectively called bacteriophytochromes (BphPs), generated further supporting evidence for such a view (Fankhauser, 2000; Vierstra and Davis, 2000). Phytochrome-like sequences were identified in the cyanobacteria *Fremyella diplosiphon*, *Synechocystis sp. PCC6803*, the purple photosynthetic bacterium *Rhodospirillum rubrum* and non-photosynthetic bacteria such as *Deinococcus radiodurans*, *Pseudomonas putida* and *Pseudomonas aeruginosa* (Kehoe and Grossman, 1996; Hughes et al., 1997; Davis et al., 1999b; Hughes and Lamparter, 1999; Jiang et al., 1999; Wu and Lagarias, 2000). Some of them, such as Cph1 of *Synechocystis sp. PCC6803*, can bind to the plant phytochrome chromophore (phytochromobilin PΦB or phycocyanobilin PCB) autocatalytically and display R/FR absorption spectra similar to plant phytochromes (Hughes et al., 1997; Lamparter et al., 1997; Yeh et al., 1997). Further, Cph1 was shown to be a light-regulated histidine kinase. Both autophosphorylation of Cph1 and transphosphorylation of Rcp1 (the response regulator for Cph1) are inhibited by R light and stimulated by FR light (Yeh et al., 1997), suggesting that in cyanobacteria phosphorylation is an important and very early step of phytochrome signal transduction.

However, higher plant phytochromes share limited sequence similarity with Cph1 at their C-termini (Figure 8). In addition, plant phytochromes have two additional domains compared with BphPs: a serine-rich NTE region, and a PRD domain located between PHY and HKRD domains (Figures 8). Moreover, mutating several critical residues required for bacterial His kinase activity does not affect the activity of plant phytochromes, suggesting that plant phytochromes are not active His kinases (Quail, 1997b). In fact, purified recombinant plant phytochromes exhibit a serine/threonine kinase activity, suggesting that eukaryotic phytochromes are histidine kinase paralogs with serine/threonine specificity (Yeh and Lagarias, 1998). Consistent with this discovery, several *in vitro* substrates of phytochrome kinase activity were subsequently discovered, such as histone H1, PHYTOCHROME KINASE SUBSTRATE 1 (PKS1), cryptochromes, AUX/IAA proteins, and FHY1 (Wong et al., 1989; Ahmad et al., 1998; Fankhauser et al., 1999; Colon-Carmona et al., 2000; Shen et al.,




Figure 8. Structural comparison of Arabidopsis phytochromes and the bacterial phytochrome Cph1 (adapted from Yeh and Lagarias, 1998).

HKD, histidine kinase domain. The percent amino acid identities between the HKD domain of Cph1 and both PRD and HKRD domains of Arabidopsis phytochromes are indicated.

Image adapted from Yeh and Lagarias (1998). Reprinted with permission from the National Academy of Sciences.

2009). However, the kinase domain of phytochrome has not been determined. It is notable that both PRD and HKRD domains show similarities to the HKD domain of Cph1 (Yeh and Lagarias, 1998; Figure 8). In addition, to date the kinase activity has only been proven for one higher plant phytochrome: oat phyA. Although a recent report indicated that phyB shows some kinase activity *in vitro* (Phee et al., 2008), more evidence is required before a firm conclusion can be reached. Whether all phytochromes have kinase activity, and whether different phytochromes behave differently as protein kinases need to be further characterized.

The claim that higher plant phytochromes function as protein kinases invites many questions, such as, is the Pfr form more active than the Pr form? What is the biological role of this kinase activity of phytochromes in plants? Answers to these questions have just begun to be unraveled. For example, autophosphorylation of recombinant oat phyA is both chromophore and light regulated, with Pfr being more active than Pr (Yeh and Lagarias, 1998). *In vitro* kinase assays also showed that the Pfr form of phytochrome more effectively phosphorylates some substrates, such as PKS1 and CRY1 (Ahmad et al., 1998; Fankhauser et al., 1999), but for some other substrates, such as AUX/IAA proteins and FHY1, the Pr and Pfr forms showed similar kinase activities (Colon-Carmona et al., 2000; Shen et al., 2009). However, *in vivo* assays are required to verify the conclusions of these experiments. For example, FHY1 phosphorylation in Arabidopsis seedlings is solely dependent on the active Pfr form of phyA, although both Pr and Pfr forms of phyA could phosphorylate FHY1 *in vitro* (Shen et al., 2009).

Recently it was shown that oat phyA autophosphorylates two serine sites in its NTE region *in vitro* (Han et al., 2010). Mutation of these two autophosphorylation sites in transgenic Arabidopsis plants caused hypersensitive light responses, indicating an increase in phyA activity (Han et al., 2010). Consistently, the degradation of the mutant phyA was significantly slower than the wild-type phyA under light conditions, suggesting that phyA autophosphorylation plays an important role in the regulation of phytochrome signaling through the control of phyA protein stability (Han et al., 2010). Another report showed that phyA is the only photoreceptor responsible for rapid R light-dependent FHY1 phosphorylation, and interestingly, this phosphorylation is R/FR light reversible, a typical LFR mode of phytochrome action (Shen et al., 2009). Notably, phosphorylated FHY1 was shown to be a preferred substrate for ubiquitin/26S proteasome-mediated degradation, suggesting that phyA-dependent FHY1 phosphorylation in R light may serve as a biochemical mechanism to desensitize FHY1-mediated phyA signaling (Shen et al., 2005b, 2009). These examples suggest that phytochrome autophosphorylation and kinase activity may play a negative role in light signal transduction.

However, the *in vivo* functional mechanism of phytochrome kinase activity is only beginning to be understood. For example, phytochrome kinase activity is stimulated in the presence of histone H1 in a Pr-specific manner, thus it is suggested that phytochrome kinase activity is activated in the nucleus that contains cationic molecules such as histones (Yeh and Lagarias, 1998; Kim et al., 2005; Han et al., 2010). Another example showing the complexity of *in vivo* phytochrome-related kinase activity comes from the study of the PHYTOCHROME-INTERACTING FACTORS (PIFs; see below). All PIFs except PIF7 are rapidly phosphorylated and then ubiquitinated in response to light *in vivo* prior to

their degradation in a phytochrome-dependent manner (Al-Sady et al., 2006; Shen et al., 2007; Shen et al., 2008). A recent study reported that both phyA and phyB mediate PIF3 phosphorylation *in vitro* (Phee et al., 2008), suggesting that phytochromes may be the protein kinases responsible for phosphorylating the PIF proteins. However, compelling evidence supporting this assumption is still lacking. Recently, it was shown that CASEIN KINASE II (CK2), a highly conserved and ubiquitous Ser/Thr kinase, directly phosphorylates PIF1 (Bu et al., 2011). Therefore, whether phytochromes (both phyA and phyB) might directly phosphorylate PIFs *in vivo*, and how CK2 functions in phytochrome-induced rapid phosphorylation of PIFs await further investigation.

PHYTOCHROME SIGNALING INTERMEDIATES

The light signals perceived by the phytochrome photoreceptors are transduced to downstream signaling intermediates, which alter the expression of target genes and ultimately lead to the modulation of the biological responses (Quail, 2002; Jiao et al., 2007). Genetic research has identified a complex and interconnecting signaling network downstream of the phytochromes, together with a considerable number of positive or negative regulators, which act either in a specific pathway (such as LAF1 and HFR1) or in all branches of phytochrome signaling pathways (such as COP1 and HY5; Figure 9).

Negative Regulators of Phytochrome Signaling

Phytochrome-Interacting Factors (PIFs). Protein-protein interactions are necessary for many signal transduction cascades. Both general screenings for phytochrome-interacting proteins and targeted protein-protein interaction studies have identified a number of phytochrome-interacting partners. Those include PIF3 and other subsequently identified PIFs (Ni et al., 1998; Leivar and Quail, 2011), PKS1 (Fankhauser et al., 1999), NDPK2 (Choi et al., 1999), cryptochromes (both CRY1 and CRY2) (Ahmad et al., 1998; Mas et al., 2000), AUX/IAA proteins (Colon-Carmona et al., 2000), FyPP (Kim et al., 2002), COP1 (Seo et al., 2004; Jang et al., 2010), PAPP5 (Ryu et al., 2005), FHY1/FHL (Hiltbrunner et al., 2005, 2006), and were summarized recently by Bae and Choi (2008). Growing evidence demonstrates that the PIF proteins, a small subset of basic helix-loop-helix (bHLH) transcription factors, play central roles in phytochrome-mediated light signaling networks (Duek and Fankhauser, 2005; Castillon et al., 2007; Leivar and Quail, 2011).

The PIF proteins belong to the 15-member Subfamily 15 of the Arabidopsis bHLH transcription factor superfamily (Bailey et al., 2003; Heim et al., 2003; Toledo-Ortiz et al., 2003). PIF3 is the foundation member of the PIF subset, initially identified in a yeast two-hybrid screen for phyB-interacting proteins (Ni et al., 1998). The second member of the PIF family, PIF4, was isolated by the convergence of both genetic and reverse-genetic approaches (Huq and Quail, 2002). Several other PIFs were then identified based on the sequence homology to PIF3, and were named PIF1, PIF5, PIF6 and PIF7 (Huq et al., 2004; Khanna et al., 2004; Oh et al., 2004; Leivar et al., 2008). As members of the bHLH superfamily transcription factors, all the PIF proteins contain a




Figure 9. A simplified model of the phytochrome signaling pathway.

phyA is the primary photoreceptor responsible for perceiving and mediating various responses to FR light, whereas phyB is the predominant phytochrome regulating responses to R light. Under light conditions, these photoreceptors act to suppress two main branches of light signaling: COP1-TFs and PIFs. COP1, whose activity is repressed by phytochromes in light conditions, is an E3 ubiquitin ligase targeting several photomorphogenesis-promoting transcription factors (such as HY5, HYH, LAF1 and HFR1) for degradation. PIFs are a subset of bHLH transcription factors required for skotomorphogenesis. Photo-activated phytochromes directly interact with PIFs, resulting in PIFs' phosphorylation and degradation, while COP1 positively regulates PIFs' protein levels. Phytochromes are targeted for degradation by COP1, and PIFs contribute to the degradation of phyB by promoting COP1/phyB interaction. Arrow, positive regulation; bar, negative regulation; solid line, direct regulation; dotted line, indirect regulation. Image adapted from Lau and Deng (2010). Reprinted with permission from Elsevier.

bHLH signature domain, consisting of a basic region (~ 15 aa) involved in DNA binding and an HLH region (~ 60 aa) involved in dimerization (Toledo-Ortiz et al., 2003; Castillon et al., 2007). The majority of the bHLH proteins bind to a *cis*-element called the E-box (CANNTG), whereas all the PIF proteins, where examined, bind specifically to a subtype of E-box, called the G-box (CACGTG) (Castillon et al., 2007; Leivar and Quail, 2011).

All PIF members contain a conserved motif in their N-termini, designated active phytochrome B-binding (APB) motif, which confers specific binding of PIFs to the biologically active Pfr form of phyB (Khanna et al., 2004; Duek and Fankhauser, 2005; Castillon et al., 2007; Leivar and Quail, 2011). However, only two PIF proteins, PIF1 and PIF3, also bind to the Pfr form of phyA, with PIF1 showing much stronger affinity for phyA than PIF3 (Ni et al., 1998; Huq et al., 2004). Accordingly, PIF1 and PIF3 each contain a motif called active phytochrome A-binding (APA) necessary for binding to phyA, but the actual sequences of these two APA motifs are not conserved (Al-Sady et al., 2006; Shen et al., 2008).

The recent finding that a quadruple mutant of PIFs, *pif1 pif3 pif4 pif5* (*pifq*), develops a *constitutively photomorphogenic* (*cop*)-like phenotype in the dark provides compelling evidence that the PIF proteins repress photomorphogenesis and promote skotomorphogenesis in etiolated seedlings (Leivar et al., 2008; Shin et

al., 2009; Quail, 2011). Consistently, microarray analysis showed that the dark-grown *pifq* mutant has a gene expression pattern similar to that of R light-grown wild-type plants (Shin et al., 2009). By comparing rapidly light-responsive genes in wild-type seedlings with those responding in darkness in the *pifq* mutant, an overlapping subset of genes were identified as potential direct targets of these bHLH transcription factors (Leivar et al., 2009). Notably, transcription factor-encoding genes are highly enriched among these genes, suggesting that they may be potential primary targets of PIF transcriptional regulation.

At the same time, evidence obtained in the last decade demonstrates that one way phytochromes promote photomorphogenesis is by inducing rapid (within minutes) phosphorylation of most, if not all PIFs, upon light exposure. This subsequently leads to their ubiquitination and degradation via the ubiquitin/proteasome system (Bauer et al., 2004; Park et al., 2004; Shen et al., 2005a; Al-Sady et al., 2006; Oh et al., 2006; Nozue et al., 2007; Shen et al., 2007; Al-Sady et al., 2008; Lorrain et al., 2008; Shen et al., 2008). The interaction between phytochromes and PIFs is necessary for the light-dependent phosphorylation of PIFs, because mutant PIF1 and PIF3 proteins that abolish interactions with the Pfr forms of phyA and phyB do not undergo light-dependent phosphorylation (Al-Sady et al., 2006; Shen et al., 2008). Moreover,

upon returning light-grown plants to darkness, PIF proteins rapidly re-accumulate to high levels. Subsequent re-exposure to light once again induces rapid degradation of PIFs (Leivar and Quail, 2011), indicating that this rapid regulation of PIFs is dynamic and controlled by phytochrome LFRs. Therefore, phytochrome-induced phosphorylation and proteolysis of PIFs may represent a major biochemical mechanism of signal transfer from the photoactivated phytochromes to their interacting signaling partners in the nucleus, which rapidly alters the gene expression profiles of the genome. In addition to controlling phytochrome-regulated gene expression, PIF proteins were recently shown to also modulate phyB abundance in R light, possibly by stimulating COP1-catalyzed ubiquitination and degradation of phyB (see below) (Khanna et al., 2007; Leivar et al., 2008; Al-Sady et al., 2008; Jang et al., 2010; Figure 9).

COP/DET/FUS Proteins. Genetic screens for Arabidopsis mutants involved in light-regulated seedling development followed by biochemical analyses have identified a group of pleiotropic Constitutive Photomorphogenic/De-etiolated/Fusca (COP/DET/FUS) proteins that are central negative regulators of photomorphogenesis (Sullivan et al., 2003; Yi and Deng, 2005). This group of COP/DET/FUS proteins defines three biochemical entities: the COP1-SPA complexes, the COP9 signalosome (CSN), and the CDD complex (COP10, DDB1, and DET1), all of which are involved in proteasomal degradation of photomorphogenesis-promoting factors (Saijo et al., 2003; Serino and Deng, 2003; Yanagawa et al., 2004; Yi and Deng, 2005; Zhu et al., 2008). Interestingly, the COP1-SPA complexes and the CDD complex were recently shown to form two groups of CUL4-based E3 ligases *in vivo* (Chen et al., 2006, 2010). Therefore, these two groups of E3 ligases regulate the degradation of downstream factors to mediate light regulation of plant development (Chen et al., 2010).

COP1 is a conserved RING finger E3 ubiquitin ligase involved in multiple processes in many different organisms, including plant development and mammalian cell survival, growth, and metabolism (Yi and Deng, 2005). COP1 was first cloned and characterized in the model plant Arabidopsis as a repressor of light-regulated plant development (Deng et al., 1991, 1992; Figure 10). COP1 contains three domains: a RING finger domain in its N-terminal region, a WD40 repeat domain in its C-terminus, and a coiled-coil domain in the middle (Deng et al., 1992; Yi and Deng, 2005). COP1 has been shown to act as an E3 ligase targeting several photomorphogenesis-promoting proteins for degradation, including HY5 (Osterlund et al., 2000), HY5 HOMOLOG (HYH; Holm et al., 2002), LAF1 (Seo et al., 2003), HFR1 (Duek et al., 2004; Jang et al., 2005; Yang et al., 2005), and the phytochromes (Seo et al., 2004; Jang et al., 2010) (Figure 9). In addition, COP1 was recently shown to regulate flowering time by directly targeting transcriptional activator CONSTANS (CO) for degradation (Jang et al., 2008; Liu et al., 2008). Moreover, COP1 can interact with the substrate adaptor EARLY FLOWERING 3 (ELF3) to modulate light input signal to the circadian clock by destabilizing GIGANTEA (GI) protein (Yu et al., 2008).

SUPPRESSOR OF PHYA-105 (SPA1) was first identified as a repressor of phyA (Hoecker et al., 1998). Subsequent studies found three additional SPA1-like proteins in the Arabidopsis genome, named SPA2, SPA3, and SPA4 (Laubinger and Hoecker, 2003; Laubinger et al., 2004). Biochemical analysis demonstrated that SPA1 interacts with COP1 (Hoecker and Quail, 2001; Saijo




Figure 10. Dark-grown *cop1* mutant seedlings phenotypically mimic light-grown wild-type seedlings.

et al., 2003; Seo et al., 2003), and interestingly, the SPA proteins can self-associate or interact with each other, forming a heterogeneous group of COP1/SPA complexes in Arabidopsis (Zhu et al., 2008). Genetic analysis showed that the four SPA genes are partially redundant in mediating light responses at both seedling and adult stages (Laubinger et al., 2004). Moreover, the quadruple *spa* mutant displays a phenotype similar to that of strong *cop1* alleles (Laubinger et al., 2004), consistent with the notion that the SPA proteins work in concert with COP1 in controlling photomorphogenesis.

Empfindlicher Im Dunkelroten Licht 1 (EID1). EID1 is an F-box protein that functions as a negative regulator in phyA-specific light signaling (Buche et al., 2000; Dieterle et al., 2001). The fact that EID1 interacts with several Arabidopsis Skp1-like (ASK) proteins and Cullin1 suggests that EID1 is a component of a SCF (SKP1/Cullin1/F-box protein) ubiquitin ligase complex targeting positively acting component(s) of phyA signaling pathway to ubiquitin-dependent proteolysis (Dieterle et al., 2001; Marrocco et al., 2006). A unique feature of the *eid1* mutant is a shift in the peak of the action spectra of phyA-mediated hypocotyl elongation from FR to R part of the spectrum (Dieterle et al., 2001; Zhou et al., 2002). Although both EID1 and SPA1 function as negatively acting components in phyA-specific light signaling, mutant analysis indicated that EID1 and SPA1 have different but overlapping functions in phyA-dependent signal transduction chains (Zhou et al., 2002). Interestingly, a L946F mutation in the HKRD domain of phyA (named *phyA-402* allele) was found to suppress the hypersensitive phenotype of the *eid1-3* mutant (Muller et al., 2009). However, when *phyA-402* is introgressed into the wild-type background, only moderate phenotype was observed, indicating that the mutation mainly alters phyA functions in an EID1-dependent signaling cascade (Muller et al., 2009).

Positive Regulators of Phytochrome Signaling

HY5 and HYH. HY5, a constitutively nuclear bZIP protein, is the first known and most extensively studied transcription factor involved in promoting photomorphogenesis under a wide spectrum of wavelengths, including FR, R, B, and UV-B (Koornneef et al.,

1980; Oyama, et al., 1997; Osterlund et al., 2000; Ulm et al., 2004). It was shown that the abundance of HY5 protein is directly correlated with the extent of photomorphogenic development (Osterlund et al., 2000). Recent chromatin immunoprecipitation (ChIP)-chip studies revealed that HY5 binds directly to a large number of genomic sites, mainly at the promoter regions of annotated genes (Lee et al., 2007; Zhang et al., 2011). It seems that HY5 directly mediates both upregulation and downregulation of gene expression by light. The gene expression regulation attributable to HY5 is included largely within genes that are regulated by light and comprises ~20% of all light-regulated genes (Ma et al., 2002). Therefore, HY5 is likely to be a high hierarchical regulator of the transcriptional cascades involved in seedling photomorphogenesis (Lee et al., 2007).

COP1 is capable of directly interacting with HY5 in the nucleus through its WD40 repeat domain and targets HY5 for proteasome-mediated degradation (Ang et al., 1998; Osterlund et al., 2000). HY5 has a homolog in the Arabidopsis genome, named HYH, and interestingly, HYH was also shown to be a target of COP1 (Holm et al., 2002). HY5 and HYH physically interact with COP1 through a COP1-interaction motif (Holm et al., 2001, 2002). Consistent with the finding that COP1 forms protein complexes with the SPA proteins, SPA1 contributes to the down-regulation of HY5 abundance (Saijo et al., 2003). Multiple photoreceptors, including phytochromes and cryptochromes, promote the accumulation of HY5 under specific light conditions, possibly by reducing the nuclear abundance of COP1 (Osterlund and Deng, 1998; Osterlund et al., 2000). However, little is known as to how the light-activated photoreceptors regulate the activities of COP1, as well as other COP/DET/FUS proteins.

HFR1 and LAF1. HFR1, an atypical bHLH protein, was originally identified as a positive regulator specific to phyA signaling (Fairchild et al., 2000; Fankhauser and Chory, 2000; Soh et al., 2000). However, subsequent studies revealed that HFR1 is also a component of cry1-mediated B light signaling (Duek and Fankhauser, 2003). Thus, HFR1 may represent a point of signal integration from phyA and cry1, either as a convergence of two independent signaling pathways or as a result of interaction of phyA and cry1 at the photoreceptor molecule level (Ahmad et al., 1998). It was demonstrated that HFR1 is capable of forming homodimers as well as heterodimers with PIF3. However, in contrast to PIF3, HFR1 does not bind directly to either phyA or phyB, although the HFR1/PIF3 complex can bind preferentially to the Pfr form of both phyA and phyB (Fairchild et al., 2000). In addition, unlike the PIF proteins, HFR1 contains an atypical basic domain which might not be functional for directly binding to DNA (Fairchild et al., 2000; Heim et al., 2003). Consistent with this proposal, HFR1 was recently shown to prevent an exaggerated shade avoidance response by forming non-DNA-binding heterodimers with PIF4 and PIF5, two bHLH transcription factors directly regulating the expression of shade-responsive marker genes (Sessa et al., 2005; Lorrain et al., 2008; Hornitschek et al., 2009; Galstyan et al., 2011). Moreover, the finding that a stabilized version of HFR1 leads to a constitutively photomorphogenic phenotype in darkness (similar to that of the *pif1 pif3 pif4 pif5* quadruple mutants) suggest that HFR1 may function to sequester all these PIF proteins (Yang et al., 2003; Leivar et al., 2008; Shin et al., 2009).

LAF1 is an R2R3-MYB transcription factor with trans-activation activity, and functions as a positive component of phyA signaling

(Ballesteros, et al., 2001). However, no direct target gene of LAF1 has been reported so far. The *hfr1 laf1* double mutant has an additive phenotype compared to the two single mutants, indicating that HFR1 and LAF1 regulate largely independent pathways (Jang et al., 2007). Interestingly, both HFR1 and LAF1 were found to be the targets of COP1's E3 ubiquitin ligase activity *in vitro*, and in both cases, genetic and physical interactions between HFR1/LAF1 and COP1 were also observed (Seo et al., 2003; Jang et al., 2005; Yang et al., 2005). Moreover, it was recently shown that HFR1 also physically interacts with LAF1, and this interaction stabilizes each other through inhibition of ubiquitination by COP1, thereby enhancing phyA photoresponses (Jang et al., 2007).

PHYTOCHROME CONTROL OF NUCLEAR GENE EXPRESSION

Microarray analyses conducted in the last decade revealed genome-wide gene expression profiles regulated by light. About 10% or so of the genes in the Arabidopsis genome display phytochrome-regulated changes in expression during the seedling de-etiolation transition triggered by initial exposure of etiolated seedlings to light (Tepperman et al., 2004, 2006; Quail, 2011). These genes include numerous photosynthetic genes related to the biogenesis of active chloroplasts, various auxin-, gibberellin-, cytokinin- and ethylene hormone pathway-related genes potentially mediating growth responses, and metabolic genes reflecting the transition from heterotrophic to autotrophic growth (Tepperman et al., 2004, 2006; Quail, 2011). It is believed that the changes in expression of this large number of light-responsive genes ultimately lead to various morphogenic changes during seedling de-etiolation. Significantly, among functionally classifiable early light-responsive genes responding within 1 hour of FR or R light exposure, 44% (for FR light) and 25% (for R light) encode transcription factors (Tepperman et al., 2001, 2004, 2006), suggesting that they may represent a master set of transcriptional regulators that orchestrate the expression of the downstream target genes in the phytochrome-directed transcriptional network.

Extensive studies have shed light on the mechanisms by which phytochromes regulate light-responsive gene expression. Firstly, phytochromes may promptly alter the expression of a large number of genes by inducing rapid phosphorylation and proteolysis of PIF transcription factors, as discussed above. Secondly, it is generally assumed that phytochromes rapidly inactivate the COP/DET/FUS proteins in response to light, which leads to the accumulation of photomorphogenesis-promoting transcription factors, such as HY5, HYH, LAF1 and HFR1, although the mechanisms governing this process are largely unknown (Figure 9). In addition, it has been proposed that the direct protein-protein interactions between phytochromes and COP1/SPA proteins might be responsible for the rapid, initial inactivation of COP1 activity, whereas long-term inactivation of COP1 is achieved by subsequent depletion of the molecule from the nucleus (Wang and Deng, 2003).

Moreover, it has been suggested that phytochromes might directly target light signals to the light-responsive gene promoters. phyB was shown to bind reversibly to G-box-bound PIF3 specifically upon light-triggered conversion of the photoreceptor to its biologically active Pfr form, thus suggesting a provocative model in which phytochromes may function as integral light-switchable

components of transcription regulator complexes, permitting direct targeting of light signals to target gene promoters (Martinez-Garcia et al., 2000; Quail, 2002). The direct interactions of PIF1 and PIF3 with phyA suggest that phyA could be targeted to gene promoters as well. However, conclusive evidence in favor of this model is still lacking. There is no evidence that phytochromes are associated with DNA *in vivo*, and that phytochromes indeed modulate transcription on the target gene promoters.

PHYTOCHROME SIGNALING AND THE CIRCADIAN CLOCK

The circadian clock controls many metabolic, developmental and physiological processes in a time-of-day-specific manner in both plants and animals (McClung, 2008; de Montaigu et al., 2010). Although circadian rhythms are endogenously generated, they can be modulated by external cues such as light and temperature, thus allowing plants to anticipate and adapt to daily and seasonal changes in their environment. Light signals perceived and transduced by phytochromes and cryptochromes ensure the clock is in tune with the external light/dark cycles. This process is known as entrainment. phyA, phyB, phyD, and phyE act as photoreceptors in R light input to the clock, while phyA and the cryptochromes cry1 and cry2 act as photoreceptors in B light input (Devlin and Kay, 2000). Interestingly, it was shown that phyA acts in low-intensity R light for circadian control, while phyB functions in high-intensity R light (Somers et al., 1998). In FR light, phyA is expected to be the only active photoreceptor transducing the light input to the circadian clock (Wenden et al., 2011). A recent report showed that Arabidopsis mutants deficient in all five phytochromes still displayed clock-controlled robust rhythmic oscillations of leaf position, indicating that phytochromes are not part of the core mechanism of the circadian clock and that other photoreceptors are sufficient for entrainment (Strasser et al., 2010). It should be noted that the LOV/F-box/Kelch-repeat family photoreceptors ZTL, LKP2 and FKF are also involved in the regulation of the circadian clock in Arabidopsis (Nelson et al., 2000; Somers et al., 2000; Schultz et al., 2001; Imaizumi et al., 2003; Mas et al., 2003; Somers et al., 2004; Kim et al., 2007; Demarsy and Fankhauser, 2009; Baudry et al., 2010).

The rhythm of leaf movement in the wild-type Arabidopsis plants can be reset by FR light, but this resetting was absent in the *phyA*, *hy1* and *hy3* mutants, suggesting that phyA signaling pathway is required for the entrainment of the circadian clock (Yanovsky et al., 2000, 2001). Interestingly, the *hy3* mutant also shows an enhanced response to R light during seedling de-etiolation, and shows disrupted rhythmicity of central-clock and clock-output gene expression in continuous R light (Allen et al., 2006). Further, FHY3 is required for the clock resetting in response to R light pulses, suggesting that FHY3 functions in gating phytochrome signaling to the circadian clock (Allen et al., 2006). Recently, FHY3 and its homolog FAR1 were shown to bind directly to the promoter of *EARLY FLOWERING 4 (ELF4)*, a component of the central oscillator of Arabidopsis circadian clock (Li et al., 2011). Interestingly, HY5, a well-characterized bZIP transcription factor involved in promoting photomorphogenesis, and CIRCADIAN CLOCK ASSOCIATED 1 (CCA1) and LATE ELONGATED HYPOCOTYL (LHY), two MYB-related transcription factors that are key components of the central oscillator, were also shown to

bind directly to the *ELF4* gene promoter via their respective *cis*-elements. FHY3, FAR1 and HY5 activate *ELF4* expression during the day, whereas CCA1 and LHY suppress *ELF4* expression periodically at dawn. Thus, this set of light- and circadian-regulated transcription factors act directly and coordinately at the *ELF4* promoter to regulate its cyclic expression (Li et al., 2011). In addition, phyA itself is essential for the expression of *ELF4* (Khanna et al., 2003), and *ELF4* is a likely target mediating phyA-transduced FR light input to the circadian clock (Wenden et al., 2011). Moreover, phyA was shown to directly associate with FHY3 *in vivo* (Saijo et al., 2008). These data suggest that FHY3 may represent a potential molecular link directly gating the environmental light signals perceived by phytochrome to the circadian clock. It is also interesting to note that FHY3 directly binds to the promoter regions of other circadian genes such as *CCA1* and *CCA1 HIKING EXPEDITION (CHE)* (Li et al., 2011; Ouyang et al., 2011), while its co-regulator HY5 binds to the promoters of *CCA1*, *LHY*, *TOC1*, *ELF3* and *GI* (Lee et al., 2007). Thus, whether FHY3 could regulate other clock genes (such as *CCA1*) together with HY5 will be an intriguing question for future studies.

Another mechanism of gating the light input to the circadian clock, called the external coincidence model, has been proposed in the recent years. In Arabidopsis, *CONSTANS (CO)* plays a central role in the induction of flowering by long days (LDs) (Putterill et al., 1995; Robson et al., 2001). *CO* encodes a nuclear protein containing zinc fingers, and activates the transcription of floral regulators such as *FLOWERING LOCUS T (FT)* in the light. The mRNA level of *CO* is tightly controlled by the circadian clock and shows a striking temporal pattern of expression. Under LDs *CO* mRNA peaks before dusk and stays high until the following dawn, whereas under short days (SDs) *CO* mRNA peaks during the night (Suarez-Lopez et al., 2001). Under these conditions, when *CO* mRNA expression coincides with the exposure of plants to light, photoreceptors enhance the level and activity of the *CO* protein (Valverde et al., 2004). *CO* then triggers flowering by activating *FT* transcription. Genetic experiments demonstrate that photoreceptors cry1, cry2 and phyA promote flowering and stabilize the *CO* protein, whereas phyB delays flowering and promotes the degradation of *CO* (Johnson et al., 1994; Guo et al., 1998; El-Assal et al., 2001; Yanovsky and Kay, 2002; Cerdan and Chory, 2003; Searle and Coupland, 2004; Valverde et al., 2004). Therefore, the coincidence between *CO* mRNA and exposure to light (thus *CO* stability could be regulated post-transcriptionally by photoreceptors) is required to promote flowering.

On the other hand, several clock-related genes are involved in phytochrome signaling. For example, *GIGANTEA (GI)* encodes a novel protein found only in plants, and regulates flowering, circadian rhythms, and seedling photomorphogenesis under continuous R and B light (Araki and Komeda, 1993; Fowler et al., 1999; Park et al., 1999; Huq et al., 2000; Tseng et al., 2004; Mizoguchi et al., 2005; Martin-Tryon et al., 2007). Interestingly, although *GI* does not affect the phyA HIRs, it does affect phyA-mediated VLFs via mechanisms that do not obviously involve its circadian functions (Oliverio et al., 2007). *ELF3* is a highly conserved plant-specific nuclear protein which has been suggested to be part of the central clock oscillator and to act as a link between light and the circadian clock (Hicks et al., 1996; Zagotta et al., 1996; McWatters et al., 2000; Covington et al., 2001; Hicks et al., 2001; Liu et al., 2001; Thines and Harmon, 2010). Like *phyB*

mutants, both *gi* and *elf3* mutants display elongated hypocotyls in R light (Zagotta et al., 1996; Huq et al., 2000; Reed et al., 2000). However, the *gi* mutants are late flowering, which is in contrast with the early flowering phenotype of the *phyB* and *elf3* mutants, suggesting that ELF3 and GI play different roles or use different mechanisms in controlling hypocotyl elongation and flowering responses. Interestingly, ELF3 was shown to interact with both *phyB* and GI (Liu et al., 2001; Yu et al., 2008). In addition, ELF3 may act as an adaptor protein allowing COP1 to interact with GI which leads to GI degradation (Yu et al., 2008).

PHOSPHORYLATION/DEPHOSPHORYLATION AND DEGRADATION OF PHYTOCHROMES

Phosphorylation and Dephosphorylation of Phytochromes

It is well established that phytochromes are phosphoproteins, because they could be readily labeled with ^{32}P isotope *in vivo* (Wong et al., 1986; Biermann et al., 1994). The phosphorylation sites of oat *phyA* have been investigated since the 1980s. There are two phosphorylation sites (Ser8 and Ser18) in the NTE region, and one site (Ser599) in the hinge region of oat *phyA*, among which, Ser8 and Ser599 were confirmed as *in vivo* phosphorylation sites (Wong et al., 1986; McMichael and Lagarias, 1990; Lapko et al., 1997, 1999). Ser8 is constitutively phosphorylated in both the Pr and Pfr forms, whereas Ser599 phosphorylation is light dependent and is preferentially phosphorylated in the Pfr form *in vivo* (Lapko et al., 1997, 1999).

The observation that phytochromes are phosphoproteins suggests the existence of protein kinase(s) and phosphatase(s) responsible for phosphorylating and dephosphorylating phytochromes. However, despite the extensive investigations of phytochrome-interacting proteins, there is no report thus far of a protein kinase that can specifically phosphorylate phytochromes. Instead, as mentioned above, two phosphorylation sites in the NTE region of oat *phyA*, i.e. Ser8 and Ser18, are autophosphorylated by *phyA* itself *in vitro* (Han et al., 2010). Mutations of these two serines to alanines effectively abolished autophosphorylation of oat *phyA* *in vitro* (Han et al., 2010). However, another phosphorylation site of oat *phyA*, Ser599, is not autophosphorylated by *phyA* (Kim et al., 2004; Han et al., 2010). Therefore, the protein kinase(s) responsible for phosphorylating this site of oat *phyA* will be a major interest for future studies.

In contrast to the failure of identifying a phytochrome-associated kinase, there are several studies reporting the discovery of protein phosphatases that could dephosphorylate phytochromes. A phytochrome-associated protein phosphatase 2A, designated FyPP, was shown to interact with and dephosphorylate *phyA* (Kim et al., 2002). The transgenic Arabidopsis plants with over-expressed or suppressed FyPP levels exhibited delayed or accelerated flowering, respectively, indicating that FyPP modulates phytochrome-mediated light signaling in the timing of flowering (Kim et al., 2002). PHYTOCHROME-ASSOCIATED PROTEIN PHOSPHATASE 5 (PAPP5) was isolated from yeast two-hybrid screen using the full-length Arabidopsis *phyA* as bait (Ryu et al., 2005). PAPP5 interacts with both *phyA* and *phyB*, and dephosphorylates all three phosphor-serine residues of oat *phyA* in a Pfr-specific manner. It was shown that PAPP5 positively influences

the protein stability of phytochrome and the interaction of phytochrome with a downstream signal transducer NDPK2 (Ryu et al., 2005). Another phosphatase, PHYTOCHROME-ASSOCIATED PROTEIN PHOSPHATASE 2C (PAPP2C), was recently shown to interact with both *phyA* and *phyB* and effectively dephosphorylate phytochromes *in vitro* (Phee et al., 2008).

What are the *in vivo* functional roles of phytochrome phosphorylation and dephosphorylation? Recent studies suggest two possible roles: controlling phytochrome stability, and regulating interactions of phytochromes with downstream signal transducers. In the last two decades, great efforts have been made to investigate the role of a stretch of serine residues (including Ser8 and Ser18 of oat *phyA*) in the first 20 amino acids of *phyA* proteins, and it has been well-documented that substitution of these serines to alanines, or deletion of this 20-aa region, results in an increased biological activity of *phyA*, suggesting that these serine residues are involved in desensitization of *phyA* signaling (Stockhaus et al., 1992; Jordan et al., 1996, 1997; Casal et al., 2002; Han et al., 2010). Recently it was shown that the increased activity of mutated *phyA* proteins is related to the increased *phyA* stability (Han et al., 2010). Therefore, it is likely that autophosphorylation of serine sites in the NTE of *phyA* controls *phyA* stability.

However, phosphorylation of another serine residue, Ser599 in the hinge region of oat *phyA*, does not affect phytochrome stability (Kim et al., 2004). Instead, it was shown that phosphorylation of Ser599 prevents the interaction of *phyA* with its signal transducers such as NDPK2 and PIF3, suggesting that phosphorylation of Ser599 serves as a signal modulating switch affecting protein-protein interactions between phytochrome and its signal transducers (Kim et al., 2004). Interestingly, an earlier study showed that if Ser599 was mutated (Ser599Lys), *phyA* autophosphorylation and phosphorylation of its substrate PKS1 were no longer regulated by light (Fankhauser et al., 1999), suggesting that Ser599 plays an important role in the regulation of phytochrome kinase activity as well.

A recent study firstly detected an *in vivo* phosphorylated form of Arabidopsis *phyA* protein (Saijo et al., 2008). It was shown that phosphorylated *phyA* preferentially associates with COP1/SPA1 complex for degradation, whereas unphosphorylated *phyA* predominantly associates with *phyA* signaling intermediates FHY1 and FHY3 for signal transduction. Interestingly, COP1 associates with unphosphorylated *phyA* in the absence of FHY1 or FHY3, suggesting that *phyA* associations with FHY1 and FHY3 protect unphosphorylated *phyA* from being recognized by COP1/SPA1 complex (Saijo et al., 2008). Therefore, these data suggest that light-induced *phyA* phosphorylation *in vivo* acts as a switch controlling differential interactions of the photoreceptor with signal propagation or attenuation machineries. However, how many phosphorylation sites contribute to produce this phosphorylated form of Arabidopsis *phyA* protein needs to be further characterized.

Degradation of Phytochromes

As mentioned above, *phyA* is a light labile phytochrome. *phyA* is most abundant in etiolated seedlings, but its level drops up to 100-fold after exposure to light (Clough and Vierstra, 1997;

Sharrock and Clack, 2002). The proteolysis of phyA following photoconversion from Pr to Pfr is rapid, as the Pr form has a half-life of approximately 1 week, whereas the Pfr form has a half-life of only 1–2 h (Clough and Vierstra, 1997). It was observed more than two decades ago that phyA is ubiquitinated *in vivo*, and is degraded by the ubiquitin/26S proteasome pathway (Shanklin et al., 1987; Jabben et al., 1989a, b; Clough and Vierstra, 1997).

However, the E3 ligase responsible for phyA ubiquitination had not been identified until 2004, when COP1 was shown to ubiquitinate phyA *in vitro* (Seo et al., 2004). COP1 and phyA colocalize in the nuclear bodies, and the two proteins physically interact with each other mediated by the PRD domain of phyA and the WD40 domain of COP1. Consistently, the degradation rate of phyA is decreased in *cop1* mutants (Seo et al., 2004). This finding was extended by a later report that the light signaling repressors, SPA proteins, contribute to COP1-mediated degradation of phyA, and that a COP1/SPA1 protein complex is tightly associated with phyA ubiquitination activity (Saijo et al., 2008). Consistent with the notion that phyA phosphorylation status controls its protein stability, the phosphorylated phyA form was suggested to be a preferred target for COP1-mediated degradation (Saijo et al., 2008).

Moreover, it is notable that light-induced phyA degradation still occurs in the null *cop1-5* allele, suggesting the presence of a COP1-independent phyA-degradation pathway (Saijo et al., 2008). Consistent with this assumption, it was observed long ago that a R light pulse promotes the rapid formation of phyA-containing cytosolic spots, referred to as sequestered areas of phytochromes (SAPs), which appear prior to the nuclear transport of phyA and is thought to be the place of phyA ubiquitination and degradation (Speth et al., 1986; Nagatani, 2004; Kevei et al., 2007). The cytosolic degradation of phyA is also supported by the recent reports that phyA is still degraded in the *fhyl fhil* double mutants, in which phyA is maintained in the cytosol, although at a slower rate than in the wild-type plants (Rosler et al., 2007; Debrieux and Fankhauser, 2010).

Although the type II phytochromes (phyB–phyE in Arabidopsis) are described as “light stable”, they are slowly degraded upon irradiation with R light (Sharrock and Clack, 2002). Recently, a report showed that phyB is stable in darkness, but in R, a fraction of phyB translocates into the nucleus and becomes degraded by the ubiquitin/26S proteasome pathway (Jang et al., 2010). Nuclear phyB degradation is also mediated by COP1 E3 ligase, which preferentially interacts with the N-terminal region of phyB. Interestingly, phyB polyubiquitination by COP1 *in vitro* can be enhanced by different PIF proteins that promote COP1/phyB interaction. Consistent with these results, nuclear phyB accumulates to higher levels in *pif* and *cop1* mutants. Moreover, COP1 was shown to interact with and ubiquitinate phyC–phyE *in vitro* (Jang et al., 2010). Therefore, COP1 is an E3 ligase for all five phytochromes in Arabidopsis.

CYTOPLASMIC EVENTS OF PHYTOCHROME SIGNALING

Besides nuclear signaling events, phytochrome signaling likely entails cytoplasmic events based on the following reasons. Firstly, as mentioned above, the nuclear import of some phytochrome

species (such as phyB) takes hours, and their nuclear actions obviously could not explain some rapid phytochrome responses which occur within minutes of light irradiation, such as the change in hypocotyl growth rate (Parks and Spalding, 1999). Secondly, phytochromes interact with a number of cytoplasmic proteins, such as PKS1 and NDPK2 (Choi et al., 1999; Fankhauser et al., 1999). Thirdly, genetic studies have identified several cytoplasmic proteins as signaling intermediates, such as PAT1 and FIN219 for phyA signaling (Bolle et al., 2000; Hsieh et al., 2000).

The fastest phytochrome response described so far is the cytoplasmic streaming of *Vallisneria gigantea*, which can be locally stimulated with pulsed R light and becomes measurable within 2.5 s, and is likely mediated by phyB-type phytochromes (Takagi et al., 2003). As FHY1/FHL are indispensable for phyA nuclear accumulation (see above), the cytoplasmic responses regulated by phyA were identified recently using the *fhyl fhil* double mutant (Rosler et al., 2007). The phyA-specific cytoplasmic responses include the R-enhanced phototropism, abrogation of gravitropism in B, and the inhibition of hypocotyl elongation in B (Rosler et al., 2007). It is noteworthy that the latter two cytoplasmic responses of phyA occur in B light, which might be explained in part by a recent observation that phyA coordinates the localization and distribution of PHOT1 (Han et al., 2008).

Previous microinjection and pharmacological studies suggested the involvement of G proteins, cGMP and Ca²⁺/calmodulin in phytochrome signaling (Bowler et al., 1994; Neuhaus et al., 1997). In particular, it was reported that transgenic Arabidopsis plants conditionally overexpressing the α subunit (encoded by *GPA1* gene) of the heterotrimeric G protein under the control of a glucocorticoid-inducible promoter exhibited a light-dependent hypersensitive response as a result of reduced hypocotyl cell elongation (Okamoto et al., 2001). However, a separate study reported that loss-of-function *gpa1* mutants display partial de-etiolation in the dark, with short hypocotyls and open apical hooks typical of light-irradiated seedlings (Ullah et al., 2001). The short hypocotyl of *gpa1* seedlings was reportedly due to a defect in cell division, but not cell elongation (Ullah et al., 2001). To resolve these discrepancies regarding GPA1 function in photomorphogenesis, Jones et al. (2003) re-evaluated the roles of the heterotrimeric G protein employing multiple alleles of *gpa1* and *agb1* (impaired in the β subunit) and their double mutants. They concluded that these mutants have wild-type sensitivity to R and FR light. In addition, there is no apparent alteration in R or FR sensitivity in transgenic plants overexpressing GPA1 or AGB1. The observed shorter hypocotyls and partially open cotyledons of *gpa1* and *agb1* mutants grown in the dark are mainly caused by a defect in cell division, rather than cell elongation (Jones et al., 2003).

Genetic studies have also provided supporting evidence for the involvement of Ca²⁺ in phytochrome signaling. The *sub1* mutant exhibits hypersensitive responses to both FR and B light (Guo et al., 2001). The *SUB1* gene was found to encode a Ca²⁺-binding protein. Genetic interaction studies suggest that SUB1 is a component of cryptochrome signaling pathway and is a modulator of phyA signaling pathway. Further, SUB1 negatively regulates HY5, a photomorphogenesis-promoting transcription factor (Guo et al., 2001).

FUTURE ISSUES

The last two decades have seen significant progress in unraveling the signaling mechanisms of plant phytochromes. However, several important questions about phytochromes still remain to be answered:

1. We still do not understand the earliest signaling events following photo-activation. Which activities of phytochromes are sufficient to trigger light responses?
2. Further to the first question, are phytochromes light-regulated kinases? If so, which domain is the kinase domain? What is the biological role of this kinase activity in inducing light responses?
3. How do phytochromes induce the phosphorylation of PIFs? How are PIFs degraded?
4. How are nuclear bodies formed? What are their precise functions in phytochrome signaling?
5. Are phytochromes associated with the target gene promoters to directly regulate their gene expression?
6. How do phytochromes inactivate the COP1/SPA protein degradation machinery upon photo-activation?
7. How are phytochromes phosphorylated? In addition to auto-phosphorylation, which kinase(s) are responsible for phosphorylating phytochromes?
8. How are phytochromes degraded in the nucleus and in the cytosol?

Further investigation and ultimate elucidation of these questions and others will undoubtedly shed more light on phytochrome signaling mechanisms and photomorphogenesis in general.

ACKNOWLEDGMENTS

We apologize to the authors whose work might have been overlooked here. We thank William Terzaghi for reading and commenting on the manuscript. This work was supported by a National Institute of Health grant (GM47850) to X.W.D. and a National Science Foundation award (IOS-1026630) to H.W. Xing Wang Deng's laboratory at Peking University was supported by grants from the Ministry of Science and Technology of China (2009DFB30030) and the Ministry of Agriculture of China (2009ZX08012-021B).

REFERENCES

- Ahmad, M., and Cashmore, A.R. (1997). The blue-light receptor cryptochrome 1 shows functional dependence on phytochrome A or phytochrome B in *Arabidopsis thaliana*. *Plant J.* **11**: 421-427.
- Ahmad, M., Jarillo, J.A., Smirnova, O., and Cashmore, A.R. (1998). The CRY1 blue light photoreceptor of *Arabidopsis* interacts with phytochrome A *in vitro*. *Mol. Cell* **1**: 939-948.
- Allen, T., Koustenis, A., Theodorou, G., Somers, D.E., Kay, S.A., Whitelam, G.C., and Devlin, P.F. (2006). *Arabidopsis* FHY3 specifically gates phytochrome signaling to the circadian clock. *Plant Cell* **18**: 2506-2516.
- Al-Sady, B., Kikis, E.A., Monte, E., and Quail, P.H. (2008). Mechanistic duality of transcription factor function in phytochrome signaling. *Proc. Natl. Acad. Sci. USA* **105**: 2232-2237.
- Al-Sady, B., Ni, W., Kircher, S., Schafer, E., and Quail, P.H. (2006). Photoactivated phytochrome induces rapid PIF3 phosphorylation prior to proteasome-mediated degradation. *Mol. Cell* **23**: 439-446.
- Andel, F., 3rd, Lagarias, J.C., and Mathies, R.A. (1996). Resonance Raman analysis of chromophore structure in the lumi-R photoproduct of phytochrome. *Biochemistry* **35**: 15997-16008.
- Ang, L.H., Chattopadhyay, S., Wei, N., Oyama, T., Okada, K., Batschauer, A., and Deng, X.W. (1998). Molecular interaction between COP1 and HY5 defines a regulatory switch for light control of *Arabidopsis* development. *Mol. Cell* **1**: 213-222.
- Araki, T., and Komeda, Y. (1993). Flowering in darkness in *Arabidopsis thaliana*. *Plant J.* **4**: 801-811.
- Aukerman, M.J., Hirschfeld, M., Wester, L., Weaver, M., Clack, T., Amasino, R.M., and Sharrock, R.A. (1997). A deletion in the *PHYD* gene of the *Arabidopsis* Wassilewskija ecotype defines a role for phytochrome D in red/far-red light sensing. *Plant Cell* **9**: 1317-1326.
- Bae, G., and Choi, G. (2008). Decoding of light signals by plant phytochromes and their interacting proteins. *Annu. Rev. Plant Biol.* **59**: 281-311.
- Bailey, P.C., Martin, C., Toledo-Ortiz, G., Quail, P.H., Huq, E., Heim, M.A., Jakoby, M., Werber, M., and Weisshaar, B. (2003). Update on the basic helix-loop-helix transcription factor gene family in *Arabidopsis thaliana*. *Plant Cell* **15**: 2497-2502.
- Ballesteros, M.L., Bolle, C., Lois, L.M., Moore, J.M., Vielle-Calzada, J.P., Grossniklaus, U., and Chua, N.H. (2001). LAF1, a MYB transcription activator for phytochrome A signaling. *Genes Dev.* **15**: 2613-2625.
- Barnes, S.A., Quaggio, R.B., Whitelam, G.C., and Chua, N.H. (1996). *thyl1* defines a branch point in phytochrome A signal transduction pathways for gene expression. *Plant J.* **10**: 1155-1161.
- Baudry, A., Ito, S., Song, Y.H., Strait, A.A., Kiba, T., Lu, S., Henriques, R., Pruneda-Paz, J.L., Chua, N.H., Tobin, E.M., Kay, S.A., and Imai-zumi, T. (2010). F-box proteins FKF1 and LKP2 act in concert with ZEITLUPE to control *Arabidopsis* clock progression. *Plant Cell* **22**: 606-622.
- Bauer, D., Viczian, A., Kircher, S., Nobis, T., Nitschke, R., Kunkel, T., Panigrahi, K.C., Adam, E., Fejes, E., Schafer, E., and Nagy, F. (2004). Constitutive Photomorphogenesis 1 and multiple photoreceptors control degradation of Phytochrome Interacting Factor 3, a transcription factor required for light signaling in *Arabidopsis*. *Plant Cell* **16**: 1433-1445.
- Biermann, B.J., Pao, L.I., and Feldman, L.J. (1994). Pre-specific phytochrome phosphorylation *in vitro* by a protein kinase present in anti-phytochrome maize immunoprecipitates. *Plant Physiol.* **105**: 243-251.
- Bolle, C., Koncz, C., and Chua, N.H. (2000). PAT1, a new member of the GRAS family, is involved in phytochrome A signal transduction. *Genes Dev.* **14**: 1269-1278.
- Borthwick, H.A., Hendricks, S.B., Parker, M.W., Toole, E.H., and Toole, V.K. (1952). A reversible photoreaction controlling seed germination. *Proc. Natl. Acad. Sci. USA* **38**: 662-666.
- Botto, J.F., Sanchez, R.A., Whitelam, G.C., and Casal, J.J. (1996). Phytochrome A mediates the promotion of seed germination by very low fluences of light and canopy shade light in *Arabidopsis*. *Plant Physiol.* **110**: 439-444.
- Bowler, C., Neuhaus, G., Yamagata, H., and Chua, N.H. (1994). Cyclic GMP and calcium mediate phytochrome phototransduction. *Cell* **77**: 73-81.
- Boylan, M.T., and Quail, P.H. (1996). Are the phytochromes protein kinases? *Protoplasma* **195**: 12-17.
- Bu, Q., Zhu, L., Dennis, M.D., Yu, L., Lu, S.X., Person, M.D., Tobin, E.M., Browning, K.S., and Huq, E. (2011). Phosphorylation by CK2 enhances the rapid light-induced degradation of Phytochrome Interacting Factor 1 in *Arabidopsis*. *J. Biol. Chem.* **286**: 12066-12074.
- Buche, C., Poppe, C., Schafer, E., and Kretsch, T. (2000). *eid1*: a new

- Arabidopsis mutant hypersensitive in phytochrome A-dependent high-irradiance responses. *Plant Cell* **12**: 547-558.
- Casal, J.J., Davis, S.J., Kirchenbauer, D., Viczian, A., Yanovsky, M.J., Clough, R.C., Kircher, S., Jordan-Beebe, E.T., Schafer, E., Nagy, F., and Vierstra, R.D.** (2002). The serine-rich N-terminal domain of oat phytochrome A helps regulate light responses and subnuclear localization of the photoreceptor. *Plant Physiol.* **129**: 1127-1137.
- Casal, J.J., and Mazzella, M.A.** (1998). Conditional synergism between cryptochrome 1 and phytochrome B is shown by the analysis of *phyA*, *phyB*, and *hy4* simple, double, and triple mutants in Arabidopsis. *Plant Physiol.* **118**: 19-25.
- Casal, J.J., Sanchez, R.A., and Botto, J.F.** (1998). Modes of action of phytochromes. *J. Exp. Bot.* **49**: 127-138.
- Castillon, A., Shen, H., and Huq, E.** (2007). Phytochrome Interacting Factors: central players in phytochrome-mediated light signaling networks. *Trends Plant Sci.* **12**: 514-521.
- Cerdan, P.D., and Chory, J.** (2003). Regulation of flowering time by light quality. *Nature* **423**: 881-885.
- Chen, H., Huang, X., Gusmaroli, G., Terzaghi, W., Lau, O.S., Yanagawa, Y., Zhang, Y., Li, J., Lee, J.H., Zhu, D., and Deng, X.W.** (2010). Arabidopsis CULLIN4-damaged DNA binding protein 1 interacts with CONSTITUTIVELY PHOTOMORPHOGENIC1-SUPPRESSOR OF PHYA complexes to regulate photomorphogenesis and flowering time. *Plant Cell* **22**: 108-123.
- Chen, H., Shen, Y., Tang, X., Yu, L., Wang, J., Guo, L., Zhang, Y., Zhang, H., Feng, S., Strickland, E., Zheng, N., and Deng, X.W.** (2006). Arabidopsis CULLIN4 forms an E3 ubiquitin ligase with RBX1 and the CDD complex in mediating light control of development. *Plant Cell* **18**: 1991-2004.
- Chen, M.** (2008). Phytochrome nuclear body: an emerging model to study interphase nuclear dynamics and signaling. *Curr. Opin. Plant Biol.* **11**: 503-508.
- Chen, M., Galvao, R.M., Li, M., Burger, B., Bugea, J., Bolado, J., and Chory, J.** (2010). Arabidopsis HEMERA/pTAC12 initiates photomorphogenesis by phytochromes. *Cell* **141**: 1230-1240.
- Chen, M., Schwab, R., and Chory, J.** (2003). Characterization of the requirements for localization of phytochrome B to nuclear bodies. *Proc Natl. Acad. Sci. USA* **100**: 14493-14498.
- Chen, M., Tao, Y., Lim, J., Shaw, A., and Chory, J.** (2005). Regulation of phytochrome B nuclear localization through light-dependent unmasking of nuclear-localization signals. *Curr. Biol.* **15**: 637-642.
- Choi, G., Yi, H., Lee, J., Kwon, Y.K., Soh, M.S., Shin, B., Luka, Z., Hahn, T.R., and Song, P.S.** (1999). Phytochrome signalling is mediated through nucleoside diphosphate kinase 2. *Nature* **401**: 610-613.
- Clack, T., Mathews, S., and Sharrock, R.A.** (1994). The phytochrome apoprotein family in Arabidopsis is encoded by five genes: the sequences and expression of *PHYD* and *PHYE*. *Plant Mol. Biol.* **25**: 413-427.
- Clack, T., Shokry, A., Moffet, M., Liu, P., Faul, M., and Sharrock, R.A.** (2009). Obligate heterodimerization of Arabidopsis phytochromes C and E and interaction with the PIF3 basic helix-loop-helix transcription factor. *Plant Cell* **21**: 786-799.
- Clough, R.C., and Vierstra, R.D.** (1997). Phytochrome degradation. *Plant Cell Environ.* **20**: 713-721.
- Colon-Carmona, A., Chen, D.L., Yeh, K.C., and Abel, S.** (2000). Aux/IAA proteins are phosphorylated by phytochrome *in vitro*. *Plant Physiol.* **124**: 1728-1738.
- Covington, M.F., Panda, S., Liu, X.L., Strayer, C.A., Wagner, D.R., and Kay, S.A.** (2001). ELF3 modulates resetting of the circadian clock in Arabidopsis. *Plant Cell* **13**: 1305-1315.
- Davis, S.J., Bhoo, S.H., Durski, A.M., Walker, J.M., and Vierstra, R.D.** (2001). The heme-oxygenase family required for phytochrome chromophore biosynthesis is necessary for proper photomorphogenesis in higher plants. *Plant Physiol.* **126**: 656-669.
- Davis, S.J., Kurepa, J., and Vierstra, R.D.** (1999a). The *Arabidopsis thaliana* *HY1* locus, required for phytochrome-chromophore biosynthesis, encodes a protein related to heme oxygenases. *Proc. Natl. Acad. Sci. USA* **96**: 6541-6546.
- Davis, S.J., Vener, A.V., and Vierstra, R.D.** (1999b). Bacteriophytochromes: phytochrome-like photoreceptors from nonphotosynthetic eubacteria. *Science* **286**: 2517-2520.
- de Lucas, M., Daviere, J.M., Rodriguez-Falcon, M., Pontin, M., Iglesias-Pedraz, J.M., Lorrain, S., Fankhauser, C., Blazquez, M.A., Titchenko, E., and Prat, S.** (2008). A molecular framework for light and gibberellin control of cell elongation. *Nature* **451**: 480-484.
- de Montaigu, A., Toth, R., and Coupland, G.** (2010). Plant development goes like clockwork. *Trends Genet.* **26**: 296-306.
- Debieux, D., and Fankhauser, C.** (2010). Light-induced degradation of phyA is promoted by transfer of the photoreceptor into the nucleus. *Plant Mol. Biol.* **73**: 687-695.
- Dehesh, K., Franci, C., Parks, B.M., Seeley, K.A., Short, T.W., Tepperman, J.M., and Quail, P.H.** (1993). Arabidopsis *HY8* locus encodes phytochrome A. *Plant Cell* **5**: 1081-1088.
- Demarsy, E., and Fankhauser, C.** (2009). Higher plants use LOV to perceive blue light. *Curr. Opin. Plant Biol.* **12**: 69-74.
- Deng, X.W., Caspar, T., and Quail, P.H.** (1991). *cop1*: a regulatory locus involved in light-controlled development and gene expression in Arabidopsis. *Genes Dev.* **5**: 1172-1182.
- Deng, X.W., Matsui, M., Wei, N., Wagner, D., Chu, A.M., Feldmann, K.A., and Quail, P.H.** (1992). *COP1*, an Arabidopsis regulatory gene, encodes a protein with both a zinc-binding motif and a G beta homologous domain. *Cell* **71**: 791-801.
- Deng, X.W., and Quail, P.H.** (1999). Signalling in light-controlled development. *Semin. Cell Dev. Biol.* **10**: 121-129.
- Desnos, T., Puente, P., Whitelam, G.C., and Harberd, N.P.** (2001). *FHY1*: a phytochrome A-specific signal transducer. *Genes Dev.* **15**: 2980-2990.
- Devlin, P.F., and Kay, S.A.** (2000). Cryptochromes are required for phytochrome signaling to the circadian clock but not for rhythmicity. *Plant Cell* **12**: 2499-2510.
- Devlin, P.F., Patel, S.R., and Whitelam, G.C.** (1998). Phytochrome E influences internode elongation and flowering time in Arabidopsis. *Plant Cell* **10**: 1479-1487.
- Devlin, P.F., Robson, P.R., Patel, S.R., Goosey, L., Sharrock, R.A., and Whitelam, G.C.** (1999). Phytochrome D acts in the shade-avoidance syndrome in Arabidopsis by controlling elongation growth and flowering time. *Plant Physiol.* **119**: 909-915.
- Dieterle, M., Zhou, Y.C., Schafer, E., Funk, M., and Kretsch, T.** (2001). *EID1*, an F-box protein involved in phytochrome A-specific light signaling. *Genes Dev.* **15**: 939-944.
- Duek, P.D., Elmer, M.V., van Oosten, V.R., and Fankhauser, C.** (2004). The degradation of HFR1, a putative bHLH class transcription factor involved in light signaling, is regulated by phosphorylation and requires COP1. *Curr. Biol.* **14**: 2296-2301.
- Duek, P.D., and Fankhauser, C.** (2003). HFR1, a putative bHLH transcription factor, mediates both phytochrome A and cryptochrome signalling. *Plant J.* **34**: 827-836.
- Duek, P.D., and Fankhauser, C.** (2005). bHLH class transcription factors take centre stage in phytochrome signalling. *Trends Plant Sci.* **10**: 51-54.
- Eichenberg, K., Baurle, I., Paulo, N., Sharrock, R.A., Rudiger, W., and Schafer, E.** (2000). Arabidopsis phytochromes C and E have different spectral characteristics from those of phytochromes A and B. *FEBS*

- Lett. **470**: 107-112.
- El-Assal, S.E.D., Alonso-Blanco, C., Peeters, A.J., Raz, V., and Koornneef, M.** (2001). A QTL for flowering time in Arabidopsis reveals a novel allele of *CRY2*. *Nat. Genet.* **29**: 435-440.
- Fairchild, C.D., Schumaker, M.A., and Quail, P.H.** (2000). *HFR1* encodes an atypical bHLH protein that acts in phytochrome A signal transduction. *Genes Dev.* **14**: 2377-2391.
- Fankhauser, C.** (2000). Phytochromes as light-modulated protein kinases. *Semin. Cell. Dev. Biol.* **11**: 467-473.
- Fankhauser, C.** (2001). The phytochromes, a family of red/far-red absorbing photoreceptors. *J. Biol. Chem.* **276**: 11453-11456.
- Fankhauser, C., and Chen, M.** (2008). Transposing phytochrome into the nucleus. *Trends Plant Sci.* **13**: 596-601.
- Fankhauser, C., and Chory, J.** (2000). *RSF1*, an Arabidopsis locus implicated in phytochrome A signaling. *Plant Physiol.* **124**: 39-45.
- Fankhauser, C., Yeh, K.C., Lagarias, J.C., Zhang, H., Elich, T.D., and Chory, J.** (1999). PKS1, a substrate phosphorylated by phytochrome that modulates light signaling in Arabidopsis. *Science* **284**: 1539-1541.
- Feng, S., Martinez, C., Gusmaroli, G., Wang, Y., Zhou, J., Wang, F., Chen, L., Yu, L., Iglesias-Pedraz, J.M., Kircher, S., Schafer, E., Fu, X., Fan, L.M., and Deng, X.W.** (2008). Coordinated regulation of *Arabidopsis thaliana* development by light and gibberellins. *Nature* **451**: 475-479.
- Fowler, S., Lee, K., Onouchi, H., Samach, A., Richardson, K., Morris, B., Coupland, G., and Putterill, J.** (1999). *GIGANTEA*: a circadian clock-controlled gene that regulates photoperiodic flowering in Arabidopsis and encodes a protein with several possible membrane-spanning domains. *EMBO J.* **18**: 4679-4688.
- Franklin, K.A., Davis, S.J., Stoddart, W.M., Vierstra, R.D., and Whitelam, G.C.** (2003a). Mutant analyses define multiple roles for phytochrome C in Arabidopsis photomorphogenesis. *Plant Cell* **15**: 1981-1989.
- Franklin, K.A., Praekelt, U., Stoddart, W.M., Billingham, O.E., Halliday, K.J., and Whitelam, G.C.** (2003b). Phytochromes B, D, and E act redundantly to control multiple physiological responses in Arabidopsis. *Plant Physiol.* **131**: 1340-1346.
- Franklin, K.A., and Quail, P.H.** (2010). Phytochrome functions in Arabidopsis development. *J. Exp. Bot.* **61**: 11-24.
- Furuya, M., and Song, P.S.** (1994). Assembly and properties of holophytochrome. In *Photomorphogenesis in plants*, R.E. Kendrick, and G.H.M. Kronenberg, eds. (Dordrecht, The Netherlands: Kluwer Academic Publishers), pp. 105-140.
- Galstyan, A., Cifuentes-Esquivel, N., Bou-Torrent, J., and Martinez-Garcia, J.F.** (2011). The shade avoidance syndrome in Arabidopsis: a fundamental role for atypical basic helix-loop-helix proteins as transcriptional cofactors. *Plant J.* **66**: 258-267.
- Garner, W.W., and Allard, H.A.** (1920). Effect of the relative length of day and night and other factors of the environment on growth and reproduction in plants. *J. Agric. Res.* **18**: 553-606.
- Genoud, T., Schweizer, F., Tscheuschler, A., Debrieux, D., Casal, J.J., Schafer, E., Hiltbrunner, A., and Fankhauser, C.** (2008). FH1 mediates nuclear import of the light-activated phytochrome A photoreceptor. *PLoS Genet.* **4**: e1000143.
- Gil, P., Kircher, S., Adam, E., Bury, E., Kozma-Bognar, L., Schafer, E., and Nagy, F.** (2000). Photocontrol of subcellular partitioning of phytochrome-B:GFP fusion protein in tobacco seedlings. *Plant J.* **22**: 135-145.
- Guo, H., Mockler, T., Duong, H., and Lin, C.** (2001). SUB1, an Arabidopsis Ca^{2+} -binding protein involved in cryptochrome and phytochrome coaction. *Science* **291**: 487-490.
- Guo, H., Yang, H., Mockler, T.C., and Lin, C.** (1998). Regulation of flowering time by Arabidopsis photoreceptors. *Science* **279**: 1360-1363.
- Hamazato, F., Shinomura, T., Hanzawa, H., Chory, J., and Furuya, M.** (1997). Fluence and wavelength requirements for Arabidopsis *CAB* gene induction by different phytochromes. *Plant Physiol.* **115**: 1533-1540.
- Han, I.S., Tseng, T.S., Eisinger, W., and Briggs, W.R.** (2008). Phytochrome A regulates the intracellular distribution of phototropin 1-green fluorescent protein in *Arabidopsis thaliana*. *Plant Cell* **20**: 2835-2847.
- Han, Y.J., Kim, H.S., Kim, Y.M., Shin, A.Y., Lee, S.S., Bhoo, S.H., Song, P.S., and Kim, J.I.** (2010). Functional characterization of phytochrome autophosphorylation in plant light signaling. *Plant Cell Physiol.* **51**: 596-609.
- Haupt, W., and Hader, D.P.** (1994). Photomovement. In *Photomorphogenesis in plants*, R.E. Kendrick, and G.H.M. Kronenberg, eds. (Dordrecht, The Netherlands: Kluwer Academic Publishers), pp. 707-732.
- Heim, M.A., Jakoby, M., Werber, M., Martin, C., Weisshaar, B., and Bailey, P.C.** (2003). The basic helix-loop-helix transcription factor family in plants: a genome-wide study of protein structure and functional diversity. *Mol. Biol. Evol.* **20**: 735-747.
- Hennig, L., Stoddart, W.M., Dieterle, M., Whitelam, G.C., and Schafer, E.** (2002). Phytochrome E controls light-induced germination of Arabidopsis. *Plant Physiol.* **128**: 194-200.
- Heschel, M.S., Selby, J., Butler, C., Whitelam, G.C., Sharrock, R.A., and Donohue, K.** (2007). A new role for phytochromes in temperature-dependent germination. *New Phytol.* **174**: 735-741.
- Hicks, K.A., Albertson, T.M., and Wagner, D.R.** (2001). *EARLY FLOWERING3* encodes a novel protein that regulates circadian clock function and flowering in Arabidopsis. *Plant Cell* **13**: 1281-1292.
- Hicks, K.A., Millar, A.J., Carre, I.A., Somers, D.E., Straume, M., Meeks-Wagner, D.R., and Kay, S.A.** (1996). Conditional circadian dysfunction of the Arabidopsis *early-flowering 3* mutant. *Science* **274**: 790-792.
- Hiltbrunner, A., Tscheuschler, A., Viczian, A., Kunkel, T., Kircher, S., and Schafer, E.** (2006). FH1 and FHL act together to mediate nuclear accumulation of the phytochrome A photoreceptor. *Plant Cell Physiol.* **47**: 1023-1034.
- Hiltbrunner, A., Viczian, A., Bury, E., Tscheuschler, A., Kircher, S., Toth, R., Honsberger, A., Nagy, F., Fankhauser, C., and Schafer, E.** (2005). Nuclear accumulation of the phytochrome A photoreceptor requires FH1. *Curr. Biol.* **15**: 2125-2130.
- Hirschfeld, M., Tepperman, J.M., Clack, T., Quail, P.H., and Sharrock, R.A.** (1998). Coordination of phytochrome levels in *phyB* mutants of Arabidopsis as revealed by apoprotein-specific monoclonal antibodies. *Genetics* **149**: 523-535.
- Hisada, A., Hanzawa, H., Weller, J.L., Nagatani, A., Reid, J.B., and Furuya, M.** (2000). Light-induced nuclear translocation of endogenous pea phytochrome A visualized by immunocytochemical procedures. *Plant Cell* **12**: 1063-1078.
- Hoecker, U., Xu, Y., and Quail, P.H.** (1998). *SPA1*: a new genetic locus involved in phytochrome A-specific signal transduction. *Plant Cell* **10**: 19-33.
- Hoecker, U., and Quail, P.H.** (2001). The phytochrome A-specific signaling intermediate SPA1 interacts directly with COP1, a constitutive repressor of light signaling in Arabidopsis. *J. Biol. Chem.* **276**: 38173-38178.
- Holm, M., Hardtke, C.S., Gaudet, R., and Deng, X.W.** (2001). Identification of a structural motif that confers specific interaction with the WD40 repeat domain of Arabidopsis COP1. *EMBO J.* **20**: 118-127.
- Holm, M., Ma, L.G., Qu, L.J., and Deng, X.W.** (2002). Two interacting bZIP proteins are direct targets of COP1-mediated control of light-dependent gene expression in Arabidopsis. *Genes Dev.* **16**: 1247-1259.
- Hornitschek, P., Lorrain, S., Zoete, V., Michielin, O., and Fankhauser, C.** (2009). Inhibition of the shade avoidance response by formation of

- non-DNA binding bHLH heterodimers. *EMBO J.* **28**: 3893-3902.
- Hsieh, H.L., Okamoto, H., Wang, M., Ang, L.H., Matsui, M., Goodman, H., and Deng, X.W.** (2000). *FIN219*, an auxin-regulated gene, defines a link between phytochrome A and the downstream regulator COP1 in light control of Arabidopsis development. *Genes Dev.* **14**: 1958-1970.
- Hudson, M., Ringli, C., Boylan, M.T., and Quail, P.H.** (1999). The *FAR1* locus encodes a novel nuclear protein specific to phytochrome A signaling. *Genes Dev.* **13**: 2017-2027.
- Hudson, M.E., Lisch, D.R., and Quail, P.H.** (2003). The *FHY3* and *FAR1* genes encode transposase-related proteins involved in regulation of gene expression by the phytochrome A-signaling pathway. *Plant J.* **34**: 453-471.
- Hughes, J., and Lamparter, T.** (1999). Prokaryotes and phytochrome. The connection to chromophores and signaling. *Plant Physiol.* **121**: 1059-1068.
- Hughes, J., Lamparter, T., Mittmann, F., Hartmann, E., Gartner, W., Wilde, A., and Borner, T.** (1997). A prokaryotic phytochrome. *Nature* **386**: 663.
- Huq, E., Al-Sady, B., Hudson, M., Kim, C., Apel, K., and Quail, P.H.** (2004). Phytochrome-Interacting Factor 1 is a critical bHLH regulator of chlorophyll biosynthesis. *Science* **305**: 1937-1941.
- Huq, E., and Quail, P.H.** (2002). PIF4, a phytochrome-interacting bHLH factor, functions as a negative regulator of phytochrome B signaling in Arabidopsis. *EMBO J.* **21**: 2441-2450.
- Huq, E., Tepperman, J.M., and Quail, P.H.** (2000). GIGANTEA is a nuclear protein involved in phytochrome signaling in Arabidopsis. *Proc. Natl. Acad. Sci. USA* **97**: 9789-9794.
- Imaizumi, T., Tran, H.G., Swartz, T.E., Briggs, W.R., and Kay, S.A.** (2003). FKF1 is essential for photoperiodic-specific light signalling in Arabidopsis. *Nature* **426**: 302-306.
- Jabben, M., Shanklin, J., and Vierstra, R.D.** (1989a). Ubiquitin-phytochrome conjugates. Pool dynamics during *in vivo* phytochrome degradation. *J. Biol. Chem.* **264**: 4998-5005.
- Jabben, M., Shanklin, J., and Vierstra, R.D.** (1989b). Red light-induced accumulation of ubiquitin-phytochrome conjugates in both monocots and dicots. *Plant Physiol.* **90**: 380-384.
- Jang, I.C., Henriques, R., Seo, H.S., Nagatani, A., and Chua, N.H.** (2010). Arabidopsis PHYTOCHROME INTERACTING FACTOR proteins promote phytochrome B polyubiquitination by COP1 E3 ligase in the nucleus. *Plant Cell* **22**: 2370-2383.
- Jang, I.C., Yang, J.Y., Seo, H.S., and Chua, N.H.** (2005). HFR1 is targeted by COP1 E3 ligase for post-translational proteolysis during phytochrome A signaling. *Genes Dev.* **19**: 593-602.
- Jang, I.C., Yang, S.W., Yang, J.Y., and Chua, N.H.** (2007). Independent and interdependent functions of LAF1 and HFR1 in phytochrome A signaling. *Genes Dev.* **21**: 2100-2111.
- Jang, S., Marchal, V., Panigrahi, K.C., Wenkel, S., Soppe, W., Deng, X.W., Valverde, F., and Coupland, G.** (2008). Arabidopsis COP1 shapes the temporal pattern of CO accumulation conferring a photoperiodic flowering response. *EMBO J.* **27**: 1277-1288.
- Janoudi, A.K., Gordon, W.R., Wagner, D., Quail, P., and Poff, K.L.** (1997). Multiple phytochromes are involved in red-light-induced enhancement of first-positive phototropism in *Arabidopsis thaliana*. *Plant Physiol.* **113**: 975-979.
- Jiang, Z., Swem, L.R., Rushing, B.G., Devanathan, S., Tollin, G., and Bauer, C.E.** (1999). Bacterial photoreceptor with similarity to photoactive yellow protein and plant phytochromes. *Science* **285**: 406-409.
- Jiao, Y., Lau, O.S., and Deng, X.W.** (2007). Light-regulated transcriptional networks in higher plants. *Nat. Rev. Genet.* **8**: 217-230.
- Johnson, E., Bradley, M., Harberd, N.P., and Whitelam, G.C.** (1994). Photoresponses of light-grown *phyA* mutants of Arabidopsis (phytochrome A is required for the perception of daylength extensions). *Plant Physiol.* **105**: 141-149.
- Jones, A.M., Ecker, J.R., and Chen, J.G.** (2003). A reevaluation of the role of the heterotrimeric G protein in coupling light responses in Arabidopsis. *Plant Physiol.* **131**: 1623-1627.
- Jordan, E.T., Cherry, J.R., Walker, J.M., and Vierstra, R.D.** (1996). The amino-terminus of phytochrome A contains two distinct functional domains. *Plant J.* **9**: 243-257.
- Jordan, E.T., Marita, J.M., Clough, R.C., and Vierstra, R.D.** (1997). Characterization of regions within the N-terminal 6-kilodalton domain of phytochrome A that modulate its biological activity. *Plant Physiol.* **115**: 693-704.
- Kami, C., Lorrain, S., Hornitschek, P., and Fankhauser, C.** (2010). Light-regulated plant growth and development. *Curr. Top. Dev. Biol.* **91**: 29-66.
- Kehoe, D.M., and Grossman, A.R.** (1996). Similarity of a chromatic adaptation sensor to phytochrome and ethylene receptors. *Science* **273**: 1409-1412.
- Kevei, E., Schafer, E., and Nagy, F.** (2007). Light-regulated nucleocytoplasmic partitioning of phytochromes. *J. Exp. Bot.* **58**: 3113-3124.
- Khanna, R., Huq, E., Kikis, E.A., Al-Sady, B., Lanzatella, C., and Quail, P.H.** (2004). A novel molecular recognition motif necessary for targeting photoactivated phytochrome signaling to specific basic helix-loop-helix transcription factors. *Plant Cell* **16**: 3033-3044.
- Khanna, R., Kikis, E.A., and Quail, P.H.** (2003). *EARLY FLOWERING 4* functions in phytochrome B-regulated seedling de-etiolation. *Plant Physiol.* **133**: 1530-1538.
- Khanna, R., Shen, Y., Marion, C.M., Tsuchisaka, A., Theologis, A., Schafer, E., and Quail, P.H.** (2007). The basic helix-loop-helix transcription factor PIF5 acts on ethylene biosynthesis and phytochrome signaling by distinct mechanisms. *Plant Cell* **19**: 3915-3929.
- Kim, D.H., Kang, J.G., Yang, S.S., Chung, K.S., Song, P.S., and Park, C.M.** (2002). A phytochrome-associated protein phosphatase 2A modulates light signals in flowering time control in Arabidopsis. *Plant Cell* **14**: 3043-3056.
- Kim, I.S., Bai, U., and Song, P.S.** (1989). A purified 124-kDa oat phytochrome does not possess a protein kinase activity. *Photochem. Photobiol.* **49**: 319-323.
- Kim, J.I., Park, J.E., Zarate, X., and Song, P.S.** (2005). Phytochrome phosphorylation in plant light signaling. *Photochem. Photobiol. Sci.* **4**: 681-687.
- Kim, J.I., Shen, Y., Han, Y.J., Park, J.E., Kirchenbauer, D., Soh, M.S., Nagy, F., Schafer, E., and Song, P.S.** (2004). Phytochrome phosphorylation modulates light signaling by influencing the protein-protein interaction. *Plant Cell* **16**: 2629-2640.
- Kim, L., Kircher, S., Toth, R., Adam, E., Schafer, E., and Nagy, F.** (2000). Light-induced nuclear import of phytochrome-A::GFP fusion proteins is differentially regulated in transgenic tobacco and Arabidopsis. *Plant J.* **22**: 125-133.
- Kim, W.Y., Fujiwara, S., Suh, S.S., Kim, J., Kim, Y., Han, L., David, K., Putterill, J., Nam, H.G., and Somers, D.E.** (2007). ZEITLUPE is a circadian photoreceptor stabilized by GIGANTEA in blue light. *Nature* **449**: 356-360.
- Kircher, S., Gil, P., Kozma-Bognar, L., Fejes, E., Speth, V., Husselstein-Muller, T., Bauer, D., Adam, E., Schafer, E., and Nagy, F.** (2002). Nucleocytoplasmic partitioning of the plant photoreceptors phytochrome A, B, C, D, and E is regulated differentially by light and exhibits a diurnal rhythm. *Plant Cell* **14**: 1541-1555.
- Kircher, S., Kozma-Bognar, L., Kim, L., Adam, E., Harter, K., Schafer, E., and Nagy, F.** (1999). Light quality-dependent nuclear import of the plant photoreceptors phytochrome A and B. *Plant Cell* **11**: 1445-1456.

- Kohchi, T., Mukougawa, K., Frankenberg, N., Masuda, M., Yokota, A., and Lagarias, J.C. (2001). The Arabidopsis *HY2* gene encodes phytylchromobilin synthase, a ferredoxin-dependent biliverdin reductase. *Plant Cell* **13**: 425-436.
- Koornneef, M., Rolff, E., and Spruit, C.J.P. (1980). Genetic control of light-inhibited hypocotyl elongation in *Arabidopsis thaliana* (L.) Heynh. *Z. Pflanzenphysiol.* **100**: 147-160.
- Kunkel, T., Neuhaus, G., Batschauer, A., Chua, N.H., and Schafer, E. (1996). Functional analysis of yeast-derived phytochrome A and B phycocyanobilin adducts. *Plant J.* **10**: 625-636.
- Lagarias, J.C., and Rapoport, H. (1980). Chromopeptides from phytochrome. The structure and linkage of the Pr form of the phytochrome chromophore. *J. Am. Chem. Soc.* **102**: 4821-4828.
- Lamparter, T., Mittmann, F., Gartner, W., Borner, T., Hartmann, E., and Hughes, J. (1997). Characterization of recombinant phytochrome from the cyanobacterium *Synechocystis*. *Proc. Natl. Acad. Sci. USA* **94**: 11792-11797.
- Lapko, V.N., Jiang, X.Y., Smith, D.L., and Song, P.S. (1997). Posttranslational modification of oat phytochrome A: phosphorylation of a specific serine in a multiple serine cluster. *Biochemistry* **36**: 10595-10599.
- Lapko, V.N., Jiang, X.Y., Smith, D.L., and Song, P.S. (1999). Mass spectrometric characterization of oat phytochrome A: isoforms and post-translational modifications. *Protein Sci.* **8**: 1032-1044.
- Lau, O.S., and Deng, X.W. (2010). Plant hormone signaling lightens up: integrators of light and hormones. *Curr. Opin. Plant Biol.* **13**: 571-577.
- Laubinger, S., and Hoecker, U. (2003). The SPA1-like proteins SPA3 and SPA4 repress photomorphogenesis in the light. *Plant J.* **35**: 373-385.
- Laubinger, S., Fittinghoff, K., and Hoecker, U. (2004). The SPA quartet: a family of WD-repeat proteins with a central role in suppression of photomorphogenesis in Arabidopsis. *Plant Cell* **16**: 2293-2306.
- Lee, J., He, K., Stolc, V., Lee, H., Figueroa, P., Gao, Y., Tongprasit, W., Zhao, H., Lee, I., and Deng, X.W. (2007). Analysis of transcription factor HY5 genomic binding sites revealed its hierarchical role in light regulation of development. *Plant Cell* **19**: 731-749.
- Leivar, P., Monte, E., Al-Sady, B., Carle, C., Storer, A., Alonso, J.M., Ecker, J.R., and Quail, P.H. (2008). The Arabidopsis phytochrome-interacting factor PIF7, together with PIF3 and PIF4, regulates responses to prolonged red light by modulating phyB levels. *Plant Cell* **20**: 337-352.
- Leivar, P., Monte, E., Oka, Y., Liu, T., Carle, C., Castillon, A., Huq, E., and Quail, P.H. (2008). Multiple phytochrome-interacting bHLH transcription factors repress premature seedling photomorphogenesis in darkness. *Curr. Biol.* **18**: 1815-1823.
- Leivar, P., Tepperman, J.M., Monte, E., Calderon, R.H., Liu, T.L., and Quail, P.H. (2009). Definition of early transcriptional circuitry involved in light-induced reversal of PIF-imposed repression of photomorphogenesis in young Arabidopsis seedlings. *Plant Cell* **21**: 3535-3553.
- Leivar, P., and Quail, P.H. (2011). PIFs: pivotal components in a cellular signaling hub. *Trends Plant Sci.* **16**: 19-28.
- Li, G., Siddiqui, H., Teng, Y., Lin, R., Wan, X.Y., Li, J., Lau, O.S., Ouyang, X., Dai, M., Wan, J., Devlin, P.F., Deng, X.W., and Wang, H. (2011). Coordinated transcriptional regulation underlying the circadian clock in Arabidopsis. *Nat. Cell. Biol.* **13**: 616-622.
- Li, J., Li, G., Gao, S., Martinez, C., He, G., Zhou, Z., Huang, X., Lee, J.H., Zhang, H., Shen, Y., Wang, H., and Deng, X.W. (2010). Arabidopsis transcription factor ELONGATED HYPOCOTYL 5 plays a role in the feedback regulation of phytochrome A signaling. *Plant Cell* **22**: 3634-3649.
- Lin, R., Ding, L., Casola, C., Ripoll, D.R., Feschotte, C., and Wang, H. (2007). Transposase-derived transcription factors regulate light signaling in Arabidopsis. *Science* **318**: 1302-1305.
- Lin, R., Teng, Y., Park, H.J., Ding, L., Black, C., Fang, P., and Wang, H. (2008). Discrete and essential roles of the multiple domains of Arabidopsis *FHY3* in mediating phytochrome A signal transduction. *Plant Physiol.* **148**: 981-992.
- Liu, L.J., Zhang, Y.C., Li, Q.H., Sang, Y., Mao, J., Lian, H.L., Wang, L., and Yang, H.Q. (2008). COP1-mediated ubiquitination of CONSTANS is implicated in cryptochrome regulation of flowering in Arabidopsis. *Plant Cell* **20**: 292-306.
- Liu, X.L., Covington, M.F., Fankhauser, C., Chory, J., and Wagner, D.R. (2001). *ELF3* encodes a circadian clock-regulated nuclear protein that functions in an Arabidopsis *PHYB* signal transduction pathway. *Plant Cell* **13**: 1293-1304.
- Lorrain, S., Allen, T., Duek, P.D., Whitelam, G.C., and Fankhauser, C. (2008). Phytochrome-mediated inhibition of shade avoidance involves degradation of growth-promoting bHLH transcription factors. *Plant J.* **53**: 312-323.
- Ma, L., Gao, Y., Qu, L., Chen, Z., Li, J., Zhao, H., and Deng, X.W. (2002). Genomic evidence for COP1 as a repressor of light-regulated gene expression and development in Arabidopsis. *Plant Cell* **14**: 2383-2398.
- Mancinelli, A.L. (1994). The physiology of phytochrome action. In *Photomorphogenesis in plants*, R.E. Kendrick, and G.H.M. Kronenberg, eds. (Dordrecht, The Netherlands: Kluwer Academic Publishers), pp. 211-269.
- Marrocco, K., Zhou, Y., Bury, E., Dieterle, M., Funk, M., Genschik, P., Krenz, M., Stolpe, T., and Kretsch, T. (2006). Functional analysis of EID1, an F-box protein involved in phytochrome A-dependent light signal transduction. *Plant J.* **45**: 423-438.
- Martinez-Garcia, J.F., Huq, E., and Quail, P.H. (2000). Direct targeting of light signals to a promoter element-bound transcription factor. *Science* **288**: 859-863.
- Martin-Tryon, E.L., Kreps, J.A., and Harmer, S.L. (2007). GIGANTEA acts in blue light signaling and has biochemically separable roles in circadian clock and flowering time regulation. *Plant Physiol.* **143**: 473-486.
- Mas, P., Devlin, P.F., Panda, S., and Kay, S.A. (2000). Functional interaction of phytochrome B and cryptochrome 2. *Nature* **408**: 207-211.
- Mas, P., Kim, W.Y., Somers, D.E., and Kay, S.A. (2003). Targeted degradation of TOC1 by ZTL modulates circadian function in *Arabidopsis thaliana*. *Nature* **426**: 567-570.
- Mateos, J.L., Luppi, J.P., Ogorodnikova, O.B., Sineshchekov, V.A., Yarovskiy, M.J., Braslavsky, S.E., Gartner, W., and Casal, J.J. (2006). Functional and biochemical analysis of the N-terminal domain of phytochrome A. *J. Biol. Chem.* **281**: 34421-34429.
- Mathews, S., and Sharrock, R.A. (1997). Phytochrome gene diversity. *Plant Cell Environ.* **20**: 666-671.
- Matsushita, T., Mochizuki, N., and Nagatani, A. (2003). Dimers of the N-terminal domain of phytochrome B are functional in the nucleus. *Nature* **424**: 571-574.
- McClung, C.R. (2008). Comes a time. *Curr. Opin. Plant Biol.* **11**: 514-520.
- McClung, C.R., Salomé, P.A., and Michael, T.P. (2002). The Arabidopsis circadian system. *The Arabidopsis Book* **1**: e0044. doi:0010.1199/tab.0044.
- McMichael, R.W., Jr., and Lagarias, J.C. (1990). Phosphopeptide mapping of Avena phytochrome phosphorylated by protein kinases *in vitro*. *Biochemistry* **29**: 3872-3878.
- McNellis, T.W., and Deng, X.W. (1995). Light control of seedling morphogenetic pattern. *Plant Cell* **7**: 1749-1761.
- McWatters, H.G., Bastow, R.M., Hall, A., and Millar, A.J. (2000). The *ELF3* *zeitnehmer* regulates light signalling to the circadian clock. *Nature* **408**: 716-720.
- Mizoguchi, T., Wright, L., Fujiwara, S., Cremer, F., Lee, K., Onouchi, H., Mouradov, A., Fowler, S., Kamada, H., Putterill, J., and Coup-

- land, G. (2005). Distinct roles of *GIGANTEA* in promoting flowering and regulating circadian rhythms in Arabidopsis. *Plant Cell* **17**: 2255-2270.
- Monte, E., Alonso, J.M., Ecker, J.R., Zhang, Y., Li, X., Young, J., Austin-Phillips, S., and Quail, P.H. (2003). Isolation and characterization of *phyC* mutants in Arabidopsis reveals complex crosstalk between phytochrome signaling pathways. *Plant Cell* **15**: 1962-1980.
- Muller, R., Fernandez, A.P., Hiltbrunner, A., Schafer, E., and Kretsch, T. (2009). The histidine kinase-related domain of Arabidopsis phytochrome A controls the spectral sensitivity and the subcellular distribution of the photoreceptor. *Plant Physiol.* **150**: 1297-1309.
- Muramoto, T., Kohchi, T., Yokota, A., Hwang, I., and Goodman, H.M. (1999). The Arabidopsis photomorphogenic mutant *hy1* is deficient in phytochrome chromophore biosynthesis as a result of a mutation in a plastid heme oxygenase. *Plant Cell* **11**: 335-348.
- Mustilli, A.C., and Bowler, C. (1997). Tuning in to the signals controlling photoregulated gene expression in plants. *EMBO J.* **16**: 5801-5806.
- Nagatani, A. (2004). Light-regulated nuclear localization of phytochromes. *Curr. Opin. Plant Biol.* **7**: 708-711.
- Nagatani, A. (2010). Phytochrome: structural basis for its functions. *Curr. Opin. Plant Biol.* **13**: 565-570.
- Nagatani, A., Chory, J., and Furuya, M. (1991). Phytochrome B is not detectable in the *hy3* mutant of Arabidopsis, which is deficient in responding to end-of-day far-red light treatments. *Plant Cell Physiol.* **32**: 1119-1122.
- Nagatani, A., Reed, J.W., and Chory, J. (1993). Isolation and initial characterization of Arabidopsis mutants that are deficient in phytochrome A. *Plant Physiol.* **102**: 269-277.
- Nagy, F., and Schafer, E. (2002). Phytochromes control photomorphogenesis by differentially regulated, interacting signaling pathways in higher plants. *Annu. Rev. Plant Biol.* **53**: 329-355.
- Neff, M.M., and Chory, J. (1998). Genetic interactions between phytochrome A, phytochrome B, and cryptochrome 1 during Arabidopsis development. *Plant Physiol.* **118**: 27-35.
- Neff, M.M., Fankhauser, C., and Chory, J. (2000). Light: an indicator of time and place. *Genes Dev.* **14**: 257-271.
- Neff, M.M., and Van Volkenburgh, E. (1994). Light-stimulated cotyledon expansion in Arabidopsis seedlings (the role of phytochrome B). *Plant Physiol.* **104**: 1027-1032.
- Nelson, D.C., Lasswell, J., Rogg, L.E., Cohen, M.A., and Bartel, B. (2000). *FKF1*, a clock-controlled gene that regulates the transition to flowering in Arabidopsis. *Cell* **101**: 331-340.
- Nemhauser, J., and Chory, J. (2002). Photomorphogenesis. The Arabidopsis Book **1**: e0054 doi: 0010.1199/tab.0054.
- Neuhaus, G., Bowler, C., Hiratsuka, K., Yamagata, H., and Chua, N.H. (1997). Phytochrome-regulated repression of gene expression requires calcium and cGMP. *EMBO J.* **16**: 2554-2564.
- Ni, M., Tepperman, J.M., and Quail, P.H. (1998). PIF3, a phytochrome-interacting factor necessary for normal photoinduced signal transduction, is a novel basic helix-loop-helix protein. *Cell* **95**: 657-667.
- Nozue, K., Covington, M.F., Duek, P.D., Lorrain, S., Fankhauser, C., Harmer, S.L., and Maloof, J.N. (2007). Rhythmic growth explained by coincidence between internal and external cues. *Nature* **448**: 358-361.
- Oh, E., Kim, J., Park, E., Kim, J.I., Kang, C., and Choi, G. (2004). PIL5, a phytochrome-interacting basic helix-loop-helix protein, is a key negative regulator of seed germination in *Arabidopsis thaliana*. *Plant Cell* **16**: 3045-3058.
- Oh, E., Yamaguchi, S., Kamiya, Y., Bae, G., Chung, W.I., and Choi, G. (2006). Light activates the degradation of PIL5 protein to promote seed germination through gibberellin in Arabidopsis. *Plant J.* **47**: 124-139.
- Oka, Y., Matsushita, T., Mochizuki, N., Suzuki, T., Tokutomi, S., and Nagatani, A. (2004). Functional analysis of a 450-amino acid N-terminal fragment of phytochrome B in Arabidopsis. *Plant Cell* **16**: 2104-2116.
- Okamoto, H., Matsui, M., and Deng, X.W. (2001). Overexpression of the heterotrimeric G-protein alpha-subunit enhances phytochrome-mediated inhibition of hypocotyl elongation in Arabidopsis. *Plant Cell* **13**: 1639-1652.
- Oliverio, K.A., Crepy, M., Martin-Tryon, E.L., Milich, R., Harmer, S.L., Putterill, J., Yanovsky, M.J., and Casal, J.J. (2007). *GIGANTEA* regulates phytochrome A-mediated photomorphogenesis independently of its role in the circadian clock. *Plant Physiol.* **144**: 495-502.
- Osterlund, M.T., and Deng, X.W. (1998). Multiple photoreceptors mediate the light-induced reduction of GUS-COP1 from Arabidopsis hypocotyl nuclei. *Plant J.* **16**: 201-208.
- Osterlund, M.T., Hardtke, C.S., Wei, N., and Deng, X.W. (2000). Targeted destabilization of HY5 during light-regulated development of Arabidopsis. *Nature* **405**: 462-466.
- Ouyang, X., Li, J., Li, G., Li, B., Chen, B., Shen, H., Huang, X., Mo, X., Wan, X., Lin, R., Li, S., Wang, H., and Deng, X.W. (2011). Genome-wide binding site analysis of FAR-RED ELONGATED HYPOCOTYL 3 reveals its novel function in Arabidopsis development. *Plant Cell* **23**: 2514-2535.
- Oyama, T., Shimura, Y., and Okada, K. (1997). The Arabidopsis *HY5* gene encodes a bZIP protein that regulates stimulus-induced development of root and hypocotyl. *Genes Dev.* **11**: 2983-2995.
- Park, D.H., Somers, D.E., Kim, Y.S., Choy, Y.H., Lim, H.K., Soh, M.S., Kim, H.J., Kay, S.A., and Nam, H.G. (1999). Control of circadian rhythms and photoperiodic flowering by the Arabidopsis *GIGANTEA* gene. *Science* **285**: 1579-1582.
- Park, E., Kim, J., Lee, Y., Shin, J., Oh, E., Chung, W.I., Liu, J.R., and Choi, G. (2004). Degradation of phytochrome interacting factor 3 in phytochrome-mediated light signaling. *Plant Cell Physiol.* **45**: 968-975.
- Parks, B.M., and Quail, P.H. (1991). Phytochrome-deficient *hy1* and *hy2* long hypocotyl mutants of Arabidopsis are defective in phytochrome chromophore biosynthesis. *Plant Cell* **3**: 1177-1186.
- Parks, B.M., and Quail, P.H. (1993). *hy8*, a new class of Arabidopsis long hypocotyl mutants deficient in functional phytochrome A. *Plant Cell* **5**: 39-48.
- Parks, B.M., Quail, P.H., and Hangarter, R.P. (1996). Phytochrome A regulates red-light induction of phototropic enhancement in Arabidopsis. *Plant Physiol.* **110**: 155-162.
- Parks, B.M., and Spalding, E.P. (1999). Sequential and coordinated action of phytochromes A and B during Arabidopsis stem growth revealed by kinetic analysis. *Proc. Natl. Acad. Sci. USA* **96**: 14142-14146.
- Pedmale, U.V., Celaya, R.B., and Liscum, E. (2010). Phototropism: Mechanism and outcomes. The Arabidopsis Book **8**: e0125 doi:0110.1199/tab.0125.
- Phee, B.K., Kim, J.I., Shin, D.H., Yoo, J., Park, K.J., Han, Y.J., Kwon, Y.K., Cho, M.H., Jeon, J.S., Bhoo, S.H., and Hahn, T.R. (2008). A novel protein phosphatase indirectly regulates phytochrome-interacting factor 3 via phytochrome. *Biochem J.* **415**: 247-255.
- Putterill, J., Robson, F., Lee, K., Simon, R., and Coupland, G. (1995). The *CONSTANS* gene of Arabidopsis promotes flowering and encodes a protein showing similarities to zinc finger transcription factors. *Cell* **80**: 847-857.
- Quail, P.H. (1997a). An emerging molecular map of the phytochromes. *Plant Cell Environ.* **20**: 657-665.
- Quail, P.H. (1997b). The phytochromes: a biochemical mechanism of signaling in sight? *BioEssays* **19**: 571-579.
- Quail, P.H. (2002). Phytochrome photosensory signalling networks. *Nat. Rev. Mol. Cell Biol.* **3**: 85-93.
- Quail, P.H. (2011). Phytochromes. *Curr. Biol.* **20**: R504-507.
- Quail, P.H., Boylan, M.T., Parks, B.M., Short, T.W., Xu, Y., and Wagner,

- D. (1995). Phytochromes: photosensory perception and signal transduction. *Science* **268**: 675-680.
- Reed, J.W., Nagatani, A., Elich, T.D., Fagan, M., and Chory, J. (1994). Phytochrome A and phytochrome B have overlapping but distinct functions in Arabidopsis development. *Plant Physiol.* **104**: 1139-1149.
- Reed, J.W., Nagpal, P., Bastow, R.M., Solomon, K.S., Dowson-Day, M.J., Elumalai, R.P., and Millar, A.J. (2000). Independent action of ELF3 and phyB to control hypocotyl elongation and flowering time. *Plant Physiol.* **122**: 1149-1160.
- Remberg, A., Ruddat, A., Braslavsky, S.E., Gartner, W., and Schaffner, K. (1998). Chromophore incorporation, Pr to Pfr kinetics, and Pfr thermal reversion of recombinant N-terminal fragments of phytochrome A and B chromoproteins. *Biochemistry* **37**: 9983-9990.
- Rizzini, L., Favory, J.J., Cloix, C., Faggionato, D., O'Hara, A., Kaiserli, E., Baumeister, R., Schafer, E., Nagy, F., Jenkins, G.I., and Ulm, R. (2011). Perception of UV-B by the Arabidopsis UVR8 protein. *Science* **332**: 103-106.
- Robson, F., Costa, M.M., Hepworth, S.R., Vizir, I., Pineiro, M., Reeves, P.H., Putterill, J., and Coupland, G. (2001). Functional importance of conserved domains in the flowering-time gene *CONSTANS* demonstrated by analysis of mutant alleles and transgenic plants. *Plant J.* **28**: 619-631.
- Rockwell, N.C., Su, Y.S., and Lagarias, J.C. (2006). Phytochrome structure and signaling mechanisms. *Annu. Rev. Plant Biol.* **57**: 837-858.
- Rosler, J., Klein, I., and Zeidler, M. (2007). Arabidopsis *fh1/fhy1* double mutant reveals a distinct cytoplasmic action of phytochrome A. *Proc. Natl. Acad. Sci. USA* **104**: 10737-10742.
- Roux, S.J. (1994). Signal transduction in phytochrome responses. In *Photomorphogenesis in plants*, R.E. Kendrick, and G.H.M. Kronenberg, eds. (Dordrecht, The Netherlands: Kluwer Academic Publishers), pp. 187-209.
- Ryu, J.S., Kim, J.I., Kunkel, T., Kim, B.C., Cho, D.S., Hong, S.H., Kim, S.H., Fernandez, A.P., Kim, Y., Alonso, J.M., Ecker, J.R., Nagy, F., Lim, P.O., Song, P.S., Schafer, E., and Nam, H.G. (2005). Phytochrome-specific type 5 phosphatase controls light signal flux by enhancing phytochrome stability and affinity for a signal transducer. *Cell* **120**: 395-406.
- Saijo, Y., Sullivan, J.A., Wang, H., Yang, J., Shen, Y., Rubio, V., Ma, L., Hoecker, U., and Deng, X.W. (2003). The COP1-SPA1 interaction defines a critical step in phytochrome A-mediated regulation of HY5 activity. *Genes Dev.* **17**: 2642-2647.
- Saijo, Y., Zhu, D., Li, J., Rubio, V., Zhou, Z., Shen, Y., Hoecker, U., Wang, H., and Deng, X.W. (2008). Arabidopsis COP1/SPA1 complex and FHY1/FHY3 associate with distinct phosphorylated forms of phytochrome A in balancing light signaling. *Mol. Cell* **31**: 607-613.
- Sakamoto, K., and Nagatani, A. (1996). Nuclear localization activity of phytochrome B. *Plant J.* **10**: 859-868.
- Salter, M.G., Franklin, K.A., and Whitelam, G.C. (2003). Gating of the rapid shade-avoidance response by the circadian clock in plants. *Nature* **426**: 680-683.
- Schultz, T.F., Kiyosue, T., Yanovsky, M., Wada, M., and Kay, S.A. (2001). A role for LKP2 in the circadian clock of Arabidopsis. *Plant Cell* **13**: 2659-2670.
- Searle, I., and Coupland, G. (2004). Induction of flowering by seasonal changes in photoperiod. *EMBO J.* **23**: 1217-1222.
- Seo, H.S., Watanabe, E., Tokutomi, S., Nagatani, A., and Chua, N.H. (2004). Photoreceptor ubiquitination by COP1 E3 ligase desensitizes phytochrome A signaling. *Genes Dev.* **18**: 617-622.
- Seo, H.S., Yang, J.Y., Ishikawa, M., Bolle, C., Ballesteros, M.L., and Chua, N.H. (2003). LAF1 ubiquitination by COP1 controls photomorphogenesis and is stimulated by SPA1. *Nature* **423**: 995-999.
- Serino, G., and Deng, X.W. (2003). The COP9 signalosome: regulating plant development through the control of proteolysis. *Annu. Rev. Plant Biol.* **54**: 165-182.
- Sessa, G., Carabelli, M., Sassi, M., Ciolfi, A., Possenti, M., Mitterpergher, F., Becker, J., Morelli, G., and Ruberti, I. (2005). A dynamic balance between gene activation and repression regulates the shade avoidance response in Arabidopsis. *Genes Dev.* **19**: 2811-2815.
- Shanklin, J., Jabben, M., and Vierstra, R.D. (1987). Red light-induced formation of ubiquitin-phytochrome conjugates: identification of possible intermediates of phytochrome degradation. *Proc. Natl. Acad. Sci. USA* **84**: 359-363.
- Sharrock, R.A., and Clack, T. (2002). Patterns of expression and normalized levels of the five Arabidopsis phytochromes. *Plant Physiol.* **130**: 442-456.
- Sharrock, R.A., and Quail, P.H. (1989). Novel phytochrome sequences in *Arabidopsis thaliana*: structure, evolution, and differential expression of a plant regulatory photoreceptor family. *Genes Dev.* **3**: 1745-1757.
- Shen, H., Moon, J., and Huq, E. (2005a). PIF1 is regulated by light-mediated degradation through the ubiquitin-26S proteasome pathway to optimize photomorphogenesis of seedlings in Arabidopsis. *Plant J.* **44**: 1023-1035.
- Shen, H., Zhu, L., Castillon, A., Majee, M., Downie, B., and Huq, E. (2008). Light-induced phosphorylation and degradation of the negative regulator PHYTOCHROME-INTERACTING FACTOR 1 from Arabidopsis depend upon its direct physical interactions with photoactivated phytochromes. *Plant Cell* **20**: 1586-1602.
- Shen, Y., Feng, S., Ma, L., Lin, R., Qu, L.J., Chen, Z., Wang, H., and Deng, X.W. (2005b). Arabidopsis FHY1 protein stability is regulated by light via phytochrome A and 26S proteasome. *Plant Physiol.* **139**: 1234-1243.
- Shen, Y., Khanna, R., Carle, C.M., and Quail, P.H. (2007). Phytochrome induces rapid PIF5 phosphorylation and degradation in response to red-light activation. *Plant Physiol.* **145**: 1043-1051.
- Shen, Y., Zhou, Z., Feng, S., Li, J., Tan-Wilson, A., Qu, L.J., Wang, H., and Deng, X.W. (2009). Phytochrome A mediates rapid red light-induced phosphorylation of Arabidopsis FAR-RED ELONGATED HYPOCOTYL1 in a low fluence response. *Plant Cell* **21**: 494-506.
- Shin, J., Kim, K., Kang, H., Zulfugarov, I.S., Bae, G., Lee, C.H., Lee, D., and Choi, G. (2009). Phytochromes promote seedling light responses by inhibiting four negatively-acting phytochrome-interacting factors. *Proc. Natl. Acad. Sci. USA* **106**: 7660-7665.
- Shinomura, T., Nagatani, A., Hanzawa, H., Kubota, M., Watanabe, M., and Furuya, M. (1996). Action spectra for phytochrome A- and B-specific photoinduction of seed germination in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* **93**: 8129-8133.
- Smith, H., and Whitelam, G.C. (1997). The shade avoidance syndrome: multiple responses mediated by multiple phytochromes. *Plant Cell Environ.* **20**: 840-844.
- Smith, H., Xu, Y., and Quail, P.H. (1997). Antagonistic but complementary actions of phytochromes A and B allow seedling de-etiolation. *Plant Physiol.* **114**: 637-641.
- Soh, M.S., Kim, Y.M., Han, S.J., and Song, P.S. (2000). REP1, a basic helix-loop-helix protein, is required for a branch pathway of phytochrome A signaling in Arabidopsis. *Plant Cell* **12**: 2061-2074.
- Somers, D.E., Devlin, P.F., and Kay, S.A. (1998). Phytochromes and cryptochromes in the entrainment of the Arabidopsis circadian clock. *Science* **282**: 1488-1490.
- Somers, D.E., Kim, W.Y., and Geng, R. (2004). The F-box protein ZEITLUPE confers dosage-dependent control on the circadian clock, photomorphogenesis, and flowering time. *Plant Cell* **16**: 769-782.
- Somers, D.E., Schultz, T.F., Milnamow, M., and Kay, S.A. (2000). *ZEIT-*

- LUPE* encodes a novel clock-associated PAS protein from Arabidopsis. *Cell* **101**: 319-329.
- Somers, D.E., Sharrock, R.A., Tepperman, J.M., and Quail, P.H.** (1991). The *hy3* long hypocotyl mutant of Arabidopsis is deficient in phytochrome B. *Plant Cell* **3**: 1263-1274.
- Speth, V., Otto, V., and Schafer, E.** (1986). Intracellular localization of phytochrome in oat coleoptiles by electron microscopy. *Planta* **168**: 299-304.
- Stockhaus, J., Nagatani, A., Halfter, U., Kay, S., Furuya, M., and Chua, N.H.** (1992). Serine-to-alanine substitutions at the amino-terminal region of phytochrome A result in an increase in biological activity. *Genes Dev.* **6**: 2364-2372.
- Strasser, B., Sanchez-Lamas, M., Yanovsky, M.J., Casal, J.J., and Cerdan, P.D.** (2010). *Arabidopsis thaliana* life without phytochromes. *Proc. Natl. Acad. Sci. USA* **107**: 4776-4781.
- Suarez-Lopez, P., Wheatley, K., Robson, F., Onouchi, H., Valverde, F., and Coupland, G.** (2001). CONSTANS mediates between the circadian clock and the control of flowering in Arabidopsis. *Nature* **410**: 1116-1120.
- Sullivan, J.A., Shirasu, K., and Deng, X.W.** (2003). The diverse roles of ubiquitin and the 26S proteasome in the life of plants. *Nat. Rev. Genet.* **4**: 948-958.
- Takagi, S., Kong, S.G., Mineyuki, Y., and Furuya, M.** (2003). Regulation of actin-dependent cytoplasmic motility by type II phytochrome occurs within seconds in *Vallisneria spiralis* epidermal cells. *Plant Cell* **15**: 331-345.
- Tepperman, J.M., Hudson, M.E., Khanna, R., Zhu, T., Chang, S.H., Wang, X., and Quail, P.H.** (2004). Expression profiling of *phyB* mutant demonstrates substantial contribution of other phytochromes to red-light-regulated gene expression during seedling de-etiolation. *Plant J.* **38**: 725-739.
- Tepperman, J.M., Hwang, Y.S., and Quail, P.H.** (2006). *phyA* dominates in transduction of red-light signals to rapidly responding genes at the initiation of Arabidopsis seedling de-etiolation. *Plant J.* **48**: 728-742.
- Tepperman, J.M., Zhu, T., Chang, H.S., Wang, X., and Quail, P.H.** (2001). Multiple transcription-factor genes are early targets of phytochrome A signaling. *Proc. Natl. Acad. Sci. USA* **98**: 9437-9442.
- Terry, M.J.** (1997). Phytochrome chromophore-deficient mutants. *Plant Cell Environ.* **20**: 740-745.
- Thines, B., and Harmon, F.G.** (2010). Ambient temperature response establishes ELF3 as a required component of the core Arabidopsis circadian clock. *Proc. Natl. Acad. Sci. USA* **107**: 3257-3262.
- Toledo-Ortiz, G., Huq, E., and Quail, P.H.** (2003). The Arabidopsis basic/helix-loop-helix transcription factor family. *Plant Cell* **15**: 1749-1770.
- Tseng, T.S., Salome, P.A., McClung, C.R., and Olszewski, N.E.** (2004). SPINDLY and GIGANTEA interact and act in *Arabidopsis thaliana* pathways involved in light responses, flowering, and rhythms in cotyledon movements. *Plant Cell* **16**: 1550-1563.
- Ulijasz, A.T., Cornilescu, G., Cornilescu, C.C., Zhang, J., Rivera, M., Markley, J.L., and Vierstra, R.D.** (2010). Structural basis for the photoconversion of a phytochrome to the activated Pfr form. *Nature* **463**: 250-254.
- Ullah, H., Chen, J.G., Young, J.C., Im, K.H., Sussman, M.R., and Jones, A.M.** (2001). Modulation of cell proliferation by heterotrimeric G protein in Arabidopsis. *Science* **292**: 2066-2069.
- Ulm, R., Baumann, A., Oravecz, A., Mate, Z., Adam, E., Oakeley, E.J., Schafer, E., and Nagy, F.** (2004). Genome-wide analysis of gene expression reveals function of the bZIP transcription factor HY5 in the UV-B response of Arabidopsis. *Proc. Natl. Acad. Sci. USA* **101**: 1397-1402.
- Valverde, F., Mouradov, A., Soppe, W., Ravenscroft, D., Samach, A., and Coupland, G.** (2004). Photoreceptor regulation of CONSTANS protein in photoperiodic flowering. *Science* **303**: 1003-1006.
- Vierstra, R.D., and Davis, S.J.** (2000). Bacteriophytochromes: new tools for understanding phytochrome signal transduction. *Semin. Cell Dev. Biol.* **11**: 511-521.
- Vince-Prue, D.** (1994). The duration of light and photoperiodic responses. In *Photomorphogenesis in plants*, R.E. Kendrick, and G.H.M. Kronenberg, eds. (Dordrecht, The Netherlands: Kluwer Academic Publishers), pp. 447-490.
- Wagner, D., Fairchild, C.D., Kuhn, R.M., and Quail, P.H.** (1996). Chromophore-bearing NH₂-terminal domains of phytochromes A and B determine their photosensory specificity and differential light lability. *Proc. Natl. Acad. Sci. USA* **93**: 4011-4015.
- Wang, H., and Deng, X.W.** (2002). Arabidopsis *FHY3* defines a key phytochrome A signaling component directly interacting with its homologous partner *FAR1*. *EMBO J.* **21**: 1339-1349.
- Wang, H., and Deng, X.W.** (2003). Dissecting the phytochrome A-dependent signaling network in higher plants. *Trends Plant Sci.* **8**: 172-178.
- Wang, H., Ma, L., Habashi, J., Li, J., Zhao, H., and Deng, X.W.** (2002). Analysis of far-red light-regulated genome expression profiles of phytochrome A pathway mutants in Arabidopsis. *Plant J.* **32**: 723-733.
- Wenden, B., Kozma-Bognar, L., Edwards, K.D., Hall, A.J., Locke, J.C., and Millar, A.J.** (2011). Light inputs shape the Arabidopsis circadian system. *Plant J.* **66**: 480-491.
- Whitelam, G.C., Johnson, E., Peng, J., Carol, P., Anderson, M.L., Cowl, J.S., and Harberd, N.P.** (1993). Phytochrome A null mutants of Arabidopsis display a wild-type phenotype in white light. *Plant Cell* **5**: 757-768.
- Wong, Y.S., Cheng, H.C., Walsh, D.A., and Lagarias, J.C.** (1986). Phosphorylation of Avena phytochrome *in vitro* as a probe of light-induced conformational changes. *J. Biol. Chem.* **261**: 12089-12097.
- Wong, Y.S., McMichael, R.W., and Lagarias, J.C.** (1989). Properties of a polycation-stimulated protein kinase associated with purified Avena phytochrome. *Plant Physiol.* **91**: 709-718.
- Wu, S.H., and Lagarias, J.C.** (2000). Defining the bilin lyase domain: lessons from the extended phytochrome superfamily. *Biochemistry* **39**: 13487-13495.
- Yanagawa, Y., Sullivan, J.A., Komatsu, S., Gusmaroli, G., Suzuki, G., Yin, J., Ishibashi, T., Saijo, Y., Rubio, V., Kimura, S., Wang, J., and Deng, X.W.** (2004). Arabidopsis COP10 forms a complex with DDB1 and DET1 *in vivo* and enhances the activity of ubiquitin conjugating enzymes. *Genes Dev.* **18**: 2172-2181.
- Yang, J., Lin, R., Sullivan, J., Hoecker, U., Liu, B., Xu, L., Deng, X.W., and Wang, H.** (2005). Light regulates COP1-mediated degradation of HFR1, a transcription factor essential for light signaling in Arabidopsis. *Plant Cell* **17**: 804-821.
- Yang, K.Y., Kim, Y.M., Lee, S., Song, P.S., and Soh, M.S.** (2003). Overexpression of a mutant basic helix-loop-helix protein HFR1, HFR1-deltaN105, activates a branch pathway of light signaling in Arabidopsis. *Plant Physiol.* **133**: 1630-1642.
- Yang, S.W., Jang, I.C., Henriques, R., and Chua, N.H.** (2009). FAR-RED ELONGATED HYPOCOTYL1 and FHY1-LIKE associate with the Arabidopsis transcription factors LAF1 and HFR1 to transmit phytochrome A signals for inhibition of hypocotyl elongation. *Plant Cell* **21**: 1341-1359.
- Yanovsky, M.J., Izaguirre, M., Wagmaister, J.A., Gatz, C., Jackson, S.D., Thomas, B., and Casal, J.J.** (2000). Phytochrome A resets the circadian clock and delays tuber formation under long days in potato. *Plant J.* **23**: 223-232.
- Yanovsky, M.J., and Kay, S.A.** (2002). Molecular basis of seasonal time measurement in Arabidopsis. *Nature* **419**: 308-312.
- Yanovsky, M.J., Mazzella, M.A., Whitelam, G.C., and Casal, J.J.** (2001). Resetting of the circadian clock by phytochromes and cryptochromes in

- Arabidopsis. *J. Biol. Rhythms* **16**: 523-530.
- Yeh, K.C., and Lagarias, J.C.** (1998). Eukaryotic phytochromes: light-regulated serine/threonine protein kinases with histidine kinase ancestry. *Proc. Natl. Acad. Sci. USA* **95**: 13976-13981.
- Yeh, K.C., Wu, S.H., Murphy, J.T., and Lagarias, J.C.** (1997). A cyanobacterial phytochrome two-component light sensory system. *Science* **277**: 1505-1508.
- Yi, C., and Deng, X.W.** (2005). COP1 - from plant photomorphogenesis to mammalian tumorigenesis. *Trends Cell Biol.* **15**: 618-625.
- Yu, J.W., Rubio, V., Lee, N.Y., Bai, S., Lee, S.Y., Kim, S.S., Liu, L., Zhang, Y., Irigoyen, M.L., Sullivan, J.A., Lee, I., Xie, Q., Paek, N.C., and Deng, X.W.** (2008). COP1 and ELF3 control circadian function and photoperiodic flowering by regulating GI stability. *Mol. Cell* **32**: 617-630.
- Yu, X., Liu, H., Klejnot, J., and Lin, C.** (2010). The cryptochrome blue light receptors. *The Arabidopsis Book* **8**: e0135. doi:0110.1199/tab.0135.
- Zagotta, M.T., Hicks, K.A., Jacobs, C.I., Young, J.C., Hangarter, R.P., and Meeks-Wagner, D.R.** (1996). The Arabidopsis *ELF3* gene regulates vegetative photomorphogenesis and the photoperiodic induction of flowering. *Plant J.* **10**: 691-702.
- Zeidler, M., Zhou, Q., Sarda, X., Yau, C.P., and Chua, N.H.** (2004). The nuclear localization signal and the C-terminal region of FHY1 are required for transmission of phytochrome A signals. *Plant J.* **40**: 355-365.
- Zhang, H., He, H., Wang, X., Yang, X., Li, L., and Deng, X.W.** (2011). Genome-wide mapping of the HY5-mediated gene networks in Arabidopsis that involve both transcriptional and post-transcriptional regulation. *Plant J.* **65**: 346-358.
- Zhou, Q., Hare, P.D., Yang, S.W., Zeidler, M., Huang, L.F., and Chua, N.H.** (2005). FHL is required for full phytochrome A signaling and shares overlapping functions with FHY1. *Plant J.* **43**: 356-370.
- Zhou, Y.C., Dieterle, M., Buche, C., and Kretsch, T.** (2002). The negatively acting factors EID1 and SPA1 have distinct functions in phytochrome A-specific light signaling. *Plant Physiol.* **128**: 1098-1108.
- Zhu, D., Maier, A., Lee, J.H., Laubinger, S., Saijo, Y., Wang, H., Qu, L.J., Hoecker, U., and Deng, X.W.** (2008). Biochemical characterization of Arabidopsis complexes containing CONSTITUTIVELY PHOTOMORPHOGENIC1 and SUPPRESSOR OF PHYA proteins in light control of plant development. *Plant Cell* **20**: 2307-2323.