

Handbook of the Birds of the World, vol. 13: Penduline-Tits to Shrikes

Author: Bowie, Rauri C. K.

Source: The Auk, 126(4) : 936-937

Published By: American Ornithological Society

URL: <https://doi.org/10.1525/auk.2009.41009>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Book Reviews

EDITED BY R. TODD ENGSTROM

The following critiques express the opinions of the individual evaluators regarding the strengths, weaknesses, and value of the books they review. As such, the appraisals are subjective assessments and do not necessarily reflect the opinions of the editors or any official policy of the American Ornithologists' Union.

The Auk 126(4):936–937, 2009
© The American Ornithologists' Union, 2009.
Printed in USA.

Handbook of the Birds of the World, vol. 13: Penduline-Tits to Shrikes.—Josep del Hoyo, Andrew Elliott, and David Christie, Eds. 2008. Lynx Edicions, Barcelona, Spain. 879 pp., 60 color plates, 536 photographs, 611 distribution maps. ISBN: 84-96553-45-0 and 978-84-96553-45-3. Hardcover, \$300.—With the publication of volume 13 (of 16, final volume out in 2011), some 174 of approximately 200 families of birds have been covered in the internationally acclaimed series *Handbook of the Birds of the World* (HBW). Volume 13 follows the classic layout of HBW by starting with a highly informative essay, which in this volume reviews the fascinating subject of bird migration (written by Ian Newton). The main body of the book covers an impressive 595 species in the following families: Remizidae (penduline-tits: 13 species), Aegithalidae (long-tailed tits: 13), Sittidae (nuthatches: 27), Tichodromidae (Wallcreeper: 1), Certhiidae (treecreepers: 10), Rhabdornithidae (*Rhabdornis*: 3), Nectariniidae (sunbirds: 132), Melanocharitidae (berrypeckers and longbills: 10), Paramythiidae (painted berrypeckers: 2), Dicaeidae (flowerpeckers: 44), Pardalotidae (pardalotes: 4), Zosteropidae (white-eyes: 98), Promeropidae (sugarbirds: 2), Meliphagidae (honeyeaters: 175), Oriolidae (orioles: 30), and Laniidae (shrikes: 31).

These 16 families, presented in a modern taxonomic framework (e.g., Barker et al. 2004, Jönsson and Fjeldså 2006, Johansson et al. 2008), span the songbird tree of life. From an ecological perspective, the sequence of families rather pleasingly encompasses nearly all the major creeper and nectarivore groups of traditional classifications, ending with the shrikes (shrike-like birds being the focus of the forthcoming vol. 14). I was particularly pleased to see that the editors had resisted attempts of some classification systems to lump traditionally recognized families such as the flowerpeckers, sugarbirds, and sunbirds together in the Nectariniidae or the painted berrypeckers with the berrypeckers and longbills in the Melanocharitidae. At the same time, the validity of some of these families is unlikely to be retained in the future. For example, there is now convincing evidence that species in the Rhabdornithidae belong in the Sturnidae (starlings; Cibois and Cracraft 2004, Zuccon et al. 2006) and that the white-eyes are part of the babbler radiation (Timaliidae; Gelang et al. 2009). One cannot fault the editors for this, for the taxonomic sequence was set several years

ago and many of these results are recently published. But this does highlight a point that readers who do not closely follow avian systematics (e.g., general bird enthusiasts and comparative biologists, among others) will find somewhat frustrating. With the advent of cheap and efficient methods for collecting DNA sequence data, together with several new methods of analysis, avian systematics is at present undergoing a revolution, with new discoveries being made almost monthly. One of the most recent and dramatic (duly noted in the honeyeater chapter) pertains to the discovery that the two genera *Moho* and *Chaetoptila*, the “Hawaiian honeyeaters,” are more closely related to waxwings, Neotropical silky flycatchers, and Palmchats (*Dulus dominicus*) than to Australasian honeyeaters (Fleischer et al. 2008). Although such large changes in taxonomy are likely to remain limited, others such as generic-level revisions are not. A simple case can be illustrated with the Nectariniidae, where use of molecular sequence data has revealed the lack of monophyly for nearly all the presently recognized 16 genera (Bowie 2003). My point is not to criticize the book—far from it: many of the authors do an excellent job of highlighting problematic taxonomy and aspects that are likely to change in the future—but rather to sound a cautionary note. The present taxonomy, at this fascinating time of massive flux within the classification of birds, is unlikely to remain stable even in the near future.

Aside from a detailed review of the systematics of each group, the beginning of each family chapter includes detailed sections on the ecology, behavior, and status of the taxa in each family, which make for fascinating reading. These texts are excellently written by experts on individual families, and in several cases they include data from studies that were published after this book (e.g., the discussion of Hawaiian honeyeaters cited above). This initial introductory overview of each family is followed by detailed species accounts together with excellent color maps of distribution ranges that are easy to interpret.

The book contains 60 color plates illustrating all species and including “all significant sexual and subspecific differences.” As a whole this goal is met, although for large families such as the sunbirds and honeyeaters not all “significant” taxa are illustrated; however, the quality of the plates is truly exceptional. A test, which I often use for my favorite group of birds (the sunbirds)

The Auk, Vol. 126, Number 4, pages 936–940. ISSN 0004-8038, electronic ISSN 1938-4254. © 2009 by The American Ornithologists' Union. All rights reserved. Please direct all requests for permission to photocopy or reproduce article content through the University of California Press's Rights and Permissions website, <http://www.ucpressjournals.com/reprintInfo.asp>. DOI: 10.1525/auk.2009.41009

when consulting a new field guide or monographic compilation, is to see how well the artist has illustrated the Rufous-winged Sunbird (*Cinnyris rufipennis*), a species first described in 1983. Despite an illustration that appeared with the original species description, its depiction in subsequent texts has varied from fantasy (e.g., Stevenson and Fanshawe 2002) to average (Cheke and Mann 2001) to exceptionally accurate (this book).

I own all the previous volumes of the HBW, and I admit that the one aspect of these books that gives me the most pleasure is the out-of-this-world photography. Volume 13 is no different, including 536 stunning photographs. This lavish illustration, together with an informative text (over 6,000 references, including a specific section devoted to bibliographic details of every genus, species, and subspecies accepted by HBW) makes this one of those very rare books that can pass as both an outstanding coffee-table book and an exceptional reference. To those who have already purchased the previous volumes: you will not be disappointed, for this one lives up to the very highest standards set by the HBW production team. To those who have not yet invested in a volume of the HBW series and have an interest in Old World birds or some of North America's more enigmatic species and families—for example, Verdin (Remizidae: *Auriparus flaviceps*), Bushtit (Aegithalidae: *Psaltiriparus minimus*), Brown Creeper (Certhiidae: *Certhia americana*), and the nuthatches (Sittidae: *Sitta* spp.)—buy this book! It will prove a fundamentally important reference and a fascinating read, and provide a great sense of pleasure when you come back to it time and time again.—RAURI C. K. BOWIE, *Museum of Vertebrate Zoology and Department of Integrative Biology, 3101 Valley Life Science Building, University of California, Berkeley, California 94720, USA. E-mail: bowie@berkeley.edu*

LITERATURE CITED

- BARKER, F. K., A. CIBOIS, P. SCHIKLER, J. FEINSTEIN, AND J. CRACRAFT. 2004. Phylogeny and diversification of the largest avian radiation. *Proceedings of the National Academy of Sciences USA* 101:11040–11045.
- BOWIE, R. C. K. 2003. Birds, molecules and evolutionary patterns among Africa's islands in the sky. Ph.D. dissertation, University of Cape Town, Cape Town, South Africa.
- CHEKE, R. A., AND C. F. MANN. 2001. Sunbirds: A Guide to Sunbirds, Flowerpeckers, Spiderhunters and Sugarbirds of the World. Christopher Helm, London.
- CIBOIS, A., AND J. CRACRAFT. 2004. Assessing the passerine "tapestry": Phylogenetic relationships of the Muscipoidea inferred from nuclear DNA sequences. *Molecular Phylogenetics and Evolution* 32:264–273.
- FLEISCHER, R. C., H. F. JAMES, AND S. L. OLSON. 2008. Convergent evolution of Hawaiian and Australo-Pacific honeyeaters from distant songbird ancestors. *Current Biology* 18:1927–1931.
- GELANG, M., A. CIBOIS, E. PASQUET, U. OLSSON, P. ALSTRÖM, AND P. G. P. ERICSON. 2009. Phylogeny of babblers (Aves: Passeriformes): Major lineages, family limits and classification. *Zoologica Scripta* 38:225–236.
- JOHANSSON, U. S., J. FJELDSÅ, AND R. C. K. BOWIE. 2008. Phylogenetic relationships within Passerida (Aves: Passeriformes): A review and a new molecular phylogeny based on three nuclear intron markers. *Molecular Phylogenetics and Evolution* 48:858–876.
- JÖNSSON, K. A., AND J. FJELDSÅ. 2006. A phylogenetic supertree of oscine passerine birds (Aves: Passeri). *Zoologica Scripta* 35:149–186.
- STEVENSON, T., AND J. FANSHAW. 2002. Field Guide to the Birds of East Africa: Kenya, Tanzania, Uganda, Rwanda, and Burundi. T & A D Poyser, London.
- ZUCCON, D., A. CIBOIS, E. PASQUET, AND P. G. P. ERICSON. 2006. Nuclear and mitochondrial sequence data reveal the major lineages of starlings, mynas and related taxa. *Molecular Phylogenetics and Evolution* 41:333–344.

The Auk 126(4):937–939, 2009

© The American Ornithologists' Union, 2009.

Printed in USA.

At-Sea Distribution and Abundance of Seabirds off Southern California: A 20-Year Comparison.

—John W. Mason, Gerard J. McChesney, William R. McIver, Harry R. Carter, John Y. Takekawa, Richard T. Golightly, Joshua T. Ackerman, Dennis L. Orthmeyer, William M. Perry, Julie L. Yee, Mark O. Pierson, and Michael D. McCrary. 2007. *Studies in Avian Biology* no. 33. Cooper Ornithological Society, Camarillo, California. ix + 101 pp., 45 figures, 17 tables. ISBN: 9780943610726. Paper, \$15.—Thirty years ago, the U.S. Department of the Interior, Minerals Management Service (MMS), organized the Outer Continental Shelf Environmental Assessment Program to gather the information necessary to meet requirements under the National Environmental Policy Act toward development of minerals extraction on the U.S. continental shelf. A portion of the assessment involved marine birds, especially their abundance and distribution in waters off California, which were deemed by industry to have significant hydrocarbon potential (in production since the 1960s). A few years of intense aerial censusing on a grid extending from the U.S.–Mexico border to the California–Oregon border resulted in an impressive data set, available electronically and summarized in *Studies in Avian Biology*, no. 11 (Briggs et al. 1987). As of 1983, therefore, the status and distribution of marine birds in California and waters within 50 miles (80 km) of the coast were well quantified. These data were used to approve lease sales and, subsequently, to assess effects of oil spills in California waters. Since completion of that project, several studies have detailed significant changes in the populations of a number of avian species in these waters (e.g., Ainley et al. 1995a, Veit et al. 1997, Ainley and Divoky 2001, Oedekoven et al. 2001, Hyrenbach and Veit 2003). Those findings apparently led to the study under review here—a reassessment, again using aerial censusing, of seabird populations in the portion of California's continental-shelf waters still available, or currently being used, for minerals extraction.

Mason et al.'s study area included waters from the coast to the outer edge of the continental shelf (3,000 m isobath, 80–100 km from the mainland beach) from Morro Bay to San Diego. Flights were flown on 102 days during three months: January, May, and September (May 1999–January 2002). Using the data obtained and reanalysis of Briggs et al.'s (1987) data, the present volume compares spatial distributions and populations of seabirds in