

## Erratum

Author: Ericson, Per G. P.

Source: The Auk, 118(2) : 563

Published By: American Ornithological Society

URL: [https://doi.org/10.1642/0004-8038\(2001\)118\[0563:E\]2.0.CO;2](https://doi.org/10.1642/0004-8038(2001)118[0563:E]2.0.CO;2)

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Erratum

PER G. P. ERICSON<sup>1</sup>

Department of Vertebrate Zoology, Swedish Museum of Natural History, P.O. Box 50007,  
SE-10405 Stockholm, Sweden

Ericson et al. (2000) reported on the taxonomic distributions of two insertions in the nuclear *c-myc* gene observed in a survey of passerine species. On the basis of an expanded set of sequences, it has become evident that the most parsimonious placement for the first insertion is between positions 789 and 790 in the published chicken (*Gallus*) sequence (Watson et al. 1983). The positional change of this insertion does not affect the phylogenetic interpretation of its occurrence. We thank Michael J. Braun and John Harshman for alerting us to the possibility that the insertion should be located at a different position than was first suggested.

---

<sup>1</sup> E-mail: per.ericson@nrm.edu

## LITERATURE CITED

- ERICSON, P. G. P., U. S. JOHANSSON, AND T. J. PARSONS. 2000. Major divisions of oscines revealed by insertions in the nuclear gene *c-myc*: A novel gene in avian phylogenetics. *Auk* 117:1077–1086.
- WATSON, D. K., E. P. REDDY, P. H. DUESBERG, AND T. S. PAPAS. 1983. Nucleotide sequence analysis of the chicken *c-myc* gene reveals homologous and unique coding regions by comparison with the transforming gene of avian myelocytomatosis viros MC29, delta *gag-myc*. *Proceedings of the National Academy of Sciences USA* 80:2146–2150.