

# A new species of antbird (Passeriformes: Thamnophilidae) from the Cordillera Azul, San Martín, Peru

Authors: Moncrieff, Andre E., Johnson, Oscar, Lane, Daniel F., Beck, Josh R., Angulo, Fernando, et al.

Source: The Auk, 135(1): 114-126

Published By: American Ornithological Society

URL: https://doi.org/10.1642/AUK-17-97.1

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



# RESEARCH ARTICLE

# A new species of antbird (Passeriformes: Thamnophilidae) from the Cordillera Azul, San Martín, Peru

# Andre E. Moncrieff,<sup>1#</sup>\* Oscar Johnson,<sup>1#</sup> Daniel F. Lane,<sup>1</sup> Josh R. Beck,<sup>2</sup> Fernando Angulo,<sup>3</sup> and Jesse Fagan<sup>4</sup>

<sup>1</sup> Louisiana State University Museum of Natural Science, Baton Rouge, Louisiana, USA

<sup>2</sup> El Cajon, California, USA

<sup>3</sup> Centro de Ornitología y Biodiversidad (CORBIDI), Lima, Peru

<sup>4</sup> Lima, Peru

<sup>#</sup> These authors contributed equally to the paper.

\* Corresponding author: amoncr2@lsu.edu

Submitted June 1, 2017; Accepted October 5, 2017; Published January 28, 2018

# ABSTRACT

We describe a distinctive new species of antbird (Passeriformes: Thamnophilidae) from humid montane forest (1,340– 1,670 m above sea level) of the Cordillera Azul, San Martín Region, Peru. Plumage, voice, and molecular evidence distinguish this species from its sister taxon *Myrmoderus ferrugineus* (Ferruginous-backed Antbird), which is found in lowland Amazonian rainforests of the Guiana Shield and Madeira-Tapajós interfluvium. The new species is presently known only from one ridge in the Cordillera Azul, and therefore we recommend further fieldwork to better estimate its distribution and population size.

Keywords: Myrmeciza, Myrmoderus, new species, outlying ridges, taxonomy

# Una nueva especie de hormiguero (Passeriformes: Thamnophilidae) de la Cordillera Azul, San Martín, Perú

#### RESUMEN

Describimos una nueva y distintiva especie de hormiguero (Passeriformes: Thamnophilidae) de los bosques montanos húmedos (1,340–1,670 metros sobre el nivel del mar) de la Cordillera Azul, región de San Martín, Perú. El plumaje, la voz, y la evidencia molecular distinguen a esta especie de su taxón hermano *Myrmoderus ferrugineus* (Hormiguero Lomirrufo), el cual se encuentra en los bosques tropicales amazónicos de llanura del Escudo Guyanés y el interfluvio Madeira-Tapajós. A esta nueva especie se le conoce en la actualidad únicamente de una cresta de la Cordillera Azul, y por lo tanto, recomendamos mayor investigación de campo para poder estimar mejor su distribución y el tamaño de su población.

Palabras clave: cadenas montañosas aisladas, Myrmeciza, Myrmoderus, nueva especie, taxonomía

#### **INTRODUCTION**

Antbirds in the genus *Myrmoderus* Ridgway, 1909 (*sensu* Remsen et al. 2017) are found in the Atlantic rainforests of eastern Brazil (White-bibbed Antbird [*M. loricatus* Lichtenstein, 1823], Squamate Antbird [*M. squamosus* Pelzeln, 1868], and Scalloped Antbird [*M. ruficauda* Wied, 1831]) and the Amazonian rainforests of the Guiana Shield and Madeira-Tapajós interfluvium (Ferruginous-backed Antbird [*M. ferrugineus* Müller, 1776]). With no representative known in western Amazonian South America, it is with some surprise that we report the discovery of a new species in this genus from the foothills of the Peruvian Andes. The new antbird inhabits forests surrounding the small coffee-

growing town of Flor de Café (7.398°S, 76.299°W), previously called "Plataforma," at 1,600 m above sea level [a.s.l.] in the western Cordillera Azul. This town has become a destination for ornithologists and birdwatchers since Todd Mark and Walter Vargas discovered an easily accessible population of Scarlet-banded Barbets (*Capito wallacei* O'Neill et al., 2000) there in May 2011 (T. Mark personal communication). This barbet species was first discovered in 1996 during an ornithological expedition to a remote and uninhabited part of the eastern Cordillera Azul (O'Neill et al. 2000).

In July 2016, J.R.B. visited Flor de Café, and on July 9, while birding in undisturbed, humid montane forest near the town at an elevation of  $\sim$ 1,570 m a.s.l., he heard a

Direct all requests to reproduce journal content to the AOS Publications Office at pubs@americanornithology.org

<sup>© 2018</sup> American Ornithological Society. ISSN 0004-8038, electronic ISSN 1938-4254

rattling alarm call of an antbird. Playback of similar antbird calls did not result in any direct response but may have excited the bird, which sang from the vicinity of the initial alarm call. After making an audio recording of the song on his HTC One M8 cell phone, J.R.B. used playback to obtain views of a ground-walking antbird that did not match any known species. The following day, as J.R.B. was preparing to leave Flor de Café to request help in documenting his find, A.E.M., D.F.L., F.A., and J.F. serendipitously arrived. Joining efforts through July 13, we documented this new species with further voice recordings, photos, video, and the collection of voucher specimens. A.E.M. remained through July 20, further documenting the species around Flor de Café at elevations of 1,390-1,670 m a.s.l. In September and October 2016, O.J., D.F.L., Emil Bautista, and Walter Vargas visited a nearby area, accessed from the town of Selva Andina (7.398°S, 76.230°W), and documented the new species at 1,340 m a.s.l.

Plumage, morphometrics, voice, behavior, and genetic evidence point to a close relationship between *Myrmoderus ferrugineus* and the new species. There are numerous characters unique to these 2 taxa, yet they are also closely allied to the other 3 members of the genus (*M. loricatus, M. squamosus,* and *M. ruficauda*). For this reason, we recognize *Myrmoderus* as the appropriate genus for the new species, which we name

#### *Myrmoderus eowilsoni,* species novum Cordillera Azul Antbird *Hormiguero de la Cordillera Azul* (Spanish)

#### Holotype

Study skin, Centro de Ornitología y Biodiversidad (CORBIDI) AV-12381; tissue, Louisiana State University Museum of Natural Science (LSUMZ) B-93477; adult male; 2.5 km west-northwest of Flor de Café, Cordillera Azul, San Martín Region, Peru (7.390°S, 76.320°W; Figure 1); elevation 1,570 m a.s.l.; collected July 12, 2016; prepared by A.E.M., original catalog number 193; audio-recorded by A.E.M., D.F.L., F.A., and J.F. (Macaulay Library of Natural Sounds [ML], Cornell Laboratory of Ornithology, Ithaca, New York, USA; audio: ML 46435931, ML 224879, ML 224880); GenBank sequences KY849953 (ND2) and KY849952 (ND3).

#### Diagnosis

We assign *Myrmoderus eowilsoni* to the genus *Myrmoderus* on the basis of its combination of black ear coverts, extensively rufous brown plumage, and blackish wing coverts with broad white or buff tips (see Isler et al. 2013). Additional characters shared with *M. ferrugineus* include bluish bare orbital skin, gray feet and tarsi, lack of white interscapular patch, and terrestrial walking behavior. Despite these similarities, it is readily distinguishable from



**FIGURE 1.** Map of the Cordillera Azul showing the type locality, which is also the only known locality, of *Myrmoderus eowilsoni* (yellow star) in relation to other surveyed sites on ridges of the Cordillera Azul (orange circles; O'Neill et al. 2000, Alverson et al. 2001, Merkord et al. 2009, J. Bates personal communication, T. Mark personal communication, LSUMZ field expeditions). The white line represents the boundary of the Cordillera Azul National Park. Areas east of the Huallaga River between 1,300 and 1,700 m a.s.l. are shaded to emphasize potential areas of occurrence of *M. eowilsoni*. Inset map shows the location of the Cordillera Azul in Peru.

*M. ferrugineus* by (1) crown and nape color, (2) supercilium color, (3) extent of bare orbital skin, (4) belly color, (5) breast color (females), and (6) song (Figures 2 and 3). Compared to *M. ferrugineus*, the new species has a colder brown crown and nape, a gray rather than white supercilium, and less extensive bluish bare orbital skin. The belly of *M. eowilsoni* is black (males) or dark brown (females), whereas both sexes of *M. ferrugineus* show substantial white in this area. Additionally, females of *M. eowilsoni* have a rufous breast, whereas females of *M. ferrugineus* have a black breast (Figure 2A). The song differs substantially from *M. ferrugineus* by having fewer notes and a slower pace.



**FIGURE 2.** Ventral (**A**) and lateral (**B**) views of both species of *Myrmoderus*. Left to right: male *M. eowilsoni* (CORBIDI 12381; holotype), female *M. eowilsoni* (CORBIDI 12380; mate of holotype), male *M. ferrugineus* (LSUMZ 178458), and female *M. ferrugineus* (LSUMZ 178456). Scale bars = 5 cm.

#### **Description of Holotype**

Capitalized color names are based on Ridgway (1912), and alphanumeric color codes in parentheses refer to Munsell (no date). Crown and nape Warm Sepia (10YR3/4) with feather tips edged slightly darker, imparting a faintly scaled appearance. Forehead and lores black. Back Brussels Brown (5YR3/6) tending toward Amber Brown (5YR4/8) on upper scapulars and upper mantle. Two mantle feathers have a small amount of white fringing toward the vane bases, suggesting a rudimentary interscapular patch, but otherwise mantle feather bases are variably dark gray to blackish. Rump and uppertail coverts Brussels Brown (5YR3/6). Dorsal side of tail Bone Brown (7.5YR2/2) tending toward Fuscous-Black (slightly darker than 10YR2/2) at tip with outer vanes fringed Brussels Brown (5YR3/6). Ventral side of tail uniformly Fuscous (10YR2/2). Undertail coverts Brussels Brown (5YR3/6). Chin, throat, auriculars, breast, sides, upper flanks, and belly black. Lower belly color blending toward Raw Umber (10YR3/4) on vent and lower flanks. Supercilium Light Gull Gray (N6.75), beginning above anterior edge of eye and extending to and broadening at nape. Lesser coverts black with narrow white tips, becoming entirely white on leading edge of wing. Greater and median secondary coverts black, broadly tipped Warm Buff (7.5YR8/8) to Ochraceous-Buff (7.5YR6/8), forming 2 obvious wing bars. Primary coverts black, narrowly tipped Warm Buff (7.5YR8/8). Underwing coverts blackish, greater coverts tipped white. Primaries and secondaries Fuscous-Black (slightly darker than 10YR2/2) fringed with Brussels Brown (5YR3/6). Tertials Fuscous (10YR2/2) with diffuse black subterminal spot on secondaries 7 and 8 and diffusely tipped with Cinnamon (5YR5/8) on outer vane. Soft-part colors recorded at the

time of collection: irides dark brown, skin of orbit bluegray, feet and tarsi dark gray, mandible blue-gray, and maxilla blackish with blue-gray tomium. Measurements are provided in Table 1; no molt or fat, stomach contained insect parts, left testis  $4.5 \times 2$  mm.

# Paratypes

There are 7 paratypes of M. eowilsoni. Six were collected near Flor de Café between July 12 and 19, 2016 (CORBIDI 12380, 12382, 12383; LSUMZ 190882, 190883, 190884), and one was collected 6 km southeast of Flor de Café on September 30, 2016 (LSUMZ 190885). Data from field labels and morphometric data for all specimens of M. eowilsoni are shown in Table 1. Female plumage differs strikingly from that of males in the following characters: throat and malar white with fine black tips, more extensively black toward chin; breast and sides between Xanthine Orange (5YR6/10) and Amber Brown (5YR4/8); center of belly Dark Gull Gray (N4.25) blending to Raw Umber (10YR3/4) on the surrounding belly and through the flanks and vent. All individuals of both sexes show an area of paler Amber Brown (5YR4/8) on the upper mantle and scapulars, but to varying degrees of intensity and extent. The tips of the lesser coverts are slightly variable in color, with 2 females (LSUMZ 190882 and 190884) showing buffy tips and all other individuals showing exclusively white tips. The wing bars formed by the tips of the median coverts are also slightly variable in color, with some individuals tending toward white. The amount of Raw Umber (10YR3/4) on the belly is variable among females, being largely replaced by gray on one individual (CORBIDI 12380). A male and a female in formative plumage (LSUMZ 190883 and 190884), aged on the basis of molt limits in the rectrices and between the greater secondary and greater primary coverts and by the presence of bursa; like adults in definitive plumage, except that the young female has a nearly pure white throat and chin with greatly reduced scaling.

## Etymology

We name *Myrmoderus eowilsoni* in honor of Dr. Edward Osborne Wilson to recognize his tremendous devotion to conservation and his patronage of the Rainforest Trust, which strives to protect the most imperiled species and habitats in the Neotropics and across the globe. We select the English name to draw attention to the little known but biogeographically important and biodiverse mountain range that contains the type locality of the species.

## REMARKS

#### **Phylogenetic Relationships**

To assess the evolutionary relationships within *Myrmoderus*, we estimated a phylogeny based on sequence data

The Auk: Ornithological Advances 135:114-126, © 2018 American Ornithological Society



FIGURE 3. Sonograms of representative vocalizations of *Myrmoderus* species. Recordings are available at Macaulay Library (ML), Xenocanto.org (XC), or Isler and Whitney (2002), with the exception of Macedo recordings, which are in the collection of the recordist and used here with permission. Specimen number, recordist, date and location recorded, and archival catalog number follow in parentheses, as available. (A) Song of holotype male *M. eowilsoni* (CORBIDI 12381; D.F.L.; July 12, 2017; ML224880). (B) Song of female *M. eowilsoni* (CORBIDI 12380; mate of holotype; D.F.L.; July 12, 2017; ML224880). (C, D) Variation among songs of female *M. eowilsoni* (D.F.L.; July 13, 2017; ML224882). (E) Song of female *M. eowilsoni* (LSUMZ 190882; D.F.L.; July 12, 2017; ML224881). (F) Chatter call of *M. eowilsoni* with insect noise between 3 and 5 kHz (CORBIDI 12380 or 12381; D.F.L.; July 12, 2017; ML224880). (G) Song of male *M. ferrugineus* (B. Whitney; Isler and Whitney 2002). (H) Song of female *M. ferrugineus* (B. Whitney; Isler and Whitney 2002). (H) Song of female *M. ferrugineus* (B. Whitney; Isler and Whitney 2002). (I) Chatter call of *M. ferrugineus* (D.F.L.; March 13, 2011; Suriname, Palumeu; XC74997). (J) Song of male *M. ruficauda* (T. Parker III; Isler and Whitney 2002). (K) Song of male *M. loricatus* (G. Macedo field recording no. 0327; November 30, 2014; Brazil, Rio de Janeiro, Estrada do Contorno). (L) Song of female *M. squamosus* (G. Macedo field recording no. 0469; September 3, 2015; Brazil, São Paulo, Paranapiacaba). (N) Song of female *M. squamosus* (G. Macedo field recording no. 0469; September 3, 2015; Brazil, São Paulo, Paranapiacaba).

from all taxa in the genus, using 2 or 3 individuals of each taxon except *M. r. ruficauda*, for which we had genetic data from only one individual (Appendix Table 3). We obtained sequences of 2 mitochondrial genes, *NADH dehydrogenase subunit 2* (ND2; 1,041 base pairs) and *NADH dehydrogenase subunit 3* (ND3; 351 base pairs) from 11 individuals sequenced previously by Bravo (2012). To augment this dataset, we generated new sequences from 2 individuals of *M. eowilsoni* and 2 individuals of *M. ferrugineus elutus* Todd, 1927. For these 4 individuals, we extracted total DNA from 25 mg of pectoral muscle using a Qiagen DNEasy kit using the manufacturer's protocol and performed polymerase chain reactions (PCR) for ND2

and ND3 following the protocol of Bravo (2012). PCR product was purified and sequenced at Eton Biosciences (Durham, North Carolina, USA) or the LSU Genomics Facility (Baton Rouge, Louisiana, USA). Four outgroup taxa from the Bravo (2012) dataset were used to root the tree: Barred Antshrike (*Thamnophilus doliatus* Linnaeus, 1764), Spotted Antbird (*Hylophylax naevioides* Lafresnaye, 1847), White-bellied Antbird (*Myrmeciza longipes* Swainson, 1825), and White-plumed Antbird (*Pithys albifrons* Linnaeus, 1766). All sequence data are deposited in GenBank (Appendix Table 3).

Each new sequence was cleaned manually and assembled into contigs in Geneious 10.1.3 (Kearse et al. 2012).

TABLE 1	. Measurements f	rom <i>Myrm</i>	oderus eowi.	<i>lsoni</i> specimens	(n = 8).												
Sex	Catalog number <sup>a</sup>	Tissue number	Preparator number	Date collected	Macaulay Library numbers	Elevation where <sup>1</sup> collected (m a.s.l.)	Wing- span V (mm)	Veight I (g)	Tail ength l (mm)	Wing l ength e (mm)	Bill le ength xposed (mm)	Bill ength from nares (mm)	Bill vidth c (mm) (	Bill T depth lo mm)	Farsus ength (mm)	Bursa c (mm)	Skull sssifi- ation (%)
Female	CORBIDI 12380	B-93476	AEM 192	July 12, 2016 4	46435931, 224879, 224880	1,570	217	29.0	57	68.7	17.3	10.8	6.2	5.4	28.8	None	100
Male <sup>b</sup>	CORBIDI 12381	B-93477	AEM 193	July 12, 2016 4	224879, 224879, 224880	1,570	NA	27.2	62	72.8	18.7	11.2	7.8	5.8	27.3	None	100
Female Male <sup>c</sup>	LSUMZ 190882 LSUMZ 190883	B-93478 R-93479	DFL 2683 DFI 2684	July 12, 2016 2 July 12, 2016 2	224881 224881	1,490 1 490	NA 800	27.5 27.3	66 67	68.4 66.0	18.5 17.6	11.0	6.7 6.6	5.9	27.8 27.4	None	100
Female <sup>c</sup>	LSUMZ 190884	B-93482 R-93483	AEM 195	July 18, 2016 2 July 18, 2016 2	46041311 46046081	1,510	225 225 215	27.6 28.7	20 60	68.0 68.0	16.7 17.1	11.0	0.0 6.6 7 7	5.7 5.7	28.0 3 28.0 3	$5 \times 2.5$ None	001
Male Male	CORBIDI 12383 LSUMZ 190885	B-93484 B-72699	AEM 197 OJ 569	September 50, 2016 2 July 19, 2016 2 September 50, 2016	46045061 51252971, 51253111	1,390 1,340	217	24.7 24.7 27.2	61 62	69.1 70.1	16.8 16.8 18.1	10.9 11.1	6.9 6.9	5.5 5.5	26.6 25.8	None None	100
<sup>a</sup> Abbrev <sup>b</sup> Holotyp <sup>c</sup> Formati	iations: CORBIDI = >e. ve plumage.	= Centro d	e Ornitologí	a y Biodiversida	d; LSUMZ = I	-ouisiana S	itate Ur	iversity	Museur	n of Nat	ural Scie	nce.					

TABLE 2. Morphon	netric data from all M	<i>yrmoderus</i> taxa ( <i>n</i> = 10	15). Mean values are g	jiven, followed by 2 st	andard deviations, wi	ith sample sizes in pare	entheses.
Trait	M. eowilsoni	M. f. ferrugineus	M. f. elutus	M. r. ruficauda	M. r. soror	M. loricatus	M. squamosus
Weight (g)	$27.4 \pm 2.6 \ (n = 8)$	$25.5 \pm 3.8 \ (n = 36)$	NA	NA	NA	NA	NA
Wing chord (mm)	$68.6 \pm 4.6 \ (n = 8)$	63.6 ± 3.3 (n = 36)	64.6 ± 3.3 (n = 5)	63.9 ± 1.6 (n = 9)	$66.6 \pm 7.9 \ (n = 5)$	59.8 ± 4.6 (n = 13)	59.1 ± 2.6 (n = 28)
Tail length (mm)	$62.1 \pm 6.4 \ (n = 8)$	54.9 ± 4.9 (n = 36)	$54.6 \pm 3.6 (n = 5)$	51.7 ± 4.1 (n = 9)	$52.4 \pm 1.8 \ (n = 5)$	$65.2 \pm 8.0 \ (n = 13)$	$64.5 \pm 6.3 \ (n = 28)$
Wing:tail ratio	$1.1 \pm 0.1 \ (n = 8)$	$1.2 \pm 0.1 \ (n = 36)$	$1.2 \pm 0.1 \ (n = 5)$	$1.2 \pm 0.1 \ (n = 9)$	$1.3 \pm 0.1 \ (n = 5)$	$0.9 \pm 0.1$ ( $n = 13$ )	$0.9 \pm 0.1 \ (n = 28)$
Tarsus (mm)	27.4 ± 1.8 (n = 8)	$26.2 \pm 2.1 \ (n = 36)$	$26.2 \pm 1.2 \ (n = 5)$	23.8 ± 1.0 ( <i>n</i> = 9)	$23.6 \pm 2.0 \ (n = 5)$	$25.6 \pm 1.8 \ (n = 12)$	26.7 ± 2.2 (n = 28)
Exposed culmen	$17.6 \pm 1.5 \ (n = 8)$	19.1 ± 2.3 ( <i>n</i> = 35)	$18.4 \pm 1.4 \ (n = 5)$	$16.2 \pm 1.6 \ (n = 9)$	$16.3 \pm 1.0 \ (n = 5)$	$14.9 \pm 2.0 \ (n = 13)$	$13.6 \pm 1.2 \ (n = 27)$
(mm) Nares to bill tip	$11.0 \pm 0.4 \ (n = 8)$	$12.1 \pm 1.6 \ (n = 35)$	$11.6 \pm 0.6 \ (n = 5)$	$10.6 \pm 0.9 \ (n = 9)$	$10.3 \pm 1.1 \ (n = 5)$	$9.2 \pm 0.6 \ (n = 13)$	8.5 ± 0.8 (n = 27)
(mm)							
Bill width (mm)	$6.8 \pm 1.0 \ (n = 8)$	$6.6 \pm 1.0 \ (n = 36)$	$6.5 \pm 1.5 \ (n = 5)$	$5.9 \pm 0.4 \ (n = 9)$	$5.6 \pm 0.8 \ (n = 5)$	$5.3 \pm 0.8 \ (n = 12)$	$5.0 \pm 1.1 \ (n = 27)$
Bill depth (mm)	$5.6 \pm 0.5 \ (n = 8)$	5.3 ± 1.1 (n = 36)	$5.0 \pm 0.4 \ (n = 5)$	5.2 ± 1.0 (n = 9)	$5.3 \pm 1.4 \ (n = 5)$	$4.4 \pm 0.7 \ (n = 13)$	$4.2 \pm 0.7 \ (n = 27)$

118 New species of antbird from Peru

The Auk: Ornithological Advances 135:114-126, © 2018 American Ornithological Society

The 2 loci were aligned separately in MUSCLE (Edgar 2004) from within Geneious using default values and then concatenated. Using PartitionFinder 2 (Lanfear et al. 2017) to compare DNA substitution models available in RAxML 8 (Stamatakis 2014) in a corrected Akaike's Information Criterion  $(AIC_c)$  framework gave a 3-partition scheme that pertained to the first, second, and third codon positions and an overall GTR+I+G model of DNA substitution. We used these settings to conduct a maximum-likelihood analysis in RAxML 8.2.10 (Stamatakis 2014) and selected the tree with the best likelihood score from 100 independent tree searches. Statistical support for the topology of this tree was then computed with the automatic majority-rule convergence criterion in RAxML 8.2.10 (Stamatakis 2014), resulting in 800 bootstrap replicates (Felsenstein 1985) and a final majority-rule consensus tree (Figure 4). When selecting among the substitution models available in MrBayes 3.2.6 (Ronquist et al. 2012), PartitionFinder 2 (Lanfear et al. 2017) gave the same partition scheme as for the maximum-likelihood analysis, but with substitution models of GTR+G, HKY+I, and GTR+I for the first, second, and third codon positions, respectively. We used these partitions and substitution models to perform the Bayesian analysis, comprising 4 independent runs of 4 MCMC chains, a chain temperature of 1.75, sampling for 20 million generations, a sampling frequency of 1,000, and a burn-in of 20%. We set a variable substitution-rate prior, an exponential alpha of 0.05, and a flat Dirichlet prior on the exchangeability parameters and base frequencies. Results from our Bayesian analyses were visualized in Tracer 1.6.0 (Rambaut et al. 2014) to assess convergence and stationarity. All runs reached convergence, and the average standard deviation of split frequencies from MrBayes 3.2.6 (Ronguist et al. 2012) was 0.002. Topologies of the maximum-likelihood and Bayesian trees are identical, so we report the maximumlikelihood tree with nodal support values from both methods (Figure 4).

We recovered *M. eowilsoni* as sister to *M. ferrugineus* with high support in all analyses (Figure 4). *Myrmoderus eowilsoni* has an uncorrected pairwise genetic distance of 8.4% in ND2 and 8.7% in ND3 from *M. f. ferrugineus* (found north of the Amazon River), and 9.2% in ND2 and 7.6% in ND3 from *M. f. elutus* (found south of the Amazon River). These distances are high when compared to those between other antbird sister-species, even within the same genus (*M. loricatus* and *M. squamosus*: ND2 uncorrected pairwise distance of 4.5%). *Myrmoderus f. elutus* has an uncorrected pairwise distance of 3.6% in ND2 and 4.3% in ND3 from *M. f. elutus*, although it differs only slightly in plumage (Todd 1927) and voice (average 5.7 notes second<sup>-1</sup> in *M. f. ferrugineus*, n = 20 individuals).



**FIGURE 4.** Phylogeny of the genus *Myrmoderus* estimated from a concatenated ND2–ND3 dataset (1,392 bp) using a maximum-likelihood analysis. The Bayesian analysis produced a tree that was topologically identical. Nodal support values are the likelihood bootstrap values and Bayesian posterior probabilities, respectively.

Isler et al. (2013) presented a taxonomic revision of the highly polyphyletic Myrmeciza antbirds in which they resurrected the genus Myrmoderus (Ridgway 1909) for M. loricatus, M. squamosus, M. ruficauda, and M. ferrugineus. Although Myrmoderus (sensu Isler et al. 2013) formed a clade supported by molecular, morphological, and some behavioral data, taxonomic relationships within this clade were unclear except for the sister relationship of M. loricatus and M. squamosus. With the discovery of M. eowilsoni, the sister relationship of this new species and M. ferrugineus becomes clear. The placement of M. ruficauda remains uncertain in our analyses, because of low support values on the subtending node. It was placed, however, as sister to *M. loricatus–M. squamosus* in all analyses (Figure 4). Adding nuclear loci to the mitochondrial DNA dataset would likely increase confidence in our placement of M. ruficauda (Maddison 1997), but we note that plumage, behavior, and vocalizations are all consistent with the genetically inferred relationships in this study.

#### Taxonomy

Myrmoderus eowilsoni and M. ferrugineus share several distinctive morphological traits, including bluish bare orbital skin, gray feet and tarsi, and white leading edges to the inner bend of the wing (proximal to the wrist), and lack the white interscapular patch shown by other species of Myrmoderus. Conversely, the other members of Myrmoderus, including the type of the genus (M. loricatus), have pink legs and feet and obvious white interscapular patches, the latter of which are regularly used in visual signaling in the forest understory (D. F. Lane personal observation). Myrmoderus eowilsoni and M. *ferrugineus* differ behaviorally from the other *Myrmoderus* by walking rather than hopping as a means of locomotion over the ground (Zimmer and Isler 2003; B. Whitney personal communication). Furthermore, M. eowilsoni and M. ferrugineus exhibit strong sexual dimorphism in song, whereas the 3 other Myrmoderus do not (B. Whitney personal communication; Figure 3). Todd (1927) noted morphometrics, extensively bare orbital skin, plumage coloration, and dark feet to justify the creation of a monotypic genus, Myrmedestes, for Myrmoderus ferrugineus. However, the name Myrmedestes Todd, 1927 fell into disuse when this genus was subsumed into a broadly conceived Myrmeciza Gray, 1841 by multiple authorities (Zimmer 1932, Peters 1951, Meyer de Schauensee 1970), prior to separation of Myrmoderus by Isler et al. (2013). We consider it appropriate, given the sister relationship of Myrmoderus ferrugineus and M. eowilsoni and the presence of numerous morphological and behavioral synapomorphies, to recognize *Myrmedestes* as a subgenus in Myrmoderus to include these 2 species.

#### **Morphological Diagnosability**

We took standard morphological measurements from 105 specimens representing all recognized taxa in the genus *Myrmoderus* (Appendix Table 4). For consistency, O.J. took all measurements. Measurement protocols followed Baldwin et al. (1931; their figure numbers follow the measurements) for tail length (fig. 120), wing chord (fig. 100), tarsus (fig. 136), exposed culmen (fig. 3), bill tip to nares (fig. 8), bill width at base (fig. 13), and bill depth at base (fig. 10; Table 2).

To evaluate the morphological data, we performed a principal component analysis with the PCA function in Scikit-learn (Pedregosa et al. 2011) using all specimens for which we had complete morphological data (n = 101). The first 2 principal components explained 91.4% of the variation in our dataset and clustered the species into 2 non-overlapping groups: one containing *M. loricatus* and *M. squamosus* and the other containing *M. eowilsoni*, both subspecies of *M. ferrugineus* (*M. f. ferrugineus* and *M. f. elutus*), and both subspecies of *M. ruficauda* (*M. r. ruficauda* and *M. r. soror* Pinto, 1940) (Figure 5). Despite



**FIGURE 5.** The first 2 principal components (PC1 and PC2) of 7 morphometric variables from all species of *Myrmoderus*. The loadings for PC1 and PC2 accounted for 73.5% and 17.9% of the variation in the dataset, respectively.

this overlap between *M. ferrugineus* and *M. ruficauda* in morphological measurements, the genetic data suggest that *M. ruficauda* is allied with *M. loricatus* and *M. squamosus* (Figure 4). In addition, *M. ruficauda* has pink legs and feet, a white interscapular patch, and hopping locomotory behavior, which all suggest a closer relationship to *M. loricatus* and *M. squamosus*.

#### Vocalizations

Songs of both male and female M. eowilsoni are distinguished from those of other members of the Thamnophilidae by a combination of a clear whistled quality, few notes, and a simple pattern. Compared to songs of M. ferrugineus, those of M. eowilsoni consist of fewer notes (2-6 vs. 8-14) that are delivered more slowly (average 2 notes second<sup>-1</sup> vs. average 7 notes second<sup>-1</sup>; Figure 3). The male song seems quite stereotyped, but with some minor variation, particularly in pitch. Males sing 4 whistled notes given as 2 couplets (songs of 2 or 3 notes heard rarely), usually with the first note highest in pitch, the second lower in pitch, the third of similar or slightly higher pitch, and the final note lowest in pitch (Figure 3A). Although it is distinctive to the ear, the song of male M. eowilsoni can be likened to the slowed-down terminal 2 couplets of the song of male M. ferrugineus. The song of female M. eowilsoni shows wider variation within and among individuals in pitch and number of notes (3-6)than in the male, and the female song is delivered more slowly and often with a raspier quality (Figure 3B-3E). The call (Figure 3F) is generally a sputtering series of notes similar to that of *M. ferrugineus* (Figure 3I), but it is delivered at a slower pace and occasionally consists of only

single or doubled notes. The songs of *M. ferrugineus* (Figure 3G, 3H), *M. loricatus* (Figure 3K, 3L), and *M. squamosus* (Figure 3M, 3N) are similar to one another in also being composed of couplets that generally descend, but differ from that of *M. eowilsoni* in having more notes. By contrast, the dry, rattled song of *M. ruficauda* (Figure 3J) is immediately distinguishable from that of the other members of the genus. The songs of *M. eowilsoni* and *M. ferrugineus* are sexually dimorphic (i.e. female songs differ consistently from those of males in pattern, note number, and note shape), whereas the songs of the other 3 *Myrmoderus* are not (B. Whitney personal communication; Figure 3).

#### Distribution

Myrmoderus eowilsoni is known currently only from the Flor de Café ridge around the type locality, between 1,340 and 1,670 m a.s.l. (Figure 1). This distribution encompasses just 15 km<sup>2</sup> (minimum convex polygon of known localities), although extrapolating the range to include habitat from 1,300 to 1,700 m a.s.l. on the same ridge increases the distribution to  $\sim$ 78 km<sup>2</sup>. We believe it is very likely that this species is found on additional ridges within the Cordillera Azul, and it should be looked for farther afield, perhaps as far as the Cordillera El Sira. Multiple previous ornithological surveys in the Cordillera Azul since the 1990s did not detect *M. eowilsoni*, yet there are  $\sim$ 2,480 km<sup>2</sup> of habitat between 1,300 and 1,700 m a.s.l. in the Cordillera Azul that may be suitable for this species, most of which are within the Parque Nacional Cordillera Azul (Figure 1). With knowledge of the elevational distribution and habitat requirements of the new species, as well as song playback, we believe that efforts to revisit previously surveyed localities within the Cordillera Azul are warranted in addition to exploring new localities.

#### Habitat, Ecology, and Behavior

Myrmoderus eowilsoni appears to be restricted to tall, montane, evergreen forest with intact understory characterized by small trees (diameter at breast height about 5-10 cm) and ferns, many moss-covered logs, and extensive dead leaf matter on the ground. We did not observe any individuals in second growth, edge, or tree-fall gap habitats, although in a few cases we could hear individuals singing from within 10 m of large plantations of sun-grown coffee. Most sightings occurred on or near level ridge tops, although this may have been biased by a lack of trails in steeper terrain. We did encounter a few individuals up to  $\sim$ 200 m downslope of ridgeline trails. Despite conducting extensive playback (at intervals of <100 m) along trails through forested habitats around Flor de Café from 900 to 1,850 m a.s.l., we did not detect the species below 1,340 or above 1,670 m a.s.l. Both members of the pair defend their territory, and most of our observations were of pairs

responding to playback of the song together, typically walking or occasionally flying toward the source of playback, and singing for as much as an hour after playback ceased. Only one female (CORBIDI 93483) had an enlarged ovary suggestive of breeding activity. Aside from the initial detection by J.R.B., we detected the species only after playback, and it seemed largely silent under natural conditions. We note, however, that we were generally not near known territories at dawn or dusk, when unprovoked singing would be most likely. Alternatively, unsolicited singing may be seasonal; other species of Myrmoderus can become quieter in the dry season, and their period of maximum singing is typically early in the morning between about 0630 and 0700 hours (B. Whitney personal communication). We observed *M. eowilsoni* only within 1 m of the ground, and individuals typically moved by short, low flights and by walking slowly along the ground. Walking involved deliberate steps accompanied by a jerking, dove-like head bob, with the tail held straight out behind. As with many terrestrial walking antbirds, antthrushes, and tapaculos, M. eowilsoni often hopped onto fallen logs or low branches in response to playback. We noted few instances of tail movements, mostly when an agitated individual was perched off the ground and responding to playback, when it would give occasional tail-raising flicks, bringing the tail slightly above horizontal from a position well below horizontal. Brief foraging observations consisted of birds picking through leaf litter to search for arthropod prey, and preserved stomach contents of specimens (CORBIDI 12380, 12381; LSUMZ 190882, 190883, 190884) contain arthropod parts.

When surveying for the species with playback, we located several adjacent territories along a ridgetop trail, which helped us estimate territory lengths of roughly 300–500 m. We revisited territories on consecutive days, and playback did not always elicit a response; this suggests that the pair may have been far enough away as to be out of earshot and that territories are quite large. We tentatively estimate a maximum territory size of 0.25 km<sup>2</sup>, by squaring our estimate of the longest territory length.

#### Conservation

*Myrmoderus eowilsoni* is presently known only from the ridge of Flor de Café, yet little fieldwork has been conducted within the nearby Parque Nacional Cordillera Azul, one of the largest national parks in the country with an area of >13,500 km<sup>2</sup>. Future surveys should target localities within this national park, which contains ~1,940 km<sup>2</sup> of area between 1,300 and 1,700 m a.s.l. that may be appropriate habitat for *M. eowilsoni* (Figure 1). Habitat alteration, primarily clear-cutting, around Flor de Café is extensive and ongoing. Sun coffee plantations, the primary agricultural land use there, do not provide habitat for *M. eowilsoni* or other forest species. Nevertheless, we found

*M. eowilsoni* to be fairly common in forest fragments around the type locality, and these fragments seem to maintain a remarkably intact avifauna, at least for the time being. Human colonists from other parts of Peru are rapidly clearing forest in the area and may soon eliminate much of the remaining forest around Flor de Café. Therefore, confirming the presence of *M. eowilsoni* in the national park is a high priority. Considering the proximity of extensive protected habitat within the national park, we are optimistic that the species will be found to be a common resident in other parts of the mountain range, and we hope that this discovery spurs further conservation and research in this unique region.

We here make a preliminary estimate of the population size of the species. Based on a territory size ranging from 0.066 km<sup>2</sup> (territory size reported for *M. ferrugineus* by Johnson et al. 2011) to 0.25 km<sup>2</sup> (our maximum estimate), 2 birds territory $^{-1}$ , territories in only 45.2% of appropriate habitat (taken from mean estimate for occupancy of M. ferrugineus by Stouffer 2007), and an area of 1,940 km<sup>2</sup> between 1,300 and 1,700 m a.s.l., the predicted population size within the national park is 7,000-27,000 individuals. If we include ridges of the Cordillera Azul found outside the park (including the Flor de Café ridge) but within the latitudinal range of the park, the area between 1,300 and 1,700 m a.s.l. increases to  $\sim$ 2,480 km<sup>2</sup>, and the predicted population size is 9,000-34,000 individuals. These estimates of population size are crude and highly preliminary because of a large number of assumptions, but they nonetheless serve as a starting point for assessment of the actual population size. Flor de Café lies on the leeward side of the Cordillera Azul, and we suspect that the lower elevational limit of *M. eowilsoni* is locally maintained by drier habitats as one descends toward the rain shadow of the Huallaga valley. In more windward and humid regions of the Cordillera Azul, appropriate habitat may extend considerably lower in elevation, perhaps as low as  $\sim$ 900 m. Survey work is urgently needed to determine the full distribution of *M. eowilsoni*, to assess territory size and occupancy, and to improve estimates of population size. Because most of the estimated distribution of *M. eowilsoni* lies within the Parque Nacional Cordillera Azul, we do not expect this species to face extensive habitat loss. Despite the restricted distribution and habitat loss in the immediate vicinity of the type locality, we consider an IUCN Red List status of "Data Deficient" as most appropriate until the distribution and population trends can be further evaluated.

## ACKNOWLEDGMENTS

We thank K. Garrett at the Natural History Museum of Los Angeles County; B. Marks at the Field Museum of Natural History; G. Graves, C. Milensky, and J. Saucier at the National Museum of Natural History, Smithsonian Institution; and M. Robbins at the University of Kansas Biodiversity Institute for access to or loans from their respective collections. CORBIDI provided essential logistical support at various stages of this project. We are indebted to our guides and field assistants, E. Vallejos, E. Bautista Obispo, and W. Vargas, for their tireless work and pleasant company. We thank T. Mark for information on his pioneering surveys in the Cordillera Azul. A. Spencer and G. Macedo kindly shared audio recordings. B. Whitney shared audio recordings and information on the natural history of related species. The Macaulay Library at the Cornell Lab of Ornithology generously provided use of audio recording equipment.

**Funding statement:** Fieldwork was conducted thanks to a grant to the Louisiana State University Museum of Natural Tropical Bird Research Fund. A.E.M. and O.J. are supported by National Science Foundation Graduate Research Fellowships under grant no. DGE-1247192. Any opinion, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation. No funding source had any influence on the content of the submitted or published manuscript, nor required approval of the final manuscript before publication.

**Ethics statement:** Specimen collecting and export permits were provided by the Servicio Nacional Forestal y de Fauna Silvestre (SERFOR) under the following permits: Resolución no. 203-2015 SERFOR-DGGSPFFS, Resolución no. 222-2015 SERFOR-DGGSPFFS, Permiso no. 003104 SERFOR, and Permiso no. 003105 SERFOR. All specimen collection was done under Louisiana State University's Institutional Animal Care and Use Committee protocol no. 15-036.

Author contributions: J.R.B. made the initial discovery. A.E.M., O.J., D.F.L., J.R.B., F.A., and J.F. collected data. A.E.M., O.J., and D.F.L. collected and prepared specimens and analyzed the data. A.E.M. and O.J. wrote the paper.

**Data deposits:** All genetic data are deposited in GenBank (see Appendix Table 3 for accession numbers). Sequences alignments and phylogenetic trees are available at TreeBASE (http://purl.org/phylo/treebase/phylows/study/TB2:S21548).

**Nomenclature:** The nomenclature in this paper has been reviewed by the Working Group on Avian Nomenclature of the International Ornithologists' Union. *Myrmoderus eowilsoni* was registered on ZooBank and it received the following LSID: urn:lsid:zoobank.org:act:A05E5983-E5BC-4189-99C7-AC9E813D1E1C

## LITERATURE CITED

- Alverson, W. S., L. O. Rodríguez, and D. K. Moskovits (2001). Perú: Biabo Cordillera Azul. Rapid Biological Inventories 2. The Field Museum, Chicago, IL, USA.
- Baldwin, S. P., H. C. Oberholser, and L. G. Worley (1931). Measurements of birds. Scientific Publications of the Cleveland Museum of Natural History 2.
- Bravo, G. A. (2012). Phenotypic and niche evolution in the antbirds (Aves, Thamnophilidae). Ph.D. dissertation, Louisiana State University, Baton Rouge, LA, USA.
- Edgar, R. C. (2004). MUSCLE: Multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Research 32:1792–1797.

The Auk: Ornithological Advances 135:114–126, © 2018 American Ornithological Society

- Felsenstein, J. (1985). Confidence limits on phylogenies: An approach using the bootstrap. Evolution 39:783–791.
- Isler, M. L., G. A. Bravo, and R. T. Brumfield (2013). Taxonomic revision of *Myrmeciza* (Aves: Passeriformes: Thamnophilidae) into 12 genera based on phylogenetic, morphological, behavioral, and ecological data. Zootaxa 3717:469–497.
- Isler, P. R., and B. M. Whitney (2002). Songs of the Antbirds [CD]. Macaulay Library of Natural Sounds, Ithaca, NY, USA.
- Johnson, E. I., P. C. Stouffer, and C. F. Vargas (2011). Diversity, biomass, and trophic structure of a central Amazonian rainforest bird community. Revista Brasileira de Ornitologia 19:1–16.
- Kearse, M., R. Moir, A. Wilson, S. Stones-Havas, M. Cheung, S. Sturrock, S. Buxton, A. Cooper, S. Markowitz, C. Duran, T. Thierer, et al. (2012). Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. Bioinformatics 28:1647–1649.
- Lanfear, R., P. B. Frandsen, A. M. Wright, T. Senfeld, and B. Calcott (2017). PartitionFinder 2: New methods for selecting partitioned models of evolution for molecular and morphological phylogenetic analyses. Molecular Biology and Evolution 34: 772–773.
- Maddison, W. P. (1997). Gene trees in species trees. Systematic Biology 46:523–536.
- Merkord, C. L., T. Mark, D. Susanibar, A. Johnson, and C. C. Witt (2009). Avifaunal survey of the Río Chipaota Valley in the Cordillera Azul region, San Martín, Peru. Ornitología Neotropical 20:535–552.
- Meyer de Schauensee, R. (1970). A Guide to the Birds of South America. Livingston, Wynnewood, PA, USA.
- Munsell, A. (No date). Munsell Book of Color, Glossy Collection. Gretag Macbeth, New Windsor, NY, USA.
- O'Neill, J. P., D. F. Lane, A. W. Kratter, A. P. Capparella, and C. F. Joo (2000). A striking new species of barbet (Capitoninae: *Capito*) from the eastern Andes of Peru. The Auk 117:569–577.
- Pedregosa, F., G. Varoquaux, A. Gramfort, V. Michel, B. Thirion, O. Grisel, M. Blondel, P. Prettenhofer, R. Weiss, V. Dubourg, J.

Vanderplas, et al. (2011). Scikit-learn: Machine learning in Python. Journal of Machine Learning Research 12:2825–2830.

- Peters, J. L. (1951). Check-list of Birds of the World, vol. 7. Museum of Comparative Zoology, Cambridge, MA, USA.
- Rambaut, A., M. A. Suchard, D. Xie, and A. J. Drummond (2014). Tracer 1.6. http://tree.bio.ed.ac.uk/software/tracer/
- Remsen, J. V., Jr., J. I. Areta, C. D. Cadena, S. Claramunt, A. Jaramillo, J. F. Pacheco, J. Pérez-Emán, M. B. Robbins, F. G. Stiles, D. F. Stotz, and K. J. Zimmer (2017). A classification of the bird species of South America. American Ornithologists' Union. http://www.museum.lsu.edu/~Remsen/SACCBaseline.htm
- Ridgway, R. (1909). New genera, species and subspecies of Formicariidae, Furnariidae, and Dendrocolaptidae. Proceedings of the Biological Society of Washington 22:69–74.
- Ridgway, R. (1912). Color Standards and Color Nomenclature. Published by the author, Washington, DC, USA.
- Ronquist, F., M. Teslenko, P. van der Mark, D. L. Ayres, A. Darling, S. Höhna, B. Larget, L. Liu, M. A. Suchard, and J. P. Huelsenbeck (2012). MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. Systematic Biology 61:539–542.
- Stamatakis, A. (2014). RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics 30:1312–1313.
- Stouffer, P. C. (2007). Density, territory size, and long-term spatial dynamics of a guild of terrestrial insectivorous birds near Manaus, Brazil. The Auk 124:291–306.
- Todd, W. E. C. (1927). New gnateaters and antbirds from tropical America, with a revision of the genus *Myrmeciza* and its allies. Proceedings of the Biological Society of Washington 40:149–178.
- Zimmer, J. T. (1932). Studies of Peruvian birds. VI. The formicarian genera *Myrmoborus* and *Myrmeciza* in Peru. American Museum Novitates 545:1–24.
- Zimmer, K. J. and M. L. Isler. (2003). Family Thamnophilidae (typical antbirds). In Handbook of Birds of the World, vol. 8: Broadbills to Tapaculos (J. del Hoyo, A. Elliott, and D. A. Christie, Editors). Lynx Edicions, Barcelona, Spain.

#### APPENDIX

# Specimen and Tissue Samples Used for Phylogenetic and Morphometric Analyses

APPENDIX TABLE 3. Individuals sampled for mitochondrial phylogeny, including outgroup taxa.

Taxon	GenBank accession numbers <sup>a</sup>	Tissue catalog number <sup>b</sup>	Locality	Latitude	Longitude
Myrmoderus eowilsoni <sup>c</sup>	KY849953, KY849952	LSUMZ B-93477	Peru: San Martín; 2.5 km WNW Flor de Café	7.39°S	76.32°W
	KY849951, KY849950	LSUMZ B-93476	Peru: San Martín; 2.5 km WNW Flor de Café	7.39°S	76.32°W
Myrmoderus f. ferrugineus	HM637280, HM637145	LSUMZ B-55285	Suriname: Sipaliwini; 78 km S Apura	4.48°N	57.04°W
, _	KY849954, KY849937	LSUMZ B-20403	Brazil: Amazonas; ~80 km N Manaus	2.37°S	60.04°W
Myrmoderus f. elutus	KY849936, KY849935	MZUSP 96106	Brazil: Amazonas; left bank of Rio Sucunduri	5.28°S	59.7°W
	KY849934, KY849933	LSUMZ B-86315	Brazil: Amazonas; Rio Sucunduri	6.77°S	59.09°W
Myrmoderus r. ruficauda	KY849943, NA	LGEMA 1527	Brazil: Bahia; Ribeirão do Largo, Fazenda Boa Esperança	15.42°S	40.63°W
Myrmoderus r. soror	KY849945, KY849944 KY849947, KY849946	FMNH 392445 FMNH 399262	Brazil: Pernambuco; Serra do Espelho Brazil: Alagoas; Ibateguara	8.7°S 8.98°S	35.83°W 35.87°W

#### **APPENDIX TABLE 3.** Continued.

Taxon	GenBank accession numbers <sup>a</sup>	Tissue catalog number <sup>b</sup>	Locality	Latitude	Longitude
Myrmoderus loricatus	KY849940, KY849939	MZUSP 85430	Brazil: Minas Gerais; Mina da Alegria, Mariana	20.38°S	43.42°W
	KY849942, KY849941	MZUSP 85433	Brazil: Minas Gerais; Mina da Alegria, Mariana	20.38°S	43.42°W
	KC715502, KY849938	LGEMA 13014	Brazil: Rio de Janeiro; PN Serra dos Órgãos (Teresópolis)	22.43°S	42.99°W
Myrmoderus squamosus	HM637281, HM637146	LSUMZ B-16940	Brazil: São Paulo; Salesopolis	23.53°S	45.85°W
	KC715574, KY849948	LGEMA 1526	Brazil: São Paulo; Biritiba Mirim	23.56°S	46.05°W
	KC715576, KY849949	MZUSP 81159	Brazil: São Paulo; E. E. Bananal	22.81°S	44.37°W
Thamnophilus doliatus <sup>d</sup>	EF030296, EF030234	UWBM 71559	Bolivia: Santa Cruz; Provincia de Cordillero, Abapo	18.9°S	63.4°W
Hylophylax naevioides <sup>d</sup>	EF640019, EF640086	LSUMZ B-2230	Panama: Darien; ~6 km NW Cana on E slope Cerro Pirre	7.8°N	77.72°W
Myrmeciza longipes <sup>d</sup> Pithys albifrons <sup>d</sup>	HM637279, HM637144 EF640047, EF640114	LSUMZ B-46533 FMNH 391430	Panama: Panama; ~7 km SE Pacora Brazil: Amapá; Fazenda Itapoã	9.04°N 2.07°N	79.25°W 50.93°W

 <sup>a</sup> For ND2 and ND3, respectively.
<sup>b</sup> Abbreviations: FMNH = Field Museum of Natural History; LGEMA = Laboratório de Genética e Evolução Molecular de Aves, São Paulo; LSUMZ = Louisiana State University Museum of Natural Science; MZUSP = Museum of Zoology of the University of São Paulo; and UWBM = University of Washington Burke Museum.

<sup>c</sup> Holotype.

<sup>d</sup> Outgroup.

#### **APPENDIX TABLE 4.** Specimens used for morphometric analyses (n = 105; measurement data are shown in Table 2).

Species	Catalog number <sup>a</sup>	Locality	Latitude	Longitude
Myrmoderus eowilsoni	CORBIDI 12380	Peru: San Martín; 2.5 km WNW Flor de Café	7.39°S	76.32°W
-	CORBIDI 12381	Peru: San Martín; 2.5 km WNW Flor de Café	7.39°S	76.32°W
	CORBIDI 12382	Peru: San Martín; 1.6 km NE Flor de Café	7.39°S	76.29°W
	CORBIDI 12383	Peru: San Martín; 1.6 km NE Flor de Café	7.39°S	76.29°W
	LSUMZ 190882	Peru: San Martín; 2.5 km WNW Flor de Café	7.39°S	76.32°W
	LSUMZ 190883	Peru: San Martín; 2.5 km WNW Flor de Café	7.39°S	76.32°W
	LSUMZ 190884	Peru: San Martín; 2.5 km WNW Flor de Café	7.39°S	76.32°W
	LSUMZ 190885	Peru: San Martín; 6 km SE Flor de Café	7.43°S	76.25°W
Myrmoderus f. ferrugineus	FMNH 108377	Guyana: Essequibo River, Rockstone	5.98°N	58.55°W
	FMNH 108378	Guyana: Essequibo River, Rockstone	5.98°N	58.55°W
	FMNH 120247	Guyana: Corentyne, Itabu Creek head	1.55°N	58.17°W
	FMNH 120248	Guyana: Corentyne, Itabu Creek head	1.55°N	58.17°W
	FMNH 120249	Guyana: Corentyne, Itabu Creek head	1.55°N	58.17°W
	FMNH 264476	Suriname: Wilhelmina Mountains, West River	3.43°N	56.75°W
	FMNH 264477	Suriname: Wilhelmina Mountains, West River	3.43°N	56.75°W
	FMNH 295737	Brazil: Amapá; mun. Macapá, Rio Amaparí	1.6°N	52.49°W
	FMNH 295738	Brazil: Amapá; mun. Macapá, Rio Amaparí	1.6°N	52.49°W
	FMNH 373073	Guyana: Essequibo; between Tapakuma and Pomeroon rivers	7.24°N	58.72°W
	FMNH 56900	Guyana: Mazaruni-Potaro, Bartica Grove	6.41°N	58.63°W
	KU 88819 <sup>b</sup>	Guyana: Iwokrama Reserve; ~41 km SW Kurupukari	4.34°N	58.85°W
	KU 89038	Guyana: North West district, Baramita	7.35°N	60.48°W
	KU 90840	Guyana: Parabara savanna	2.16°N	59.3°W
	LSUMZ 178456	Suriname: Sipaliwini; $\sim$ 70 km SE Apura	4.66°N	56.76°W
	LSUMZ 178457 <sup>b</sup>	Suriname: Sipaliwini; $\sim$ 70 km SE Apura	4.66°N	56.76°W
	LSUMZ 178458	Suriname: Sipaliwini; $\sim$ 70 km SE Apura	4.66°N	56.76°W
	LSUMZ 67334	Brazil: Amapá; alto Rio Araguari	1.52°N	52.02°W
	LSUMZ 67335	Brazil: Amapá, Estrada de Ferro Amapá	0.67°N	51.83°W
	LSUMZ 67336	Brazil: Amapá, Estrada de Ferro Amapá	0.67°N	51.83°W
	LSUMZ 67337	Brazil: Amapá, Fóz do Rio Falcino	0.93°N	51.58°W
	USNM 147368	Guyana: Muzaruni-Potaro, Bartica Grove	6.41°N	58.63°W
	USNM 147369	Guyana: Muzaruni-Potaro, Bartica Grove	6.41°N	58.63°W

The Auk: Ornithological Advances 135:114–126, © 2018 American Ornithological Society

# **APPENDIX TABLE 4.** Continued.

	Catalog			
Species	number <sup>a</sup>	Locality	Latitude	Longitude
Myrmoderus f ferruaineus	USNM 514992	Brazil: Amaná Serra do Navio Amaná	0.98°N	52 05°₩
nijinioacias n terragineas	USNM 514993	Brazil: Amapá, Serra do Navio Amapá	0.98°N	52.05°W
	USNM 515594	Brazil: Amapá, Serra do Navio Amapá	0.98°N	52.05°W
	USNM 586395	Guvana: North West district, Baramita	7.35°N	60.48°W
	USNM 625221	Guvana: North side Acari Mountains	1.38°N	58.93°W
	USNM 625526	Guyana: upper Essequibo River	1.58°N	58.63°W
	USNM 625527	Guyana: upper Essequibo River	1.58°N	58.63°W
	USNM 627058	Guyana: Courantyne River, 8 km S Siparuta	5.17°N	57.29°W
	USNM 637017	Guyana: Upper Takutu, upper Rewa River	3.45°N	58.58°W
	USNM 637075	Guyana: Upper Takutu, upper Rewa River	3.45°N	58.58°W
	USNM 84093	Guyana: Demerara	6.77°N	58.15°W
	USNM 90584	Guyana: Demerara	6.77°N	58.15°W
	USNM 92006	Guyana: Demerara	6.77°N	58.15°W
	USNM 92016	Guyana: Demerara	6.77°N	58.15°W
Myrmoderus f. elutus	FMNH 254862	Brazil: Pará; Villa Braga	4.42°S	56.29°W
	LACM 31977	Brazil: Pará; Rio Tapajós, Urucurituba	3.8°S	55.53°W
	LACM 31978	Brazil: Pará; Rio Tapajós, Urucurituba	3.8°S	55.53°W
	LACM 31979	Brazil: Pará; Rio Tapajós, Urucurituba	3.8°S	55.53°W
	LACM 31980	Brazil: Pará; Rio Tapajós, Urucurituba	3.8°S	55.53°W
Myrmoderus r. ruficauda	LSUMZ 113498	Brazil: Espírito Santo, Lagoa Juparanã	19.27°S	40.11°W
	LACM 28048	Brazil: Espírito Santo, Pau Gigante	19.83°S	40.37°W
	LACM 28049	Brazil: Espírito Santo, Pau Gigante	19.83°S	40.37°W
	LACM 28050	Brazil: Espírito Santo, Pau Gigante	19.83°S	40.37°W
	USNM 368286	Brazil: Espírito Santo, Mun. Santa Cruz	NA	NA
	USNM 368287	Brazil: Espírito Santo, Mun. Colatina, Linhares	19.4°S	40.07°W
	USNM 368288	Brazil: Espirito Santo, Mun. Colatina, Linhares	19.4°S	40.07°W
	USNM 368289	Brazil: Espirito Santo, Mun. Colatina, Linhares	19.4°S	40.07°W
	USNM 44799	Brazil: Bahia	NA	NA
Myrmoaerus r. soror	FMINH 311031	Brazil: Alagoas, Usina Sinimpu	9.92'S	36.13°W
Myrmoderus loricatus		Brazil: Alagoas, Usina Sinimbu Brazil: Alagoas, Usina Sinimbú	9.92'S	36.13°W
	LACIVI 20939	Didzii: Aidgods, Usina Sinimbu Prazil: Alagoas, Usina Sinimbu	9.92 S	26.12 VV
	LACINI 20940	Brazili, Alagoas, Usina Sinimbu Brazili, Alagoas, Usina Sinimbu	9.92 3	26 12°\N/
Myrmoderus Ioricatus	EACIN 20941 EMNH 311032	Brazil: Rio de Janeiro, Parque Nacional do Itatiaia	9.92 J 22 1/1°S	14.61°W
Mynnouerus Ioncutus	EMNH 311032	Brazil: Rio de Janeiro, Miguel Pereira	22. <del>44</del> 5 22.45°S	/13 37°\\/
	ΙΔCM 27950	Brazil: Rio de Janeiro, Terezonolis, Fazenda Rôa Fe	22. <del>4</del> 5 5 22.46°S	42 93°\\/
	LACM 66626	Brazil: Rio de Janeiro, Parque Nacional do Itatiaia	22.30 S	44 61°W
	LACM 74070	Brazil: Rio de Janeiro, Terezopolis, Fazenda Bôa Fe	22.11 S	42 93°W
	LACM 74071	Brazil: Rio de Janeiro, Miguel Pereira	22.45°S	43.37°W
	LSUMZ 113499	Brazil: Bahia. Itirussú	13.53°S	40.15°W
	USNM 14441 b	South America: unknown locality	NA	NA
	USNM 32856	Brazil: unknown locality	NA	NA
	USNM 42613	South America: unknown locality	NA	NA
	USNM 525966	South America: unknown locality	NA	NA
	USNM 84094	Brazil: Bahia	NA	NA
	USNM 84095	Brazil: Bahia	NA	NA
Myrmoderus squamosus	FMNH 258114	Brazil: São Paulo, Trapandé, Pasto Grande	25.07°S	47.93°W
	FMNH 265197	Brazil: São Paulo, Rio Ipiranga, Bôa Vista	24.37°S	47.83°W
	FMNH 265198	Brazil: São Paulo, Rio Juquiá, Pousinho	24.37°S	47.55°W
	FMNH 265199	Brazil: São Paulo, Rio Ipiranga, Laranja Azeda	24.32°S	47.85°W
	FMNH 265200	Brazil: São Paulo, Rio Ipiranga, Laranja Azeda	24.32°S	47.85°W
	FMNH 265201	Brazil: São Paulo, Barra do Rio Juquiá	24.37°S	47.82°W
	FMNH 344536	Brazil: São Paulo, Barra do Icapara	24.67°S	47.42°W
	FMNH 344537	Brazil: São Paulo, Barra do Icapara	24.67°S	47.42°W
	FMNH 344538	Brazil: São Paulo, Barra do Icapara	24.67°S	47.42°W
	FMNH 344539	Brazil: São Paulo, Barra do Icapara	24.67°S	47.42°W
	FMNH 344540	Brazil: São Paulo, Barra do Icapara	24.67°S	47.42°W
	FMINH 344541	Brazil: Sao Paulo, Estacao Engenheiro Ferraz	23.98°S	46.6°W
	FMINH 344542	Brazii: Sao Paulo, Barra do Icapara	24.67°S	47.42°W

#### **APPENDIX TABLE 4.** Continued.

Species	Catalog number <sup>a</sup>	Locality	Latitude	Longitude
Myrmoderus squamosus	FMNH 344543	Brazil: São Paulo, Costão dos Engenhos	24.67°S	47.5°W
	FMNH 344544	Brazil: São Paulo, Costão dos Engenhos	24.67°S	47.5°W
	FMNH 50741	Brazil: São Paulo, Fazenda Cajoa	22.95°S	49.98°W
	FMNH 56960	Brazil: São Paulo, Victoria	22.78°S	48.4°W
	FMNH 73615	Brazil: Santa Catarina, Joinville	26.3°S	48.83°W
	FMNH 73616	Brazil: Santa Catarina, Joinville	26.3°S	48.83°W
	FMNH 73617 <sup>b</sup>	Brazil: Santa Catarina, Joinville	26.3°S	48.83°W
	KU 67585	Brazil: São Paulo, Salesopolis, Casa Grande	23.54°S	45.85°W
	LSUMZ 52763	Brazil: São Paulo, Salesopolis, Casa Grande	23.54°S	45.85°W
	LSUMZ 65179	Brazil: São Paulo, Salesopolis, Casa Grande	23.54°S	45.85°W
	LSUMZ 68029	Brazil: São Paulo, Salesopolis, Casa Grande	23.54°S	45.85°W
	LSUMZ 68030	Brazil: São Paulo, Salesopolis, Casa Grande	23.54°S	45.85°W
	LSUMZ 68259	Brazil: São Paulo, Salesopolis, Casa Grande	23.54°S	45.85°W
	USNM 108244	Brazil: Santa Catarina	NA	NA
	USNM 177730	Brazil: Iguape	NA	NA

<sup>a</sup> Abbreviations: LSUMZ = Louisiana State University Museum of Natural Science; CORBIDI = Centro de Ornitología y Biodiversidad; FMNH = Field Museum of Natural History; USNM = National Museum of Natural History, Smithsonian Institution; KU = University of Kansas Biodiversity Institute; and LACM = Natural History Museum of Los Angeles County.

<sup>b</sup> Excluded from PCA because of incomplete data.