

# **KEYS TO THE FAMILIES OF FULGOROMORPHA WITH EMPHASIS ON PLANTHOPPERS OF POTENTIAL ECONOMIC IMPORTANCE IN THE SOUTHEASTERN UNITED STATES (HEMIPTERA: AUCHENORRHYNCHA)**

Author: Wilson, Stephen W.

Source: Florida Entomologist, 88(4) : 464-481

Published By: Florida Entomological Society

URL: [https://doi.org/10.1653/0015-4040\(2005\)88\[464:KTTFOF\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2005)88[464:KTTFOF]2.0.CO;2)

---

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## KEYS TO THE FAMILIES OF FULGOROMORPHA WITH EMPHASIS ON PLANTHOPPERS OF POTENTIAL ECONOMIC IMPORTANCE IN THE SOUTHEASTERN UNITED STATES (HEMIPTERA: AUCHENORRHYNCHA)

STEPHEN W. WILSON

Department of Biology, Central Missouri State University, Warrensburg, MO 64093

### ABSTRACT

Planthoppers (Hemiptera: Fulgoromorpha) are of concern to agriculture in the southeastern United States because of their potential to cause significant damage to economically important plants. These keys are intended to aid diagnosticians attempting to identify planthopper crop pests. This work includes lists of pathogen vectors and those species considered pests of economically important plants grown in the region, brief descriptions of planthopper collecting and curating techniques, a summary of morphological features used for identification, a revised key for identification of planthoppers to family, a key to the economically important delphacids, and a list of literature useful for planthopper identification.

Key Words: Insecta, Homoptera, Delphacidae, Cixiidae, pests

### RESUMEN

Las cigarritas (Hemiptera: Fulgoromorpha) son de mucho interés a la agricultura en el suroeste de los Estados Unidos por su potencial para causar daño significativo a plantas de importancia económica. Estas claves tienen la intención de ayudar a las personas que hacen el diagnóstico para identificar las cigarritas que son plagas en los cultivos. Este trabajo incluye una lista de los vectores patógenos y las especies consideradas como plagas de plantas de importancia económica sembradas en la región, una descripción breve de las técnicas para recolectar y montar las cigarritas, una clave de los delphácidos de importancia económica, y finalmente una lista de la literatura útil para la identificación de las cigarritas.

The planthoppers are a cosmopolitan group of insects placed in the infraorder Fulgoromorpha (=superfamily Fulgoroidea) which includes approximately 9,000 described species in 20 families. The vast majority of planthoppers feed on the phloem tissues of woody or herbaceous plants; however, some planthoppers feed on fungi, mosses, horsetails, or ferns (Wilson et al. 1994; Wheeler 2003). Their life histories range from highly polyphagous species that feed on numerous taxa of unrelated plants to strictly monophagous species that feed on only a single host plant species. Many planthoppers are univoltine and overwinter in the temperate zones in the egg stage; others are multivoltine where the number of generations per year depends on temperature and, perhaps, photoperiod, and many overwinter as nymphs or adults.

In this paper, I provide a brief survey of planthoppers of concern to agriculture in the southeastern United States. This survey includes lists of pathogen vectors and those species considered pests of economically important plants grown in the region, brief descriptions of planthopper collecting and curating techniques, a summary of morphological features used for identification, a revised key for identification of planthoppers to family, a key to the economically important delphacids, and a list of literature useful for planthopper identification.

### THE ECONOMICALLY IMPORTANT PLANTHOPPERS

Planthoppers are of concern to agriculture because several species are significant pests of a number of important crop plants. A list of economically important species and their host plants was provided by Wilson & O'Brien (1987); many of the species included in their list are of relatively minor economic importance in that they have been reported as pests on only one or a few occasions and/or attack plants of minor economic importance. Several planthopper species are of major importance because they reach high population densities thereby damaging crops and/or they are vectors of bacterial, phytoplasma, or viral pathogens. Planthoppers found to serve as pathogen vectors include 36 species in 4 families (Table 1). As well, 8 species in 7 families are not known to be pathogen vectors but are considered pests due to population eruptions on economically important plants; 5 other species have been introduced to the United States and could ultimately prove to be of economic significance (Table 2). The planthopper species considered pests of crops of importance in the southeastern United States—corn, rice, sugarcane, and palms—and selected studies on their biology are listed in Table 3.

TABLE 1. PLANTHOPPER VECTORS AND PATHOGENS OF ECONOMICALLY IMPORTANT PLANTS.<sup>1</sup>

Planthopper Taxon	Pathogen <sup>2</sup>	Host plants	Region <sup>3</sup>
<b>CIXIIDAE</b>			
<i>Hyalesthes obsoletus</i> Signoret	StP, GBNP	polyphagous	PA-W
<i>Myndus crudus</i> Van Duzee	LYP, CFDV (?)	coconut palm	NA, NT
<i>Myndus taffini</i> Bonfils	CFDV	coconut palm	AP
<i>Oliarus atkinsoni</i> Myers	PYP	flax lily	AP
<i>Pentastiridius beieri</i> (Wagner)	StP	sugar beet	PA-W
<b>DELPHACIDAE</b>			
<i>Caenodelphax teapae</i> (Fowler)	UHBV	plantain signalgrass	NT
<i>Chilodelphax albifascia</i> (Mats.)	RBSDV, RSV	grasses (rice, etc.)	PA-E
<i>Dicranotropis hamata</i> (Boheman)	MRDV, OSDV, PGSV	grasses (oats, etc.)	PA-W
<i>Javesella discolor</i> (Boheman)	OSDV	grasses (oats, etc.)	HO
<i>Javesella dubia</i> (Kirschbaum)	EWSMV, OSDV	grasses (wheat, etc.)	HO
<i>Javesella obscurella</i> (Boheman)	EWSMV, OSDV	grasses (wheat, etc.)	HO
<i>Javesella pellucida</i> (Fabricius)	EWSMV, MRDV, OSDV	grasses (wheat, etc.)	HO
<i>Laodelphax striatellus</i> (Fallen)	BYSMV, MRDV, NCMV, RSV, RBSDV, WCSV, WSTV	grasses (wheat, etc.)	OR, PA
<i>Megadelphax sordidulus</i> (Stål)	PGSV	grasses (timothy, etc.)	PA-W
<i>Muellerianella fairmairei</i> (Perris)	NCMV	grasses (oats, etc.)	PA-W
<i>Nilaparvata bakeri</i> (Muir)	RGSV, RRSV	rice	OR
<i>Nilaparvata lugens</i> (Stål)	RGSV, RRSV	rice	AP, OR
<i>Nilaparvata muiri</i> (Ashmead)	RGSV	rice	OR
<i>Peregrinus maidis</i> (China)	FMMV, IMMV, MSSV, MSV	maize, sorghum	AP, ET, OR, NA, NT
<i>Perkinsiella saccharicida</i> Kirkaldy	FDV	sugarcane	AP, ET
<i>Perkinsiella vastatrix</i> (Bredin)	FDV	sugarcane	AP, ET
<i>Perkinsiella vitiensis</i> Kirkaldy	FDV	sugarcane	AP
<i>Ribautodelphax notabilis</i> (Logv.)	IMMV, MRDV, RBSDV	grasses (maize, etc.)	PA-W
<i>Sogatella furcifera</i> (Horvath)	PSV	grasses	AP, OR
<i>Sogatella kolophon</i> (Kirkaldy)	BWSV, DSV, MSSV	grasses (maize, barley, etc.)	AP, ET, OR, NA, NT
<i>Sogatella vibix</i> (Haupt)	FMMV, MRDV, MSSV	grasses (maize, etc.)	AP, OR, PA
<i>Tagosodes cubanus</i> (Crawford)	EHBV, RHBV	maize, coconut	ET, NA,
<i>Tagosodes orizicolus</i> (Muir)	RHBV	grasses (rice, barnyard grass)	NA, NT
<i>Tarophagus prosperina</i> (Kirkaldy)	CBV	taro	AP, OR
<i>Terthron albovittatum</i> (Matsumura)	RSV, RBSDV	rice, barnyard grass	OR, PA-E

<sup>1</sup>Sources: Cumber 1952, 1953; Wilson & O'Brien 1987; Maixner 1994; Nault 1994; Liefing et al. 1998; Sforza et al. 1999; Ponnamma & Babjan 2000; Gatineau et al. 2001; Mpunami et al. 2002; Rajan et al. 2002; International Committee on Taxonomy of Viruses. Universal Virus Database 2004; Randles et al. 2004.

<sup>2</sup>Pathogen Abbreviations:

<b>Pathogen abbreviations:</b>			
<b>Bacteria:</b>	<i>Bacillus amylovorus</i> (Burr.) DeToni (apple tree canker)		
<b>Phytoplasmas:</b>	ANYLP      Areca nut yellow leaf phytoplasma	LYP	Lethal yellowing phytoplasma
	CP          Coconut phytoplasma (unidentified)	PYP	Phormium yellow leaf phytoplasma
	CRWP        Coconut root wilt phytoplasma	StP	Stolbur phytoplasma
	GBNP        Grapevine bois noir phytoplasma		
<b>Viruses:</b>	BWSV        Brazilian wheat spike disease	MSV	Maize stripe virus
	BYSMV       Barley yellow striate mosaic virus	NCMV	Northern cereal mosaic virus
	CBV          Colocasia babone virus	OSDV	Oat sterile dwarf virus
	CCSV        Cynodon chlorotic streak virus	PGSV	Phleum green stripe virus
	CFDV        Coconut foliar decay virus	PSV	Pangola stunt virus
	DSV          Digitaria striate virus	RBSDV	Rice black-streaked dwarf virus
	EHBV        Echinochloa hoja blanca virus	RGSV	Rice grassy stunt virus
	EWSMV      European wheat striate mosaic virus	RHBV	Rice hoja blanca virus
	FDV          Fiji disease virus	RRSV	Rice ragged stunt virus
	FMMV        Finger millet mosaic virus	RSV	Rice stripe virus
	IMMV        Iranian maize mosaic virus	UHBV	Urochloa hoja blanca virus
	MIRDV       Maize rough dwarf virus	WCSV	Wheat chlorotic streak virus
	MSSV        Maize sterile stunt virus	WRSV	Wheat rosette stunt virus

<sup>b</sup>Biogeographic Regions Abbreviations: AP—Australo-Pacific, ET—Ethiopian, HO—Holarctic, NA—Nearctic, NT—Neotropical, OR—Oriental, PA—Palaeartic, PA-E—Eastern Palaeartic, PA-W—Western Palaeartic.

TABLE 1. (CONTINUED) PLANTHOPPER VECTORS AND PATHOGENS OF ECONOMICALLY IMPORTANT PLANTS.<sup>1</sup>

Planthopper Taxon	Pathogen <sup>2</sup>	Host plants	Region <sup>3</sup>
<i>Toya propinqua</i> (Fieber)	CCSV, MRDV	grasses (Bermuda grass, maize, etc.)	ET, NA
<i>Unkanodes sapporonus</i> (Mats.)	NCMV, RSV, RBSDV	rice	OR, PA-E
<i>Unkanodes tanasijevici</i> (Dlabola)	IMMV	grasses (maize, etc.)	PA-W
DERBIDAE			
<i>Diosctrombus mkusangai</i>	CP (?)	coconut	ET
<i>Proutista moesta</i> (Westwood)	CRWP, ANYLP	coconut, areca nut	ET
FLATIDAE			
<i>Anzora unicolor</i> (Walker)	<i>B. amylovorus</i>	polyphagous	AP

<sup>1</sup>Sources: Cumber 1952, 1953; Wilson & O'Brien 1987; Maixner 1994; Nault 1994; Liefting et al. 1998; Sforza et al. 1999; Ponnamma & Babjan 2000; Gatineau et al. 2001; Mpunani et al. 2002; Rajan et al. 2002; International Committee on Taxonomy of Viruses, Universal Virus Database 2004; Randles et al. 2004.

<sup>2</sup>Pathogen Abbreviations:  
Bacteria: *Bacillus amylovorus* (Burr.) DeToni (apple tree canker)  
Phytoplasmas: ANYLP Areca nut yellow leaf phytoplasma LYP Lethal yellowing phytoplasma  
CP Coconut phytoplasma (unidentified) PYP Phormium yellow leaf phytoplasma  
CRWP Coconut root wilt phytoplasma StP Stolbur phytoplasma  
GBNP Grapevine bois noir phytoplasma  
Viruses: BWSV Brazilian wheat spike disease MSV Maize stripe virus  
BYSMV Barley yellow striate mosaic virus NCMV Northern cereal mosaic virus  
CBV Colocasia babone virus OSDV Oat sterile dwarf virus  
CCSV Cynodon chlorotic streak virus PGSV Phleum green stripe virus  
CFDV Coconut foliar decay virus PSV Pangola stunt virus  
DSV Digitaria striate virus RBSDV Rice black-streaked dwarf virus  
EHBV Echinochloa hoja blanca virus RGSV Rice grassy stunt virus  
EWSMV European wheat striate mosaic virus RHBV Rice hoja blanca virus  
FDV Fiji disease virus RRSV Rice ragged stunt virus  
FMMV Finger millet mosaic virus RSV Rice stripe virus  
IMMV Iranian maize mosaic virus UHBV Urochloa hoja blanca virus  
MRDV Maize rough dwarf virus WCSV Wheat chlorotic streak virus  
MSSV Maize sterile stunt virus WRSV Wheat rosette stunt virus

<sup>3</sup>Biogeographic Regions Abbreviations: AP—Australo-Pacific, ET—Ethiopian, HO—Holarctic, NA—Nearctic, NT—Neotropical, OR—Oriental, PA—Palaeartic, PA-E—Eastern Palaeartic, PA-W—Western Palaeartic.

PLANTHOPPER COLLECTING  
AND CURATING TECHNIQUES

Collecting Techniques. The method employed for most efficiently collecting planthoppers largely depends upon the structure of the host plant. Those on woody plants can be collected individually by knocking specimens into a vial, by use of an aspirator with a relatively large opening, or by putting a sheet or large net under the foliage and striking the branches above. Planthoppers on herbaceous plants can be collected with a sweep net with a very deep net bag which is slowly opened and the planthoppers collected with an aspirator as they crawl toward the opening. Another type of sweep net is constructed of a fine woven translucent plastic netting that retains its open shape; after sweeping, this type of net is held up toward the sun, the planthoppers tend to congregate at the back of the net and can be collected with an aspirator. As many planthoppers tend to live within the thatch near the ground, collecting them requires sitting on the ground, pushing the plants aside, then waiting for the insects to crawl up the plants where they

can be sucked up in an aspirator (Beamer, L. 1946). A more efficient means of collecting in this type of habitat is to employ a vacuum, either a "D-vac" which is very heavy and has a somewhat unreliable engine or a modified leaf blower/vacuum which is much lighter and more reliable (Wilson et al. 1993). The net bag at the end of the vacuum can be placed in a plastic jar containing cotton balls soaked in ethyl acetate and the specimens sorted out later, or the bag can be emptied into a sweep net and the specimens collected with an aspirator. Planthoppers also can be collected with Malaise traps, Sante canopy traps, window traps (Weber & Wilson 1981), and blacklights. Blacklights are useful for collecting achilids, cixiids, and some dictyopharids but are of limited value in collecting certain delphacids and caliscelids as the vast majority of their populations are composed of brachypters.

Curating and Specimen Preparation. Collected planthoppers may be killed by placing them directly into 70% ethyl or isopropyl alcohol, or putting specimens in killing jars containing a cotton ball or plaster bottom soaked in ethyl acetate, or by freezing specimens. Nymphal specimens must

TABLE 2. NON-VECTOR PLANTHOPPERS OF ECONOMICALLY IMPORTANT PLANTS OF CONCERN IN THE SOUTHEASTERN UNITED STATES.

Planthopper Taxon	Host plants	Native region <sup>1</sup>	Introduced to
CALISCELIDAE			
<i>Asarcopus palmarum</i> Horvath	date palm	PA-W	California <sup>2</sup>
<i>Caliscelis bonellii</i> (Latreille)	grasses (?)	PA-W	California <sup>3</sup>
DELPHACIDAE			
<i>Eoerysa flavocapitata</i> Muir	sugarcane	OR	—
<i>Harmalia anacharsis</i> Fennah	rice (?)	OR	Florida <sup>4</sup>
<i>Saccharosydne saccharivora</i> (West.)	sugarcane	NA, NT	
FLATIDAE			
<i>Melormenis basalis</i> (Walker)	polyphagous (?)	NT	Florida <sup>5</sup>
<i>Metcalfa pruinosa</i> (Say)	polyphagous	NA	Europe
<i>Ormenaria rufifascia</i> (Walker) <sup>6</sup>	palms	NA	—
<i>Siphanta acuta</i> (Walker)	polyphagous	AP	California <sup>5</sup>
LOPHOPIDAE			
<i>Pyrilla perpusilla</i> (Walker)	sugarcane, rice	OR	—
MEENOPLIDAE			
<i>Nisia nervosa</i> (Motschulsky)	rice	OR	—
TETTIGOMETRIDAE			
<i>Hilda patruelis</i> (Stål) <sup>7</sup>	peanuts	ET	—
TROPIDUCHIDAE			
<i>Numicia viridis</i> Muir <sup>8</sup>	sugarcane	ET	—
<i>Ommatissus lybicus</i> DeBerg	date palm	PA-W	—

<sup>1</sup>See Table 1 for abbreviations; <sup>2</sup>Stickney et al. 1950; <sup>3</sup>O'Brien 1967; <sup>4</sup>Wooten et al. 1993; <sup>5</sup>Wilson, S. W. pers. obs.; <sup>6</sup>Mead 1965, Wilson & Tsai 1984; <sup>7</sup>Weaving 1980; <sup>8</sup>Carnegie 1980.

be stored in vials, preferably with caps with plastic cone inserts, filled with 70% ethyl or isopropyl alcohol; adults may be stored this way but colors will usually bleach out. Adult specimens are usually pinned or glued to a point on the right side of the specimen. Label data accompanying the specimen should include locality (country, state or province, county or district, nearest town, specific location, and if available, GPS coordinates), date, collector, and, if possible, host plant.

Accurate species identification in most planthopper taxa requires examination of the structures of the male genitalia; unfortunately, for many taxa, it is not possible to identify females to species, and in some cases, to genus level. Identification of males to species usually requires removing the abdomen with a pair of fine forceps or the point of an insect pin, then placing the abdomen in a 10% potassium hydroxide (KOH) solution for 12-24 h at room temperature (small, lightly sclerotized specimens may require as little as 4-8 h or less in the KOH solution). An efficient means of processing a relatively large number of specimens is to fill wells of a 96-well tissue culture plate with water, place a KOH pellet in each well containing water, then place an abdomen in a well. The pinned specimen is then placed in a Schmidt box or foam bottom insect drawer that has been

marked with grid locations corresponding to those on the 96-well tissue culture plate (A1-H12). To examine the genitalia, remove the abdomen from the KOH solution and place it in a deep depression slide filled with water; let it soak in the water for a few minutes then place it in a depression slide filled with glycerine. If the genitalic structures are not clearly visible, hold the pygofer with a pair of fine forceps or the point of an insect pin then gently move the anal tube caudoventrally, or push the connective attached to the anterior base of the aedeagus caudally. This will usually force the aedeagus, anal tube, and styles out of the pygofer. After examination of the genitalia, fill a genital vial with glycerine, place the abdomen into the vial, cap the vial, then pin the vial through the cap under the specimen.

MORPHOLOGY

This brief review of planthopper external morphology should provide enough basic information to allow effective use of the keys for identification. More complete treatments are provided by O'Brien & Wilson (1985) for all families, Asche (1985) for delphacids, Freund & Wilson (1995) for acanaloniids, and McPherson & Wilson (1995) for dictyopharids. Overviews of the anatomy of the male

TABLE 3. SELECTED BIOLOGICAL STUDIES ON THE PRINCIPAL PLANTHOPPER PESTS OF THE MAJOR ECONOMICALLY IMPORTANT PLANTS IN THE SOUTHEASTERN UNITED STATES.

Host plant	Planthoppers	Biological studies
Corn <sup>1</sup>	<i>Peregrinus maidis</i> (Ashmead)	Tsai & Wilson 1986
	<i>Sogatella vibix</i> (Haupt)	Ammar 1977
Sugarcane <sup>2</sup>	<i>Eoerysa flavocapitata</i> Muir	Chatterjee 1971, Wilson & Tsai 1992
	<i>Numicia viridis</i> Muir	Carnegie 1980
	<i>Perkinsiella saccharicida</i> Kirkaldy	Bull 1972, Chang & Ota 1978, Sosa 1985
	<i>Pyrilla perpusilla</i> (Walker)	Brar & Bains 1979, Rahman & Nath 1940
	<i>Saccharosydne saccharivora</i> (West.)	Metcalfe 1969
Rice <sup>3</sup>	<i>Laodelphax striatellus</i> (Fallen)	Ruan et al. 1981
	<i>Nilaparvata lugens</i> (Stål)	Bae & Pathak 1970, Mochida & Okada 1979, Wilson & Claridge 1991
	<i>Nilaparvata meander</i> Fennah	Wilson & Claridge 1991
	<i>Sogatella furcifera</i> (Horvath)	Ammar et al. 1980
	<i>Tagosodes cubanus</i> (Crawford)	Wilson & Claridge 1991
	<i>Tagosodes orizicolus</i> (Muir)	Atkins et al. 1958
Palms <sup>4</sup>	<i>Asarcopus palmarum</i> Horvath	Stickney et al. 1950
	<i>Myndus crudus</i> Van Duzee	Howard et al. 1983, Wilson & Tsai 1982
	<i>Ommatissus lybicus</i> DeBerg	Hussain 1963, Klein & Venezian 1985, Asche & Wilson 1989

<sup>1</sup>*Zea mays* L.; <sup>2</sup>*Saccharum officinarum* L.; <sup>3</sup>*Oryza sativa* L.; <sup>4</sup>*Cocos nucifera* L., *Phoenix dactylifera* L.

genitalia for selected taxa are given by Muir (1926a); Fennah (1945); Bourgoin (1990); and Yang & Chang (2000). Descriptions and illustrations of the anatomy of the female genitalia are provided by Heady & Wilson (1990) and Bourgoin (1993).

Planthoppers can be separated from members of the Cicadomorpha by the location of the antennae which are ventral to the compound eyes (except in one cixiid subfamily in which the compound eyes are positioned anteroventrally to the compound eyes but are partially surrounded by cup-like concavities); the enlarged antennal pedicel bearing wart-like sensilla; the presence, in most taxa, of pad-like tegulae at the base of the forewings; the presence, again, in most taxa, of a “Y-vein”, that is, two anal veins coalescing on the clavus of the forewing to form a “Y-shaped” set of veins; the elongate, widely separated base of the mesothoracic coxae; and the lack of rows of setae along the shaft of each metathoracic tibia as in Cicadellidae.

Head. The head in planthoppers ranges from a broad, short head to those that are strongly laterally compressed or some that have an elongate straight or curved projection extending well beyond the compound eyes. The dorsal aspect of the head is termed the vertex which may be separated from the anterior or anteroventral frons by a transverse carina or, in those planthoppers with broadly rounded heads, not be demarcated from it. The frons, in all but one family, is separated from the gena on each side by a longitudinal external lateral carina; as well, the frons may bear a median longitudinal carina and/or internal longitudi-

nal lateral carinae. Posteroventral to the frons is the clypeus from which the three-segmented beak extends. Some planthoppers have a median ocellus on the frons; most have a lateral ocellus between the frons and gena. The compound eyes are generally large and may be strongly notched ventrally; in some subterranean species the eyes may be reduced or absent. The three segmented antennae in almost all planthoppers are situated ventral to the compound eyes; in some cixiids the antennae are in a cup-like concavity located anterior to the eyes. The basal antennal antennal segment is termed the scape, the middle segment which may be elongate and/or foliose is the pedicel, the terminal segment consists of a bulbous base and elongate, seta-like flagellum.

Thorax. The pronotum is generally short and collar-like and may bear a median longitudinal carina and a pair of lateral carinae. The mesonotum is generally subtriangular and may bear a median longitudinal carina and a pair of lateral carinae; a transverse suture demarcates a small triangular piece at the apex of the mesonotum in tropiduchids and some lophopids. The fore wings may be very large with numerous veins or greatly reduced as in brachypters and may be opaque, concolorous with the body or brightly colored, or completely or partially translucent. The subtriangular anal region of the forewings, or clavus, is set off from the rest of the wing by the postcubital vein which may or may not reach the edge of the wing. The clavus typically bears a “Y-shaped” vein and, in flatids, has numerous pustules on its sur-

TABLE 4. KEYS FOR IDENTIFICATION OF PLANTHOPPERS IN ECONOMICALLY SIGNIFICANT TAXA.

Taxa	Region or Topic	Reference
Families	World	O'Brien & Wilson 1985
Families	World, immatures	O'Brien et al. 1998
Families, Genera, Species	Central Europe	Holzinger et al. 2003
Families, Genera, Species	Fennoscandia	Ossiannilsson 1978
Families, Genera, Species	Eastern Russia	Anufriev & Emeljanov 1988
Families, Genera, Species	Eastern U.S.	Metcalf 1923
Families, Genera, Species	Illinois	Wilson & McPherson 1980c
Families, Genera, Species	Puerto Rico	Caldwell & Martorell 1951
Families, Genera, Species	Rice pests	Wilson & Claridge 1991
Acanaloniidae		
<i>Acanalonia</i>	U.S.	Freund & Wilson 1995
Caliscelidae		
Genera	U.S.	Doering 1938
<i>Aphelonema</i>	U.S.	Doering 1941
<i>Bruchomorpha</i>	U.S.	Doering 1940
<i>Fitchiella</i>	U.S.	Doering 1941
Cixiidae		
Genera	U.S.	Kramer 1983
<i>Cixius</i>	U.S.	Kramer 1981
<i>Hyalesthes</i>	Europe	Hoch & Remane 1985
<i>Myndus</i>	New World	Kramer 1979
<i>Oecleus</i>	U.S.	Kramer 1977
<i>Oliarus</i>	U.S.	Mead & Kramer 1982
Delphacidae		
Genera	Mississippi	Coley 1970
<i>Achorotile</i>	U.S.	Beamer 1954
<i>Bakerella</i>	U.S.	Beamer 1950
<i>Bostaera</i>	U.S.	Penner 1952
<i>Delphacodes</i>	North Carolina	DuBose 1960
<i>Eurysa</i>	U.S.	Beamer 1952
<i>Kelisia</i>	U.S.	Beamer 1951
<i>Laccocera</i>	U.S.	Penner 1945
<i>Megamelanus</i>	U.S.	McDermott 1952
<i>Megamelus</i>	U.S.	Beamer 1955
<i>Neomegamelanus</i>	U.S.	McDermott 1952
<i>Pentagramma</i>	U.S.	Penner 1947
<i>Phyllodinus</i>	U.S.	Morgan & Beamer 1949
<i>Pissonotus</i>	U.S.	Bartlett & Deitz 2000
<i>Prokelisia</i>	U.S.	Wilson 1982, Heady & Wilson 1990
<i>Sogatella</i>	World	Asche & Wilson 1990
<i>Stobaera</i>	U.S.	Kramer 1973
<i>Stenocranus</i>	U.S.	Beamer 1946
<i>Tagosodes</i>	World	Asche & Wilson 1990
<i>Tumidagena</i>	U.S.	McDermott 1952
Flatidae		
Subfamilies, Tribes	Panama	Metcalf 1938
Species	U.S.	Shepard 1939
<i>Siphanta</i>	Australasia	Fletcher 1985
Issidae		
Subfamilies, Tribes	World	Fennah 1954
Genera	U.S.	Doering 1938
<i>Hysteropterum</i>	U.S.	Doering 1938
<i>Thionia</i>	U.S.	Doering 1938
Tropiduchidae		
Subfamilies, Tribes	World	Fennah 1982
Genera, Species	U.S.	O'Brien 1992
<i>Ommatissus</i>	Old World	Asche & Wilson 1989

face. In some taxa, some or all of the forewing veins may bear enlarged pustules. The hind wings may be almost as large as the forewings or greatly reduced or absent. The hindwings of fulgorids have a characteristic reticulate venation in the anal region. The metathoracic or hind legs have numerous features useful for identification. The tibia has a row of apical spines on the plantar surface and usually bears large lateral spines along the tibial shaft. Delphacids have a characteristic metatibial spur which may be spike-like or foliose and with or without teeth on its inner margin. The tarsus consists of three tarsomeres; the basal or first tarsomere bears a row of apical teeth, the second tarsomere may have a row of apical teeth, one tooth on each side or may be lacking teeth; the third tarsomere terminates in a pair of claws and a central arolium.

**Abdomen.** Some planthoppers have sound producing tymbals which consist of internal apodemes in the basal abdominal segments and a subrectangular piece of the first abdominal tergite. The sounds made by planthoppers are species specific courtship signals detected via substrate vibrations.

The structures of the male external genitalia are necessary for accurate species identification in most groups of planthoppers. The male genital capsule is a subcylindrical pygofer and represents the modified 9th abdominal segment. The pygofer may have lateral and/or ventral caudally-directed lobes or spines and, in delphacids, may have a partial dorsoventral wall or diaphragm. The diaphragm may bear lobes, spines or other forms of diaphragm armature on its median ventral margin. The anal tube, the 10th abdominal segment, is usually moveable and often bears spines or lobes on its caudal margin. The styles are a pair of clasper-like structures emanating from the ventrocaudal aspect of the pygofer. The styles may be elongate and can have spines or teeth or bear lobes or keels or, in some taxa, they can be relatively small and club-shaped. A "Y-shaped" connective attaches the base of the styles to the base of the aedeagus. The intromittent organ, or aedeagus, ranges in shape from a simple tube-like structure to a complex elaborate entity bearing numerous spines and, in some taxa, inflatable translucent lobes.

Terminology used in describing female genitalia follows Heady & Wilson (1990). The external female genitalia, collectively referred to as the ovipositor, may be "orthopteroid" or "fulgoroid". In the "orthopteroid ovipositor" the fused median gonopophyses (=valvula) of the 9th abdominal segment often have a serrated dorsal aspect and serve as a saw-like structure for depositing eggs, usually into plant tissues. The lateral gonopophyses of the 9th abdominal segment and the gonopophyses of the 8th segment serve in egg deposition and may coat the eggs with wax. In some taxa, these structures are reduced and/or modified and

very large wax plates are present. The "fulgoroid ovipositor" may be modified for raking particles prior to egg deposition in the soil or on the ground or for depositing eggs in plant tissue. In those that insert eggs into plant tissue the fused gonopophyses of the 8th abdominal segment serve as the saw-like deposition structure and the lateral gonopophyses of the 9th abdominal segment as protective covers and/or egg guide.

#### IDENTIFICATION OF PLANTHOPPERS

There are 196 species in 11 families of planthoppers recorded from the southeastern United States (Wilson & McPherson 1980a; Wilson, unpubl. data). Most of these species are of no known economic significance; in fact, many are rare and little is known about their life cycles and hosts. A few species present in the region are of economic importance. As well, there are several species that could be imported by accident or on purpose into the region. To aid in identifying taxa of potential economic importance, keys to the 20 planthopper families and to the economically significant delphacids are included. The key to families has been adapted from O'Brien & Wilson (1985). Since publication of their key, the following family-level changes have been made: 18 genera of Issidae were transferred to Nogodinidae (Fennah 1984); the family Caliscelidae was established to include the former issid subfamilies Caliscelinae and Ommatidiotinae (Emeljanov 1999); the family Acanaloniidae was reestablished and includes the former issid subfamilies Acanaloniinae, Tonginae, and Trenopinae (Emeljanov 1999); and the family Achilixiidae was subsumed within Cixiidae (Liang 2001). Planthopper families with economically important species are designated with an asterisk (\*).

The family Acanaloniidae is identified as one that includes species of potential economic importance because *Acanalonia conica* (Say), while not a significant pest in the United States, has been introduced to Sicily (D'Urso, personal communication). This species has a similar life history and feeding habits to *Metcalfa pruinosa* (Say), also not a significant pest in the United States, which was introduced to Italy from the United States in the late 1970s and has spread from Spain and southern France to Austria, Slovenia, and Greece (Wilson & Lucchi 2001). *Acanalonia conica* is often found in mixed nymphal feeding assemblages with *M. pruinosa* (Wilson & McPherson 1980b) and could become a pest in Europe. As such, it joins the native US species, *M. pruinosa* and *Prokelisia marginata* Van Duzee (Seljak 2004) that have recently been introduced to Europe and may be intercepted at United States ports of entry. The life histories and descriptions and illustrations of the immatures of *A. conica*, *M. pruinosa*, and similar associated species are provided by Wilson & McPherson (1981a, b).




There are no recent comprehensive keys to the genera and species of a number of families that include economically significant species (e.g., Flatidae). A list of keys that identify species in these taxa and those that survey taxa on selected economically significant crops is provided in Table 4. The following key to delphacids of potential economic importance includes species present in the United States, whether native or introduced, and known to be of economic significance, viral or phytoplasma vectors that could be introduced to the US, and those that are significant pests of crops grown in the United States. Several species recorded as transmitters of viral pathogens based only on laboratory or greenhouse studies with no apparent field evidence are not considered significant pests and, thus, are not included. As well, species that are vectors of pathogens of crop plants of little significance to agriculture in the southeastern United States are not included (e.g., *Javesella* spp. except for the ubiquitous *J. pellucida* (Fabricius)). A few common species that are likely to be frequently collected and those that

can easily be confused with major pests are included in the key (e.g., *Caenodelphax teapae* (Fowler), *Nilaparvata wolcottii* Muir and Giffard). Finally, some taxa, such as *Sogatella* and *Togodes*, include numerous species, some of which could be introduced to the southeastern United States. If specimens fail to fit the key and illustrations given below, then the taxonomic literature on these taxa should be consulted (e.g., Asche & Wilson 1990; also, see Table 4). In order to ensure accurate identification of these and other delphacid taxa, species identifiers should always compare structures of the male genitalia with either the illustrations provided in this paper or with those in the works listed in Table 4. Figures accompanying the keys were redrawn from Muir & Giffard (1924); Muir (1926b); Doering (1940); Zimmerman (1948); Caldwell & Martorell (1951); Wilson & McPherson (1980c); Asche (1985); O'Brien & Wilson (1985); Tsai & Wilson (1986); Wilson & Claridge (1991); Wilson & Tsai (1992); Wooten et al. (1993); Freund & Wilson (1995); and Holzinger et al. (2003).


#### KEY TO THE FAMILIES OF PLANTHOPPERS

1. Hind tibia with a large movable spur at apex (Fig. 1) ..... Delphacidae\*  
—Hind tibia without a movable spur at apex (Figs. 2-4) ..... 2
2. Second tarsomere of hind legs with a row of apical spines (Fig. 2) ..... 3  
—Second tarsomere of hind legs with one apical spine on each side (Fig. 3) or spines absent (Fig. 4) ..... 9
3. Hindwings with numerous cross veins near apex and in anal area; head usually with a transverse suture separating vertex from frons (Fig. 5); aedeagus with inflatable lobes ..... Fulgoridae  
—Hindwings without cross veins near apex or in anal area; head without a transverse suture separating vertex from frons; aedeagus variable ..... 4
4. Forewings overlapping posteriorly (Fig. 6); body usually flattened ..... Achilidae  
—Forewings not overlapping posteriorly; body variable ..... 5
5. Forewings with tubercles on one or both claval veins (Fig. 7); beak with apical segment longer than wide ..... Meenoplidae\*  
—Forewings usually without tubercles on claval veins; if so, then all veins with tubercles or beak with apical segment subequal in length and width ..... 6
6. Forewings with tubercles on claval veins and beak with apical segment subequal in length and width; styles much longer than pygofer; anal tube not moveable, attached to pygofer ..... Derbidae (in part)  
—Forewings without tubercles on claval veins or with tubercles on all veins; beak with apical segment longer than wide; styles shorter than length of pygofer; anal tube moveable ..... 7
7. Frons with two or three median carinae and/or head with elongate anterior projection (Fig. 8); if not, then tegulae absent ..... Dictyopharidae  
—Frons with one median carina (Fig. 9); head usually without elongate anterior projection; tegulae present ..... 8
8. Forewings usually with tubercles on veins; abdominal tergites 6-8 subrectangular; females with caudally directed wax plates on tergite 9 and a spike-like ovipositor, or wax plates absent and ovipositor sword-shaped ..... Cixiidae\*  
—Forewings without tubercles on veins; abdominal tergites 6-8 chevron shaped, females with wax plates on tergites 7-9; female genitalia greatly reduced ..... Kinnaridae
9. Second tarsomere of hind legs with one apical spine on either side (Fig. 3) ..... 10  
—Second tarsomere of hind legs without apical spines on either side (Fig. 4) ..... 17




Figs. 1-13. Planthopper key structures. 1. Delphacid hind leg, dorsal view. 2. Cixiid hind leg, ventral view. 3. Flatid hind leg, ventral view. 4. Lophopid hind leg, ventral view. 5. Fulgorid head. 6 Achilid habitus. 7. Meenoplid forewing. 8. Dictyopharid head. 9. Cixiid head. 10. Tettigometrid head. 11. Tropiduchid head and thorax. 12. Flatid habitus. 13. Nogodinid head and pronotum.

- 10. Frons with lateral carinae ..... 11  
—Frons without lateral carinae (Fig. 10).....Tettigometridae\*
- 11. Mesonotum with a posterior transverse suture (Fig. 11); forewings with few or  
no cross veins anterior to nodus, with numerous veins posterior to nodus ..... Tropiduchidae\*  
—Mesonotum usually without a posterior transverse suture; forewings with sparse  
venation or dense venation throughout ..... 12
- 12. Beak with apical segment subequal in length and width; styles much longer than pygofer;  
anal tube not moveable, attached to pygofer ..... Derbidae (in part)  
—Beak with apical segment longer than wide; styles shorter than length of pygofer;  
anal tube moveable ..... 13




Figs. 14-18. Planthopper key structures. 14. Acanaloniid female genitalia, lateral view. 15. Issid female genitalia, ventral view. 16. Caliscelid head. 17. Issid head. 18. Ricaniid habitus.

13. Forewings with tubercles between veins on clavus (Fig. 12) and with numerous costal crossveins ..... Flatidae  
 —Forewings without tubercles on clavus; usually without numerous costal crossveins ..... 14
14. Ovipositor laterally compressed, gonopophyses of the 8th abdominal segment bearing apical teeth (Fig. 14); pronotum extending anteriorly beyond middle of eyes (Fig. 13), or forewings usually with reticulate venation and hind tibiae without lateral spines ..... 15  
 —Ovipositor not laterally compressed, gonopophyses of the 8th abdominal segment without apical teeth (Fig. 15); pronotum not extending anteriorly beyond middle of eyes; forewings of macropters with normal venation; hind tibiae with lateral spines ..... 16
15. Pronotum extending anteriorly beyond middle of eyes (Fig. 13); clypeus with lateral carinae; macropterous or brachypterous; hind tibia with lateral spines ..... Nogodinidae  
 —Pronotum extending not much beyond posterior aspect of eyes; clypeus without lateral carinae; macropterous, often with reticulate venation; hind tibia usually without lateral spines ..... Acanaloniidae



Figs. 19-34. *Nilaparvata* spp. 19. Hind leg, ventral view. 20-34. Male genitalia. 20-23. *N. bakeri*. 24-27. *N. muii*. 28-30. *N. lugens*. 31-32. *N. meander*. 33-34. *N. wolcottii*. 20, 24: Diaphragm armature of pygofer, caudal view. 21, 25, 28, 31, 33: Anal tube. 22, 26, 29, 32, 34: Aedeagus. 23, 27, 30: Styles.

16. Frons with lateral carinae bordering a large disc-like or elongate areolet (Fig. 16);  
usually brachypterous . . . . . Caliscelidae\*  
—Frons with median carina, with or without lateral carina (Fig. 17);  
if lateral carinae present, not enclosing a central areolet; macropterous . . . . . Issidae
17. Brachypterous, or forewings barely exceeding beyond the apex of the abdomen and with apical  
venation strongly carinate (southern Africa) . . . . . 18  
—Macropterous, forewings almost always exceeding well beyond the apex of the abdomen,  
venation not carinate. . . . . 19




Figs. 35-46. *Sogatella* spp. 35-38. *S. furcifera*. 39-42. *S. vibix*. 43-46. *S. kolophon*. 35, 39, 43: Head. 36, 40, 44: Anal tube. 37, 41, 45: Aedeagus. 38, 42, 46: Styles.

18. Eyes reduced, body white, brachypterous, adults subterranean ..... Hypochthonellidae  
 —Eyes not reduced, body pigmented, macropterous with strongly carinate  
 apical veins, adults not subterranean ..... Gengidae
19. Forewings at least twice as long as wide; claval vein extending ca. 2/3 to 3/4 distance  
 to caudal aspect of wing ..... 20  
 —Forewings broadly triangular, almost as wide as long; claval vein extending  
 almost to caudal aspect of wing (Fig. 18) ..... Ricaniidae\*
20. Clypeus with lateral carinae; vertex with length more than 1/3 width, frons usually  
 longer than wide and with 1-3 longitudinal carinae ..... Lophopidae\*  
 —Clypeus without lateral carinae; vertex with length less than 1/3 width,  
 frons wider than long with obscure median carina or carina absent ..... Eurybrachidae


#### KEY TO THE ECONOMICALLY IMPORTANT DELPHACIDAE\*

1. Hind legs with lateral spines on first (basal) tarsomere (Fig. 19) ..... *Nilaparvata*, 2  
 —Hind legs without lateral spines on first tarsomere (Fig. 1) ..... 6
2. Pygofer with process on ventral margin (Figs. 20, 24) ..... 3  
 —Pygofer without processes on ventral margin ..... 4




Figs. 47-53. *Perkinsiella* and *Tagosodes* spp. 47-49. *P. saccharicida*. 50-51. *T. orizicolus*. 52-53. *T. cubanus*. 47: Head. 48: Anal tube. 49, 50, 52: Aedeagus. 51, 53: Styles.

3. Pygofer process with lateral spines (Fig. 20); genitalia as in Figs. 21-23 ..... *N. bakeri* (Muir)  
—Pygofer with three small processes without lateral spines (Fig. 24);  
genitalia as in Figs. 25-27 ..... *N. muiri* China
4. Anal tube with elongate spines; styles not bifurcate; genitalia as in Figs. 28-30 ..... *N. lugens* (Stål)  
—Anal tube with very small upturned spines or lacking spines (Figs. 31, 33) ..... 5
5. Anal tube with very small upturned spines (Fig. 31), aedeagus as in Fig. 32 ..... *N. meander* Fennah  
—Anal tube lacking spines (Fig. 33), aedeagus as in Fig. 34 ..... *N. wolcotti* Muir & Giffard
6. Mesonotum light brown to black with median longitudinal pale stripe;  
genitalia as in Figs. 36-46, 50-53 ..... 7  
—Mesonotum without median longitudinal pale stripe;  
genitalia as in Figs. 48-49, 54-75 ..... 13
7. Pygofer with diaphragm margin thickened and U-shaped, aedeagus with dorsal  
and ventral elongate rows of teeth (e.g., Fig. 37) ..... *Sogatella*, 8  
—Pygofer with diaphragm margin not thickened and U-shaped, aedeagal teeth,  
if present, not arranged as above (e.g., Fig. 50) ..... 11
8. Frons dark brown, carinae often pale (Fig. 35); genitalia as in Figs. 36-38 ..... *S. furcifera* (Horvath)  
—Frons pale yellow (Figs. 39, 43); if frons brown then styles as in Fig. 46 ..... 9
9. Sides of head beneath eyes dark brown (Fig. 39); genitalia as in Figs. 40-42 ..... *S. vibix* (Haupt)  
—Sides of head beneath eyes pale yellow (Fig. 43) or dark brown;  
genitalia as in Figs. 44-46 ..... 10
10. Sides of head beneath eyes pale yellow (Fig. 43); forewing without dark marking  
at apex of clavus ..... *S. kolophon* (Kirkaldy)  
—Sides of head beneath eyes dark brown; frons pale or light brown;  
forewing with dark marking at apex of clavus ..... *S. molina* (Fennah)
11. Frons with median carina forked near ventral aspect of eyes, dorsal half  
of frons dark with pale spots, ventral half pale (Fig. 47);  
genitalia as in Figs. 48, 49 ..... *Perkinsiella saccharicida* Kirkaldy



Figs. 54-63. *Saccharosydne*, *Eoeurysa*, *Caenodelphax*, and *Peregrinus* spp. 54-56. *S. saccharivora*. 57-58. *E. flavocapitata*. 59-60. *C. teapae*. 61-63. *P. maidis*. 54, 57, 59, 62: Anal tube. 55, 58, 60, 63: Aedeagus. 56, 61: Styles.

- Frons with median carina forked near dorsal aspect of eyes,  
frons not darker dorsally and lighter ventrally. . . . . *Tagosodes*, 12
- 12. Styles broad at apex, aedeagus with numerous ventral teeth (Figs. 50-51) . . . . . *T. orizicolus* (Muir)  
—Styles slender at apex, aedeagus with one row of ventro-lateral  
teeth on each side (Figs. 52-53). . . . . *T. cubanus* (Crawford)
- 13. Head narrow, produced well beyond eyes; body green to yellow;  
anal tube without spines, styles with small, apical lobes,  
aedeagus elongate, ribbon-like (Figs. 54-56) . . . . . *Saccharosydne saccharivora* (Westwood)  
—Head broad, not produced much beyond eyes; body not green . . . . . 14
- 14. Forewings almost entirely dark black . . . . . 15  
—Forewings clear or with dark markings . . . . . 16
- 15. Forewings dark with pale transverse stripe; anal tube with bifurcate  
posterocaudal spine on each side (Figs. 57-58) . . . . . *Eoeurysa flavocapitata* Muir  
—Forewings dark without pale transverse stripe; anal tube  
with minute teeth on posterocaudal aspect (Figs. 59-60) . . . . . *Caenodelphax teapae* (Fowler)
- 16. Forewings with dark markings in apical half; aedeagus slender  
with spine-like dorsal projections (Figs. 61-63) . . . . . *Peregrinus maidis* (Ashmead)  
—Forewings without dark markings in apical half; aedeagus not as above . . . . . 17
- 17. Styles splayed laterally; aedeagus bent strongly ventrally (Figs. 64-66) . . . . . *Javesella pellucida* (Fabricius)  
—Styles oriented dorsoventrally; aedeagus directed caudad . . . . . 18



Figs. 64-75. *Javesella*, *Harmalia*, *Laodelphax*, and *Toya* spp. 64-66. *J. pellucida*. 67-69. *H. anacharsis*. 70-71. *L. striatellus*. 72-75. *T. propinqua*. 64, 67, 72: Anal tube. 65, 68, 70, 73: Aedeagus. 66, 69, 71, 74: Styles. 75: Diaphragm armature of pygofer, caudal view.

18. Styles each with elongate rounded process at apex; aedeagus without dorsal teeth (Figs. 67-69). . . . . *Harmalia anacharsis* Fennah  
—Styles each without elongate rounded process at apex . . . . . 19
19. Mesonotum black; aedeagus curved ventrally (Figs. 70-71) . . . . . *Laodelphax striatellus* (Fallen)  
—Mesonotum light brown to straw yellow; aedeagus straight with minute dorsal teeth near apex; pygofer with “y-shaped” dorsal armature (Figs. 72-75) . . . . . *Toya propinqua* (Fieber)

Note—to ensure accurate species identification always compare the dissected male genitalia with illustrations in the appropriate publications.

ACKNOWLEDGMENTS

I thank Dr. Amanda Hodges, Entomology and Nematology Department, University of Florida, Gainesville, for encouragement and for efforts spent organizing the “Homoptera Workshop”; Dr. Susan Halbert, Division of Plant Industry, Florida Department of Agriculture and Consumer Services, Gainesville, for enthusiasm and very fruitful discussions about some of the planthoppers introduced into Florida; the anonymous reviewers whose suggestions greatly improved the manuscript;

and the participants of the “Homoptera Workshop” who asked all the right questions and, by doing so, helped improve the keys immensely.

REFERENCES CITED

AMMAR, E. 1977. Biology of the planthopper *Sogatella vibix* (Haupt) in Giza, Egypt (Hom., Delphacidae). Deutsche Entomol. Zeitschrift 24: 151-158.  
AMMAR, E., O. LAMIE, AND I. A. KHODEIR. 1980. Biology of the planthopper *Sogatella furcifera* Horv. in Egypt



- (Homoptera: Delphacidae). Deutsche Entomol. Zeitschrift 27: 21-27.
- ANUFRIEV, G. A., AND A. F. EMELJANOV. 1988. Cicadinea (Auchenorrhyncha)—Leafhoppers, pp. 12-495 *In* P. A. Lera [ed.], Identification Guide to the Insects of the Far Eastern USSR. Vol. 2. Homoptera and Hemiptera.
- ASCHE, M. 1985. Zur Phylogenie der Delphacidae Leach, 1815 (Homoptera Cicadina Fulgoromorpha). Marburger Entomol. Publ. 2(1): 1-910.
- ASCHE, M., AND M. R. WILSON. 1989. The palm-feeding planthopper genus *Ommatissus* (Homoptera: Fulgoroidea: Tropiduchidae). Syst. Entomol. 14: 127-147.
- ASCHE, M., AND M. R. WILSON. 1990. The delphacid genus *Sogatella* and related groups: a revision with special reference to rice-associated species (Homoptera: Fulgoroidea). Syst. Entomol. 15: 1-42.
- ATKINS, J. G., J. P. KRAMER, AND S. D. HENSLEY. 1958. Hoja blanca and its insect vector found in a second area in the United States. Plant Disease Rep. 42: 1414.
- BAE, S. H., AND M. D. PATHAK. 1970. Life history of *Nilaparvata lugens* (Homoptera: Delphacidae) and susceptibility of rice varieties to its attacks. Ann. Entomol. Soc. America. 63: 149-155.
- BARTLETT, C., AND L. L. DEITZ. 2000. Revision of the New World delphacid planthopper genus *Pissonotus* (Hemiptera: Fulgoroidea). Thomas Say Publ. Entomol. Monogr. 234 pp.
- BEAMER, L. 1946. I'll tell you how. J. Kansas Entomol. Soc. 19: 127-129.
- BEAMER, R. H. 1946. The genus *Stenocranus* in America north of Mexico (Homoptera-Delphacinae). J. Kansas Entomol. Soc. 19: 1-11.
- BEAMER, R. H. 1950. The genus *Bakerella* in North America north of Mexico (Homoptera-Fulgoridae-Delphacinae). J. Kansas Entomol. Soc. 23: 102-109.
- BEAMER, R. H. 1951. A review of the genus *Kelisia* in America north of Mexico with four new species (Homoptera-Fulgoridae-Delphacinae). J. Kansas Entomol. Soc. 24: 117-121.
- BEAMER, R. H. 1952. The genus *Eurysa* in America north of Mexico (Homoptera: Fulgoridae, Delphacinae). Pan-Pacific Entomol. 28: 51-55.
- BEAMER, R. H. 1954. A revision of the genus *Achorotile* in America north of Mexico (Homoptera-Fulgoridae-Delphacinae). J. Kansas Entomol. Soc. 27: 143-147.
- BEAMER, R. H. 1955. A revision of the genus *Megamelus* in America north of Mexico (Homoptera-Fulgoridae-Delphacinae). J. Kansas Entomol. Soc. 28: 29-40, 41-46.
- BOURGOIN, T. 1990. Morphologie comparée des genitalia mâles des Trypetimorphini et remarques phylogénétiques (Hemiptera Fulgoromorpha). Ann. Soc. Entomol. Français (N. S.) 26: 555-564.
- BOURGOIN, T. 1993. Female genitalia in Hemiptera Fulgoromorpha, morphological and phylogenetic data. Ann. Soc. Entomol. Français (N. S.) 29: 225-244.
- BRAR, R. S., AND S. S. BAINS. 1979. Population dynamics of *Pyrilla perpusilla* Walker. Mortality factors. Indian J. Ecology 6: 110-121.
- BULL, R. M. 1972. A study of the sugar-cane leafhopper *Perkinsiella saccharicida* Kirk. (Homoptera, Delphacidae) in the Bundaberg District of southeastern Queensland. Proc. Queensland Soc. Sugar Cane Technol. 39: 173-183.
- CALDWELL, J. S., AND L. F. MARTORELL. 1951. Review of the auchenorrhynous Homoptera of Puerto Rico. Part II. The Fulgoroidea except Kinnaridae. J. Agric. Univ. Puerto Rico 34: 133-269.
- CARNEGIE, A. J. M. 1980. Egg mortality on *Numicia viridis* Muir (Homoptera: Tropiduchidae) in sugarcane and in indigenous hosts. J. Entomol. Soc. South Africa 43: 215-222.
- CHANG, V. C. S., AND A. K. OTA. 1978. Feeding activities of *Perkinsiella* leafhoppers and Fiji disease resistance of sugarcane. J. Econ. Entomol. 21: 297-300.
- CHATTERJEE, P. B. 1971. Occurrence of *Eoeurysa flavo-capitata* Muir (Fam. Delphacidae) on sugarcane in India. Indian J. Entomol. 33: 220.
- COLEY, J. D. 1970. The family Delphacidae (Homoptera) in Mississippi. Unpubl. MS Thesis, Mississippi State Univ., State College, Mississippi, 106 pp.
- CUMBER, R. A. 1952. Studies on *Oliarus atkinsoni* Myers (Hem. Cixiidae), vector of the "yellow-leaf" disease of *Phormium tenax* Forst. I. Habits and environment with a note on natural enemies. New Zealand J. Sci. Technol. 34(B): 92-98.
- CUMBER, R. A. 1953. Investigations into yellow-leaf disease of *Phormium*. IV. Experimental induction of yellow leaf condition in *Phormium tenax* Forst. by the insect vector *Oliarus atkinsoni* Myers (Hom., Cixiidae). New Zealand J. Sci. Technol. 34(A): 31-46.
- DOERING, K. C. 1938. A contribution to the taxonomy of the subfamily Issinae in America north of Mexico (Fulgoridae, Homoptera). Part II. Univ. Kansas Sci. Bull. 25: 447-575.
- DOERING, K. C. 1940. A contribution to the taxonomy of the subfamily Issinae in America north of Mexico (Fulgoridae, Homoptera). Part III. Univ. Kansas Sci. Bull. 26: 83-167.
- DOERING, K. C. 1941. A contribution to the taxonomy of the subfamily Issinae in America north of Mexico (Fulgoridae, Homoptera). Part IV. Univ. Kansas Sci. Bull. 27: 185-233.
- DUBOSE, W. P. 1960. The genus *Delphacodes* Fieber in North Carolina (Homoptera: Delphacidae). J. Elisha Mitchell Sci. Soc. 76: 36-63.
- EMELJANOV, A. F. 1999. Notes on delimitation of families of the Issidae group with description of a new species of Caliscelidae belonging to a new genus and tribe (Homoptera, Fulgoroidea). Zoosyst. Rossica 8: 61-72.
- FENNAH, R. G. 1945. The external male genitalia of Fulgoroidea (Homoptera). Proc. Entomol. Soc. Washington 47: 217-229.
- FENNAH, R. G. 1954. The higher classification of the family Issidae (Homoptera: Fulgoroidea) with descriptions of new species. Trans. Royal Entomol. Soc. London 105: 455-474.
- FENNAH, R. G. 1982. A tribal classification of the Tropiduchidae (Homoptera: Fulgoroidea), with the description of a new species on tea in Malaysia. Bull. Entomol. Res. 72: 631-643.
- FENNAH, R. G. 1984. Revisionary notes on the classification of the Nogodinidae (Homoptera, Fulgoroidea), with descriptions of a new genus and a new species. Entomol. Mon. Mag. 120: 81-86.
- FLETCHER, M. J. 1985. Revision of the genus *Siphanta* Stål (Homoptera: Fulgoroidea: Flatidae). Australian J. Zool. Suppl. Ser. No. 110: 1-94.
- FREUND, R., AND S. W. WILSON. 1995. The planthopper genus *Acanalonia* in the United States (Homoptera: Issidae): morphology of male and female external genitalia. Insecta Mundi 9: 195-215.
- GATINEAU, F., J. LARRUE, D. CLAIR, F. LORTON, M. RICHARD-MOLARD, AND E. BOUDON-PADIEU. 2001. A new natural planthopper vector of stolbur phyto-

- plasma in the genus *Pentastiridius* (Hemiptera: Cixiidae). European J. Plant Pathol. 107(3): 263-271.
- HEADY, S. E., AND S. W. WILSON. 1990. The planthopper genus *Prokelisia* (Homoptera: Delphacidae) in the United States: Descriptions of female genitalia and copulatory behavior. J. Kansas Entomol. Soc. 63: 267-278.
- HOCH, H., AND R. REMANE. 1985. Evolution und Speziation der Zikaden-Gattung *Hyalesthes* Signoret, 1865 (Homoptera Auchenorrhyncha Fulgoroidea Cixiidae). Marburger Entomol. Publ. 2(2): 1-427.
- HOLZINGER, W. E., I. KAMMERLANDER, AND H. NICKEL. 2003. The Auchenorrhyncha of Central Europe. Vol. 1: Fulgoromorpha, Cicadomorpha excl. Cicadellidae. Brill, Boston.
- HOWARD, F. W., R. C. NORRIS, AND D. L. THOMAS. 1983. Evidence of transmission of palm lethal yellowing agent by a planthopper, *Myndus crudus* (Homoptera: Cixiidae). Tropical Agric. 60: 168-171.
- HUSSAIN, A. A. 1963. Biology and control of the dubas bug, *Ommatissus binotatus lybicus* De Berg. (Homoptera, Tropiduchidae), infesting date palms in Iraq. Bull. Entomol. Res. 53: 737-745.
- INTERNATIONAL COMMITTEE ON TAXONOMY OF VIRUSES. Universal Virus Database (ICTVdB). www.ncbi.nlm.nih.gov/ICTVdb/index (accessed 29 Oct 2004).
- KLEIN, M., AND A. VENEZIAN. 1985. The dubas bug tropiduchid, *Ommatissus binotatus lybicus*, a threat to date palms in Israel. Phytoparasitica 13: 95-101.
- KRAMER, J. P. 1973. Revision of the American planthoppers of the genus *Stobaera* (Homoptera: Delphacidae) with new distributional data and host plant records. Proc. Entomol. Soc. Washington 75: 379-402.
- KRAMER, J. P. 1977. Taxonomic study of the planthopper genus *Oecleus* in the United States (Homoptera: Fulgoroidea: Cixiidae). Trans. American Entomol. Soc. 103: 379-449.
- KRAMER, J. P. 1979. Taxonomic study of the planthopper genus *Myndus* in the Americas (Homoptera: Fulgoroidea: Cixiidae). Trans. American Entomol. Soc. 105: 301-389.
- KRAMER, J. P. 1981. Taxonomic study of the planthopper genus *Cixius* in the United States and Mexico (Homoptera: Fulgoroidea: Cixiidae). Trans. American Entomol. Soc. 107: 1-68.
- KRAMER, J. P. 1983. Taxonomic study of the planthopper family Cixiidae in the United States (Homoptera: Fulgoroidea). Trans. American Entomol. Soc. 109: 1-58.
- LIANG, A.-P. 2001. Morphology of antennal sensilla in *Achilixius sandakanensis* Muir (Hemiptera: Fulgoromorpha: Achilixiidae) with comments on the phylogenetic position of the Achilixiidae. Raffles Bull. Zool. 49: 221-226.
- LIEFTING, L. W., R. E. BEEVER, C. J. WINKS, M. N. PEARSON, AND R. L. S. FORSTER. 1998. Planthopper transmission of *Phormium* yellow leaf phytoplasma. Australasian Plant Pathol. 26(3): 148-154.
- MAIXNER, M. 1994. Transmission of German grapevine yellows (Vergilbungskrankheit) by the planthopper *Hyalesthes obsoletus*. Vitis 3: 103-104.
- MCDERMOTT, B. T. 1952. A revision of the genus *Megamelanus* and its allies (Homoptera, Fulgoridae, Delphacinae). J. Kansas Entomol. Soc. 25: 41-49.
- MCPHERSON, K. R., AND S. W. WILSON. 1995. The planthopper genus *Phylloscelis* in the United States (Homoptera: Dictyopharidae). Insecta Mundi 9: 177-188.
- MEAD, F. W. 1965. *Ormenaria rufifascia* (Walker), a planthopper pest of palms (Homoptera: Flatidae). Florida Dep. Agric. Entomol. Circ. 37.
- MEAD, F. W., AND J. P. KRAMER. 1982. Taxonomic study of the planthopper genus *Oliarus* in the United States (Homoptera: Fulgoroidea: Cixiidae). Trans. American Entomol. Soc. 107: 381-569.
- METCALF, Z. P. 1923. A key to the Fulgoridae of eastern North America. J. Elisha Mitchell Sci. Soc. 38: 139-230.
- METCALF, Z. P. 1938. The Fulgorina of Barro Colorado and other parts of Panama. Bull. Mus. Comp. Zool. 82: 276-433, plates 1-23.
- METCALFE, J. R. 1969. Studies on the biology of the sugar cane pest *Saccharosydne saccharivora* (Westw.) (Hom., Delphacidae). Bull. Entomol. Res. 59: 393-408.
- MOCHIDA, O., AND T. OKADA. 1979. Taxonomy and biology of *Nilaparvata lugens* (Hom., Delphacidae) In O. Mochida and T. Okada [eds.], Brown Planthopper: Threat to Rice Production in Asia. Los Baños, Philippines.
- MORGAN, L. W., AND R. H. BEAMER. 1949. A revision of three genera of delphacine fulgorids from America north of Mexico (Homoptera-Fulgoroidea-Delphacinae). J. Kansas Entomol. Soc. 22: 97-120, 21-142.
- MPUNAMI, A. A., K. KULLAYA, AND J. MUGINI. 2002. The status of lethal yellowing-type diseases in east Africa, pp. 161-168 In Proc. Expert Consultation on Sustainable Coconut Production through Control of Lethal Yellowing Disease, Kingston, Jamaica. CFC/FAO.
- MUIR, F. 1926a. The morphology of the aedeagus in Delphacidae. Trans. Royal Entomol. Soc. London 2:371-380.
- MUIR, F. 1926b. Contributions to our knowledge of South American Fulgoroidea (Homoptera). Part I. The family Delphacidae. Bull. Hawaiian Sugar Planter's Assoc. Div. Entomol. 18: 2-38.
- MUIR, F., AND W. M. GIFFARD. 1924. Studies in North American Delphacidae. Bull. Hawaiian Sugar Planter's Assoc. Div. Entomol. 15: 1-53.
- NAULT, L. R. 1994. Transmission biology, vector specificity and evolution of planthopper transmitted plant viruses (Chapter 12), pp. 429-448 In R. F. Denno and T. J. Perfect [eds.], Planthoppers: Their Ecology and Management. Chapman and Hall, Inc., New York.
- O'BRIEN, L. B. 1967. *Caliscelis bonellii* (Latreille), a European genus of Issidae new to the United States (Homoptera: Fulgoroidea). Pan-Pacific Entomol. 43: 130-133.
- O'BRIEN, L. B. 1992. Tropiduchidae (Homoptera: Fulgoroidea) of the United States. Ann. Entomol. Soc. America 85:121-126.
- O'BRIEN, L. G., M. B. STOETZEL, AND D. R. MILLER. 1998. Order Homoptera, pp. 66-111 In F. W. Stehr [ed], Immature Insects. Vol. 2. Kendall Hunt Publ. Co. Dubuque, IA.
- O'BRIEN, L. R., AND S. W. WILSON. 1985. The systematics and morphology of planthoppers (Fulgoroidea), pp. 61-102 In L. Nault and R. Rodriguez [eds.], The Leafhoppers and Planthoppers. John Wiley & Sons, Inc., N.Y.
- OSSIANNILSSON, F. 1978. The Auchenorrhyncha (Homoptera) of Fennoscandia and Denmark. Part 1: Introduction, infraorder Fulgoromorpha. Fauna Entomol. Scandinavica 7: 5-222.
- PENNER, L. R. 1945. The genus *Laccocera* Van Duzee (Homoptera: Delphacidae). J. Kansas Entomol. Soc. 18: 30-47.

- PENNER, L. R. 1947. Some notes on the genus *Pentagramma* and four new species (Homoptera-Delphacidae-Asiracinae). *J. Kansas Entomol. Soc.* 20: 30-39.
- PENNER, L. R. 1952. A new *Bostaera* from Florida (Homoptera-Delphacidae). *J. Kansas Entomol. Soc.* 25: 1-3.
- PONNAMMA, K. N., AND B. BABJAN. 2000. Biocontrol agents associated with *Proutista moesta*, a vector of phytoplasma diseases of palms, with particular reference to *Halictophagus palmi*. XXI International Entomol. Congr., Iguassu, Brasil.
- RAHMAN, K. A., AND R. NATH. 1940. Bionomics and control of the Indian sugar-cane leafhopper, *Pyrilla perpusilla* Walk. (Rhynchota, Fulg.) in the Punjab. *Bull. Entomol. Res.* 31: 179-190.
- RAJAN, P., C. P. R. NAIR, J. J. SOLOMON, AND L. GEETHA. 2002. Identification of phytoplasma in the salivary gland of *Proutista moesta* (Westwood), a putative vector of coconut root (wilt) disease. *J. Plantation Crops* 30(2): 55-57.
- RANDLES, J. W., R. A. J. HODGSON, D. HANOLD, D. C. MILLER, J. P. MORIN, E. WEFELS, G. WALL, AND W. ROHDE. 2004. Diagnosis of virus, viroids and viroid-like sequences in coconut. [www.mpiz-koeln.mpg.de/~rohde/randles](http://www.mpiz-koeln.mpg.de/~rohde/randles) (accessed 29 Oct 2004).
- RUAN, Y. L., W. L. CHIANG, AND R. F. LIN. 1981. Studies on the rice virus vector small brown planthopper *Laodelphax striatella* Fallen. *Acta Entomol. Sinica* 24: 283-290.
- SFORZA, R., T. BOURGOIN, S. W. WILSON, AND E. Boudon-Padieu. 1999. Field observations, laboratory rearing, and descriptions of immatures of *Hyalesthes obsoletus* (Hemiptera: Cixiidae). *European J. Entomol.* 96: 409-418.
- SELJAK, G. 2004. *Prokelisia marginata* (Van Duzee, 1897)—a Nearctic planthopper new to Slovenia and Europe (Auchenorrhyncha: Delphacidae). *Acta Entomologica Slovenica* 12: 260-263.
- SHEPARD, C. L. 1939. The genus *Ormenis* (Fulgoroidea) in North America north of Mexico. Unpubl. M. A. Thesis, Univ. Kansas. 60 pp. 8 plates. [keys to species now placed in *Anormenis*, *Metcalfa*, *Ormenaria*, *Ormenis* (s. l.), and *Ormenoides*]
- SOSA, O., JR. 1985. The sugarcane delphacid, *Perkinsiella saccharicida* (Homoptera: Delphacidae), a sugarcane pest new to North America detected in Florida. *Florida Entomol.* 68: 357-358.
- STICKNEY, F. S., D. F. BARNES, AND P. SIMMONS. 1950. Date palm insects in the United States. United States Dep. Agric. Circ. 846: 1-57.
- TSAL, J. H., AND S. W. WILSON. 1986. Biology of *Peregrinus maidis* with descriptions of immature stages. *Ann. Entomol. Soc. America* 79: 395-401.
- WEAVING, A. J. S. 1980. Observations on *Hilda patruelis* Stål (Homoptera: Tettigometridae) and its infestation of the groundnut crop in Rhodesia. *J. Entomol. Soc. South Africa* 43: 151-167.
- WEBER, B. C., AND S. W. WILSON. 1981. Seasonal and vertical distributions of planthoppers (Homoptera: Fulgoroidea) within a black walnut plantation. *Great Lakes Entomol.* 14: 71-75.
- WHEELER, JR., A. G. 2003. Bryophagy in the Auchenorrhyncha: seasonal history and habits of a moss specialist, *Javesella opaca* (Beamer) (Fulgoroidea: Delphacidae). *Proc. Entomol. Soc. Washington* 105: 599-610.
- WILSON, M. R., AND M. R. CLARIDGE. 1991. Handbook for the Identification of Leafhoppers and Planthoppers of Rice. CAB International, Oxon, UK. 142 pp.
- WILSON, S. W. 1982. The planthopper genus *Prokelisia* in the United States (Homoptera: Fulgoroidea: Delphacidae). *J. Kansas Entomol. Soc.* 55: 532-546.
- WILSON, S. W., R. F. DENNO, C. MITTER, AND M. R. WILSON. 1994. Evolutionary patterns of host plant use by delphacid planthoppers and their relatives (Chapter 1), pp. 7-113 *In* R. F. Denno and T. J. Perfect [eds.]. *Planthoppers: Their Ecology and Management*. Chapman and Hall, Inc., New York.
- WILSON, S. W., AND A. LUCCHI. 2001. Distribution and ecology of *Metcalfa pruinosa* and associated planthoppers in North America (Homoptera: Fulgoroidea). *Atti dell' Acad. Naz. Italiana di Entomol. Rendiconti* 49: 121-130.
- WILSON, S. W., AND J. E. MCPHERSON. 1980a. The distribution of the Fulgoroidea of the eastern United States (Homoptera). *Trans. Illinois Acad. Sci.* 73(4): 7-20.
- WILSON, S. W., AND J. E. MCPHERSON. 1980b. Mixed species feeding assemblages of planthopper nymphs (Homoptera: Fulgoroidea). *Great Lakes Entomol.* 13: 185-187.
- WILSON, S. W., AND J. E. MCPHERSON. 1980c. Keys to the planthoppers, or Fulgoroidea, of Illinois (Homoptera). *Trans. Illinois Acad. Sci.* 73(2): 1-61.
- WILSON, S. W., AND J. E. MCPHERSON. 1981a. Life histories of *Acanalonia bivittata* and *A. conica* with descriptions of immature stages. *Ann. Entomol. Soc. Amer.* 74: 289-298.
- WILSON, S. W., AND J. E. MCPHERSON. 1981b. Life histories of *Anormenis septentrionalis*, *Metcalfa pruinosa*, and *Ormenoides venusta* with descriptions of immature stages. *Ann. Entomol. Soc. Amer.* 74: 299-311.
- WILSON, S. W., AND L. B. O'BRIEN. 1987. A survey of planthopper pests of economically important plants (Homoptera: Fulgoroidea). *Proc. 2nd Inter. Workshop on Leafhoppers and Planthoppers of Economic Importance*. Commonwealth Inst. Entomol., London, pp. 343-360.
- WILSON, S. W., J. L. SMITH, AND A. H. PURCELL III. 1993. An inexpensive vacuum collector for insect sampling. *Entomol. News* 104: 203-208.
- WILSON, S. W., AND J. H. TSAI. 1982. Descriptions of the immature stages of *Myndus crudus* (Homoptera: Fulgoroidea: Cixiidae). *J. New York Entomol. Soc.* 90: 166-175.
- WILSON, S. W., AND J. H. TSAI. 1984. *Ormenaria rufifascia* (Homoptera: Fulgoroidea: Flatidae): Descriptions of nymphal instars and notes on biology. *J. New York Entomol. Soc.* 92: 307-315.
- WILSON, S. W., AND J. H. TSAI. 1992. Descriptions of immatures of *Eoeurysa flavocapitata* (Homoptera: Delphacidae). *Pan-Pacific Entomol.* 68: 133-139.
- WOOTEN, C. M., S. W. WILSON, AND J. H. TSAI. 1993. Descriptions of nymphs of *Harmalia anacharsis* Fennah, a species new to the United States (Homoptera: Delphacidae). *J. New York Entomol. Soc.* 101: 567-573.
- YANG, C.-T., AND T.-Y. CHANG. 2000. The External Male Genitalia of Hemiptera (Homoptera-Heteroptera). Shih Way Publ. Taichung, Taiwan. 746 pp.
- ZIMMERMAN, E. C. 1948. Homoptera: Auchenorrhyncha. *Insects of Hawaii* 4: 1-268.