

## **JACKFRUIT, ARTOCARPUS HETEROPHYLUS, IS NOT A HOST OF DIAPHORINA CITRI (HOMOPTERA: PSYLLIDAE) IN FLORIDA**

Authors: Peña, J. E., Mannion, C. M., Ulmer, B. J., and Halbert, S. E.

Source: Florida Entomologist, 89(3) : 412-413

Published By: Florida Entomological Society

URL: [https://doi.org/10.1653/0015-4040\(2006\)89\[412:JAHINA\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2006)89[412:JAHINA]2.0.CO;2)

---

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## JACKFRUIT, *ARTOCARPUS HETEROPHYLUS*, IS NOT A HOST OF *DIAPHORINA CITRI* (HOMOPTERA: PSYLLIDAE) IN FLORIDA

J. E. PEÑA<sup>1</sup>, C. M. MANNION<sup>1</sup>, B. J. ULMER<sup>1</sup> AND S. E. HALBERT<sup>2</sup>

<sup>1</sup>University of Florida, IFAS, Tropical Research and Education Center, 18905 SW 280<sup>th</sup> Street, Homestead, FL 33031

<sup>2</sup>Florida Department of Agriculture and Consumer Services, Division of Plant Industry, P.O. Box 147100, Gainesville, FL 32611-7100

The Asian citrus psyllid, *Diaphorina citri* Kuwayama 1907 (Rhynchoeae: Psyllidae), was discovered in south Florida in 1998 and is now established in the state (Halbert & Manjunath 2005). It is considered one of the most serious pests of citrus when the pathogens that cause citrus greening, which can vector, are also present (Halbert & Manjunath 2005). *Diaphorina citri* is an oligophagous species with a host range within genera in the family Rutaceae, e.g., *Aegle*, *Aeglopsis*, *Afraegle*, *Atalantia*, *Balsamocitrus*, *Citropsis*, *Citrus*, *Clausena*, *Murraya*, *Fortunella*, *Linonia*, *Merrillia*, *Microcitrus*, *Pamburus*, *Poncirus*, *Severinia*, and *Swinglea* (Viraktamath & Bhuvanavar 2002; Tirtawidjaja 1981; Koizumi et al. 1996; Chavan & Summanwar 1993; Aubert 1990a,b; Lim et al. 1990; Garnier & Bové 1993; Halbert & Manjunath 2005). In addition to these host plant associations, Shivankar et al. (2000) reported *D. citri* on jackfruit, *Artocarpus heterophyllus* Lamarek, which is in the family Moraceae (Crane et al. 2002). Because there have been no other reports of *D. citri* infesting *A. heterophyllus* in Florida, nor anywhere *D. citri* has been observed, we investigated *A. heterophyllus* as a host plant of *D. citri*.

Field surveys were initiated in the fall of 2005. On Sep 30, 2005 samples of *A. heterophyllus* and orange jasmine, *Murraya paniculata*, a known host of *D. citri*, were collected in Miami-Dade County. Ten 10-cm shoots were collected from each of two jackfruit trees located approximately 28 m from a hedge of orange jasmine. Twenty 10-cm shoots also were collected from the orange jasmine hedge. Shoots were placed in individual bags and examined under a microscope to determine the numbers of eggs, nymphs, and adults per sample.

The mean numbers of *D. citri* collected from *Murraya* shoots were  $8.85 \pm 2.24$  ( $n = 20$ ). An average  $3.45 \pm 2$  *Toxoptera aurantii* (Boyer de Fonscolombe) (Homoptera: Aphididae) were also recorded from *Murraya* shoots. No *D. citri* were found on the shoots collected from jackfruit; however, 10 nymphs of *T. aurantii* were found on the jackfruit terminals. Means for other arthropods recorded from jackfruit were  $0.5 \pm 0.22$  *Anascirtothrips arorai* Bhatti (Thysanoptera: Thripidae),  $0.05 \pm 0.05$  *T. aurantii*,  $51.35 \pm 0.22$  *Ditrymaceus*

*integrifoliae* Mohanasundaran (Acari: Eriophyidae), and  $0.15 \pm 0.15$  unidentified leafhoppers (Homoptera: Cicadellidae).

On Dec 12, 2005, five 10-cm shoots ( $n = 55$ ) were collected from 11 randomly selected jackfruit trees from the National Germplasm Repository Collection at the USDA, ARS, SHRS in Miami, FL. No other common hosts of *D. citri* were located within a 100-m radius from the site; therefore, no samples were collected from known hosts on that date. Samples were inspected as mentioned above. No *D. citri* were recorded from 55 shoots of jackfruit; the mean numbers of other arthropods per shoot were  $0.4 \pm 0.17$  *A. arorai*,  $2.46 \pm 1.90$  *D. integrifoliae*,  $0.04 \pm 0.04$  *T. aurantii*, and  $0.02 \pm 0.02$  unidentified Diaspididae.

On Feb 22, 2006, 15 shoots of jackfruit were randomly collected from the Fruit and Spice Park, Homestead, FL. Citrus species were located ca. 30 m from the jackfruit trees but no samples were collected from the citrus spp. No *D. citri* were collected from jackfruit; but  $0.13 \pm 0.09$  *A. arorai* and  $29.86 \pm 15.85$  *D. integrifoliae* were collected from the terminals.

In addition to the field surveys, no-choice trials were conducted in the greenhouse. Four 0.8-m potted jackfruit trees with suitable new growth were introduced each into individual nylon mesh screen cages ( $91 \times 91 \times 122$  cm), each supported on a PVC frame. Depending on the date, ten to 20 adults of *D. citri* were introduced into each cage on Oct 5, 13, 19, Nov 1, 29 and Dec 22, 2005. *D. citri* densities were recorded 7 d after the introduction of each psyllid cohort. The jackfruit trees were inspected for *D. citri* eggs, nymphs and adults with aid of a 10 $\times$  hands lens 7 d after the introduction of each cohort. No psyllid eggs, nymphs or adults were collected on Oct 11 (7-10 shoots/plant), Oct 18 (6-20 shoots/plant), Oct 26 (6-20 shoots/plant), Nov 9 (7-14 shoots/plant), Dec 5 (9-17 shoots/plant), or Dec 30 (6-11 shoots/plant).

On Feb 13, 2006, 4 potted jackfruit trees were placed in close contact with a *D. citri* infested hedge of *M. paniculata*. The shoots of each jackfruit plant were inspected with a 10 $\times$  hand lens on Feb 16, 20, 22, 2006. One adult psyllid was observing resting, but not feeding on, a shoot on Feb 16. No eggs or nymphs of *D. citri* were observed

from 6 - 13 inspected shoots per plant. On Feb 20, 1 second to third instar *D. citri* nymph was observed crawling on one shoot, from a range of 9-14 shoots inspected per plant. No adults or eggs were recorded. On Feb 22, 2006, no adults, nymphs, or eggs of *D. citri* were observed from these plants.

Based on the field collections of several varieties of jackfruit, our no-choice greenhouse experiment, and our field two-choice experiment, it does not appear that *A. heterophyllus* is an acceptable host plant for *D. citri*. The present study refutes the report by Shivankar et al. (2000) listing jackfruit as a host of *D. citri*.

#### SUMMARY

The status of jackfruit, *A. heterophyllus* as a host plant of *D. citri* was investigated. Field surveys, a no-choice greenhouse test, and a field study showed that jackfruit is not an acceptable host for *D. citri*.

#### REFERENCES CITED

- AUBERT, B. 1990a. Integrated activities for the control of hunaglungbin-greening and its vector *Diaphorina citri* Kuwayama in Asia, pp. 133-144 *In* B. Aubert, S. Tontyaporn, and D. Buangsuwon [eds.], Rehabilitation of Citrus Industry in the Asia Pacific Region. Proc. Asia Pacific International Conference on Citriculture, Chiang Mai, Thailand, 4-10 February 1990. UNDP-FAO, Rome.
- AUBERT, B. 1990b. High density planting (HDP) of Jiaogan mandarin in the lowland area of Shantou (Guandong China) and implications for greening control, pp. 149-157 *In* B. Aubert, S. Tontyaporn, and D. Buangsuwon [eds.], Rehabilitation of Citrus Industry in the Asia Pacific Region. Proc. Asia Pacific International Conference on Citriculture, Chiang Mai, Thailand, 4-10 February 1990. UNDP-FAO, Rome.
- CHAVAN, V. M., AND A. S. SUMMANWAR. 1993. Population dynamics and aspects of the biology of citrus psylla, *Diaphorina citri* Kuw., in Maharashtra, pp. 286-290 *In* P. Moreno, J. V. da Graca, and L. W. Timmer [eds.], Proc. 12<sup>th</sup> Conference of the International Organization of Citrus Virologists, University of California, Riverside.
- CRANE, J., C. F. BALERDI, AND R. CAMPBELL. 2002. The Jackfruit (*Artocarpus heterophyllus* Lam.) in Florida. University of Florida, FL Coop. Ext. Service, Inst. Food Agr. Sci, HS-882. 8 p.
- GARNIER, S. M., AND J. M. BOVÉ. 1993. Production of monoclonal antibodies recognizing most Asian strains of the greening BLO by vitro immunization with an Antigenic protein purified from the BLO, pp. 244-249 *In* P. Moreno, J. V. da Graca, and L. W. Timmer [eds.], Proc. 12<sup>th</sup> Conference of the International Organization of Citrus Virologists, University of California, Riverside.
- HALBERT, S., AND K. MANJUNATH. 2005. Asian citrus psyllids (Sternorrhyncha: Psyllidae) and greening disease of citrus: a literature review and assessment of risk in Florida. Florida Entomol. 87: 330-353.
- KOIZUMI, M., M. PROMMINTARA, AND Y. OHTSU. 1996. Wood apple, *Limonia acidissima*: A new host for the huanglongbing (greening) vector, *Diaphorina citri*, pp. 271-275 *In* J. V. da Graca, P. Moreno, and R. K. Yokomi [eds.], Proc. 13<sup>th</sup> Conference of the International Organization of Citrus Virologists (IOBC). University of California, Riverside.
- LIM, W. H., O. SHAMUDIN, AND W. KO. 1990. Citrus greening disease in Malaysia, pp. 100-105 *In* B. Aubert, S. Tontyaporn, and D. Buangsuwon [eds.], Rehabilitation of Citrus Industry in the Asia Pacific Region. Proc. Asia Pacific International Conference on Citriculture, Chiang Mai, Thailand, 4-10 February 1990. UNDP-FAO, Rome.
- SHIVANKAR, V. J., C. N. RAO, AND S. SINGH. 2000. Studies on citrus Psylla, *Diaphorina citri* Kuwayama: A review. Agricultural Reviews (Karnal, India) 21: 100-204.
- TIRTAWIDJAJA, S. 1981. Insect, dodder and seed transmissions of citrus vein phloem Degeneration (CVPD). Proc. International Soc. Citriculture 1: 469-471.
- VIRAKTAMATH, C., AND B. BHUMANAVAR. 2002. Biology and ecology and management of *Diaphorina citri* Kuwayama (Hemiptera: Psyllidae): Pest management in Horticultural Ecosystems. 7:1-27.