



Survey of the Termites (Isoptera: Kalotermitidae, Rhinotermitidae, Termitidae) of Lanyu Island, Taiwan

Authors: Li, Hou-Feng, Scheffrahn, Rudolf H., Su, Nan-Yao, Kanzaki, Natsumi, and Yang, Rou-Ling

Source: Florida Entomologist, 91(3) : 472-473

Published By: Florida Entomological Society

URL: [https://doi.org/10.1653/0015-4040\(2008\)91\[472:SOTTIK\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2008)91[472:SOTTIK]2.0.CO;2)

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

SURVEY OF THE TERMITES (ISOPTERA: KALOTERMITIDAE, RHINOTERMITIDAE, TERMITIDAE) OF LANYU ISLAND, TAIWAN

HOU-FENG LI, RUDOLF H. SCHEFFRAHN, NAN-YAO SU, NATSUMI KANZAKI AND ROU-LING YANG
 Department of Entomology and Nematology, Fort Lauderdale Research and Education Center,
 University of Florida, Institute of Food and Agricultural Sciences,
 3205 College Avenue, Fort Lauderdale, FL 33314, U.S.A.

Lanyu Island (= Botel Tobago Island; Orchid Island) is a 48-km² volcanic island, ~90 km south-east of Taiwan Island. Oshima (1912) and Tu (1955) conducted a termite survey on Lanyu Island, and recorded 6 species (3 families, 6 genera) in total, but the voucher specimens were neither mentioned nor available. During 10-12 Jun 2007, we collected 24 termite colony samples from 11 localities along roadsides, in structures, and on boats on the island (Fig. 1). Our survey also yielded 6 species in 6 genera and 3 families (Table 1). Identifications were made by Li based on original descriptions and references listed in Table 1. The voucher specimens are deposited in the University of Florida Termite Collection, Fort Lauderdale Research and Education Center, and National Taiwan University, Department of Entomology, Taipei, Taiwan. The 6 species with brief description are the following.

Cryptotermes domesticus (Haviland): Alates were taken from a sport-fishing yacht, and 4 other colonies were found in the dead branches of living trees. This distribution pattern indicates that *Cr. domesticus* is a structural pest and also a woodland species of Lanyu Island. In Taiwan *Cr. domesticus* was only found in structural wood (Li, unpublished data). Miller & Paton (1983) reported that *Cr. domesticus* is indigenous and only found in mangroves in northern Australia near Darwin, but is a domestic pest in northern Queensland. We assume that Lanyu Island and North Australia are the northern and southern limits of

the indigenous range of *Cr. domesticus*. *Cryptotermes kotoensis* (Oshima) collected in Lanyu Island by Oshima (1912) is a junior synonym of *Cr. domesticus* (Chung & Chen 1994; Huang 2000).

Neotermes koshunensis (Shiraki): *Neotermes koshunensis*, a common species in Lanyu Island, is usually found in dead trees or dead branches of living trees with high moisture and is not known to be a structural pest.

Coptotermes formosanus Shiraki: *Coptotermes formosanus* is the major structural pest on Lanyu Island and found in houses, discarded wood material, wood retaining walls, fence posts, and boats. The yacht infested by *Cr. domesticus* also was infested by *Co. formosanus*, and the latter obtained fresh water by building shelter tubes to the bilge tank. We speculate that *Co. formosanus* is an anthropogenically dispersed species from Taiwan Island due to frequent shipping.

Prorhinotermes japonicus (Holmgren): *Prorhinotermes japonicus* was collected by Hozawa in 1911 and is known only from Lanyu Island. Holmgren (1912) described and named this species as *japonicus* due to the fact that Taiwan and its surrounding islands belonged to Japan between 1895 and 1945. In this survey, we collected only 1 sample in a central dead part of a living *Casuarina* sp. tree, along with 2 other termite species, *Co. formosanus* and *Nasutitermes takasagoensis* (Shiraki).

Reticulitermes flaviceps (Oshima): Oshima (1912) collected a *Reticulitermes* species on Lanyu Island and identified it as *Reticulitermes flaviceps* from specimens collected in Taipei, Taiwan (Oshima 1911). In the termite list of Taiwan and Lanyu Islands, *Reticulitermes speratus* (Kolbe) is listed instead of *R. flaviceps* (Tu 1955). Morimoto (1968) suggested that the *Reticulitermes* sp. collected at Lanyu Island is closer morphologically to *R. fukiensis* Light or *R. speratus* rather than *R. flaviceps* collected in Taipei. Recently, *R. speratus* from Japan, *R. flavipes* from U.S.A., and *R. flaviceps* from Taiwan Island have been supported as 3 valid species based on DNA sequences (Yashiro & Matsuura 2007). However, the taxonomic status of *Reticulitermes* sp. on Lanyu Island has not been validated. Chung & Chen (1994) also pointed out the uncertain identification of *Reticulitermes* sp. on Lanyu Island. In this survey, samples of *Reticulitermes* sp. from 2 colonies were collected at the interface between wet soil and fallen dead trees, and identified as *R. flaviceps* based upon soldier morphology. This result was confirmed by the nucleotide sequences of 3 mitochondrial genes, cytochrome oxidase subunit




Fig. 1. Collection sites (dark circles) represented in the current survey.

TABLE 1. TERMITE SPECIES RECORDS FROM LANYU ISLAND LISTED ALPHABETICALLY BY FAMILY.

Taxon	Collected location ^a	Original species description	Previous locality reference
Kalotermitidae			
<i>Cryptotermes domesticus</i> (Haviland)	A, B, F, J	Haviland (1898)	Oshima (1912), Tu (1955)
<i>Neotermes kosshunensis</i> (Shiraki)	B, D, I	Shiraki (1909)	Oshima (1912), Tu (1955)
Rhinotermitidae			
<i>Coptotermes formosanus</i> Shiraki	C, E, F, H, K	Shiraki (1909)	Oshima (1912), Tu (1955)
<i>Prorhinotermes japonicus</i> (Holmgren)	K	Holmgren (1912)	Lo et al. (2004), Oshima (1912), Tu (1955)
<i>Reticulitermes flaviceps</i> (Oshima)	A, B	Oshima (1911)	Oshima (1912), Tu (1955)
Termitidae			
<i>Nasutitermes takasagoensis</i> (Shiraki)	B, D, E, G, K	Nawa (1911)	Oshima (1912), Tsai (2003), Tu (1955)

^aLocation codes are indicated in Fig. 1.

II (COII), ribosomal RNA small subunit 12S and ribosomal RNA large subunit 16S, and a nuclear gene, Internal Transcribed Spacer (ITS) region. The sequences determined in the present study were deposited in the GenBank database with accession numbers EU627778-EU627785.

Nasutitermes takasagoensis (Shiraki): *Nasutitermes takasagoensis* was the most dominant termite species on Lanyu Island. This species builds narrow foraging tubes and sheeting on wood poles, fence posts, discarded construction material, and trees. *Odontotermes formosanus* (Shiraki) common on Taiwan Island builds similar shelters, but is not found in Lanyu Island.

We thank Wen-Jer Wu (Dept. of Entomology), Ai-Chi Lin (Dept. of Library and Information Science), and Hsiu-Chuan Liang (Dept. of Pharmacology) from National Taiwan University for providing literature references. This study was supported in part by a grant from USDA-ARS under the grant agreement No. 58-6435-2-0075.

SUMMARY

A field survey of the termites of Lanyu Island resulted in confirmation of the island's fauna to 6 species in 6 genera and 3 families. The voucher specimens are maintained at the University of Florida, Fort Lauderdale Research and Education Center, Ft. Lauderdale, FL, U.S.A., and National Taiwan University, Department of Entomology, Taipei, Taiwan.

REFERENCES CITED

CHUNG, C. H., AND C. S. CHEN. 1994. A review of Taiwanese termites (Insecta, Isoptera) with keys to

adults and soldiers. *Yushania* 11: 193-203 (In Chinese with English abstract).

HAVILAND, G. D. 1898. Observations on termites, with descriptions of new species. *Linn. Soc. J. Zool.* 26: 358-442.

HOLMGREN, N. 1912. Die Termiten Japans. *Annot. Zool. Japan.* 8: 107-136 (In German).

HUANG, F. 2000. *Cryptotermes domesticus*, pp. 170-172. In F. Huang, S. Zhu, Z. Ping, X. He, G. Li, and D. Gao [eds.], *Fauna Sinica, Insecta vol. 17: Isoptera*. Science Press, Beijing, China (In Chinese).

LO, N., O. KITADE, T. MIURA, R. CONSTANTINO, AND T. MATSUMOTO. 2004. Molecular phylogeny of the Rhinotermitidae. *Insect. Soc.* 51: 365-371.

MILLER, L. R., AND R. PATON. 1983. *Cryptotermes* in mangroves in the northern territory (Isoptera: Kalotermitidae). *J. Australian Entomol. Soc.* 22: 189-190.

MORIMOTO, K. 1968. Termites of the genus *Reticulitermes* of Japan and Taiwan. *Bull. Govt. Forest Exp. Sta.* 217: 43-73.

NAWA, U. 1911. Two termite species from Taiwan. *Insect World* 15: 413-417 (In Japanese).

OSHIMA, M. 1911. Discrimination between *Termes flaviceps* Oshima and *Termes speratus* Kolbe, and several remarks on the scientific names of Japanese termites. *Insect World* 15: 355-363 (In Japanese).

OSHIMA, M. 1912. The taxonomy and distribution of termites in Taiwan, pp. 54-94. In M. Oshima [ed.], *The Third Official Report on Termites, Taiwan Sotokufu, Taihoku, Japan* (In Japanese).

SHIRAKI, T. 1909. Japanese termites. *Trans. Entomol. Soc. Japan* 2: 229-242 (In Japanese).

TSAI, C.-C. 2003. A taxonomic study of termite (Isoptera) from Taiwan. Ph. D. dissertation. Tunghai University, Taichung, Taiwan.

TU, T. 1955. The termites of Formosan. *Formos. Sci.* 9: 80-87.

YASHIRO, T., AND K. MATSUURA. 2007. Distribution and phylogenetic analysis of termite egg-mimicking fungi "termite balls" in *Reticulitermes* termites. *Ann. Entomol. Soc. America* 100: 532-538.