



**Steinernema diaprepesi (Nematoda: Steinernematidae):
Its Occurrence in Western Mexico and Susceptibility of
Engorged Cattle Ticks Boophilus microplus (Acari:
Ixodidae)**

Authors: Molina-Ochoa, Jaime, Nguyen, Khuong B., González-Ramírez, Martín, Quintana-Moreno, Miguel Guadalupe, Lezama-Gutiérrez, Roberto, et al.

Source: Florida Entomologist, 92(4) : 661-663

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.092.0423>

BioOne Complete (complete.bioone.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

STEINERNEMA DIAPREPESI (NEMATODA: STEINERNEMATIDAE): ITS OCCURRENCE IN WESTERN MEXICO AND SUSCEPTIBILITY OF ENGORGED CATTLE TICKS *BOOPHILUS MICROPLUS* (ACARI: IXODIDAE)

JAIME MOLINA-OCHOA^{1,4}, KHUONG B. NGUYEN², MARTÍN GONZÁLEZ-RAMÍREZ¹,

MIGUEL GUADALUPE QUINTANA-MORENO³, ROBERTO LEZAMA-GUTIÉRREZ¹ AND JOHN E. FOSTER⁴

¹Universidad de Colima, Facultad de Ciencias Biológicas y Agropecuarias, Laboratorio de Nematología Entomopatológica, Km. 40 autopista Colima-Manzanillo, Apartado Postal 36, Tecmán, Colima 28930, México
E-mail: jmolina@ucol.mx, jmolina18@hotmail.com, mglezr@ucol.mx, rlezama32@hotmail.com

²University of Florida Gainesville, Department of Entomology and Nematology, Gainesville, Florida 32611-0620, USA
E-mail: kbn@ufl.edu

³Universidad de Colima, Facultad de Medicina Veterinaria y Zootecnia, Apartado Postal 36, Km. 40 autopista Colima-Manzanillo, Apartado Postal 36, Tecmán, Colima 28930, México
E-mail: quintana_moreno@hotmail.com

⁴University of Nebraska Lincoln, Department of Entomology, 312F Plant Industry Building, Lincoln, NE 68583-0816, USA
E-mail: jfoster1@unl.edu

Entomopathogenic nematodes (EPN) belong to the families Steinernematidae and Heterorhabditidae. They infect a wide range of insect hosts and are used to control numerous species of soil borne insect pests (Grewal 2002). We identified EPN genera based on the color of cadavers of the wax moth, *Galleria mellonella* L., in which *Heterorhabditis* and *Steinernema*, produce red and tan colors, respectively, according to Woodring & Kaya (1988). We obtained an isolate of an EPN from sandy loam soil with the *Galleria* technique (Bedding & Akhurst, 1975) from grasslands in the Campus Tecmán of the Universidad de Colima, and we designated it as JMO94. In 2008, the morphometry and molecular characterization of JMO94 was conducted at the University of Florida Gainesville, Department of Entomology and Nematology under the supervision of Dr. K. B. Nguyen, and our JMO94 isolate was identified as *Steinernema diaprepesi* by morphological, morphometrical, and molecular results. Our strain is now labeled as *S. diaprepesi* Colimense strain, and represents a new record of occurrence of *S. diaprepesi* outside of the United States. DNA extraction, PCR amplification, sequencing, multiple alignments and other processes were conducted by methodology reported by Nguyen et al. (2001, 2004, 2006). PCR amplification was conducted by methodology reported by Nguyen et al. (2004) with the following exceptions: for ITS regions, the 2 internal primers for ITS regions suggested by Nguyen et al. (2001): KN58 = 5'-GTATGTTTGGTTGAAG-GTC-3' and KNRV = 5'-CACGCTCATACAACT-GCTC-3', were used; and for region D2/D3 regions, the primers D2F, 5'-CCTTAGTAACGGC-GAGTAAA-3' (forward) and 536, 5'-CAGC-

TATCCTGAGGAAAC-3' (reverse), were used as external primers to amplify the D2/D3 regions; primers 502, 5'-CAAGTACCGTGAGGGAAAGT-TGC-3' (forward) and 503, 5'-CCTTGGTCCGT-GTTTCAAGACG-3' (reverse), were used as internal primers for sequencing. Molecular phylogenetic relationships were obtained by maximum parsimony (MP) with PAUP, 4.0b8 (Swofford 2002). For phylogenetic analysis of the ITS regions, *Steinernema intermedium* was treated as the outgroup taxon for resolving relationships among the rest of *Steinernema* species (Nguyen et al. 2001). For D2/D3 regions, *Panagrellus redivivus* was used as the outgroup taxon (Stock et al. 2001). Branch support was estimated by bootstrap analysis (1000 replicates) based on the same parameters as the original search.

Steinernema diaprepesi has been isolated from citrus groves in Polk County, Florida, naturally attacking the citrus root weevil, *Diaprepes abbreviatus* L. (Coleoptera: Curculionidae) (Nguyen & Duncan 2002), and has been isolated from other orchards located on the central ridge of Florida, which is characterized by deep sandy soils similar to our soils in the Agricultural Experimental Station in the Tecmán Campus.

We used the Petri dish bioassay procedure (Freitas et al. 2005; Reis-Menini et al. 2008) to evaluate the susceptibility of engorged ticks (*Boophilus=Rhipicephalus microplus* Canestrini) to concentrations of 0, 500, 1000, 2000, 4,000, 8,000, and 16,000 infective juveniles (IJ) of *S. diaprepesi* Colimense strain applied in 1 mL of sterile distilled water dispensed on the surface of a couple of moistened filter papers (Whatman No. 1) in Petri

dishes. For each treatment concentration, 10 ticks were placed in the Petri dishes (60 × 15 mm) and each treatment was replicated 4 times.

Tick mortality was recorded every 24 h for 9 d (de Oliveira-Vasconcelos et al. 2004). Dead ticks were incubated and later examined to confirm the presence of nematodes. To compare tick mortality among different nematode concentrations, analysis of variance was used. Angular transformation was performed on the percent of mortality by square-root transformation before the analysis. Means were separated by the Tukey test ($P = 0.05$) (SAS Institute 1985).

The ANOVA showed statistical differences for all the sources of variance including the interaction concentrations × times ($P < 0.001$). The highest mortality (approx. 40%) was obtained with the concentration of 2000 IJ after 9 d post-exposure (Quintana-Moreno et al. 2007).

We observed a positive correlation between mortality of engorged ticks and the concentration of EPN and exposure time. Similar results were reported by Freitas-Rivero et al. (2005) with *Boophilus microplus*, and Samish et al. (1999) with other species of ticks. Because the cattle ticks are not specific hosts of *Steinernema diaprepesi*, we speculate that this is one of the reasons for the low cumulative mortality exhibited in our Petri dish experiment. Another possible reason is the body dimensions of the IJs; *S. diaprepesi* has a long body length of 1002 μm and a width of 34 μm. These dimensions could reduce the opportunities to penetrate the natural openings of the engorged ticks.

The potential of our strain could be determined in the future with other insect pest species in western Mexico such as the agave weevil, *Scyphophorus acupunctatus* Gyllenhal (Coleoptera: Curculionidae), and the coconut weevil, *Rhynchophorus palmarum* (L.) (Coleoptera: Curculionidae).

The authors thank MSc. Miguel Ángel Aguayo-López, Rector of the Universidad de Colima, and Secretaría de Educación Pública-PROMEP-PIFI 3.3 for supporting the senior author with grants to conduct research in Mexico and the University of Florida, Gainesville, and Dr. John J. Hamm (United States Department of Agriculture, Agricultural Research Service, Crop Protection and Management Research Unit, 2747 Davis Road, Tifton, GA 31793-0748, retired) for critical review of the manuscript.

SUMMARY

A steinernematid nematode was isolated from soil of grasslands in Tecomán Colima, Mexico; it was initially designated as JMO94, and determined to be *Steinernema diaprepesi* Colimense strain based upon morphometrics and molecular data; the susceptibility of cattle engorged ticks,

Boophilus microplus, to the nematode strain is reported. The highest tick cumulative mortality was approximately 40% after 9 d post-exposure to the nematode.

REFERENCES CITED

- BEDDING, R. A., AND AKHURST, R. J. 1975. A simple technique for the detection of insect parasitic nematodes in soil. *Nematologica* 21: 109-110.
- DE OLIVEIRA-VASCONCELOS, V., FURLONG, J., MARQUES DE FREITAS, G., DOLINSKI, C., MENDONA-AGUILERA, M., DEVITTE-RODRIGUES, R. C., AND PRATA, M. 2004. *Steinernema glaseri* Santa Rosa strain (Rhabditida: Steinernematidae) and *Heterorhabditis bacteriophora* CCA strain (Rhabditida: Heterorhabditidae) as biological control agents of *Boophilus microplus* (Acari: Ixodidae). *Parasitol. Res.* 94: 201-206.
- FREITAS-RIVERO, G. M., FURLONG, J., OLIVEIRA-VASCONCELOS, V., DOLINSKI, C., AND LOURENS-RIVERO, A. 2005. Analysis of biological parameters of *Boophilus microplus* Canestrini, 1887, exposed to entomopathogenic nematodes *Steinernema carpocapse* Santa Rosa and All strain (Steinernema: Rhabditida). *Braz. Arch. Biol. Technol.* 48: 911-919.
- GREWAL, P. S. 2002. Formulation, and application technology, pp. 265-287 *In* R. Gaugler (ed.), *Entomopathogenic Nematology*, Springer, Berlin Germany.
- NGUYEN, K. B., AND DUNCAN, L. W. 2002. *Steinernema diaprepesi* n. sp. (Rhabditida: Steinernematidae) a parasite of the citrus root weevil *Diaprepes abbreviatus* (L.) (Coleoptera: Curculionidae). *J. Nematol.* 34: 159-170.
- NGUYEN, K. B., MARUNIAK, J., AND ADAMS, B. J. 2001. Diagnostic and phylogenetic utility of the rDNA internal transcribed spacer sequences of *Steinernema*. *J. Nematol.* 33: 73-82.
- NGUYEN, K. B., SHAPIRO-ILAN, D. J., STUART, R. J., JAMES, R. R., AND ADAMS, B. J. 2004. *Heterorhabditis mexicana* n. sp. (Heterorhabditidae: Rhabditida) from Tamaulipas, Mexico, and morphological studies of the bursa of *Heterorhabditis* spp. *Nematology* 6: 231-244.
- NGUYEN, K. B., MALAN, A. P., AND GOZEL, U. 2006. *Steinernema khoisanense* n. sp. (Rhabditida: Steinernematidae) a new entomopathogenic nematode from South Africa. *Nematology* 8: 157-175.
- QUINTANA-MORENO, M. G., MOLINA-OCHOA, J., GONZÁLEZ-RAMÍREZ, M., AND GOVEA-CASAREZ, R. 2007. Susceptibilidad de la garrapata *Boophilus microplus* Canestrini, al nematodo entomopatógeno *Steinernema* sp. (Rhabditida: Steinernematidae) bajo condiciones de laboratorio. Universidad de Colima, Facultad de Medicina Veterinaria, Seminario de Investigación II, Tecomán, Colima, México, Agosto de 2007, 15 p.
- REIS-MENINI, C. M. R., PRATA, M. C. A., FURLONG, J., AND SILVA, E. R. 2008. Compatibility between the entomopathogenic nematode *Steinernema glaseri* (Rhabditida: Steinernematidae) and an acaricide in the control of *Rhipicephalus (Boophilus) microplus* (Acari: Ixodidae). *Parasitol. Res.* 103: 1391-1396.
- SAMISH, M., ALEKSEEV, E., AND GLAZER, I. 1999. Efficacy of entomopathogenic nematode strains against

- engorged females (Acari: Ixodidae) under simulated field conditions. *J. Med. Entomol.* 36: 727-732.
- SAS INSTITUTE. 1985. SAS user's guide: Statistics. SAS Institute, Cary, NC.
- STOCK, S. P., CAMPBELL, J. F., AND NADLER, S. A. 2001. Phylogeny of *Steinernema* Travassos, 1927 (Cephalobina: Steinernematidae) inferred from ribosomal DNA sequences and morphological characters. *J. Parasitol.* 87: 877-889.
- SWOFFORD, D. L. 2002. PAUP* Phylogenetic analysis using parsimony (*and other methods). Sunderland, MA, USA, Sinauer Associates.
- WOODRING, J. L., AND KAYA, H. A. 1988. Steinernematid and Heterorhabditid Nematodes. A Handbook of Techniques. Arkansas Agricultural Experiment Station, Fayetteville, Arkansas, Southern Cooperative Series Bulletin 331, 30 p.