



## **New Records of Thrips from Mesoamerica and Comments Regarding Specific Characters (Tubulifera: Phlaeothripidae)**

Authors: Valenzuela-García, Rita Dinora, Retana-Salazar, Axel P., García-Martínez, Oswaldo, and Cazola, Carlos Carvajal

Source: Florida Entomologist, 94(2) : 372-373

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.094.0242>

---

BioOne Complete ([complete.bioone.org](https://complete.bioone.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## NEW RECORDS OF THRIPS FROM MESOAMERICA AND COMMENTS REGARDING SPECIFIC CHARACTERS (TUBULIFERA: PHLAEOTHRIPIDAE)

RITA DINORA VALENZUELA-GARCÍA<sup>1</sup>, AXEL P. RETANA-SALAZAR<sup>2</sup>, OSWALDO GARCÍA-MARTÍNEZ<sup>1</sup>  
AND CARLOS CARVAJAL CAZOLA<sup>3</sup>

<sup>1</sup>Departamento de Parasitología, Universidad Autónoma Agraria Antonio Narro, Calzada Antonio Narro 1923, Buenavista, Saltillo, Coahuila, México, C.P. 25315

<sup>2</sup>Programa Universitario de Biología Aplicada (PUA), Centro de Investigación en Estructuras Microscópicas (CIEMIC), Ciudad de la Investigación, Universidad de Costa Rica 2060

<sup>3</sup>Unidad Académica de Agricultura, Universidad Autónoma de Nayarit, Xalisco, Nayarit, México, Tepic-Compostela Km 9

Several genera in the Tubulifera are monotypic, and in some cases species were described based on a sole original holotype (Mound, 1976). This scenario complicates the study of morphological variation in species, and in several cases, intraspecific variation may be greater than variation between species (Retana-Salazar & Mound 1994). Reports of taxa from new localities and descriptions of specific character variation are fundamental in the study of these groups. Modest publications of this sort advance the construction of a more complete landscape of morphological variation on an interspecific and intraspecific level.

A new locality (Nayarit, Mexico) for two known species is reported in this paper. Comments about variations of characters are included.

### Phlaeothripinae

#### *Eurythrips* Hinds 1902

*Eurythrips* is a New World genus, particularly common in the Neotropical region. The most recent revision of this genus placed in synonymy 16 of the 54 species included in this group (Mound 1976). The latest work concerning the thrips of Central and South America listed 37 species for this region (Mound & Marullo 1996). Several of these species are not well known and in some cases they are only known from the type material.

#### *Eurythrips longilabris* Watson

#### *Eurythrips longilabris* Watson 1921

#### *Eurythrips harti* Hood 1925

Material: Holotype female, USA, Florida State Collection of Arthropods (FSCA) and material from Texas (USNM) (Holotype of *E. harti*).

New Record. Collected on weeds, Santiago Ixcuintla County, Nayarit, México. 1 specimen.

Comments. Mound (1976), after of examination of the holotypes, considered both *longilabris* and *harti* species as synonymous; he considered that the morphological differences were not

enough to consider these as two different species. Indeed, Mound (1976) showed that main differences are in color details and in the postocular setae. Each of these setae is acuminate in the *longilabris* holotype but softly pointed in the *harti* holotype. The color pattern of this specimen collected in Mexico is congruent with the most comprehensive key published to date (Mound, 1976). The postocular setae in the material collected in Mexico are similar to those of the *longilabris* holotype described by Mound (1976), that is to say, an acuminate shaped apex is present.

### Idolothripinae

#### *Ethirothrips* Karny 1925

This is complex genus. Mound and Palmer (1983) considered 7 genera as synonyms of *Ethirothrips*. There are considerable differences between several species included in this genus. Mound and Marullo (1996) considered that some of these species are widely distributed and variable in color as well as in structure. *Ethirothrips* is a common member of the fauna of thrips associated with dead branches and leaf-litter in Brazil (Monteiro 2002).

#### *Ethirothrips firmus* Hood 1952

Material: Identified solely from a type series (5 specimens) collected in Sao Paulo, Brazil.

New Record. Collected on weeds, Santiago Country, Nayarit, Mexico. 1 specimen

Comments. Although described as a species of *Gastrothrips*, it is currently considered to be included in *Ethirothrips*. There are no reports of this species found anywhere else but at its original type locality. The structural variation in the species of this genus (Mound & Marullo 1996) presents a complex scenario for the study of this group. New material collected in México is congruent with all the morphological details described in the key presented by Mound and Marullo (1996). The presence of this species only in Brazil and Mexico suggests that Mound's hypothesis regarding distribution, related to human activity, holds true.

These two species have become identified through descriptions based on a single holotype, in the first case, and from a single type series in the second case. The new material presented allows new localities to be defined and the variability of morphological characters used in species definition to be explored further.

#### SUMMARY

Two Mexican species, apparently absent in other places in Mesoamerica are briefly described and recorded in this paper. It is very important to note that *Eurythrips longilabris* is recognized solely from the holotype collected in Florida, USA.

#### REFERENCED CITED

- MONTEIRO, R. C. 2002. The Thysanoptera fauna of Brazil, pp. 325-340 *In* R. Marullo and L. A. Mound [eds.], Thrips and Tosspoviruses: Proceedings of the 7th International Symposium on Thysanoptera, Reggio Calabria, Italy, 2-7 July 2001. Canberra, Australian National Insect Collection.
- MOUND, L. A., AND MARULLO, R. 1996. The Thrips of Central and South America: An Introduction. *Mem. Entomol. International* 6: 1-488.
- MOUND, L. A., AND PALMER, J. 1983. The generic and tribal classification of spore-feeding thrips Thysanoptera (Thysanoptera: Idolothripinae). *Bull. British Mus. Nat. Hist. Entomol.* 46: 1-174.
- MOUND, L. A. 1976. American leaf-litter Thysanoptera of the genera *Erkosothrips*, *Eurythrips* and *Terthrotrips* (Phlaeothripidae: Phlaeothripinae). *Bull. British Mus. Nat. Hist. Entomol.* 35(2): 27-64.
- RETANA, S. A. P., AND MOUND, L. A. 1994. Thrips of the *Frankliniella minuta* group (Insecta: Thysanoptera) in Costa Rican Asteraceae flowers. *Rev. Biol. Trop.* 42: 639-648.