



## **Biological Activity of Hyphomycete Entomopathogenic Fungi Against *Gynaikothrips uzeli* (Thysanoptera: Phlaeothripidae)**

Authors: Rios-Velasco, Claudio, Cambero-Campos, Jhonathan, Valenzuela-García, Rita, Gallegos-Morales, Gabriel, Cazola, Carlos Carvajal, et al.

Source: Florida Entomologist, 94(4) : 1060-1062

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.094.0447>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# BIOLOGICAL ACTIVITY OF HYPHOMYCETE ENTOMOPATHOGENIC FUNGI AGAINST *GYNAIKOTHRIPS UZELI* (THYSANOPTERA: PHLAEOTHIRIPIDAE)

CLAUDIO RIOS-VELASCO<sup>1</sup>, JHONATHAN CAMBERO-CAMPOS<sup>2</sup>, RITA VALENZUELA-GARCÍA<sup>3</sup>, GABRIEL GALLEGOS-MORALES<sup>3</sup>, CARLOS CARVAJAL CAZOLA<sup>2</sup> AND LUIS AGUIRRE-URIBE<sup>3</sup>

<sup>1</sup>Centro de Investigación en Alimentación y Desarrollo, A.C. Unidad Cuauhtémoc, Chihuahua. Av. Río Conchos S/N Parque Industrial, Apdo. Postal 781. P. 31570 Cd. Cuauhtémoc, Chihuahua, México

<sup>2</sup>Unidad Académica de Agricultura, Universidad Autónoma de Nayarit, Xalisco, Nayarit, México  
Tepic-Compostela Km 9

<sup>3</sup>Department of Parasitology, Universidad Autónoma Agraria Antonio Narro, Calzada Antonio Narro 1923, Saltillo, Coahuila, México, C.P. 25315

Weeping fig, *Ficus benjamina* L. (Moraceae) is an ornamental plant species grown worldwide. In Mexico, it is one of the most widely used ornamentals in urban green areas. Leaves of the tree are damaged by the thrips *Gynaikothrips uzeli* Zimmerman (Cambero et al. 2010). Thrips feed on leaf terminals and inject toxins causing deformed leaves and gall formation (Held et al. 2005; Retana-Salazar & Sánchez-Chacón 2009). Chemical control of *Gynaikothrips* spp. is possible (Held & Boyd 2008), but insecticides should be used wisely in order to preserve associated natural enemies (Wheeler et al. 2007). In Mexico, *Montandoniola confusa* Streito & Matocq and *Androthrips ramachandrai* (Karny) have been found to be predators of *Gynaikothrips* spp (Cambero et al. 2010). In North America north of Mexico, *A. ramachandrai*, *Montandoniola moraguesi* Puton and *Thripastichus gentilei* (Del Guercio) are reported natural enemies (Held et al. 2005; Held & Boyd 2008).

Microbial control could be integrated with natural enemies as an alternative to insecticides. Entomopathogenic fungi (Hyphomycetes) infecting thrips have been described, especially on *Frankliniella occidentalis* (Pergande) and *Thrips tabaci* (Lindeman) (Gouli et al. 2008; Thungrabeab et al. 2006). The Hyphomycetes are common and well known soil-borne pathogens in nature, with a wide range of insect hosts, and are consid-

ered excellent biological control agents (Gouli et al. 2008). Evaluations of entomopathogenic fungi noticeably are absent from previous research on control of *G. uzeli* (Held & Boyd 2008) or recent reviews (Held et al. 2005). The objective of this study was to test isolates of *Beauveria bassiana* (Balsamo) Vuillemin, *Metarhizium anisopliae* (Metschnikoff), and *Paecilomyces fumosoroseus* (Wize) against larvae and adults of *G. uzeli* under laboratory conditions.

Thrips used in bioassays were collected from trees of *F. benjamina* located in gardens of the Universidad Autónoma de Nayarit, in Tepic, Nayarit, Mexico, located at 21°29'18.73"N, 104°53'25"W and 944 masl, and were transferred to the Parasitology Department of the Universidad Autónoma Agraria Antonio Narro, Mexico. Thrips were identified by M. S. Jhonathan Cambero Campos. Entomopathogenic fungi were obtained from different hosts (Table 1) and were maintained and propagated on potato dextrose agar with 2% yeast extract (PDAY) that was supplemented with corn liquor to obtain pH 6.0. Purified fungi were identified according to their micro- and macroscopic characteristics (Humber 1997). Spores were stored under aseptic conditions in 0.05% Tween 80 sterile distilled water solution at pH 6.0. Spore viability ranged from 98 to 100% and was evaluated one day before bioassay setup by spreading 10 µL of each spore suspen-




Fig. 1. *Gynaikothrips uzeli* specimens infected with entomopathogenic fungi; a) *Metarhizium anisopliae* Ma-C;

TABLE 1. PERCENT MORTALITY (MEANS ± SD) OF *GYNAIKOTHRIPS UZELI* 9 DAYS AFTER TREATMENT WITH ISOLATES OF ENTOMOPATHOGENIC FUNGI IN MEXICO.

Entomopathogen	Original host	Concentration of spores per ml	% Mortality ± SD <sup>1</sup>	
			Larvae	Adult
<i>B. bassiana</i> Bb-S <sub>1</sub>	<i>Pogonomyrmex</i> sp. (Hymenoptera: Formicidae)	7.5 × 10 <sup>9</sup>	40.05 ± 5.08 b*	54.30 ± 7.8 a**
<i>B. bassiana</i> Bb-S <sub>2</sub>	<i>Spodoptera frugiperda</i> (Lepidoptera: Noctuidae)		11.30 ± 1.20 f	9.08 ± 0.94 c
<i>B. bassiana</i> Bb-S <sub>3</sub>	<i>Melanoplus bivittatus</i> (Orthoptera: Acrididae)		17.02 ± 1.06 e	7.40 ± 1.1 c
<i>M. anisopliae</i> Ma-A	<i>Amphides latrifons</i> (Coleoptera: Curculionidae)	1.5 × 10 <sup>9</sup>	64.80 ± 3.01 a*	60.50 ± 6.4 a**
<i>M. anisopliae</i> Ma-C	<i>Atta mexicana</i> (Hymenoptera: Formicidae)		36.04 ± 3.43c*	14.09 ± 0.60 c
<i>P. fumosoroseus</i> Pf-4a	Unknown host	2.5 × 10 <sup>10</sup>	20.10 ± 2.76 d	38.30 ± 4.8 b**
Control		0	0.10 ± 0.007	0.016 ± 0.002

<sup>1</sup>Means followed by different letters are significantly different according to Tukey's test on arcsine transformed data ( $P < 0.05$ ). Untransformed means are presented.  
\* Highly pathogenic (> 64.49%).  
\*\* moderately pathogenic (64.49-30.99%); according to Thungrabeab et al. (2006).

sion onto PDAY in a Petri dish. After 24 h of incubation at 25 ± 2 °C, percentage of spore germination was evaluated. A spore was considered viable if the germ tube was twice the length of the spore. Prior to thrips treatments, original spore suspensions were stirred for 2 min, and spore concentrations were determined with a hemacytometer (Blau Brand, Germany) and adjusted to the concentration used for each isolate (Table 1).

Leaves containing thrips (larvae and adults) were dipped in a spore suspension (5 × 10<sup>7</sup> to 2.5 × 10<sup>10</sup> spores per mL) for 10 s (Lewis 1997) and left to dry at ambient temperature (28 °C). Controls were treated with 0.05% Tween 80 water solution. Treated thrips (either 1st and 2nd instar larvae or adults) were then transferred to uninfested leaves that had been washed in 0.5% sodium hypochlorite water solution to eliminate contaminants and fixed to a base of plaster (4 mm thick) on Petri dishes. Each dish contained one leaf and 20 larvae or 20 adults. Dishes were sealed with parafilm, placed on a sponge saturated with water, and stored at 25 ± 2 °C, a photoperiod of 14:10 h L:D, and 95% relative humidity. Mortality of *G. uzeli* was recorded daily for 9 days post-inoculation, and fungal infection was confirmed by the presence of mycelium and conidia on the insect cuticle, as observed under a microscope.

Percent mortality was corrected using Abbott's formula (Abbott 1925) before statistical analysis. The experiment was conducted in 3 replicates using a randomized complete block design with 6 treatments, where each treatment was an isolate of entomopathogenic fungi (a total of 180 larvae and 180 adults per treatment were tested). Mortality data were normalized using arcsine transformation and were analyzed using Statistical Analysis System (SAS 2002) for balanced Analysis of Variance (ANOVA). Means were separated by Tukey's test ( $P < 0.05$ ).

Results showed that the 6 tested isolates were pathogenic to both stages of *G. uzeli*, and mycelial growth of fungi was observed on the cuticle of larvae and adults (Fig. 1). Mortality differed significantly ( $F = 72.09$ ,  $df = 5$ ,  $P < 0.001$  for adults and  $F = 3.09$ ,  $df = 5$ ,  $P < 0.0001$  for larvae) with isolates of *B. bassiana* (Bb-S1) and *M. anisopliae* (Ma-A) causing the highest mortality (Table 1). *B. bassiana* and *M. anisopliae* also have been reported as effective pathogens against *F. occidentalis* (Vestergaard et al. 1995) and *T. tabaci* (Thungrabeab et al. 2006; Gouli et al. 2008). We chose to test *M. anisopliae*, *B. bassiana*, and *P. fumosoroseus* due to their geographical distribution, host range, and potential as biological control agents. Although all tested isolates were pathogenic to *G. uzeli*, further research is needed to test methods and application equipment under field conditions. Thrips cause leaves to fold, which serves as protection, but may also complicate control.

## SUMMARY

The biological activity of *Beauveria bassiana* (isolates Bb-S1, Bb-S2, and Bb-S3), *Metarhizium anisopliae* (Ma-A and Ma-C), and *Paecilomyces fumosoroseus* (Pf-4a) as biological control agents of the thrips *Gynaikothrips uzeli* was evaluated under laboratory conditions. Isolates Bb-S1 and Ma-A caused the highest mortality in both larvae and adults. Further research is needed to test methods and equipment for application under field conditions.

## REFERENCES CITED

- ABBOTT, W. S. 1925. A method of computing the effectiveness of an insecticide. *J. Econ. Entomol.* 18: 265-267.
- CAMBERO, C. J., VALENZUELA, G. R., CARVAJAL, C. C., RIOS, V. C., AND GARCÍA, M. O. 2010. New records for Mexico: *Gynaikothrips uzeli*, *Androthrips ramachandrai* (Thysanoptera: Phlaeothripidae) and *Montandoniella confusa* (Hemiptera: Anthocoridae). *Florida Entomol.* 93: 470-472.
- GOULI, S., GOULI, V., SKINNER, M., PARKER B., MARCELINO, J., AND SHTERNISH, M. 2008. Mortality of western flowers thrips, *Frankliniella occidentalis*, under influence of single and mixed fungal inoculations. *J. Agr. Techn.* 4: 37-47.
- HELD, D. W., BOYD, D., LOCKLEY, T., AND EDWARDS, G. B. 2005. *Gynaikothrips uzeli* Zimmerman (Thysanoptera: Phlaeothripidae) in the southeastern United States: Distribution and review of biology. *Florida Entomol.* 88: 538-540.
- HELD, D. W., AND BOYD, D. W. 2008. Evaluation of sticky traps and insecticides to prevent gall induction by *Gynaikothrips uzeli* Zimmerman (Thysanoptera: Phlaeothripidae) on *Ficus benjamina*. *Pest. Manag. Sci.* 64: 133-140.
- HUMBER, R. A. 1997. Fungi. identification *In*: L. A. Lacey [ed.], *Manual of Techniques in Insect Pathology*. Academic Press, New York, USA.
- LEWIS, T. 1997. Pest thrips in perspective. *In* T. Lewis [ed.], *Thrips as Crop Pests*. CAB International, Wallingford, UK: 1-14.
- RETANA-SALAZAR, A. P., AND SÁNCHEZ-CHACÓN, E. 2009. Anatomía de las agallas en *Ficus benjamina* (Moraceae) asociada a "thrips" (Tubulifera: Phlaeothripidae). *Rev. Biol. Trop.* 57: 179-186.
- SAS INSTITUTE. 2002. *SAS User's Guide*. Version 9.0. SAS Institute, Cary, North Carolina, USA.
- THUNGRABEAB, M., BLAESER, P., AND SENGONCA, C. 2006. Possibilities for biocontrol of the onion thrips *Thrips tabaci* Lindeman (Thys., Thripidae) using different entomopathogenic fungi from Thailand. *Mitt. Dtsch. Ges. Allg. Angew. Entomol.* 15: 299-304.
- VESTERGAARD, S., GILLESPIE, A. T., BUTT, T. M., SCHREITER, G., AND EILENBERG, J. 1995. Pathogenicity of the Hyphomycetes fungi *Verticillium lecanii* and *Metarhizium anisopliae* to the western flower thrips, *Frankliniella occidentalis*. *Biocontrol Sci. Tech.* 5: 185-192.
- WHEELER, C., HELD, D. W., AND BOYD, D. 2007. Mortality of *Montandoniella moraguesi* following treatment of *F. benjamina* for control of weeping fig thrips. *Proc. South. Nurs. Assoc.* 52: 47-49.