



## **The Redbay Ambrosia Beetle (Coleoptera: Curculionidae) Prefers Lauraceae in Its Native Range: Records from the Chinese National Insect Collection**

Authors: Hulcr, Jiri, and Lou, Qiao-Zhe

Source: Florida Entomologist, 96(4) : 1595-1596

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.096.0444>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# THE REDBAY AMBROSIA BEETLE (COLEOPTERA: CURCULIONIDAE) PREFERS LAURACEAE IN ITS NATIVE RANGE: RECORDS FROM THE CHINESE NATIONAL INSECT COLLECTION

JIRI HULCR<sup>1,2,\*</sup> AND QIAO-ZHE LOU<sup>3,4</sup>

<sup>1</sup>School of Forest Resources and Conservation, University of Florida-IFAS, Gainesville, FL 32611-0410, USA

<sup>2</sup>Department of Entomology and Nematology, University of Florida-IFAS, Gainesville, FL 32611-0620, USA

<sup>3</sup>State Key Laboratory of Integrated Management of Pest Insects and Rodents, Institute of Zoology (IOZ), Chinese Academy of Sciences (CAS), Beijing, 100101, China

<sup>4</sup>University of Chinese Academy of Sciences, Beijing, 100049, China

\*Corresponding author; E-mail: hulcr@ufl.edu

The redbay ambrosia beetle *Xyleborus glabratus* is one of the most destructive invasive insect species in North America. The beetle is associated with a symbiotic fungus *Raffaelea lauricola* T.C. Harr., Fraedrich & Aghayeva (Ophiostomatales: Ophiostomataceae) which is extremely pathogenic to North American trees in the family Lauraceae (Fraedrich et al. 2008) and causes a disease known as laurel wilt. Since the introduction of this symbiotic couple in the early 2000's, laurel wilt killed millions of trees, mostly in the genus *Persea*, throughout most of southeastern USA (J. A. Smith, pers. comm.).

One of the important unknown aspects of the biology of this invasive species is its conspicuous specificity to Lauraceae. Most ambrosia species are broad host generalists (Hulcr et al. 2007). Current data on the original host specificity of *X. glabratus* are ambiguous. Published records of tree species from which the beetle was collected includes mostly Lauraceae, but also occasionally unrelated plant families (Wood & Bright 1992; Rabaglia et al. 2006), and it is not possible to confirm the accuracy of many of these records. The current hypotheses can be summarized as follows:

H0: The beetle is an unusual ambrosia species even in its native region, specialized on Lauraceae.

H1: As most ambrosia beetle species, the redbay ambrosia beetle is a host generalist in its native region. The specificity to Lauraceae in North America is a rapid evolutionary-ecological change.

Determining the correct hypothesis is important for many reasons. For example, knowing the original host specificity of the beetle may explain its unusual semiochemical ecology (Kendra et al. 2011). Importantly, it may also help us predict whether the symbiotic couple requires Lauraceae for its survival in invaded ecosystems, or if it is likely to persist in other tree families when Lauraceae are eliminated.

Here we present previously unpublished host records of *Xyleborus glabratus* deposited in the National Zoological Museum of China, Institute of Zoology, Chinese Academy of Sciences, Beijing. These are the first published records of this beetle occurring on mainland China. Identity of all specimens identified as *X. glabratus* was checked for correctness. Chinese host tree names were associated with scientific names according to Chang (1998), Cheng & Fu (1978) and Li (1982).

The collection at the Institute of Zoology in Beijing contains 45 specimens of *Xyleborus glabratus*, mostly collected by Dr. Fu-sheng Huang and identified by Dr. Hui-fen Yin. This is more than the number of records from the rest of Asia by Wood & Bright (1992). Specimens originated in the Sichuan, Hunan and Fujian provinces. This collection displays greater morphological variation than the American non-native population, particularly in terms of beetle size, but the larger or smaller beetles are not confined to specific host species or locations. The majority of specimens were collected from trees in the family Lauraceae (Table 1).

The host list indicates that *Xyleborus glabratus* is not a strict specialist to trees from the family Lauraceae, but it strongly prefers them. While this is an unusual behavior for an ambrosia beetle in the tribe Xyleborini, there are several other xyleborines for which preference for a particular host group has been reported (Kirkendall 2006; Hulcr et al. 2007). This also confirms that the apparent bias to Lauraceae in the host tree summary in Wood & Bright (1992) is correct, and it explains the observation of Dr. J. Peña (UF Tropical Research and Education Center, pers. comm.) who found *X. glabratus* in Taiwan only in *Cinnamomum osmophloeum* Kanehira. The 2 collections from non-lauraceous trees may represent uncommon behavior of the species.

The CAS collection labels do not contain any information on whether the trees were killed by the beetle. However, no records of this behavior in Southeast Asia have ever been reported. Healthy populations of the host tree species reported here

TABLE 1. HOSTS TREES OF *XYLEBORUS GLABRATUS* SPECIMENS IN THE NATIONAL ZOOLOGICAL MUSEUM OF CHINA.

Tree species	Family	Province(s)	Number of specimens
<i>Phoebe zhennan</i>	Lauraceae	Sichuan	21
<i>Machilus nanmu</i>	Lauraceae	Sichuan	6
<i>Cinnamomum camphora</i>	Lauraceae	Sichuan, Hunan	7
<i>Phoebe neurantha</i>	Lauraceae	Fujian	4
<i>Schima superba</i>	Theaceae	Fujian	1
<i>Picea asperata</i>	Pinaceae	Sichuan	6 (all the same collection)

occur in the areas where the *X. glabratus* specimens were collected, thus the authors doubt that *X. glabratus* displays tree-killing behavior in its native range.

The records explain the strong attraction of *X. glabratus* to phoebe oil, extracted from trees of the genus *Phoebe* (Kendra et al. 2011) and to many North American Lauraceae (Hanula et al. 2008). It also appears that occasionally the beetle is capable of colonizing non-lauraceous hosts, and it remains unknown whether or not it may eventually disappear from American ecosystems when all lauraceous of suitable size hosts have been eliminated. The origin of several *X. glabratus* from a pine tree (Table 1) is unclear, it may be an unusual or erratic behavior of those particular specimens, or an erroneous record.

We thank Drs. Jiang-hua Sun, Min Lu and Ji-an Yao for facilitating access to the collection. JH was funded by the University of Florida Emerging Pathogens Institute, the USDA Forest Service, and the National Science Foundation.

SUMMARY

The redbay ambrosia beetle *Xyleborus glabratus* is unusual among ambrosia beetles because of its host specificity to Lauraceae, but it isn't clear whether this is only a feature of the invasive American population. Our examination of the extensive collection in the Chinese Academy of Sciences suggests that *Xyleborus glabratus* strongly prefers Lauraceae also in its native Asia. These are also the first published records of the species from continental China, which highlights the value of entomological collections.

Key Words: host specificity, laurel wilt, *Persea*, symbiotic fungus, Theaceae, tree-killing behavior

RESUMEN

El escarabajo ambrosia *Xyleborus glabratus* es una especie insólita entre los escarabajos ambrosia, debido a su especificidad de hospederos de la familia Lauraceae, pero no está claro si esto es sólo una característica de esta población invasiva en los EE. UU. Nuestro examen de la extensa colección de la Academia de Ciencias de China sugiere que *Xyleborus glabratus*

prefiere fuertemente hospederos de la familia Lauraceae también en su nativa Asia. Estos son también los primeros registros publicados de las especies en China continental, lo que destaca el valor de las colecciones entomológicas. Palabras Clave: especificidad de hospedero, marchitez del laurel, *Persea*, hongo simbiótico, Theaceae, comportamiento de matar árboles

REFERENCES CITED

CHANG, H. 1998. Angiospermae: Dicotyledoneae: Theaceae (1): Theoideae *In* Flora Republicae Popularis Sinicae. Agendae Academiae Sinicae ed. 49 (3). Science Press, Beijing. (in Chinese).

CHENG, W., AND FU, L. 1978. Gymnospermae *In* Flora Republicae Popularis Sinicae. Agendae Academiae Sinicae ed. vol. 7. (in Chinese)

FRAEDRICH, S. W., HARRINGTON, T. C., RABAGLIA, R. J., ULYSHEN, M. D., MAYFIELD, A. E., HANULA, J. L., EICKWORT, J. M., AND MILLER, D. R. 2008. A fungal symbiont of the redbay ambrosia beetle causes a lethal wilt in redbay and other Lauraceae in the southeastern United States. *Plant Disease* 92: 215-224.

HANULA, J. L., MAYFIELD III, A. E., FRAEDRICH, S. W., AND RABAGLIA, R. J. 2008. Biology and Host Associations of the Redbay Ambrosia Beetle, (Coleoptera: Curculionidae: Scolytinae), exotic vector of laurel wilt killing redbay trees in the southeastern United States. *J. Econ. Entomol.* 101(4): 1276-1286.

HULCR, J., MOGIA, M., ISUA, B., AND NOVOTNY, V. 2007. Host specificity of ambrosia and bark beetles (Col., Curculionidae: Scolytinae and Platypodinae) in a New Guinea rain forest. *Ecol. Entomol.* 32: 762-772.

KENDRA, P. E., MONTGOMERY, W. S., NIOGRET, J., PEÑA, J. E., CAPINERA, J. L., BRAR, G., EPSKY, N. D., AND HEATH, R. R. 2011. Attraction of redbay ambrosia beetle (Coleoptera: Curculionidae: Scolytinae) to avocado, lychee, and essential oil lures. *J. Chem. Ecol.* 37: 932-942.

KIRKENDALL, L. R. 2006. A New Host-Specific, *Xyleborus vochysiae* (Curculionidae: Scolytinae), from Central America Breeding in Live Trees. *Ann. Entomol. Soc. America* 99: 211-217.

LI, H. 1982. Angiospermae: Dicotyledoneae: Lauraceae and Hernandiaceae *In* Flora Republicae Popularis Sinicae. Agendae Academiae Sinicae ed. vol. 31. Science Press, Beijing. (in Chinese).

WOOD, S. L., AND BRIGHT, D. E. 1992. A catalog of Scolytidae and Platypodidae (Coleoptera), Part 2: Taxonomic Index. *Great Basin Nat. Mem.*