

## **First Survey of Collembola (Hexapoda: Entognatha) Fauna in Soil of Archipelago Fernando de Noronha, Brazil**

Authors: Araujo De Lima, Estevam C., and Zeppelini, Douglas

Source: Florida Entomologist, 98(1) : 368-369

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.098.0161>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# First survey of Collembola (Hexapoda: Entognatha) fauna in soil of Archipelago Fernando de Noronha, Brazil

Estevam C. Araujo De Lima<sup>1,\*</sup> and Douglas Zeppelini<sup>1,2</sup>

Collembola (Hexapoda: Entognatha) is one of the most abundant and widely distributed taxa among terrestrial Hexapoda (Hopkin 1997). Collembola specimens are found in almost all habitats, excluding aquatic environments below the surface firm where their occurrence is rare or accidental. The greatest diversity and abundance of these species occurs in soil and in adjacent microhabitats, especially where there is much organic matter (Zeppelini et al. 2008). The potential value of Collembola as biological indicators of soil health and ecosystem quality is increasingly recognized and therefore knowledge of the diversity of Collembola becomes useful in the development of conservation strategies and environmental monitoring (Stork & Eggleton 1992; Zeppelini et al. 2008).

The growing human impact on biodiversity requires large efforts in developing conservation strategies (Myers et al. 2000). In island environments these efforts should be a priority, as these environments have major barriers to dispersal and colonization, especially for terrestrial fauna, which usually shows very particular diversity in islands. Analyses of the distribution of terrestrial species may be useful as a means to evaluate environmental quality (Rusek 1998; Kumssa et al. 2004).

In this work we present the first survey of Collembola fauna in the soil of the Brazilian archipelago Fernando de Noronha, which expands knowledge of the distribution of these animals. The collections were made in Jul and Aug 2012 on the 2 largest islands of the archipelago, Fernando de Noronha island and Rata island. The archipelago (S 3°50' W 32°24') is approximately 345 km away from the nearest point of the Brazilian coast. Collecting was done by soil sample on 7 different beaches on Fernando de Noronha. At each site we collected on the sand beach, on the cliff slope and in the forest at the hill top, with 4 samples in each collection site. Collembola were captured with Berlese-Tullgren funnels. Preparation and mounting followed Christiansen & Bellinger (1998). World distribution of species and nomenclature in Table 1 were based on Bellinger et al. (1996–2014), Culik & Zeppelini (2003), Abrantes et al. (2009) and original new records. Biogeographical distribution regions in Table 1 are according to Good (1974), as modified by Christiansen & Bellinger (1995). Species with distribution restricted to Brazil are assigned to biogeographic regions in Table 1 that include Brazil as in Culik & Zeppelini (2003). We collected 20,308 specimens of Collembola belonging to the following orders: Poduromorpha, Börner, 1913; Entomobryomorpha Börner, 1913; Symphypleona, Börner 1901 and Neelipleona Massoud, 1971. These specimens were distributed in 12 families, 28 genera and 38 species. Nine of these species appear to be restricted entirely to Fernando de Noronha. Fifteen of these species on this archipelago been observed at one or more other

**Table 1.** Collembola recorded on the Fernando de Noronha archipelago, Brazil. Collection localities were: a sandy beach (SB), soil on the slope of a cliff (SC) and the Soil in the forest at the hilltop (SF). World distribution was summarized for each species as follows: Boreal (Bor) include regions 1–8, Neotropical (Neo) regions 24–30, South African (Saf) region 31, Palearctic (Pal) regions 9–23, Australian (Aus) regions 32–34, and Antarctic (Ant) regions 35–37. Species distributed in at least, in 4 of the major regions (Neo, Pal, etc.) are considered to be cosmopolitan (Cos). Species distribution restricted to Northeast and Central Brazil (NCB), restricted to Fernando de Noronha (RFN) and doubtful distribution Record (?).

Taxa	Localities			World distribution
	SB	SC	SF	
<b>Neanuridae Börner, 1901</b>				
<i>Arlesia</i> sp. nov.		x		RFN
<i>Friezea cubensis</i> Potapov & Banasko, 1985		x		Neo
<b>Brachystomellidae Stach, 1949</b>				
<i>Brachystomella agrosa</i> Wray, 1953		x	x	Neo
<b>Hypogastruridae Börner, 1906</b>				
<i>Acherontiella</i> sp. nov.		x		RFN
<i>Willemia</i> sp. nov.		x		RFN
<i>Xenylla yucatanica</i> Mills, 1938		x	x	Cos
<b>Isotogastruridae Thibaud J-M &amp; Najt J, 1992</b>				
<i>Isotogastrura mucrospatulata</i> Palacios et al., 2013	x			RFN
<b>Isotomidae Schäffer, 1896</b>				
<i>Cryptopygus</i> sp.1			x	?
<i>Cryptopygus</i> sp.2		x	x	?
<i>Hemisotoma thermophila</i> Axelson, 1900		x	x	Cos
<i>Isotomiella</i> sp.		x	x	?
<i>Isotomodes</i> sp.		x	x	?
<i>Desoria</i> sp.		x		?
<i>Folsomides</i> sp.		x	x	?
<i>Folsomina</i> sp.		x	x	?
<i>Proisotoma</i> sp.1		x	x	?
<i>Proisotoma</i> sp.2		x	x	?
<i>Proisotoma</i> sp.3		x	x	?
<i>Proisotoma</i> sp.4			x	?
<b>Entomobryidae Schäffer, 1896</b>				
<i>Lepidocyrtus violaceus</i> Fourcroy, 1785			x	Bor, Neo, Pal
<i>Lepidocyrtus (Setogaster)</i> sp. nov.		x	x	RFN
<i>Pseudosinella flatua</i> Christiansen & Bellinger, 1996		x	x	Bor
<i>Pseudosinella aera</i> Christiansen & Bellinger, 1980		x	x	Bor, Neo
<i>Pseudosinella</i> sp.		x	x	?
<i>Entomobrya atrocincta</i> Schött, 1896			x	Cos
<i>Seira musarum</i> Ridley, 1890		x		RFN

<sup>1</sup>Laboratório de Sistemática de Collembola e Conservação. Depto. Biologia, Centro de Ciências Biológicas e Sociais Aplicadas. Programa de Pós Graduação em Ecologia e Conservação, Universidade Estadual da Paraíba Campus V, João Pessoa, Paraíba, Brazil

<sup>2</sup>Associação Guajiru-Ciência-Educação-Meio Ambiente Scientific Board, João Pessoa, Paraíba, Brazil

\*Corresponding author; E-mail: estevam.araujo@gmail.com

**Table 1.** (Continued) Collembola recorded on the Fernando de Noronha archipelago, Brazil. Collection localities were: a sandy beach (SB), soil on the slope of a cliff (SC) and the Soil in the forest at the hilltop (SF). World distribution was summarized for each species as follows: Boreal (Bor) include regions 1–8, Neotropical (Neo) regions 24–30, South African (Saf) region 31, Palearctic (Pal) regions 9–23, Australian (Aus) regions 32–34, and Antarctic (Ant) regions 35–37. Species distributed in at least, in 4 of the major regions (Neo, Pal, etc.) are considered to be cosmopolitan (Cos). Species distribution restricted to Northeast and Central Brazil (NCB), restricted to Fernando de Noronha (RFN) and doubtful distribution Record (?).

Taxa	Localities			World distribution
	SB	SC	SF	
<b>Paronellidae Börner, 1913</b>				
<i>Cyphoderus agnotus</i> Börner, 1906		x		Bor, Neo
<i>Cyphoderus inominatus</i> Mills, 1938		x	x	Neo
<i>Cyphoderus javanus</i> Börner, 1906		x	x	Neo, Pal
<i>Cyphoderus</i> <b>sp. nov.</b>		x	x	RFN
<i>Lepidonella</i> sp.		x	x	?
<b>Neelidae Folsom JW, 1896</b>				
<i>Megalothorax minimus</i> Willem, 1900		x	x	Cos
<b>Dicyrtomidae Börner, 1906</b>				
<i>Calvatomina</i> sp. nov.1 ( <i>rufescens</i> , group)		x	x	RFN
<i>Calvatomina</i> sp. nov.2 ( <i>rufescens</i> , group)		x	x	RFN
<b>Collophoridae Bretfeld, 1999</b>				
<i>Collophora terrabasilis</i> <b>nom. nov.</b>			x	NCB
<b>Bourletiellidae Börner, 1912</b>				
<i>Prorastriopes</i> sp.		x		?
<b>Sminthurididae Börner, 1906</b>				
<i>Sminthurides aquaticus</i> Bourlet, 1842			x	Bor, Aus
<i>Sphaeridia pumilis</i> Krausbauer, 1898		x	x	Cos

localities and 14 have questionable distribution record (Table 1). The only previous records of Collembola from the archipelago were those of *Seira musarum* Ridley, 1890 and *Isotogastrura mucrospatulata* Palacios et al. 2013. Furthermore, the subgenus *Setogaster* Salmon, 1951 and the genus *Calvatomina* Yosii are new records for Brazil.

The species *Collophora brasilis*, originally described in Zeppelini & Brito (2013), was renamed to make the epithet consistent with the etymology. The etymology does not refer to the native tree 'Pau-Brasil', but to 'Terra Brasiliis', the location. The original etymology refers to the location twice: (1) in the first sentence as 'Brazil' and (2) in the second sentence as 'Terra Brasiliis' which is an ancient name for Brazil. So the etymology explicitly refers to the location Brazil, therefore the epithet 'brasiliis' is not consistent with the etymology and thus invalid. To make it consistent it is being emended to *Collophora terrabasilis* nom. nov. for *Collophora brasilis* Zeppelini & Brito 2013.

## Summary

We provide the first survey of collembolan fauna of the archipelago Fernando de Noronha (Brazil) with 36 new records of species for the archipelago and a genus *Calvatomina* Yosii, 1966) and a subgenus (*Setogaster* Salmon, 1951) new for Brazil. Moreover the following is proposed replacement name: *Collophora terrabasilis* nom. nov. for *Collophora brasilis* Zeppelini & Brito (2013).

Key Words: distribution, microarthropods, springtails, oceanic island, replacement name

## Sumário

Apresentamos o primeiro levantamento da fauna de collembola do arquipélago de Fernando de Noronha com 36 novos registros de espécies para o Arquipélago e o registro de um gênero (*Calvatomina* Yosii, 1966) e um subgênero (*Setogaster* Salmon, 1951) para o Brasil. Além disso o seguinte nome de substituição é proposto: *Collophora terrabasilis* nom. nov. em substituição à *Collophora brasilis* Zeppelini & Brito (2013).

Palavras Chave: distribuição, microartrópodes, colêmbolos, ilhas oceânicas, nome de substituição

## References Cited

- Abrantes EA, Bellini BC, Bernardo AN, Fernandes LH, Mendonça MC, Oliveira EP, Queiroz GC, Sautter KD, Silveira TC, Zeppelini D. 2010 Synthesis of Brazilian Collembola: An update to the species list. *Zootaxa* 2388: 1-22.
- Abrantes EA, Bellini BC, Bernardo AN, Fernandes LH, Mendonça MC, Oliveira EP, Queiroz GC, Sautter KD, Silveira TC, Zeppelini D. 2012 Errata corrigenda and up date for the "Synthesis of Brazilian Collembola: An update to the species list." Abrantes et al. (2010), *Zootaxa*, 2388: 1-22; *Zootaxa* 3168: 1-21.
- Bellinger PF, Christiansen KA, Janssens F. 1996-2014. Checklist of the Collembola of the World. <http://www.collembola.org>
- Christiansen K, Bellinger P. 1998. The Collembola of North America. North of Rio Grande. Grinnell College, Grinnell, Iowa, USA. 1520 pp.
- Culik MP, Zeppelini D. 2003. Diversity and distribution of Collembola (Arthropoda: Hexapoda) of Brazil. *Biodiversity and Conservation* 12: 1119-1143.
- Good R. 1974. The geography of flowering plants. London, Longman Group, 4th ed., 574 pp.
- Jacquemart S. 1980. Collembolles nouveaux du Pérou. *Bulletin de l'Institut royal des Sciences naturelles de Belgique. Entomologie* 52(15): 1-27.
- Hopkin SP. 1997. Biology of Springtails: Collembola (Insecta). Oxford University Press, Oxford, UK.
- Kumssa DB, Van Aarde RJ, Wassenaar TD. 2004. The regeneration of soil microarthropod assemblages in a rehabilitating coastal dune forest at Richards Bay, South Africa. *African Journal of Ecology* 42: 346-354.
- Myers N, Mittermeier R, Mittermeier CG, Fonseca AB, Kent J. 2000. Biodiversity hotspots for conservation priorities. *Nature* 403: 853-858.
- Rusek J. 1998. Biodiversity of Collembola and their functional role in the ecosystem. *Biodiversity and Conservation* 7: 1207-1219.
- Stork NE, Eggleton P. 1992. Invertebrates as indicators of soil quality. *American Journal of Alternative Agriculture* 7: 38-47.
- Zeppelini D, Bellini BC. 2004. Introdução ao estudo dos Collembola. João Pessoa: Editora Universitária da UFPB.
- Zeppelini D, Bellini BC, Duarte AJC, Hernandez MIM. 2008. Collembola as bio-indicators of restoration in mined sand dunes of Northeastern Brazil. *Biodiversity and Conservation* 18: 1161-1170.