



## **Expression Profile of Two HSP70 Chaperone Proteins in Response to Extreme Thermal Acclimation in *Xestia c-nigrum* (Lepidoptera: Noctuidae)**

Authors: Wang, Ling, Yang, Shuai, Han, Lanlan, Zhao, Kuijun, and Ye, Lefu

Source: Florida Entomologist, 98(2) : 506-515

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.098.0218>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Expression profile of two HSP70 chaperone proteins in response to extreme thermal acclimation in *Xestia c-nigrum* (Lepidoptera: Noctuidae)

Ling Wang<sup>1</sup>, Shuai Yang<sup>1,2</sup>, Lanlan Han<sup>1</sup>, Kuijun Zhao<sup>1,\*</sup>, and Lefu Ye<sup>1</sup>

---

## Abstract

Heat shock proteins 70 (HSP70s) are highly evolutionarily conserved and play important roles in helping insects survive under extreme temperatures. In this study, *Xestia c-nigrum* (L.) (Lepidoptera: Noctuidae) was stressed at various temperatures, and the impacts on thermotolerance and cold endurance were examined. Two complementary deoxyribonucleic acid (cDNA) clones encoding heat shock cognate 70 (HSC70) and inducible heat shock protein 70 (HSP70), named *Xc-HSC70* and *Xc-HSP70*, respectively, were isolated from *X. c-nigrum* using reverse transcriptase polymerase chain reaction (RT-PCR) and rapid amplification of cDNA ends (RACE). Amino acid sequence analysis indicated that *Xc-HSC70* and *Xc-HSP70* contained the signature sequences of the HSP70 family, and phylogenetic analysis showed that *Xc-HSP70s* were consistent with the known insect classification. Genomic DNA structure analysis revealed that the *Xc-HSC70* gene contained 8 introns that all conformed to the GT/AG rule, whereas the *Xc-HSP70* gene did not contain any introns in the coding region. Fluorescent real-time quantitative RT-PCR revealed that both genes were significantly up-regulated under thermal stress and cold stress. The expression patterns indicated that *Xc-HSP70* was heat inducible and *Xc-HSC70* was constitutively expressed. The temperature response was more intense for *Xc-HSP70* than for *Xc-HSC70*, regardless of the temperature or developmental stage; however, the expression of *Xc-HSC70* and *Xc-HSP70* was not altered in the pupa. Taken together, the expression profiles of both genes (*Xc-HSP70s*) provide good support to the insect for coping with adverse conditions.

Key Words: heat shock proteins 70 (HSP70s); expression profile; thermal stress; *Xestia c-nigrum*

## Resumen

Proteínas de choque térmico 70 (Hsp70s) están evolutivamente muy conservadas y juegan un papel importante en ayudar a los insectos a sobrevivir las temperaturas extremas. En este estudio, se puso *Xestia c-nigrum* (L.) (Lepidoptera: Noctuidae) bajo el estrés de diferentes temperaturas, y se examinó el impacto sobre la termotolerancia y resistencia al frío. Se aislaron dos clones de ácido desoxirribonucleico (ADNc) complementarios que codifican el cognado choque térmico 70 (HSC70) y la proteína de choque térmico inducible 70 (HSP70), llamado *Xc-HSC70* y *Xc-HSP70*, respectivamente, a partir de *X. c-nigrum* mediante la cadena de polimerasa con transcriptasa inversa reacción (RT-PCR) y la amplificación rápida de extremos de ADNc (RACE). El análisis de la secuencia de aminoácidos indicó que *Xc-HSC70* y *Xc-HSP70* contenían las secuencias de la firma de la familia HSP70, y el análisis filogenético demostró que *Xc-Hsp70s* es consistente con la clasificación conocida del insecto. El análisis genómico de la estructura del ADN reveló que el gen *Xc-HSC70* contenía 8 intrones que todos se ajustaban a la regla GT/AG, mientras que el gen *Xc-HSP70* no contenía intrones en la región codificante. El análisis de RT-PCR fluorescente cuantitativa en tiempo real reveló que ambos genes fueron significativamente regulados hacia arriba bajo estrés térmico y estrés por frío. Los patrones de expresión indicaron que *Xc-HSP70* fue inducible por el calor y *Xc-HSC70* se expresó constitutivamente. La respuesta de la temperatura fue más intensa para *Xc-HSP70* que para *Xc-HSC70*, independientemente de la temperatura o el estadio de desarrollo; sin embargo, la expresión de *Xc-HSC70* y *Xc-HSP70* no se alteró en la pupa. Tomados en conjunto, los perfiles de expresión de ambos genes (*Xc-Hsp70s*) proveen un buen apoyo para el insecto para hacer frente a las condiciones adversas.

Palabras Clave: proteínas de choque térmico 70 (Hsp70s); perfil de expresión; estrés térmico; *Xestia c-nigrum*

---

Cellular stress response is involved in protecting organisms from damage caused by exposure to a variety of stressors, including temperature, heavy metals, and other xenobiotics. The stress response entails the rapid synthesis of heat shock proteins (HSPs) to protect cellular proteins against denaturation (Feder & Hofmann 1999; Boutet et al. 2003; Rinehart et al. 2007). In normal conditions, heat shock proteins primarily act as molecular chaperones involved in protein metabolism,

cell cycle regulation, and apoptosis (Welch 1993; Ming et al. 2010). On the basis of molecular weight and sequence similarity, HSPs can be divided into several families, including HSP90, HSP70, HSP60, HSP40, and small heat shock proteins (molecular weights ranging from 12 to 43 kDa) (Sørensen et al. 2003; Huang et al. 2008; Shen et al. 2014). Among the HSPs, HSP70s are the most conserved and important protein families and have been studied extensively. The HSP70 family is

---

<sup>1</sup>Northeast Agricultural University, College of Agriculture, Heilongjiang Province, Harbin 150030, China

<sup>2</sup>Virus-Free Seedling Research Institute, Heilongjiang Academy of Agricultural Sciences, Heilongjiang Province, Harbin, 150086, China

\*Corresponding author; E-mail: kjzhao@163.com

encoded by 2 different genes, constitutive type *HSC70* (heat shock cognate 70) and inducible type *HSP70* (Boutet et al. 2003; Deane & Woo 2005; Franzellitti & Fabbri 2005). In normal conditions, *HSC70* is constitutively expressed in all cells, whereas *HSP70* is not expressed or is expressed only at a low level. Under stress conditions, *HSC70* expression remains unchanged or is slightly up-regulated, whereas *HSP70* expression is highly induced (Lindquist & Craig 1988; Franzellitti & Fabbri 2005). These 2 heat shock proteins play key roles in the cell as molecular chaperones. However, *HSC70* is mainly involved in physiological processes, such as cell division, multiplication, and development (Park et al. 2001; Kregel 2002).

In insects, it is well known that HSP70s increase heat tolerance and provide protection against thermal injury and death. Many types of *HSP70* genes are up-regulated in response to heat or cold stress in various insects, such as Diptera (Huang & Kang 2007; Huang et al. 2009; Gray 2013), Lepidoptera (Jiang et al. 2012; Choi et al. 2014; Shen et al. 2014), and Coleoptera (Mahroof et al. 2005). Because of the important roles of HSP70s in thermal stress, it is important to investigate the function and expression characteristics of *HSP70* genes under temperature stress. In *Drosophila melanogaster* Meigen (Diptera: Drosophilidae), thermotolerance was found to be significantly improved in a strain with extra copies of *HSP70s* (Bettencourt et al. 2008; Jensen et al. 2014). The suppression of *HSP70* mRNA levels by RNAi decreased the heat and cold tolerance in *Pyrrhocoris apterus* (L.) (Heteroptera: Pyrrhocoridae) and *Spodoptera exigua* Hübner (Lepidoptera: Noctuidae) (Kostal & Tollarova-Borovanska 2009; Choi et al. 2014). In addition, HSP70s may be involved in the developmental processes of some insects and in fecundity (Jensen et al. 2014), longevity (Zhang et al. 2010; Choi et al. 2014), diapause (Rinehart et al. 2007), development (Huang et al. 2009), and metamorphosis (Zheng et al. 2010). These studies have provided direct evidence of the roles of HSP70s in cellular activity and development and have elucidated important biological functions.

*Xestia c-nigrum* L. (Lepidoptera: Noctuidae) is an important polyphagous pest of vegetables, cotton, wheat, maize, soybean, and ornamental plants. It is a worldwide pest that occurs in tropical, subtropical, and temperate regions (Mukawa & Goto 2011). In South China, including Guangdong, and Taiwan, *X. c-nigrum* is adapted to high temperatures that occur throughout the year. Severe crop damage occurs when the temperature is high in the summer and autumn, indicating that *X. c-nigrum* has a significantly positive response to high temperature (Zheng & Wang 2010; Zhang et al. 2013). In addition to heat tolerance, *X. c-nigrum* exhibits cold resistance, overwintering as larvae and pupae in temperate zones without entering diapause. The pupae have an average supercooling point of approximately  $-17^{\circ}\text{C}$  with a low of  $-21^{\circ}\text{C}$  in some individuals (Mukawa & Goto 2010; Landolt et al. 2011). In the southern region of North America and in Central and East Asia, overwintering pupae have been observed in fields (Xi et al. 2002). In China, the infesting population reaches northward to Heilongjiang Province in Northeast China. Notably, this species undertakes a long-distance migration between or within tropical, subtropical, and temperate regions, which significantly increases the geographic range over which infestations occur (Jiang et al. 2012). Furthermore, *X. c-nigrum* is known for its rapid development of resistance to many chemical pesticides and its lack of susceptibility to transgenic Bt crops (Landolt et al. 2011; Liu et al. 2011). Therefore, it can be presumed that *X. c-nigrum* has a high potential to tolerate or acquire tolerance to various environmental stresses. However, the biochemical and molecular mechanisms of such tolerance are unknown, and no information is available regarding stress-induced *HSP70* expression in *X. c-nigrum*; therefore, the mechanisms underlying this species' ecological adaptability and stress tolerance remain unclear.

This study examined basal gene expression and thermal responses among different developmental stages, such as the larvae and pupae of *X. c-nigrum*, using expression profiling of *Xc-HSC70* and *Xc-HSP70*. Furthermore, we evaluated *Xc-HSP70* expression at the mRNA level under heat or cold stress in larvae and pupae. We explored the functions of these genes in the context of thermotolerance and development and provided information needed to explore the mechanism of environmental tolerance and ecological adaptation in *X. c-nigrum*.

## Materials and Methods

### INSECTS

Adult *X. c-nigrum* specimens were collected from Xiangfang Farm of Harbin, Heilongjiang Province, China. After oviposition, the eggs were incubated at  $25 \pm 1^{\circ}\text{C}$  with a 12:12 h L:D photoperiod for hatching. Newly hatched larvae were reared at  $25 \pm 1^{\circ}\text{C}$  and 70% relative humidity (RH) under a photoperiod of 14:10 h L:D in a climatic cabinet (RXZ-308B, Jiangsu, China), and cabbage leaf was used to feed the different instars. For diapause induction, 6th instars were placed in an  $18 \pm 1^{\circ}\text{C}$  controlled-temperature room with an 8:16 h L:D photoperiod, and the diapause status of the pupae was ascertained as previously described (Liu et al. 2011).

### TEMPERATURE EXPOSURE

Based on previous studies of thermotolerance in insects (Jiang et al. 2012; Lyytinen et al. 2012; Franke et al. 2014), developing larvae from the 2nd to 6th instar were chosen for heat and cold shock treatments. For each treatment, 5 individuals were placed into a cryogenic tube and shocked at high temperatures of 28, 31, 34, 37, 40, 43, 45, and  $47^{\circ}\text{C}$  for 2 h, then allowed to recover at  $25^{\circ}\text{C}$  for 1 h. Similarly, larvae from the 2nd to 6th instar and 4-d-old pupae were shocked at low temperatures of 5, 0,  $-4$ , and  $-7^{\circ}\text{C}$  for 2 h, then transferred to  $25^{\circ}\text{C}$  for 1 h of recovery. Corresponding untreated larvae and pupae were used as controls. The larvae did not survive at above  $47^{\circ}\text{C}$  and below  $-7^{\circ}\text{C}$ , and therefore the mRNA was not analyzed. Each treatment was replicated 3 times. To detect the developmental regulation of the *HSC70* and *HSP70* genes, 2-, 4-, and 6-d-old pupae were reared at  $25^{\circ}\text{C}$ , with 3 replications. All of the samples were frozen quickly in liquid nitrogen and stored at  $-80^{\circ}\text{C}$ . Temperature control was achieved using environmental chambers (Sanyo, Tokyo, Japan).

### CLONING OF THE FULL-LENGTH CDNAS OF *XC-HSC70* AND *XC-HSP70*

Total RNA isolation and first-strand cDNA synthesis: RNA was isolated using an RNA isolation kit (Omega Bio-Tek, Norcross, Georgia, USA) according to the manufacturer's instructions. The concentration and quality of the RNA were verified by spectrophotometry and electrophoresis on a 1.0% agarose gel. The cDNA was synthesized using a cDNA kit (TaKaRa, Dalian, China) according to the manufacturer's instructions.

Primer design: The *HSC70* and *HSP70* genes of other insects were aligned in GenBank to identify the conserved region of the gene family. Primer Premier 5.0 (Premier, Canada) was used to design the primers, and the sequences of the primers are listed in Table 1. The PCR protocol was as follows: 3 min at  $94^{\circ}\text{C}$  and 35 cycles of 30 s at  $94^{\circ}\text{C}$ , 30 s at  $57.8^{\circ}\text{C}$  for *Xc-HSC70* or 30 s at  $58.0^{\circ}\text{C}$  for *Xc-HSP70*, and 1 min at  $72^{\circ}\text{C}$ , followed by a 10 min extension at  $72^{\circ}\text{C}$ . Then, the amplified fragment was isolated using a 1.0% agarose gel and ligated into the pMD18-T vector (TaKaRa, Dalian, China) for sequencing. RACE amplification was

**Table 1.** Primers for the experiment in this study.

Primer type	Sequence (5' to 3')	Use of primers
P1	GAGGGMRTCGACTTCTAYACBTCCATCAC	PCR for <i>Xc-HSC70</i>
P2	CTTGTA CTCTCTG CNTCYTGBACCAT	
P3	AACGARCCBACYGCGYCGC	PCR for <i>Xc-HSP70</i>
P4	CGGTC SATSTCRAADGTGAC	
P5	CGTGAAAAGAAGCCGTC AATCG	For full-length and gDNA sequence of <i>Xc-HSC70</i>
P6	GAGTTTCAATCATTTATTGGAGAGACTGTAC	
P7	CACCTTTGCTGAGTTACTCTACGAGTTAAG CTGAAG	For full-length and gDNA sequence of <i>Xc-HSP70</i>
P8	CAATATCAGTAAAGCTTGCTGTAATGTTG	
5' <i>HSC70</i> GSP1	CGTTACGTTCTCCATCCCTTCTTGTC AAGGCCG	5' RACE for <i>Xc-HSC70</i>
5' <i>HSC70</i> GSP2	GTTGGAAGCGAGGTCTCTCTTG	5' RACE for <i>Xc-HSP70</i>
5' <i>HSP70</i> GSP1	ATCAACCCCGACGAGGCCGTAGCTTA	
5' <i>HSP70</i> GSP2	AGGTACAGGATCTGCTGCTGCTCGA	3' RACE for <i>Xc-HSC70</i>
3' <i>HSC70</i> GSP1	GAACGCTCTGGAATCTTACTGCTTCAACATG	
3' <i>HSC70</i> GSP2	GAAGTCTACCAACAAGGAGAACAAAG	3' RACE for <i>Xc-HSP70</i>
3' <i>HSP70</i> GSP1	GTCGGTGTTCTACCCGTATTCC	
3' <i>HSP70</i> GSP 2	TACCGTCGTTCCACTAGTGATT	RT-qPCR for <i>Xc-HSC70</i>
<i>HSC70</i> F	CTTGACCGGTGACAAGTCTGAG	
<i>HSC70</i> R	GAATGTCTGGGTTTGCTTG GTG	RT-qPCR for <i>Xc-HSP70</i>
<i>HSP70</i> F	GTGACGCGAAGATGGACAAGTC	
<i>HSP70</i> R	CACCGTAAGCTACAGCCTCGTC	RT-qPCR for $\beta$ -actin
$\beta$ -actin F	CGCGACCTCACAGACTACCTG	
$\beta$ -actin R	CGTAGGACTTCTCCAGGGAGC	

Note: All primers were synthesized at Shanghai Generey Biotech Co., Ltd. (Shanghai, China).

performed based on the sequenced fragment of *Xc-HSC70* and *Xc-HSP70* using the 3'Full and 5'Full RACE kit (TaKaRa, Dalian, China). We designed the 5'RACE and 3'RACE primers 5' *HSC70* GSP1, 5' *HSC70* GSP2, 5' *HSP70* GSP1, 5' *HSP70* GSP2, 3' *HSC70* GSP1, 3' *HSC70* GSP2, 3' *HSP70* GSP1, and 3' *HSP70* GSP2 (Table 1) and combined the 3 single fragments of *Xc-HSC70* and *Xc-HSP70* to design the full-length primer pairs P5, P6 and P7, P8, respectively.

CLONING OF THE *XC-HSC70* AND *XC-HSP70* GENES FROM GENOMIC DNA

Genomic DNA was isolated from 5th instars using the Universal Genomic DNA Extraction kit (Takara, Dalian, China) according to the manufacturer's instructions. The concentration and quality of the DNA were verified by spectrophotometry and electrophoresis on a 1.0% agarose gel.

To determine whether the *Xc-HSC70* and *Xc-HSP70* genes contained introns in their coding regions, the genomic DNA fragments for *Xc-HSC70* and *Xc-HSP70* were amplified from genomic DNA (100 ng) as a template, and their nucleotide sequences were analyzed. For *Xc-HSC70* intron analysis, the amplification was performed using a pair of specific primers (P5, P6; Table 1), and the amplification conditions were 94 °C for 3 min, followed by 35 cycles of 94 °C for 30 s, 58 °C for 45 s, and 72 °C for 4 min, with a final elongation step at 72 °C for 10 min. For *Xc-HSP70* intron analysis, the genomic DNA fragment was amplified using a pair of specific primers (P7, P8; Table 1), and the amplification conditions were as described above. The DNA fragments were then cloned and sequenced.

SEQUENCE ANALYSIS

The bioinformatic software DNASTar (DNASTar, USA) was used to split joint the full-length sequences of *Xc-HSC70* and *Xc-HSP70*. BLAST software was used to analyze the homology of the sequences (<http://www.ncbi.nlm.nih.gov/blast/>). Biology WorkBench ([\[workbench.sdsc.edu/\]\(http://workbench.sdsc.edu/\)\) was used to identify the open reading frames \(ORFs\). ProtParam \(\[http:// web.expasy.org/protparam/\]\(http://web.expasy.org/protparam/\)\), Scan-Prosit e \(<http://prosite.expasy.org/scanprosite/>\), and SWISS-MODEL \(<http://www.swissmodel.expasy.org/SWISS-MODEL.html>\) were used to analyze the protein sequence characteristics. PSORT II \(<http://psort.hgc.jp/form2.html>\) was used to calculate the nuclear localization signal fragment. CLUSTALW and MEGA 4.0 were used to draw the cladogram. Spidey \(<http://www.ncbi.nlm.nih.gov/spidey/>\) was used to analyze the genomic introns.](http://</a></p></div><div data-bbox=)

QUANTITATIVE ANALYSIS OF *XC-HSC70* AND *XC-HSP70* MRNA EXPRESSION

Total RNA isolation and first-strand cDNA synthesis were isolated as above. Real-time quantitative PCR (qRT-PCR) primers were designed using the *Xc-HSC70* (*HSC70* F, *HSC70* R) and *Xc-HSP70* (*HSP70* F, *HSP70* R) sequences, and the resulting products had lengths of 145 bp and 141 bp, respectively. The reference gene  $\beta$ -actin was used as an endogenous control to quantify the expression of the target genes. This gene is an appropriate control for studies on *HSP70s* in insects during this wide developmental window as determined in our previous studies (Jiang et al. 2012; Yu et al. 2012), with a resulting product of 156 bp. *Xc-HSC70*, *Xc-HSP70*, and  $\beta$ -actin sequences were amplified from each of the instars under each treatment condition using specific primers, Thunderbird® Sybr qPCR Mix (ToYoBo, <http://www.toyobo-global.com>, Japan), and a Chromo4™ Real-Time PCR instrument (Bio-Rad, Hercules, California, USA). All of the amplifications were performed in triplicate. The final volume of each qRT-PCR reaction was 20  $\mu$ l, which contained 10  $\mu$ l of 2  $\times$  SYBR Mix (ToYoBo, <http://www.toyobo-global.com>, Japan), 1  $\mu$ l of diluted cDNA template, 7.8  $\mu$ l of PCR-grade water, and 0.6  $\mu$ l of each primer at 10  $\mu$ M. PCR conditions were as follows: 95 °C for 30 s and 40 cycles of 95 °C for 10 s and 60 °C for 30 s.

*Xc-HSC70* and *Xc-HSP70* expression levels were calculated using the 2<sup>- $\Delta\Delta$ CT</sup> comparative threshold cycle (CT) method (Livak & Schmittgen 2001). The mean and standard deviation were calculated from experi-



ments performed in triplicate and presented as *n*-fold differences in expression. Differences in the transcriptional features of *Xc-HSC70* and *Xc-HSP70* in different developmental stages were analyzed using SPSS 16.0. Statistical significance was determined using one-way analysis of variance (ANOVA) and post-hoc Duncan multiple range tests. Significance was defined as  $P < 0.05$ .

## Results

### NUCLEOTIDE ANALYSIS AND DEDUCED AMINO ACID SEQUENCES OF *XC-HSC70* AND *XC-HSP70*

Total RNAs of *Xc-HSC70* and *Xc-HSP70* were isolated from larvae of *X. c-nigrum* which reared at room temperature and treated at 37 °C respectively. Full-length cDNAs of *Xc-HSC70* and *Xc-HSP70* were 2,152 bp (published in Wang et al. 2014) and 2,213 bp (published in Wang et al. 2015), respectively. The sequence of *Xc-HSC70* contains a 5'-terminal untranslated region (UTR) of 101 bp, a 3'-terminal UTR of 86 bp, and an ORF of 1,965 bp that encoded a protein of 654 amino acids with a calculated molecular weight of 71.59 kDa (Fig. 1). The sequence of

*Xc-HSP70* also contains a 5'-terminal UTR of 147 bp, a 3'-terminal UTR of 86 bp, and an ORF of 1,965 bp that encoded a protein of 654 amino acids with a calculated molecular weight of 71.62 kDa (Fig. 2). Amino acid sequence analysis indicated that *Xc-HSC70* and *Xc-HSP70* contained the cytoplasmic characteristic motif EEVD (Boutet et al. 2003) and 3 signature sequences of the HSP70 family (Figs. 1 and 2). At the carboxyl terminal region, *Xc-HSC70* contained 3 consecutive repeats of the tetrapeptide motif GGMP (617 to 628 amino acids), whereas *Xc-HSP70* contained 2 tetrapeptides of GGMP (617 to 620 amino acids, 625 to 628 amino acids). Putative bipartite nuclear localization signals (NLS) were also observed in the two HSP70s using the online software PSORT II (Figs. 1 and 2).

### ANALYSIS OF THE GENOMIC DNA STRUCTURE OF *XC-HSC70* AND *XC-HSP70*

For *Xc-HSC70* intron analysis, a genomic DNA fragment of 3,710 bp was isolated by specific primers (P5, P6; Table 1). Comparison of the sequence isolated from genomic DNA with the cDNA sequence of *Xc-HSC70* revealed 9 exons and 8 introns in the *Xc-HSC70* fragment. The coding region of the *Xc-HSC70* gene contained 8 exons

```

1      GAAAAATTTTTCGTGAAAGAACGCCGTAATCGCGCAGTTTGTCTGAGTTACTCAACGAGTTAAGTCAACGACTGAGATAGTTATACAAAAAATTAATAATGGCAGCACTAAAGCAC
1      M A A T K A P
121    CCGCTGTAGGTATCGACTTGGGTACCACTTACTCATGTGTGGGAGTTTCCAGCATGGTAAAGTGGAGATCATCGCCAACGACAGGGCAACAGGACCACGCCCTCATATGTCCGGTTCA
8      A V G I D L G T T Y S C V G V F Q H G K V E I I A N D Q G N R T T P S Y V A F T
241    CAGACACCGAGCGTCTCATCGGAGATGCCCAAGAACAGGTGGCGATGAACCTTAACAACACAATTTTCGATGCCAAACGTTCTCATCGGCCGTAAATTCGAAGATGCCACTGTACAAG
48      D T E R L I G D A A K N Q V A M N P N N T I F D A K R L I G R K F E D A T V Q A
361    CCGACATGAAGCACTGGCCCTTCGAGGTAGTCAGTGACGGTGGCAAGCAAGATCAAGGTCTCATACAAAGTGAAGACAAACCTTCTCCCTGAGGAAGTAGCTCAATGGTGCTCA
88      D M K H W P F E V V S D G G K P K I K V S Y K G E D K T F F P E E V S S M V L T
481    CAAAAATGAAGAAACAGCCGAGGCTTACCTCGGCAAACTGTGCAGATGCAGTAATCACGGTTCAGCGTACTTCAATGACTCACAGACAGCAAGACAAAAGATGCGGGTACCATCT
128     K M K E T A E A Y L G K T V Q N A V I T V P A Y F N D S Q R Q A T K D A G T I S
601    CTGGCTTGAATGTTCTCGTATTATCAATGAACGACTGCTGCTGCGATTGCGTACGGCCTTGACAAGAGGGGAGTGGAGAAGCTAACGTATTGATTTTCATCTCGGCGCGGTACCT
168     G L N V L R I I N E P T A A A I A Y G L D K K G S G E R N V L I F D L G G G T F
721    TCGACGTTTCCATCTGACCATCGAGGATGGTATCTTTGAAGTAAAGTCCACGCGGTGACGCGATTGGGAGGAGAGGACTTCGACAACCGAATGGTAAACCACTTTGTGCAAGAGT
208     D V S I L T I E D G I F E V K S T A G D T H L G G E D F D N R M V N H F V Q E F
841    TCAAGAGGAAATACAAGAAGGACCTCGCTTCCAACAGAGGGCCCTTCGTGGTTGCGCACTGCGCTGTGAAGGGCGAAGAGAACTCTCTCTCGTCCACCAGGCTAGCATTGAATCG
248     K R K Y K K D L A S N K R A L R R L R T A C E R A K R T L S S S T Q A S I E I D
961    ACTCTCTGTTGAGGGTATGACTTCTATACCTCCATCACCAGGGCTGCTTTGAAGAACTGAAGCTGACCTGTTGAGTCCATCTGGAGCCTGTGGAGAAGTCCCTTCGTGACGCGA
288     S L F E G I L D F Y T S I T R A R F E E L N A D L F R S T M E P V E K S L R D A K
1081   AGATGGACAAGTCTCAATCCACGACATCGTACTTGTGGTGGTCTACCGGTATCCCAAAGTGCAAGGCTCTTCAAGACTTCTTAAATGGCAAGGAGCTTAACAAATCCATCAACC
328     M D K S Q I H D I V L V G G S T R I P K V Q K L L Q D F F N G K E L N K S I N P
1201   CTGACGAGGCGGTAGCTTACGGTGTGCGGTCCAGGCTGCCATCTTGACCGGTGACAAGTCTGAGGAGGTGACGAGTCTTCTGCTGCTCGACGTGACTCCGTTGTGCTCGGTATCGAAA
368     D E A V A Y G A A V Q A A I L H G D K S E E V Q D L L L L D V T P L S L G I E T
1321   CTGCGGCGGTGTAATGACCACACTCATCAAGCGCAACACCACCATCCCACCAAGCAAAACCCAGACATTACCACCTACTCTGACAACAGCCTGGAGTACTTATCCAGGTGTTGGAAG
408     A G G V M T T L I K R N T T I P T K Q T Q T F T T Y S D N Q P G V L I Q V F E G
1441   GTGAGCGGCCATGACCAAGGATAACAACCTTCTCGGAAAGTTCGAGTGCAGCGCATCCCTCCCGCGCGCGTGGCGTGCCTCCAGATCGAAGTCACTTTGACATTGACGCCAACGGCA
448     E R A M T K D N N L L G K F E L T G I P P A P R G V P Q I E V T F D I D A N G I
1561   TTCTTAACTGTGCGCCGTGAGAGTCTACCAACAGGAGAACAGATCACCATCACCACGCAAGGGCCGTCTCTCAAGGAGGAGATCGAGCGCATGGTCAACGAGGCCGAGAAAT
488     L N V S A V E K S T N K E N K I T I T N D K G R L S K E E I E R M V N E A E K Y
1681   ACAGGACTGAGGATGAGAAGCAAGAGACCATCCAGGCTAAGAACGCTCTGGAATCTTACTGCTTCAACATGAAGTCCACATGGAAGACGAGAAGCTCAAGGACAAAATCTCGACT
528     R T E D E K Q K E T I Q A K N A L E S Y C F N M K S T M E D E K L K D K I S D S
1801   CCGACAAACAGACCATCTGGACAAGTGCAACGACACCATTAATGGCTAGACTCCAATCAGTTGGCTGACAAGAAGAATACGAGCATTAAGCAGAAGGAATGGAAGGCATCTGCAACC
568     D K Q T I L D K C N D T I K W L D S N Q L A D K E E Y E H K Q K E L E G I C N P
1921   CTATTATACCAAAATGTACAGGAGCAGGTGGTATGCCCGGCGGTATGCCCGGAGGTATGCCCGGCTTCCCGGCGGCGTCCCGGCGGCGGTGGCGGCGTCCCGGTGGCGGTGCCG
608     I I T K M Y Q G A G G M P G G M P G G M P G F P G G A P G A G G A A P G G G A G
2041   GACCTACCATCGAGGAGTTCGACTAAACATTTCAACAAATATACAGCAAGCTTTTACTGATATTGCAGTATTCACTACAGTACAGTCTCTCAATAAATGATTGAAACTCAAAAAA
648     P T I E E V D *
2161   AAAA

```

**Fig. 1.** Nucleotide and deduced amino acid sequences of *Xc-HSC70*. The signature sequences of the HSP70 family are shown in boxes, the nuclear localization signal sequence is underlined, the consensus sequence EEVD at the C-terminus is indicated in italics, and the start and stop codons are in bold. The nucleotides and amino acids are numbered along the left and right margins. The sequence encoding *Xc-HSC70* has been deposited in GenBank under accession no. KC844151.



1 ATTCGCGGATCCACAGCCTACTGATG  
 27 ATCAGTCGATGAAAAATTTTTTTTTTCGTGAAAAAGCCGTCATCGCGCACTTTGCTGAGTTACTCTACGAGTTAAGTCAACGACTGAGATAGTTATACATAAAATTAATAAAA  
 147 ATGGCAGCACTAAAGCACCCGCTGTAGGTATCGACTTGGGTACCACTTACTCATGTGTGGGAGTTTCCAGCATGGTAAGGTGGAGATCATCGCAACGACAGGGCAACAGGACCAAGC  
 1 M A A T K A P A V G I D L G T T Y S C V G V F Q H G K V E I I A N D Q G N R T T  
 I  
 267 CCCTCATATGTCGCGTTTACCGACACTGAGCGTCTCATCGGAGATGCCCAAGAACAGGTGGCGGTGAACCTTAACAACAATTTTCGATGCCAACGCTCATCGGCCGCAAAATTC  
 41 P S Y V A F T D T E R L I G D A A K N Q V A V N P N N T I F D A K R L I G R K F  
 387 GAAGATGCCACCGTACAAGCTGATGAAGCACTGGCCCTTCGAGGTAGTCAGTGACGGTGGCAACAAAGATCAAGGTCTCGTACAAGGCGAAGACAAAACCTTCTCCCTGAGGAA  
 81 E D A T V Q A D M K H W P F E V V S D G G K P K I K V S Y K G E D K T F F P E E  
 507 GTTAGCTCAATGGTCTCACAAAAATGAAGGAAACAGCCGAGGCTACCTCGGCAAAACGGTGCAGAAATGCAGTAATCACGGTTCAGCGTACTTCAATGACTACAGAGACAAGCCACA  
 121 V S S M V L T K M K E T A E A Y L G K T V Q N A V I T V P A Y F N D S Q R Q A T  
 627 AAAGATCGGGTACCATCTCTGGCTTGAATGTCTCGTATATCAATGAACCGACTGCTGCTGCGATTGCATACGGCTTGACAAAAGGGGAGTGGAGAACGTAACGATTGATTTC  
 161 K D A G T I S G L N V L R I I N E P T A A A I A Y G L D K K G S G E R N V L I F  
 747 GATCTCGGCGGCTACCTTCGACGTTTCCATCTGACCATCGAGGATGGTATCTTGAAGTAAAGTCCACCGCTGGTGACACGCAATTGGGAGGAGAGGACTTCGACACCGAATGGTA  
 201 D L G G G T F D V S I L T I E D G I F E V K S T A G D T H L G G E D F D N R M V  
 II  
 867 AACCCTTTGTCAGGAGTTCAAGAGGAAATACAAGAAGGACCTCGCTTCCAACAAGAGGGCCCTTCGTCGGTTGCGCACTGCGCTGTGAAGGGCGAAGAGAACTCTCTCTCGTCCACC  
 241 N H F V Q E F K R K Y K K D L A S N K R A L R R L R T A C E R A K R T L S S S T  
 987 CAGGCTAGCATGAAATCGACTCTCTGTTCGAGGGTATTGACTTCTATACCTCCATCACCAGGGCTCGTTTGAAGAACTGAACGCTGACCTGTTTCAGATCCACTATGGAGCCTGTGGAG  
 281 Q A S I E I D S L F E G I D F Y T S I T R A R F E E L N A D L F R S T M E P V E  
 1107 AAGTCCCTTCGTGACGCGAAGATGGACAAGTCTCAATCCACGACATCGTACTTGTGCGTGGTTCTACCCGTATTCCCAAAGTGCAGAAGCTCTTCAAGACTTCTTAAATGGAAGGAG  
 321 K S L R D A K M D K S Q I H D I V L V G G S T R I P K V Q K L L Q D F F N G K E  
 III  
 1227 CTTAACAATCCATCAACCTGACGAGGCTGTAGCTTACGGTGTCCGCTCCAGGCTGCCATCTTGCACGGTGACAAGTCTGAGGAGGTGACGATCTGCTGCTGCTCGACGTGACTCCG  
 361 L N K S I N P D E A V A Y G A A V Q A A I L H G D K S E E V Q D L L L L D V T P  
 1347 TTGTGCTCGGTATCGAAACCGCGCGGTGTATGACCACTCATCAAGCGCAACACCACTCCCAAGCAAGCAACCCAGACATTACCACTACTCTGACAACCCAGCCTGGAGTA  
 401 L S L G I E T A G G V M T T L I K R N T T I P T K Q T Q T F T T Y S D N Q P G V  
 1467 CTTATCAGGTGTTCGAGGGGAGCGGCCATGACCAAGGAAAGAACACATCGGAAAGTTCGAGTGAACCGCATCCCTCCGCGCCCCGTGGCGTGGCCAGATCGAAGTACCTTC  
 441 L I Q V F E G E R A M T K E R N T L G K F E L T G I P P A P R G V P Q I E V T F  
 1587 GACATCGACGCTAACGGCATTTCAACGTGTCTGCGTTCGAGAAGTGCAGTAAACAGGAGAAAGATCACCATCCACGACAAGGGCCGCTCTCCAAGGAGGAGATCGAGCGCATG  
 481 D I D A N G I L N V S A V E K S T N K E N K I T I T H D K R G K E E I E R M  
 1707 GTCAACGAGGCGGAGAAGTACAGGACTGAAGACGAGAAGCAGAAGGAGACCATCCAGGCTAAGAAGCGCCCTGGAATCTTACTGCTTCAACATGAAGTCCACCATGGAGGACGAGAAGCTC  
 521 V N E A E K Y R T E D E K Q K E T I Q A K N A L E S Y C F N M K S T M E D E K L  
 1827 AAGGACAAGATCTCAGACTCCGACAAGCAGACCATCTGGACAAGTGAACGATACCATCAAAATGGCTGGACTCCAACGAGTGGCTGACAAGGAAGATACGAGCCCAAGCAGAAGGAA  
 561 K D K I S D S D K Q T I L D K C N D T I K W L D S N Q L A D K E E Y E P K Q K E  
 1947 TTGGAAGGCATCTGCAACCCCAATCATCAAGATGTACCAGGAGCAGGTGGCATGCCAGTGGTATGTCGGGAGGCATGCGCGCTTCCCGGGGGCGCTCCCGGCGCGGGGGCGCA  
 601 L E G I C N P I I T K M Y Q G A G G M P S G M P G G M P G F P G G A P G A G G A  
 2067 GCCCGGGGGCGGTGCGGACCTACCATCGAGGAGTTCGACTAAACAAACATTCACCAATTACAGCAAGCTTTACTGATATTGCACTATTCACTACAGTACAGTGTCTCCAATAAAAT  
 641 A P G G G A G P T I E E V D \*  
 2187 GATTGAACTCAAAAAAAAAAAAAA

Fig. 2. Nucleotide and deduced amino acid sequences of the *Xc-HSP70* gene. The signature sequences of the *HSP70* family are shown in boxes, the nuclear localization signal sequence is underlined, the consensus sequence EEVD at the C-terminus is indicated in italics, and the start and stop codons are in bold. The nucleotides and amino acids are numbered along the left and right margins. The sequence encoding *Xc-HSP70* has been deposited in GenBank under accession no. HQ698836.


and 7 introns (Fig. 3). The first intron (561 bp) was located in the 5' non-coding region, whereas the other introns were in the coding region, and their lengths were as follows: 157, 213, 87, 86, 135, 101, and 218 bp. All of the donor and acceptor sites of these introns were GT and AG, respectively, following the GT/AG rule as described by Breathnach & Chambon (1981) and Ming et al. (2010). For *Xc-HSP70* intron analysis, a 2,043 bp genomic DNA fragment was isolated by specific primers (P7, P8; Table 1), with a sequence identical to that of the *Xc-HSP70* cDNA, indicating that there was no intron in the coding region of the *Xc-HSP70* gene. Otherwise, compared with the cDNA and gDNA sequences of the cloned *Xc-HSC70* gene, the *Xc-HSC70* gene contained 8 introns (Fig. 3), wherein the longest intron lay in the 5' UTR, which also contained a heat shock element (HSE)-like core sequence (gaatatgCaGAAtgTTCaGaa) and other introns (with different lengths from 86 to 218 bp). This is the first report on the specific amount and sites of introns in *HSC70* of *X. c-nigrum*.

#### SIMILARITY ANALYSIS AND PHYLOGENETIC TREE CONSTRUCTION

Homology analysis revealed that 2 *Xc-HSP70s* were highly conserved in insects. The deduced amino acid sequences were highly simi-

lar to those of other known *HSP70s*. *Xc-HSC70* had higher than 88.1% similarity with other insect *HSC70* genes, 97.8% similarity with *HSC70* of *Mamestra brassicae* L. (Lepidoptera: Noctuidae) and 75% similarity with *HSC70* of vertebrates. Through the similarity analysis, we found that the *Xc-HSP70* genes had higher than 76.1% similarity with other insect *HSP70* genes, e.g., 88.8% identity with *HSP70* of *M. brassicae* and 86.3% identity with *HSP70* of *Helicoverpa zea* Boddie (Lepidoptera: Noctuidae), and 71.5% similarity with *HSP70* of vertebrates. The similarity between *Xc-HSC70* and *Xc-HSP70* was 83.5%. We also found that the sequences of *Xc-HSC70* and *Xc-HSP70* were more closely related to those of other *HSC70* and *HSP70* genes, respectively, in insects than to each other.

Phylogenetic tree construction showed that the *HSP* gene family could be divided into 2 main clusters: *HSC70* and *HSP70*. This classification was supported by a high degree of confidence, as shown in the evolutionary tree cluster (Fig. 4). *Xc-HSC70* belonged to the *HSC70* cluster, which also contained all 11 other *Xc-HSC70* genes. *Xc-HSP70* belonged to the *HSP70* cluster, which only contained all of the other *HSP70* genes. Therefore, this finding supports that *Xc-HSC70* is a constitutively expressed gene and that *Xc-HSP70* is an inducible gene, with a bootstrap value of 100% in 1,000 replicates. The relationships displayed in the phylogenetic tree were in general agreement with traditional taxonomy.




**Fig. 3.** Schematic structure of the *Xc-HSC70* gene. Exons are shown as boxes in which white boxes represent untranslated regions, whereas the black boxes are the protein-coding exons; introns are indicated as lines between the boxes. The numbers above and below the drawing represent the sizes (base pairs) of each exon and intron, respectively. The start codon (ATG) and stop codon (TAA) are also indicated. The genomic DNA sequence of *Xc-HSC70* has been deposited in GenBank under accession no. KF731994.

### EXPRESSION OF *XC-HSC70* AND *XC-HSP70* RNA IN RELATION TO THERMAL STRESS DURING DEVELOPMENT


The relative mRNA expression levels of *Xc-HSC70* induced by low and high temperatures during development were quantified by qRT-PCR (Fig. 5). *Xc-HSC70* expression was significantly up-regulated at temperatures of  $\leq -4$  or  $\geq 40$  °C in all developmental stages ( $P < 0.05$ ), and the *Xc-HSC70* expression level induced by a temperature of 37 °C in 5th instars

was significantly different from that in the controls ( $P < 0.05$ ). Therefore, the onset temperatures at which the expression level of *Xc-HSC70* was significantly higher than that of the controls during cold and heat shock in most developmental stages were  $-4$  and  $40$  °C, respectively. The intensity of the temperature response of *Xc-HSC70* varied among the developmental stages. Higher increases of 2.69 and 4.61 fold appeared in 2nd instars and pupae at  $-4$  and  $-7$  °C, respectively, under cold shock, whereas the greatest increases were observed in 2nd instars under heat shock, with a maximum increase of 3.52 fold (Fig. 5).

No significant increase in relative mRNA expression levels of *Xc-HSP70* were observed when larvae and pupae were stressed by temperatures of  $\geq 5$  °C or when larvae were stressed by temperatures of  $\leq 34$  °C (Fig. 6,  $P > 0.05$ ). However, *Xc-HSP70* was significantly up-regulated at temperatures of  $\leq 0$  °C or  $\geq 37$  °C ( $P < 0.05$ ). Therefore, the onset temperatures for *Xc-HSP70* up-regulation during cold and heat shock in all of the tested developmental stages were 0 and 37 °C, respectively. Higher increases of 6.54 and 12.47 fold appeared in 5th instars at  $-4$  °C and pupae at  $-7$  °C, respectively, under cold shock, whereas the greatest increases were observed in 5th instars under heat shock, with a maximum increase of 8.98 fold (Fig. 6). The intensity of the temperature response of *Xc-HSP70* varied among developmental stages,



**Fig. 4.** Phylogenetic tree of *Xc-HSC70* and *Xc-HSP70* amino acid sequences from different species. A 3-letter code has been included to indicate the order name of the corresponding insect and vertebrate orders (COL = Coleoptera, LEP = Lepidoptera, DIP = Diptera, HYM = Hymenoptera, and VER = Vertebrata). The values indicated on the branches correspond to bootstrap percentages (BP).



**Fig. 5.** *Xc-HSC70* mRNA expression profiles induced by cold ( $-7$  to  $5$  °C) and heat ( $37$  to  $47$  °C) in 2nd, 3rd, 4th, 5th, and 6th instars and pupae of *Xestia c-nigrum*. The relative quantities indicate the levels of the *HSC70* gene transcript normalized against transcript levels of  $\beta$ -actin as an internal standard and compared with the transcript levels of the untreated control at  $25$  °C. An asterisk indicates a significant difference between the control and heat shock conditions (significant, \*  $P < 0.05$ ). The data are denoted as the mean  $\pm$  SEM (error bar).




showing expression profiles similar to those of *Xc-HSC70* during development. *Xc-HSC70* was significantly up-regulated at temperatures of  $\leq -4$  °C or  $\geq 40$  °C ( $P < 0.05$ ). And the onset temperatures for *Xc-HSC70* up-regulation during cold and heat shock in all of the tested developmental stages were  $-4$  and  $40$  °C, respectively. Higher increases of 2.7 and 4.6 fold appeared in 2th instars at  $-4$  °C and pupae at  $-7$  °C, respectively, under cold shock, whereas the greatest increases were observed in 5th instars under heat shock, with a maximum increase of 3.5 fold (Fig. 5).

Comparative mRNA expression profiles of *Xc-HSC70* and *Xc-HSP70* induced by cold and heat during development indicated 3 obvious characteristics (Figs. 5 and 6). First, the onset temperature varied between *Xc-HSC70* and *Xc-HSP70*, depending on development stage in the case of heat shock but not cold shock, with a value of  $40$  °C for *Xc-HSC70* in most developmental stages and  $37$  °C for *Xc-HSP70* in all tested stages. Second, the temperature responses of *Xc-HSP70* at any given temperature or developmental stage were more intense than those of *Xc-HSC70*.


The expression levels of the 2 *X. c-nigrum* heat shock protein genes in different developmental stages at  $25$  °C were determined relative to the expression levels in 2nd instars. *Xc-HSC70* and *Xc-HSP70* expression decreased with the development of the larva and pupa without thermal stress (Fig. 7). Expression decreased substantially with larval age, and the lowest expression level was observed in 4th instars and then increased in 5th instars and 6th instars again. Expression of *Xc-HSC70* and *Xc-HSP70* did not change significantly in 4d and 6d-old pupa.

## Discussion

Cluster analysis supported the finding that *Xc-HSC70* and *Xc-HSP70* belong to 2 different branches of *HSP70* and *HSC70*, respectively. Some



**Fig. 6.** *Xc-HSP70* mRNA expression profiles induced by cold ( $-7$  to  $5$  °C) and heat ( $37$  to  $47$  °C) in 2nd, 3rd, 4th, 5th, and 6th instars and pupae of *Xestia c-nigrum*. The relative quantities indicate the levels of the *HSP70* gene transcript normalized against transcript levels of  $\beta$ -actin as an internal standard and compared with the transcript levels of the untreated control at  $25$  °C. An asterisk indicates a significant difference between the control and heat shock conditions (significant, \*  $P < 0.05$ ). The data are denoted as the mean  $\pm$  SEM (error bar).



**Fig. 7.** Expression levels of 2 *HSP70*s at different developmental stages relative to expression levels in 2nd instars at  $25$  °C. The data are denoted as the mean  $\pm$  SEM (error bar), and the different lowercase or uppercase letters indicate a significant difference in the means as assessed using multi-comparison tests ( $P < 0.05$ ).

studies have found that expression of *HSP70* increased significantly after the induction of environmental stress, but *HSC70* expression was lower than *HSP70* expression (Deane & Woo 2005). It has been suggested that the lack of introns in inducible *HSP70* genes could help to circumvent blocks in RNA splicing and enable preferential expression of heat shock proteins during periods of stress (Huang et al. 1999), thus protecting cells against harmful insults. In this study, genomic DNA sequence analysis revealed 8 introns in the coding region of the *Xc-HSC70* gene, representative of constitutive features, whereas the *Xc-HSP70* gene did not contain any introns, representative of inducible features. All of the intron borders of the *Xc-HSC70* gene start and end with the consensus GT and AG splicing signals (Breathnach & Chambon 1981). *Xc-HSP70* was significantly different from *Xc-HSC70* with regard to expression amount and expression time, and it displayed an increased rate of expression. The qRT-PCR results suggested that *Xc-HSC70* was constitutively expressed and that *Xc-HSP70* was induced, possibly because the *Xc-HSP70* gene had no introns and *Xc-HSC70* had introns, which affect the splicing and joining of mRNA (Yost & Lindquist 1991; Chuang et al. 2007).

The heat shock response is characterized by the induction of numerous HSPs (Balch et al. 2008) and is mostly regulated at the transcription level by heat shock transcription factors (HSFs), which can specifically bind to heat shock elements (HSEs) in the promoters of heat shock genes (Morimoto 1998). HSEs are composed of at least 3 inverted repeats of the consensus sequence nGAAn (Fernandes et al. 1994). The longest intron of *Xc-HSC70* lies in the 5' UTR, which also contains an HSE-like core sequence (aGAATatgcaGAAtgttccaGAAa). The HSFs bind to the HSEs and initiate the transcription of the *HSP70*s (Morimoto 1993). Although we did not measure the *HSC70* or *HSP70* protein level in this experiment, the precondition of higher levels of *HSC70* or *HSP70* is the higher abundance of *HSC70* or *HSP70* mRNA, respectively, because the synthesis of *HSP70*s is primarily regulated at the transcriptional level (Molina et al. 2000). To sustain a proper level of *HSP* mRNA, the transcription of *HSP* genes can be controlled by negative regulation of HSPs. When HSPs reach high levels, HSFs bind to the HSPs, which blocks DNA binding to *HSE* genes (Morimoto 1993). The difference in mRNA expression between the *Xc-HSC70* and *Xc-HSP70* genes may be related to unique structures (Ali et al. 2003), functions, and regulatory characteristics (Park et al. 2001; Kregel 2002).

Although heat shock was the first stress shown to elicit synthesis of HSPs, it is now evident that various environmental, physiological, and



genetic factors (Sørensen et al. 2003), even cold shock (Goto & Kimura 1998), can regulate the expression of these same proteins. Therefore, the regulation of inducible *HSP70* genes plays an important role in cellular responses. The function of *HSP* genes in thermotolerance has been confirmed in various organisms (Sonna et al. 2002; Huang & Kang 2007). In *X. c-nigrum*, *Xc-HSC70* and *Xc-HSP70* were positively induced during development when the insect was stressed, regardless of heat or cold, suggesting that up-regulation of these genes is essential for thermotolerance in *X. c-nigrum*.

The level of *Xc-HSP70* expression and transcription was increased after thermal stress treatments in different developmental stages, which suggests that *Xc-HSP70s* are closely associated with insect resistance to thermal stress (Sonoda et al. 2006; Zhang & Denlinger 2010). The intensity of the temperature responses also varies among *HSP70* genes in different insects. Heat induction resulted in an expression increase of approximately 6 fold for the *Xc-HSP70* gene (Figs. 5 and 6). This finding is peculiar because in many other insects, including some Lepidoptera species, heat up-regulates *HSP70* expression by more than 100 fold. Regarding the 6-fold induction, we consider that the experimental materials, which were taken from the northernmost province of China in Heilongjiang Province (Habin, 130°10'N, 46°40'E), may be an important factor. Some studies have shown that production of *HSP70* in response to temperature shock is less intense in organisms that are more frequently exposed to unfavorable temperatures in their habitat than in organisms in benign conditions (Sørensen et al. 2001; Lyytinen et al. 2012). As the synthesis of *HSP70s* requires considerable energy and may thus occur at the cost of the synthesis of other proteins (Krebs & Loeschcke 1994), northern populations might have evolved a less costly way to resist cold stress (Lansing et al. 2000; Sørensen et al. 2003). *HSP70s* may not be the only proteins involved in heat and cold resistance; other proteins such as *HSP20*, *HSP60*, and *HSP90* may also be involved in heat resistance (Zhang & Denlinger 2010; Wang et al. 2012) or in the accumulation of low-molecular-weight cryoprotectants (Crowe et al. 1988), synthesis of antifreeze proteins (Duman 2001), and remodeling of the structure of the cell membrane (Tomcala et al. 2006). Some studies have found *HSP70* to be up-regulated 2–5 fold in insects (Colinet et al. 2010; Morales et al. 2011; Shu et al. 2011; Wang et al. 2012; Yu et al. 2012; Luo et al. 2014).

The onset temperature at which *HSP70* gene expression is up-regulated, whether low or high, varies among organisms (Garbuz et al. 2003). Tomanek & Somero (1999) suggested that the threshold temperature may be a useful way to identify temperature tolerance limits and that a higher onset temperature is associated with high heat tolerance, and vice versa. Findings in *Drosophila* (Garbuz et al. 2003) and leaf miner species (Huang & Kang 2007) support this idea. In *X. c-nigrum*, the relatively high onset temperature of 40 and 37 °C for *Xc-HSC70* and *Xc-HSP70*, respectively, in response to heat shock may represent an indicator of heat tolerance. In contrast, the values of –4 and 0 °C for *Xc-HSC70* and *Xc-HSP70*, respectively, under cold shock may suggest only weak cold tolerance for this species. Furthermore, the difference in onset temperature between the 2 genes under heat shock, i.e., 40 °C in most developmental stages for *Xc-HSC70* and 37 °C in all of the tested developmental stages for *Xc-HSP70*, may reflect different functions. That is, inducible expression of *HSP70* but not *HSC70* at “normal” temperatures in organisms may play a negative role in cell growth and division. Expression of *Xc-HSP70* was not significantly different across a wide temperature range from 26 to 40 °C, which is consistent with the negative effect of *HSP70* overexpression on growth, survival, and fecundity observed in other insects (Krebs & Loeschcke 1994).

Finally, the basal *HSP70* gene expression profiles varied with developmental stage, suggesting that these genes may be involved in development (Huang et al. 2009). Interestingly, the relative accumulated

amounts of mRNA from large *HSP70s*, including *HSC70* and *HSP70*, may increase or decrease as larval and pupal development progress, depending on the species (Sonoda et al. 2006). In *X. c-nigrum*, the relative transcript levels of *Xc-HSC70* and *Xc-HSP70* decreased with the developmental progress of the pupa. These results are consistent with research conducted on *Pteromalus puparum* (L.) (Hymenoptera: Pteromalidae) and the pyrrhocorid *P. apterus* (Kostal & Tollarova-Borovanska 2009; Wang et al. 2012). Furthermore, gene expression profiles for induced *HSP70s* also varied after heat or cold shock during development. Mahroof et al. (2005) suggested that increased mRNA abundance induced under thermal stress during different developmental stages may contribute to the increased thermotolerance in those stages. Increased thermotolerance in young larvae of *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae) was suggested to result from increased expression of *HSP70s* (Mahroof et al. 2005). In *X. c-nigrum*, higher *Xc-HSC70* and *Xc-HSP70* expression were observed in heat and cold shocked young larvae and pupae, consistent with high heat resistance in larvae and stronger cold tolerance in young larvae and pupae. Therefore, our findings that correspond to thermal acclimation for *X. c-nigrum* during development based on *HSP70* gene expression levels are consistent with temperature tolerances observed in field populations.

## Acknowledgments

This work was supported by the Special Fund for Modern Agro-industry Technology Research System: Soybean Technology Research System (CARS-04) and the Fund of Common Wealth Industry (Agriculture) Special Research (201103002).

## References Cited

- Ali KS, Dorgai L, Ábrahám M, Hermesz E. 2003. Tissue- and stressor-specific differential expression of two hsc70 genes in carp. *Biochemical and Biophysical Research Communications* 307: 503-509.
- Balch WE, Morimoto RI, Dillin A, Kelly JW. 2008. Adapting proteostasis for disease intervention. *Science* 319: 916-919.
- Bettencourt BR, Hogan CC, Nimali M, Drohan BW. 2008. Inducible and constitutive heat shock gene expression responds to modification of Hsp70 copy number in *Drosophila melanogaster* but does not compensate for loss of thermotolerance in Hsp70 null flies. *BMC Biology* 6: 5.
- Boutet I, Tanguy A, Rousseau S, Auffret M, Moraga D. 2003. Molecular identification and expression of heat shock cognate 70 (*hsc70*) and heat shock protein 70 (*hsp70*) genes in the Pacific oyster *Crassostrea gigas*. *Cell Stress Chaperones* 8: 76-85.
- Breathnach R, Chambon P. 1981. Organization and expression of eukaryotic split genes coding for proteins. *Annual Review of Biochemistry* 50: 349-383.
- Choi BG, Hapat R, Kim Y. 2014. RNA interference of a heat shock protein, Hsp70, loses its protection role in indirect chilling injury to the beet armyworm, *Spodoptera exigua*. *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology* 168: 90-95.
- Chuang K, Ho S, Song Y. 2007. Cloning and expression analysis of heat shock cognate 70 gene promoter in tiger shrimp (*Penaeus monodon*). *Gene* 405: 10-18.
- Colinet H, Lee SF, Hoffmann A. 2010. Temporal expression of heat shock genes during cold stress and recovery from chill coma in adult *Drosophila melanogaster*. *FEBS Journal* 277: 174-185.
- Crowe JH, Crowe LM, Carpenter JF, Rudolph AS, Wistrom CA, Spargo BJ, Anchordoguy TJ. 1988. Interactions of sugars with membranes. *Biochimica et Biophysica Acta* 947: 367-384.
- Deane EE, Woo NYS. 2005. Cloning and characterization of the hsp70 multigene family from silver sea bream: modulated gene expression between warm and cold temperature acclimation. *Biochemical and Biophysical Research Communications* 330: 776-783.
- Duman JG. 2001. Antifreeze and ice nucleator proteins in terrestrial arthropods. *Annual Review of Physiology* 63: 327-357.
- Feder ME, Hofmann GE. 1999. Heat-shock proteins, molecular chaperones, and the stress response: evolutionary and ecological physiology. *Annual Review of Physiology* 61: 243-282.

- Fernandes M, Xiao H, Lis JT. 1994. Fine structure analysis of the *Drosophila* and *Saccharomyces* heat shock factor–heat shock element interactions. *Nucleic Acids Research* 22: 167-173.
- Franke K, Heitmann N, Tobner A, Fischer K. 2014. Fitness costs associated with different frequencies and magnitudes of temperature change in the butterfly *Bicyclus anynana*. *Journal of Thermal Biology* 41: 88-94.
- Franzellitti S, Fabbri E. 2005. Differential HSP70 gene expression in the Mediterranean mussel exposed to various stressors. *Biochemical and Biophysical Research Communications* 336: 1157-1163.
- Garbuz D, Evgenev MB, Feder ME, Zatzepina OG. 2003. Evolution of the thermotolerance and heat-shock response: evidence from inter/intraspecific comparison and interspecific hybridization in the *virilis* species group of *Drosophila*. I. Thermal phenotype. *The Journal of Experimental Biology* 206: 2392-2408.
- Goto SG, Kimura MT. 1998. Heat- and cold-shock responses and temperature adaptations in subtropical and temperate species of *Drosophila*. *Journal of Insect Physiology* 44: 1233-1239.
- Gray EM. 2013. Thermal acclimation in a complex life cycle: the effects of larval and adult thermal conditions on metabolic rate and heat resistance in *Culex pipiens* (Diptera: Culicidae). *Journal of Insect Physiology* 59: 1001-1007.
- Huang LH, Kang L. 2007. Cloning and interspecific altered expression of heat shock protein genes in two leafminer species in response to thermal stress. *Insect Molecular Biology* 16: 491-500.
- Huang LH, Wang HS, Kang L. 2008. Different evolutionary lineages of large and small heat shock proteins in eukaryotes. *Cell Research* 18: 1074-1076.
- Huang LH, Wang CZ, Kang L. 2009. Cloning and expression of five heat shock protein genes in relation to cold hardening and development in the leafminer, *Liriomyza sativa*. *Journal of Insect Physiology* 55: 279-285.
- Huang Y, Wimler KM, Carmichael GG. 1999. Intronless mRNA transport elements may affect multiple steps of pre-mRNA processing. *EMBO Journal* 18: 1642-1652.
- Jensen P, Overgaard J, Loeschcke V, Schou MF, Malte H, Kristensen TN. 2014. Inbreeding effects on standard metabolic rate investigated at cold, benign and hot temperatures in *Drosophila melanogaster*. *Journal of Insect Physiology* 62: 11-20.
- Jiang XF, Zhai HF, Wang L, Luo LZ, Thomas WS, Zhang L. 2012. Cloning of the heat shock protein 90 and 70 genes from the beet armyworm, *Spodoptera exigua*, and expression characteristics in relation to thermal stress and development. *Cell Stress Chaperones* 17: 67-80.
- Kostal V, Tollarova-Borovanska M. 2009. The 70 kDa heat shock protein assists during the repair of chilling injury in the insect *Pyrrhocoris apterus*. *PLOS ONE* 4: e4546.
- Krebs RA, Loeschcke V. 1994. Costs and benefits of activation of the heat-shock response in *Drosophila melanogaster*. *Functional Ecology* 8: 730-737.
- Kregel KC. 2002. Heat shock proteins: modifying factors in physiological stress responses and acquired thermotolerance. *Journal of Applied Physiology* 92: 2177-2186.
- Landolt PJ, Guédot C, Zack RS. 2011. Spotted cutworm, *Xestia c-nigrum* (L.) (Lepidoptera: Noctuidae) responses to sex pheromone and blacklight. *Journal of Applied Entomology* 135: 593-600.
- Lansing E, Justesen J, Loeschcke V. 2000. Variation in the expression of Hsp70, the major heat-shock protein, and thermotolerance in larval and adult selection lines of *Drosophila melanogaster*. *Journal of Thermal Biology* 25: 443-450.
- Lindquist S, Craig EA. 1988. The heat-shock proteins. *Annual Review of Genetics* 22: 631-677.
- Liu XY, Ma XH, Lei CF, Xiao YZ, Zhang ZX, Sun XL. 2011. Synergistic effects of *Cydia pomonella* granulovirus GP37 on the infectivity of nucleopolyhedroviruses and the lethality of *Bacillus thuringiensis*. *Archives of Virology* 156: 1707-1715.
- Livak KJ, Schmittgen TD. 2001. Analysis of temporal gene expression data using real-time quantitative PCR and the 2<sup>-ΔΔC<sub>T</sub></sup> (T) Method. *Methods* 25: 402-408.
- Luo SQ, Wong SC, Xu CR, Hanski I, Wang JR, Lehtonen R. 2014. Phenotypic plasticity in thermal tolerance in the Glanville fritillary butterfly. *Journal of Thermal Biology* 42: 33-39.
- Lyytinen A, Mappes J, Lindström L. 2012. Variation in Hsp70 levels after cold shock: signs of evolutionary responses to thermal selection among *Leptinotarsa decemlineata* populations. *PLOS ONE* 7: e31446.
- Mahroof R, Zhu KY, Subramanyam B. 2005. Changes in expression of heat shock proteins in *Tribolium castaneum* (Coleoptera: Tenebrionidae) in relation to developmental stage, exposure time, and temperature. *Annals of the Entomological Society of America* 98: 100-107.
- Ming JH, Xie J, Xu P, Liu WB, Ge XP, Liu B, He YJ, Cheng YF, Zhou QL, Pan LK. 2010. Molecular cloning and expression of two HSP70 genes in the Wuchang bream (*Megalobrama amblycephala* Yih). *Fish and Shellfish Immunology* 28: 407-418.
- Molina A, Biemar F, Müller F, Iyengar A, Prunet P, Maclean N, Martial JA, Muller M. 2000. Cloning and expression analysis of an inducible HSP70 gene from tilapia fish. *FEBS Letters* 474: 5-10.
- Morales M, Planelló R, Martínez-Paz P, Herrero O, Cortés E, Martínez-Guitarte JL, Morcillo G. 2011. Characterization of Hsp70 gene in *Chironomus riparius*: expression in response to endocrine disrupting pollutants as a marker of ecotoxicological stress. *Comparative Biochemistry and Physiology Part C: Toxicology and Pharmacology* 153: 150-158.
- Morimoto RI. 1993. Cell in stress: transcriptional activation of heat shock genes. *Science* 259: 1409-1410.
- Morimoto RI. 1998. Regulation of the heat shock transcriptional response: cross talk between a family of heat shock factors, molecular chaperones, and negative regulators. *Genes and Development* 12: 3788-3796.
- Mukawa S, Goto C. 2010. *Mamestra brassicae* nucleopolyhedrovirus infection and enhancing effect of proteins derived from *Xestia c-nigrum* granulovirus on larvae of *Mamestra brassicae* and *Helicoverpa armigera* (Lepidoptera: Noctuidae) on cabbage. *Journal of Economic Entomology* 103: 257-264.
- Mukawa S, Goto C. 2011. Enhancing effect of proteins derived from *Xestia c-nigrum* granulovirus on *Mamestra brassicae* nucleopolyhedrovirus infection in larvae of *Autographa nigrisigna* (Lepidoptera: Noctuidae) on cabbage. *Applied Entomology and Zoology* 46: 55-63.
- Park JH, Lee JJ, Yoon S, Lee JS, Choe SY, Choe J, Park EH, Kim CG. 2001. Genomic cloning of the Hsc71 gene in the hermaphroditic teleost *Rivulus marmoratus* and analysis of its expression in skeletal muscle: identification of a novel muscle-preferred regulatory element. *Nucleic Acids Research* 29: 3041-3050.
- Rinehart JP, Li A, Yocum GD, Robich RM, Hayward SAL, Denlinger DL. 2007. Up-regulation of heat shock proteins is essential for cold survival during insect diapause. *Proceedings of the National Academy of Sciences of the USA* 104: 11130-11137.
- Shen Y, Gong YJ, Gu J, Huang LH, Feng QL. 2014. Physiological effect of mild thermal stress and its induction of gene expression in the common cutworm, *Spodoptera litura*. *Journal of Insect Physiology* 61: 34-41.
- Shu YH, Du Y, Wang JW. 2011. Molecular characterization and expression patterns of *Spodoptera litura* heat shock protein 70/90, and their response to zinc stress. *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology* 158: 102-110.
- Sonna LA, Fujita J, Gaffin SL, Lilly CM. 2002. Invited review: effects of heat and cold stress on mammalian gene expression. *Journal of Applied Physiology* 92: 1725-1742.
- Sonoda S, Fukumoto K, Izumi Y, Yoshida H, Tsumuki H. 2006. Cloning of heat shock protein genes (*hsp90* and *hsc70*) and their expression during larval diapause and cold tolerance acquisition in the rice stem borer, *Chilo suppressalis* Walker. *Archives of Insect Biochemistry and Physiology* 63: 36-47.
- Sørensen JG, Dahlgraad J, Loeschcke V. 2001. Genetic variation in thermal tolerance among natural populations of *Drosophila buzzatii*: down regulation of Hsp70 expression and variation in heat stress resistance traits. *Functional Ecology* 15: 289-296.
- Sørensen JG, Kristensen TN, Loeschcke V. 2003. The evolutionary and ecological role of heat shock proteins. *Ecology Letters* 6: 1025-1037.
- Tomanek L, Somero GN. 1999. Evolutionary and acclimation-induced variation in the heat-shock responses of congeneric marine snails (genus *Tegula*) from different thermal habitats: implications for limits of thermotolerance and biogeography. *The Journal of Experimental Biology* 202: 2925-2936.
- Tomcala A, Tollarová M, Overgaard J, Simek P, Kostál V. 2006. Seasonal acquisition of chill tolerance and restructuring of membrane glycerophospholipids in an overwintering insect: triggering by low temperature, desiccation and diapause progression. *The Journal of Experimental Biology* 209: 4102-4114.
- Wang H, Li K, Zhu JY, Fang Q, Ye GY, Wang H, Li K, Zhu JY. 2012. Cloning and expression pattern of heat shock protein genes from the endoparasitoid wasp, *Pteromalus puparum* in response to environmental stresses. *Archives of Insect Biochemistry and Physiology* 79: 247-263.
- Wang L, Yang S, Zhu MH, Xu ZX, Zhao KJ, Han LL. 2014. Analysis of transcription differences of the heat shock cognate 70 gene in different tissues in response to heat stress in *Xestia c-nigrum* Linnaeus (Lepidoptera: Noctuidae). *Chinese Journal of Applied Entomology* 51: 1014-1025. (In Chinese)
- Wang L, Yang S, Zhao KJ, Han LL. 2015. Expression Profiles of the Heat Shock Protein 70 Gene in Response to Heat Stress in *Agrotis c-nigrum* (Lepidoptera: Noctuidae). *Journal of Insect Science* DOI: 10.1093/jisesa/ieu169.
- Welch WJ. 1993. How cells respond to stress. *Scientific American* 268(5): 56-64.

- Xi JH, Pan HY, Chen YJ, Guo YL. 2002. Influence of four food plants on development and fecundity of the spotted cutworm, *Agrotis c-nigrum*. Chinese Journal of Applied Entomology 39: 428-429. (In Chinese)
- Yost HJ, Lindquist S. 1991. Heat shock proteins affect RNA processing during the heat shock response of *Saccharomyces cerevisiae*. Molecular and Cellular Biology 11: 1062-1068.
- Yu H, Wan FH, Guo JY. 2012. cDNA cloning of heat shock protein genes and their expression in an indigenous cryptic species of the whitefly *Bemisia tabaci* complex from China. Journal of Integrated Agriculture 11: 293-302.
- Zhang Q, Denlinger DL. 2010. Molecular characterization of heat shock protein 90, 70 and 70 cognate cDNAs and their expression patterns during thermal stress and pupal diapause in the corn earworm. Journal of Insect Physiology 56: 138-150.
- Zhang SJ, Qian XJ, Li CJ, Pan FJ, Xu YL. 2013. Pathogenicity of entomopathogenic nematode on *Xestia c-nigrum* in soybean field. Soybean Science 32: 63-67. (In Chinese)
- Zheng WW, Yang DT, Wang JX, Song QS, Gilbert LI, Zhao XF. 2010. Hsc70 binds to ultraspiracle resulting in the upregulation of 20-hydroxyecdysone-responsive genes in *Helicoverpa armigera*. Molecular and Cellular Endocrinology 315: 282-291.
- Zheng YR, Wang WS. 2010. Biological characteristics were observed of *Agrotis c-nigrum* in the north Chinese. JI LIN Agricultural 12: 109. (In Chinese)