

Thrips (Thysanoptera) Collected from Solanum dulcamara (Solanales: Solanaceae) in Washington and Idaho

Authors: Carrillo, Carmen I. Castillo, Funderburk, Joseph, and Snyder, William E.

Source: Florida Entomologist, 99(2): 306-307

Published By: Florida Entomological Society

URL: https://doi.org/10.1653/024.099.0225

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Thrips (Thysanoptera) collected from Solanum dulcamara (Solanales: Solanaceae) in Washington and Idaho

Carmen I. Castillo Carrillo^{1,*}, Joseph Funderburk², and William E. Snyder¹

Bittersweet nightshade, *Solanum dulcamara* L. (Solanales: Solanaceae), is a native plant in parts of Europe and Asia. It was introduced and is established in the eastern, north-central, and Pacific Northwest regions of the USA. It is commonly found in grasslands and meadows and occurs most frequently in riparian areas, wetlands, and deciduous forests (Waggy 2009). We report here various thrips (Thysanoptera) species collected from samples of *S. dulcamara* in Washington and Idaho.

Patches of *S. dulcamara* were found along irrigation canals and ponds near cultivated areas. Samples were taken with a D-Vac sucking device (Model 24, Rincon-Vitova Insectaries, Inc., Ventura, California) from the entire above-ground portions of the plants from Jun to Nov in 2012 and in Aug and Oct in 2013 (Table 1). *Solanum dulcamara* grows using other plants as support. Seventeen species of surrounding plants were identified, including *Elaeagnus angustifolia* L. (Rosales: Elaeagnaceae), *Typha* sp. (Poales: Typhaceae), *Asclepias syriaca* L. (Gentianales: Apocynaceae), and *Salix* sp. (Malpighiales: Salicaceae). The D-Vac tube was placed over the plant patch for approximately 10 s. Subsamples were taken depending on the size of the patch. Insects collected in the mesh bag were placed on dry ice for transport to the laboratory. Only adults were sorted and identified to species. However, thrips larvae of unidentified species were observed on the plants from Colfax in Jun 2013, Moses Lake in Jul and Aug, Mesa (new) in Jun, and Mattawa in Jun and Sep. Thrips were extracted under stereomicroscopy in 90% ethyl alcohol. Adults were mounted in Canada balsam for identification using the procedures in Wirth & Marston (1968) and the keys in Hoddle et al. (2012). Vouchers of each species were deposited at the North Florida Research and Education Center, University of Florida, Quincy, Florida.

Adults of 8 species of thrips from 3 families were collected (Table 2). Species of Thripidae included the phytophagous *Caliothrips fasciatus* (Pergande), *Chirothrips aculeatus* Bagnall, *Frankliniella occidentalis* (Pergande), *Thrips hawaiiensis* (Morgan), and *Thrips tabaci* Lindeman. Both *F. occidentalis* and *T. tabaci* are worldwide pests of many crops, and they are vectors of the plant viruses in the genus *Tospovirus* (Bunyaviridae) (Hoddle et al. 2012). *Solanum dulcamara* is a confirmed host for *Tomato spotted wilt virus* (Parrella et al. 2003), and *F. occidentalis* is a confirmed vector of this tospovirus species to *S. dulcamara* (Stobbs et al. 1992).

Adults of 2 species of Phlaeothripidae were collected (Table 2). *Haplothrips verbasci* (Osborn) reproduces on the stems and flowers of *Verbascum thapsus* L. (Lamiales: Scrophulariaceae) (Comegys & Schmitt 1965; Hoddle et al. 2012; Wilbur et al. 2013). This plant species was identified growing near *S. dulcamara* in eastern Washington.

Site name	Coordinates	2012	Plant stage ^a	2013	Plant stage [®]	
Twin Falls, ID	42.4992583°N, 114.1540361°W	14 Jun	Flo			
	42.4978139°N, 114.1535000°W	10 Jul	Flo & Fru			
Mesa (old), WA	46.5882556°N, 119.0003111°W	28 Jun	Flo	24 Aug	Flo & Fru	
		17 Jul	Flo			
		16 Aug	Flo			
		4 Nov	Flo & Fru			
Mesa (new), WA	46.5764861°N, 119.0092806°W	2 Aug	Flo	26 Oct	Flo & Fru	
		4 Nov	Fru			
Colfax, WA	46.8475280°N, 117.4788700°W	27 Sep	Flo & Fru			
		6 Oct	Fru			
Moses Lake, WA	46.9928306°N, 119.6851833°W			24 Aug	Fru	
	46.9986389°N, 119.6847917°W					
Mattawa, WA	46.7089806°N, 119.9451500°W	4 Sep	Flo	24-Aug	Flo & Fru	
				26 Oct	Flo & Fru	
Sacajawea, WA	46.2035470°N, 119.0471290°W	4 Nov	Fru			

 Table 1. Sample locations, coordinates, sampling dates, and plant stage.

^aFlo = flowering, Fru = fruiting

¹Washington State University, Department of Entomology, Pullman, Washington 99164-1120, USA

²University of Florida, North Florida Research & Education Center, 155 Research Rd., Quincy, Florida 32351-5677, USA

^{*}Corresponding author; E-mail: c.castillocarrillo@wsu.edu

Scientific Notes

 Table 2. Species of thrips and total numbers of adults found sampling Solanum dulcamara patches in Idaho and Washington.

	Sampling places ^a							
Thrips species	TF	MO	MN	СХ	ML	MT	SJ	
Aeolothrips fasciatus	3	2		1				
Caliothrips fasciatus	1	4	1	1	1	46		
Cephalothrips monilicornis	4	7	1	1			1	
Chirothrips aculeatus		1						
Frankliniella occidentalis	9	2				2	2	
Haplothrips verbasci			2					
Thrips hawaiiensis		1						
Thrips tabaci						1		

*TF = Twin Falls, MO = Mesa (old), MN = Mesa (new), CX = Colfax, ML = Moses Lake, MT = Mattawa, SJ = Sacajawea

Cephalothrips monilicornis (Reuter) reproduces on the leaves of various Poaceae (Hoddle et al. 2012). Four species of Poaceae were identified growing close to *S. dulcamara*. *Aeolothrips bicolor* Hinds in the family Aeolothripidae was collected. Insects in the order Thysanoptera are mainly phytophagous or mycophagous, and obligate predation is limited to only several lineages (Mound 2005). Species of *Aeolothrips* are predatory on small insects including other species of thrips.

A host plant is one on which an insect can reproduce (Mound 2013), and more research is needed to determine which species of thrips utilize *S. dulcamara* as a host plant. Our results suggest that the plant is potentially a source of economically important thrips invading crop fields.

We thank the Northwest Potato Research Consortium for supporting this research, and Marcus Hooker for identifying the plants (Marion Ownbey Herbarium, School of Biological Science, Washington State University).

Summary

Bittersweet nightshade, *Solanum dulcamara* L. (Solanales: Solanaceae), was sampled at numerous locations in Washington and Idaho. Adults of 8 species of thrips (Thysanoptera) from 3 families were collected, including the worldwide plant pests *Frankliniella occidentalis* (Pergande) and *Thrips tabaci* Lindeman (Thripidae), which are vectors of the serious plant viruses in the genus *Tospovirus*. *Aeolothrips fasciatus* Hinds (Aeolothripidae), a predator of small insects, also was collected.

Key Words: bittersweet nightshade; Pacific Northwest; Thripidae; Phlaeothripidae; Aeolothripidae

Sumario

Un muestreo de plantas hierba mora o dulcamara, Solanum dulcamara L. (Solanales: Solanaceae), fue realizado en varias localidades de Washington y Idaho. Adultos de ocho especies de trips (Thysanoptera) fueron colectados, incluidas las especies Frankliniella occidentalis (Pergande) y Thrips tabaci Lindeman (Thripidae) que son vectores de virus del género Tospovirus en cultivos agrícolas a nivel mundial. También se colectó Aeolothrips fasciatus (L.), especie que es depredadora de pequeños insectos.

Palabras Clave: hierba mora; dulcamara; Noroeste del Pacífico; Thripidae; Phlaeothripidae; Aeolothripidae

References Cited

- Comegys GR, Schmitt JB. 1965. A list of the Thysanoptera or thrips of New Jersey. Journal of the New York Entomological Society 1: 195–222.
- Hoddle MS, Mound LA, Paris DL. 2012. Thrips of California. CBIT Publishing, Queensland, Australia.
- Mound LA. 2005. Thysanoptera diversity and interactions. Annual Review of Entomology 50: 247–269.
- Mound LA. 2013. Homologies and host-plant specificity: recurrent problems in the study of thrips. Florida Entomologist 96: 318–322.
- Parrella G, Gognalons P, Gebre-Selassie K, Marchouz G, Vovlas C. 2003. An update of the host range of *Tomato spotted wilt virus* (*Lycopersicon esculentum* Mill.). Journal of Plant Pathology 85: 227–264.
- Stobbs LW, Broadbent AB, Allen WR, Stirling AL. 1992. Transmission of *Tomato spotted wilt virus* by the western flower thrips to weeds and native plants found in southern Ontario. Plant Disease 76: 23–29.
- Waggy M. 2009. Solanum dulcamara. In Fire Effects Information System [online]. United States Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). http://www. fs.fed.us/database/feis/ (last accessed 16 Dec 2015).
- Wilbur H, Hufbauer RA, Alba C, Norton A. 2013. The effect of insect herbivory on the growth and fitness of introduced *Verbascum thapsus* L. Neobiota 19: 21–44.
- Wirth WW, Marston N. 1968. A method for mounting small insects on microscope slides in Canada balsam. Annals of the Entomological Society of America 61: 783–784.