

# Sampling Plan for Bemisia tabaci (Hemiptera: Aleyrodidae) in Melon Crops

Authors: Macêdo, Ruan Victor Borges Tavares de, Sarmento, Renato Almeida, Pereira, Poliana Silvestre, Lima, Carlos Henrique Oliveira, Deus, Tânia Lígia Lopes Barbosa de, et al.

Source: Florida Entomologist, 102(1): 16-23

Published By: Florida Entomological Society

URL: https://doi.org/10.1653/024.102.0103

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Sampling plan for *Bemisia tabaci* (Hemiptera: Aleyrodidae) in melon crops

Ruan Victor Borges Tavares de Macêdo<sup>1</sup>, Renato Almeida Sarmento<sup>2,\*</sup>, Poliana Silvestre Pereira<sup>2</sup>, Carlos Henrique Oliveira Lima<sup>2</sup>, Tânia Lígia Lopes Barbosa de Deus<sup>2</sup>, Arthur Vieira Ribeiro<sup>1</sup>, and Marcelo Coutinho Picanço<sup>1</sup>

#### Abstract

Melon (*Cucumis melo* L.; Cucurbitaceae) is one of the 10 most-consumed fruits in the world. The whitefly *Bemisia tabaci* (Gennadius) (Hemiptera: Aleyrodidae) is one of the most important pests in melon crops worldwide. Conventional sampling plans are the starting point to establish decision-making systems for integrated pest management (IPM) programs. The purpose of this study was to determine a conventional sampling plan for *B. tabaci* in melon crops with plants at the vegetative, flowering, and fruiting stages. The best sampling units for *B. tabaci* were the 5th and 6th most apical leaves of the plant vine. The best sampling technique was direct counting of adult whiteflies. The most appropriate frequency distribution to describe *B. tabaci* densities in melon fields was the negative binomial. Whiteflies on melon fields with plants at different phenological stages showed a common aggregation parameter (Kcommon = 0.9134). The optimal number of samples from the sampling plan was 72 samples per field with a maximum error of 15% in population determination. The sampling plan determined by this study can be used by farmers because it is a low-cost (US\$5.27 per sampling), fast (39 minutes per sampling) and feasible (15% maximum evaluation error). The same sampling plan can be used with melon plants at the vegetative, flowering, and fruiting stages.

Key Words: Cucumis melo L.; sampling unit; sampling technique; negative binomial; decision-making

#### Resumen

El melón (*Cucumis melo* L.; Cucurbitaceae) es una de las 10 frutas más consumidas en el mundo. La mosca blanca *Bemisia tabaci* (Gennadius) (Hemiptera: Aleyrodidae) es una de las plagas más importantes en los cultivos de melón en todo el mundo. Los planes de muestreo convencionales son el punto de partida para establecer sistemas de toma de decisiones para los programas de manejo integrado de plagas (MIP). El propósito de este estudio fue determinar un plan de muestreo convencional para *B. tabaci* en cultivos de melón con plantas durante de las etapas vegetativa, de floración y de fructificación. Las mejores unidades de muestreo para *B. tabaci* fueron la quinta y la sexta hojas más apicales de la planta enredadera. La mejor técnica de muestreo fue el conteo directo de adultos. La distribución de frecuencia más adecuada para describir las densidades de *B. tabaci* en los campos de melón fue el binomio negativo. Las moscas blancas en los campos de melón con plantas en diferentes estados fenológicos mostraron un parámetro de agregación común (Kcommon = 0.9134). El número óptimo de muestras del plan de muestreo fue de 72 muestras por campo con un error máximo del 15% en la determinación de la población. El plan de muestreo determinado por este estudio puede ser utilizado por los agricultores po su bajo costo (US\$5,27 por muestreo), rápido (39 minutos por muestreo) y factible (15% de error de evaluación máximo). El mismo plan de muestreo se puede usar con plantas de melón en las etapas vegetativa, de floración y de fructificación.

Palabras Clave: Cucumis melo L.; unidad de muestreo; técnica de muestreo; binomio negativo; toma de decisiones

Melon (*Cucumis melo* L.; Cucurbitaceae), often called muskmelon or honeydew melon, is one of the 10 most-consumed fruits in the world (3.6% of global fruit consumption). Global production is 29.6 million tons per year (FAO 2014). In Brazil, cultivation of *C. melo* varies from 55 to 75 days and during this period it can generate a profit of US\$6,364 per ha, as well as 1 direct and 3 indirect jobs per ha of cultivation (Freitas et al. 2009).

One of the principal pests occurring in melon crops is the whitefly *Bemisia tabaci* (Gennadius) (Hemiptera: Aleyrodidae) (Picanço et al. 2002; Braga Sobrinho et al. 2011; Pessarakli 2016), which is a polyphagous species that occurs in all cultivated regions worldwide (Stansly & Naranjo 2010; CABI 2016). This pest has approximately 700 reported

host plants, mainly from the families Asteraceae, Brassicaceae, Cucurbitaceae, Fabaceae, Malvaceae, and Solanaceae (Greathead 1986; Li et al. 2011). It causes damage to crops due to sap sucking, injection of toxins into the vascular system, and transmission of viruses, especially geminiviruses (Blackmer & Byrne 1999; Toscano et al. 2004; Varma et al. 2011). Additionally, opportunistic fungi develop on the sweet excretions (honeydew) of *B. tabaci*, forming a dark layer of fungi that covers the plants (sooty mold) resulting in a reduction of photosynthesis and depreciation of the fruit value (Gusmão et al. 2005; Stansly & Naranjo 2010).

Sampling plans are the starting point for decision-making systems in integrated pest management (IPM) programs because they are

<sup>&</sup>lt;sup>1</sup>Universidade Federal de Viçosa, Departamento de Entomologia, Viçosa, Minas Gerais, 36570-900, Brazil; E-mail: ruanvictorufv@gmail.com (R. V. B. T); arthurvieira3@gmail.com (A. V. R.); picanco@ufv.br (M. C. P.)

<sup>&</sup>lt;sup>2</sup>Universidade Federal do Tocantins, Campus de Gurupi, Gurupi, Tocantins, 77402-970, Brazil; E-mail: rsamento@uft.edu.br (R. A. S.);

poliana\_silvestre@yahoo.com.br (P. S. P.); agrochol@yahoo.com.br (C. H. O. L.); taniacko17@gmail.com (T. L. B. D.)

<sup>\*</sup>Corresponding author; E-mail: rsamento@uft.edu.br

used to determine economic damage levels and to validate sequential sampling plans (Binns et al. 2000; Gusmão et al. 2006; Pedigo & Rice 2014; Rosado et al. 2014; Pereira et al. 2017). In conventional sampling plans, the number of samples is fixed, whereas in sequential plans this number is variable (Pereira et al. 2017; Pinto et al. 2017). Conventional sampling plans consist of a sample unit, sampling technique, and the number of samples (Gusmão et al. 2005; Moura et al. 2007; Bacci et al. 2008). Optimal sampling units and techniques must be accurate, fast, low-cost, and representative. Accurate sampling units and techniques have low variability. Thus, for units and techniques to be precise they must have a relative variance (RV) of less than 25% (Southwood 1978). On the other hand, representative units and sampling techniques are those that have relative densities (individuals per sample) that present the same variation as the absolute density (individuals per plant or vine). The number of samples in the sampling plans should be economically feasible. Therefore, sampling plans must be simple, fast, and cost effective in order to be viable for farmers (Gusmão et al. 2005; Moura et al. 2007; Bacci et al. 2008; Pinto et al. 2017).

Despite the importance of *B. tabaci* as a pest in melon crops, a sampling plan for use in IPM programs for this species has yet to be developed for melons. Therefore, the objective of this study was to determine a conventional sampling plan for *B. tabaci* in melon crops with plants during vegetative, flowering, and fruiting stages.

## **Materials and Methods**

#### EXPERIMENTAL CONDITIONS

This study was carried out during 2015 and 2016 in 12 different commercial melon fields (variety 'Valenciano') in Formoso do Araguaia, Tocantins State, Brazil (11.9021055°S, 49.5616027°W, with an altitude of 240 masl, and a tropical climate with a dry winter). Each field consisted of approximately 20 ha. The plants were spaced by 1.5 × 2.0 m and established according to Braga Sobrinho et al. (2008) and Pessarakli (2016). The cultural treatments used were plant and fruit thinning, weed control, and fruit turning.

The research was divided into 3 parts. In the first part, the optimal sampling unit to assess *B. tabaci* was selected using the direct counting technique. In the second part, the optimal sampling technique (direct counting or leaf beating on a white tray) was selected. In the third part, the appropriate number of samples for the sampling plan was determined.

#### SELECTION OF THE OPTIMAL SAMPLING UNIT TO ASSESS BEMISIA TABACI

This part of the study was carried out in 3 melon fields at different stages in 2015. The plants were evaluated in the vegetative, flowering, and fruiting stages (Figs. 1A, B, C). In each field, 100 plants were selected at random and in each plant, the largest vine was evaluated. The leaves were numbered according to their position on the branch. Leaf number 1 was the most apical of the vine, number 2 the 2nd most apical leaf, and so on (Fig. 1D). On each leaf, the number of *B. tabaci* adults was evaluated by direct count. In this evaluation, the leaf was handled carefully (to avoid insect escape) and the number of *B. tabaci* adults present was counted (Fig. 1E). The adults were evaluated because this is the insect stage that causes most damage to plants (Gusmão et al. 2006) and they are easier to count due to their larger size (Chu et al. 2003).

The sampling unit was selected based on the criteria of leaf occurrence frequency in a vine, precision, and representativeness. In the first criterion, only leaves with an occurrence frequency (frequency with which a leaf in a certain position appears in a vine) above 80% were selected. This was done due to the long time taken to locate leaves that have a low occurrence frequency on the vines, which can increase sampling time (Rosado et al. 2014; Pinto et al. 2017). The occurrence frequency of each leaf on the vines of melon plants was calculated using formula (1):

$$Fi = (100 \times Ni)/Nr \tag{1}$$

where Fi = frequency of occurrence of leaf i (%) in the 100 vines evaluated, i = position of the leaf on the vine (left to right), Ni = number of times that the leaf i was present on the 100 evaluated vines, and Nr = total number of evaluated vines.

The precision criterion used to identify the sampling unit was the relative variances of *B. tabaci* densities on each leaf of the vine. Only relative variances of less than 25% were selected because samples and techniques with relative variance higher than 25% generate unfeasible sampling plans (Southwood 1978). The relative variances of *B. tabaci* densities on each leaf of the vine were calculated using formula (2) (Moura et al. 2003; Bacci et al. 2006):

$$RV = [100 \times S(x)]/x \tag{2}$$

where RV = relative variance (%), S(x) = standard error of the mean densities, and x = mean densities.

Under the representativeness criterion, the relative densities (adults per leaf<sup>1</sup>) and the absolute densities (adults per vine<sup>-1</sup>) were calculated, and the sample units considered were those with relative densities that were significantly correlated (P < 0.05) with absolute densities (Rosado et al. 2014; Pinto et al. 2017). The final selection was made based on the sample unit being amongst those suitable for the evaluation of pests according to all criteria during the vegetative, flowering, and fruiting stages.

#### SELECTION OF THE OPTIMAL SAMPLING TECHNIQUE

This part of the study was performed in 3 melon fields in 2015. During the evaluations, the plants of each field were in the vegetative, flowering, or fruiting stages (Figs. 1A, B, C). Twenty plants were randomly selected from each field. Two vines were evaluated in each plant. The evaluated leaves were those previously selected (fifth and sixth leaves). With 1 of these vines, *B. tabaci* density was evaluated by the direct count technique (Fig. 1E). With the other vine of the plant, whitefly densities were evaluated by leaf beating onto a white tray (40 × 25 × 3 cm) and counting *B. tabaci* adults (Fig. 1F). These techniques were used because they are suitable for the evaluation of sucking insects such as *B. tabaci* (Moura et al. 2007; Pinto et al. 2017). The time required to execute each sampling technique also was evaluated.

The relative variance of *B. tabaci* densities obtained by each sampling technique was calculated. The calculation of relative variance was performed using formula (2). The data for *B. tabaci* densities and sampling times for each sampling technique were analyzed by analysis of variance at P < 0.05. The technique that provided a relative variance of less than 25% and a shorter sampling time was selected (Southwood 1978).

# DETERMINATION OF THE NUMBER OF SAMPLES FROM THE SAMPLING PLAN

This stage of the study was divided into 2 parts. In the first part, the most appropriate frequency distribution was determined to describe the *B. tabaci* densities in the melon fields. In the second part, the maximum error to be used in pest sampling was determined.



Fig. 1. Crops with melon plants in phases (A) vegetative, (B) flowering, and (C) fruiting. (D) Leaves used as sample units (number 1 represents the most apical leaf of the vine and the largest number the most basal leaf of the vine). Sampling techniques of (E) direct count and (F) leaf beating on a white tray.

## DETERMINATION OF THE FREQUENCY DISTRIBUTION THAT BEST DESCRIBES THE DENSITIES OF *BEMISIA TABACI* IN CROPS

This stage of the research was carried out in 12 melon fields during 2015 and 2016. During the evaluations, the plants of each field were in the vegetative, flowering, or fruiting stages (4 fields at each phenological stage). In each field, 300 plants were evaluated. These plants were located equidistantly to eliminate directional errors (Bacca et al. 2006; Rosado et al. 2014; Pereira et al. 2017; Pinto et al. 2017). The sampling techniques used were those previously selected (direct counting of *B. tabacci* adults on the fifth or sixth most apical leaves of the vine).

The observed and expected frequencies of *B. tabaci* densities were calculated according to the negative binomial, Poisson, and positive binomial distributions through Microsoft Excel spreadsheets. These frequencies were compared using the chi-square test ( $\chi^2$ ) (Young & Young 1998). The data for the *B. tabaci* densities in a field were adjusted to a certain frequency distribution when the difference between the observed and expected frequencies was not significant (*P* > 0.05) (Young & Young 1998). A given frequency distribution was considered descriptive of *B. tabaci* densities in crops when the whitefly distribution conformed to this frequency distribution in most fields (more than 70%) (Rosado et al. 2014; Pinto et al. 2017).

The values of the aggregation parameter for each field were calculated using formula (3) (Young & Young 1998; Moura et al. 2003):

$$k = x^2/(S^2 - x)$$
 (3)

where k = aggregation parameter in the field, x = average density of the pest in the field, and  $S^2$  = variance of the pest densities.

The aggregation parameters (k) of each of the 12 fields were assessed with simple linear regression analysis according to Bliss and Owen (1958). In this regression, it is considered that the fields present a common aggregation parameter (kc) when their intercept is not significant and the slope is significant by the F test at P < 0.05 (Rosado et al. 2014; Pinto et al. 2017).

# Determination of the Maximum Allowed Error for Calculating the Number of Samples

Initially, the sample numbers of the sampling plan for melon crops in the vegetative, flowering, and fruiting stages were calculated using formula (4) (Young & Young 1998; Gusmão et al. 2005; Moura et al. 2007):

$$NA = \frac{1}{c^2} x \left( \frac{1}{x} + \frac{1}{kc} \right)$$
(4)

where NA = number of sample units, C = permitted error, x = population mean, and kc = common parameter of aggregation of the negative binomial frequency distribution (0.9134) determined previously.

In these determinations, we used error values of 0.05, 0.10, 0.15, 0.20, and 0.25. These errors were used because they are the standard range of errors used in these calculations (Moura et al. 2007; Pinto et al. 2017). Given the number of samples in these situations, the sampling time for each of these plans was evaluated in 3 fields. The plants of each field were in the vegetative, flowering, or fruiting stages.

The sampling cost was calculated using the following formula (5):

$$CS = CM + (TS \times CL)$$
(5)

where CS = cost of 1 sampling,  $CM = \text{cost of sampling material (pencil, rubber, paper, and clipboard: US$0.07 per sample), <math>TS = \text{sampling time}$  (h), and CL = labor costs (salary of a rural worker and non-wage labor costs: US\$8.00 per h) (Moura et al. 2007; Rosado et al. 2014).

The selected sampling plan was the plan with the lowest level of error and with sampling time of less than 1 h for 3 fields in the vegetative, flowering, or fruiting stages. This was done because sampling plans with this duration are feasible (Gusmão et al. 2005). In addition, these criteria allow these sampling plans to be standardized because they can be used in any melon plantation irrespective of phenological stage or pest density (Pinto et al. 2017).

#### Results

### SELECTION OF THE PLANT COMPONENTS TO BE USED FOR BEMISIA TABACI SAMPLING

In plants during the vegetative stage, only the 7 most apical leaves displayed frequencies of occurrence on the vines higher than 80%. From those 7 leaves, only the fifth and sixth leaves had whitefly density relative variances lower than 25%. From the third to seventh leaves, the relative densities (adults per leaf<sup>-1</sup>) presented significant correlations (P < 0.05) with absolute densities (adults per vine<sup>-1</sup>) (Table 1). Therefore, in the vegetative stage, the fifth and sixth leaves were suitable to sample *B. tabaci* adults because they had frequency of occurrence above 80%, relative variance lower than 25%, and significant correlations with absolute densities.

In the flowering plants, only the 10 most apical leaves presented frequencies of occurrence on the vines higher than 80%. From the third to tenth leaves, the relative variance of *B. tabaci* densities was lower than 25%. From the fourth to tenth leaves, the relative densities (adults per leaf<sup>1</sup>) presented significant correlations (P < 0.05). Thus, the fourth to tenth leaves of flowering melon plants are sampling units that may be used to sample *B. tabaci* adults.

In fruiting plants, the 15 most apical leaves presented frequencies of occurrence on the vines higher than 80%. From the fifth to fifteenth leaves the relative variance of *B. tabaci* densities was lower than 25% and the relative densities (adults per leaf<sup>1</sup>) presented significant correlations (P < 0.05) with absolute densities (adults per vine<sup>1</sup>) (Table 1). Thus, the fifth to fifteenth leaves of fruiting melon plants are sampling units that may be used to sample *B. tabaci* adults.

Therefore, the sampling units that were suitable in vegetative, flowering, and fruiting stages were the fifth and sixth most apical leaves of the vine. These were the ideal sampling units to generate a standardized sampling plan to be used for all phenological stages.

#### SELECTION OF THE OPTIMAL SAMPLING TECHNIQUE

In plants at vegetative and fruiting stages, higher densities of *B. tabaci* adults were detected using the leaf beating on a white tray technique than with the direct counting technique. The opposite occurred when the melon plants were at the flowering stage (Fig. 2A).

In plants at the vegetative stage, the relative variance of *B. tabaci* densities was lower than 25% when using both the leaf beating on a white tray technique and the direct counting technique. In the flowering and fruiting plants, the relative variance of *B. tabaci* densities was lower than 25% only when the direct counting technique was used (Fig. 2B).

Sampling time using the direct counting technique was shorter (average of 7 min per stage<sup>-1</sup>) than with the leaf beating on a white tray technique (average of 14 min per stage<sup>-1</sup>) in melon plants for all 3 phenological stages (vegetative, flowering, and fruiting) (Fig. 2C).

Thus, the best technique to sample *B. tabaci* adults in melon plants in the 3 phenological stages (vegetative, flowering, and fruiting) was direct counting.

#### 20

Table 1. Selection of leaf position to be used in the sampling of *Bemisia tabaci* adults in melon plants during vegetative, flowering, and fruiting stages: leaf occurrence frequency on the vine, density (mean ± standard error), relative variance (RV), Pearson correlation coefficient (r) between relative densities (adults per leaf<sup>3</sup>) and absolute density (adults per vine<sup>-1</sup>).

		B. t	<i>abaci</i> sampling variab	les
Leaf position in the vine <sup>a</sup>	Frequency (%)	Density	RV (%)	r
Vegetative phase plants: absolute density = $0.74 \pm 0.10$ adults per vine <sup>-1</sup>				
1	100	$0.00 \pm 0.00$	0.00	0.00
2	100	$0.00 \pm 0.00$	0.00	0.00
3	100	$0.01 \pm 0.01$	100.00	0.23*
4	100	$0.15 \pm 0.04$	25.73	0.55*
5	100	$0.28 \pm 0.06$	20.36	0.68*
6	100	$0.18 \pm 0.04$	24.18	0.48*
7	87	$0.11 \pm 0.04$	33.13	0.42*
Flowering plants: absolute density = 33.19 ± 2.54 adults per vine <sup>-1</sup>				
1	100	$0.01 \pm 0.01$	100.00	0.23*
2	100	$0.22 \pm 0.06$	25.50	0.11
3	100	$0.66 \pm 0.13$	19.29	0.06
4	100	2.07 ± 0.27	13.07	0.40*
5	100	$5.02 \pm 0.63$	12.63	0.69*
6	100	4.86 ± 0.53	10.90	0.66*
7	100	5.38 ± 0.63	11.78	0.76*
8	100	4.39 ± 0.46	10.43	0.72*
9	100	4.70 ± 0.53	11.22	0.80*
10	97	4.45 ± 0.47	10.57	0.71*
Fruiting plants: absolute density = 5.27 $\pm$ 0.55 adults per vine <sup>-1</sup>				
1	100	$0.00 \pm 0.00$	0.00	0.00
2	100	$0.00 \pm 0.00$	0.00	0.00
3	100	$0.04 \pm 0.02$	49.24	-0.01
4	100	$0.09 \pm 0.03$	39.00	0.29*
5	100	$0.14 \pm 0.03$	24.93	0.29*
6	100	$0.16 \pm 0.04$	24.70	0.26*
7	100	$0.35 \pm 0.07$	21.27	0.49*
8	100	$0.44 \pm 0.12$	28.60	0.57*
9	100	$0.44 \pm 0.07$	16.60	0.52*
10	100	$0.53 \pm 0.09$	16.97	0.63*
11	100	$0.61 \pm 0.09$	15.20	0.55*
12	100	$0.62 \pm 0.11$	17.74	0.57*
13	100	$0.50 \pm 0.09$	18.62	0.59*
14	100	$0.71 \pm 0.13$	18.45	0.59*
15	100	0.63 ± 0.12	18.68	0.62*

\*1, 2, and n = first, second, and nth leaf from the vine apex, respectively. Only leaves with an occurrence frequency on the vine higher than 80% were included in the table. \*Significant correlation according to the t test at P < 0.05.

## DETERMINATION OF THE NUMBER OF SAMPLES FOR *BEMISIA* TABACI SAMPLING IN MELON FIELDS

In 9 of the 12 fields evaluated (75%), the differences between observed and expected frequencies of *B. tabaci* densities according to the negative binomial distribution were not significant (P > 0.05). Therefore, in these fields, pest densities followed the negative binomial frequency distribution. *Bemisia tabaci* densities followed the Poisson frequency distribution in only 3 of the melon fields. On the other hand, *B. tabaci* densities that followed a positive binomial frequency distribution were not observed in any of the fields (Table 2). Therefore, the ideal formula to calculate samples for the sampling plan is the negative binomial frequency distribution.

The simple linear regression of the common aggregation parameter of the *B. tabaci* densities in 12 melon fields (Kcommon) as a function of the values of the aggregation parameter of each crop showed a significant curve (P < 0.05) and a non-significant intercept (P > 0.05) (Table 3). Therefore, the *B. tabaci* densities in 12 melon fields showed a common aggregation parameter (Kcommon).

The sampling time for sampling plans with errors of 5 and 10% was higher than 1 h. When the sampling error was 15, 20, and 25%, the sampling time was less than 1 h. Therefore, 15% was the smallest error that generated a sampling plan with sampling time less than 1 h. Using this error (15%), the number of samples from the sampling plan was 72 per crop with a duration of 39 min and cost of US\$5.27 per sample (Table 4).

# Discussion

The fifth and sixth most apical leaves of the plant were the optimal sample units for the evaluation of *B. tabaci* in melon crops because they allowed more precise sampling of the pest for all phenological stages. In this context, these units allowed faster sampling because they were present in 100% of plant vines. These sample units also enabled accurate sampling, because *B. tabaci* densities displayed a relative variance of less than 25%. According to Southwood (1978), these



**Fig. 2.** (A) Adult density of *Bemisia tabaci* (means ± standard errors), (B) relative variance of densities, and (C) sampling times of *Bemisia tabaci* adults using direct counting and leaf beating on a white tray methods in melon crops in the vegetative, flowering, and fruiting stages. For each phenological stage, the histograms with lowercase letters indicate significantly different means according to the F test (P < 0.05).

values generate feasible sampling plans. In addition, the fifth and sixth most apical leaves of the branch allowed representative samplings of pest attacks on the vine. This is shown by the fact that relative *B. tabaci* 

densities (adults per leaf<sup>1</sup>) showed a positive and significant correlation with the absolute density of the pest (adults per vine<sup>-1</sup>). Moreover, these fast, precise, and representative samplings were observed for all phenological stages (vegetative, flowering, and fruiting) of melon plants. The preference of whitefly to attack younger leaves can be explained by the fact that these leaves present better nutritional content, besides having soft, fine cuticles, and a higher concentration of water available to the insect (Van Lenteren & Noldus 1990).

The best sampling technique to sample *B. tabaci* adults in melon crops was direct counting due to the possibility of sampling the pest at all the phenological stages of plant growth. One of the characteristics that led to the choice of the direct counting technique was the fact that it enabled accurate sampling (relative variance lower than 25%) in all situations. In addition, direct counting was faster than the leaf beating on a white tray technique for all phenological stages of the plant. The fact that a technique has a shorter sampling time indicates that its use will generate more feasible sampling plans (Southwood 1978; Gusmão et al. 2005; Moura et al. 2007).

The frequency distribution that best described *B. tabaci* adult densities in melon crops was the negative binomial. This was probably due to variance of pest densities being higher than average for this characteristic (Tonhasca et al. 1994; Moura et al. 2003). Additionally, we observed that in melon crops with plants at vegetative, flowering, and fruiting stages, the variance in *B. tabaci* densities was up to 74.65 and 104.00 times higher than the averages, respectively.

In this context, some authors misinterpret the frequency distribution data, stating that it represents the spatial distribution of insects in fields (Maruyama et al. 2006; Costa et al. 2010; Souza et al. 2013). However, this is not the case, because there is no relationship between frequency distributions of insect densities and the spatial distribution of these organisms in fields (Young & Young 1990; Barrigossi et al. 2001). To evaluate spatial distribution of insect populations in fields, it is necessary to locate the geographical coordinates of each sampling point and to analyze the data using a suitable statistical tool such as geostatistics (Young & Young 1998; Rijal et al. 2014; Rosado et al. 2015). However, frequency distribution data is an appropriate tool to select the formula to use in the calculation of samples for sampling plans (Young & Young 1998; Gusmão et al. 2005; Bacci et al. 2008) and population estimates (Pedigo & Buntin 1993; Heersink et al. 2016).

**Table 2.** Chi-squared test ( $\chi^2$ ) results between the frequencies observed and expected by the negative binomial, Poisson, and positive binomial *Bemisia tabaci* density distributions (means ± standard errors) in 12 fields of melon.

		Negative binomial		Poisson	Poisson		Positive binomial	
Fields	Density	$\chi^2$	df	χ <sup>2</sup>	df	χ <sup>2</sup>	df	
Vegetative stage plants								
1	$0.11 \pm 0.02$	1.07 <sup>NS</sup>	1	7.62*	1	892.24*	1	
2	$0.04 \pm 0.01$	$1.00 \times 10^{-4}$ NS	1	0.01 <sup>NS</sup>	1	172.84*	1	
3	4.49 ± 0.25	24.25 <sup>NS</sup>	15	12,552.71*	16	2.97 × 10 <sup>32</sup> *	16	
4	$3.29 \pm 0.17$	33.42*	13	7,415.55*	14	$2.05 \times 10^{15*}$	14	
Flowering plants								
5	0.09 ± 0.03	0.81 <sup>NS</sup>	2	1,657.02*	3	317.00*	3	
6	$0.07 \pm 0.02$	2.23 <sup>NS</sup>	1	653.95*	2	232.35*	2	
7	$2.99 \pm 0.16$	9.11 <sup>NS</sup>	8	378.26*	9	5.88 × 10 <sup>18</sup> *	9	
8	$4.95 \pm 0.22$	19.96 <sup>NS</sup>	12	1,038.96*	13	$1.01 \times 10^{27*}$	13	
Fruiting plants								
9	$0.03 \pm 0.01$	9.81 × 10 <sup>-5 NS</sup>	1	0.01 <sup>NS</sup>	1	135.89*	1	
10	$0.02 \pm 0.01$	5.50 × 10 <sup>-5 NS</sup>	1	$1.40 \times 10^{-3}$ NS	1	71.42*	1	
11	4.72 ± 0.35	28.14*	12	3,012.50*	13	2.99 × 10 <sup>30</sup> *	13	
12	$2.35 \pm 0.24$	55.94*	10	13,134.12*	11	7.10 × 10 <sup>23</sup> *	11	

<sup>NS</sup>Not significant at 5% probability. \*Significant at 5% probability. df = Degrees of freedom.

Downloaded From: https://complete.bioone.org/journals/Florida-Entomologist on 27 Apr 2024 Terms of Use: https://complete.bioone.org/terms-of-use

**Table 3.** Analysis of variance of *Bemisia tabaci* adult density, sampled in 12 melon fields, to assess the dispersion parameter of the common negative binomial distribution (Kcommon).

85 277.85 6.77*
13 7.13 0.17 <sup>NS</sup>
18 41.02

 $^{\rm NS}Not$  significant at 5% probability. \*Significant at 5% probability. df = Degrees of freedom.

The existence of a common aggregation parameter (Kcommon) for *B. tabaci* densities in melon crops indicates that it is possible to determine a sampling plan for this pest that is applicable to all sampled crops (Bliss & Owen 1958). This is even more relevant because the fields sampled in this project were in different phenological stages (vegetative, flowering, or fruiting). Therefore, the sampling plan generated here has great potential to be adopted by farmers due to its capacity for being used throughout all melon crop stages.

For a *B. tabaci* sampling plan to be viable, it should present a sampling time of less than 1 h and require a small number of samples (Bacci et al. 2006; Rosado et al. 2014). The smallest error used in the calculation to sample this pest was 15%. Therefore, it is a feasible system. According to Southwood (1978), the permitted error for this type of study can be up to 25%. The number of samples obtained was 72 samples per crop with a collection time of 39 min and a cost of US\$5.27 per sample. The advantage of these systems is that producers are able to obtain more reliable (smaller sampling error), representative, faster, and less expensive samples. According to Bacci et al. (2008), these sampling techniques can reduce pesticide use, cost, and application time, and preserve the ecosystem and human health. In addition to providing the farmer with environmentally friendly management options, IPM ensures greater profitability by reducing pesticide spraying (Picanço et al. 2004; Pedigo & Rice 2014).

In conclusion, *B. tabaci* sampling in melon crops should be performed by direct counting of pest adults on the fifth and sixth most apical leaves of the plant vine. The sampling plan is composed of 72 samples per crop. This sampling plan is feasible with a duration of 39 min, a cost of US\$5.27 per sample, and can be used in plants during vegetative, flowering, and fruiting stages.

# Acknowledgments

We thank the National Council for Scientific and Technological Development - CNPq - Brazil (Projects: 458946/2014-1 and 304178/2015-2) for financial support. We also thank the Coordination for the Im-

**Table 4.** Number of samples, time involved, and cost of sampling *Bemisia tabaci* adults in melon plants as a function of the maximum error admitted in this process.

		Sampling Effort		
Error (%)	Number of Samples	Time	Cost (US\$)	
5	644	5 h 51 min	46.87	
10	161	1 h 28 min	11.80	
15	72	39 min	5.27	
20	40	22 min	3.00	
25	26	14 min	1.94	

Downloaded From: https://complete.bioone.org/journals/Florida-Entomologist on 27 Apr 2024 Terms of Use: https://complete.bioone.org/terms-of-use

provement of Higher Education Personnel (CAPES), Brazil (Project: PROCAD-2013), and the Minas Gerais State Research Foundation (FAPEMIG) for the scholarships and resources provided. The Universidade Federal de Viçosa, Brazil, and the Federal University of Tocantins, Brazil, are thanked for providing logistical support.

# **References Cited**

- Bacca T, Lima ER, Picanço MC, Guedes RNC, Viana JHM. 2006. Optimum spacing of pheromone traps for monitoring the coffee leaf miner *Leucoptera coffeella*. Entomologia Experimentalis et Applicata 119: 39–45.
- Bacci L, Picanço MC, Moura MF, Della Lucia TMC, Semeão AA. 2006. Sampling plan for *Diaphania* spp. (Lepidoptera: Pyralidae) and for hymenopteran parasitoids on cucumber. Journal of Economic Entomology 99: 2177–2184.
- Bacci L, Picanço MC, Moura MF, Semeão AA, Fernandes FL, Morais EGF. 2008. Sampling plan for thrips (Thysanoptera: Thripidae) on cucumber. Neotropical Entomology 37: 582–590.
- Barrigossi JAF, Young LJ, Crawford CAG, Hein GL, Higley LG. 2001. Spatial and probability distribution of Mexican bean beetle (Coleoptera: Coccinellidae) egg mass populations in dry bean. Environmental Entomology 30: 244–253.
- Binns MR, Nyrop JP, van der Werf W. 2000. Sampling and Monitoring in Crop Protection: The Theoretical Basis for Developing Practical Decision Guides. CABI, Wallingford, United Kingdom.
- Blackmer JL, Byrne DN. 1999. The effect of *Bemisia tabaci* on amino acid balance in *Cucumis melo*. Entomologia Experimentalis et Applicata 93: 315–319.
- Bliss Cl, Owen ARG. 1958. Negative binomial distributions with a common k. Biometrika 45: 37–58.
- Braga Sobrinho R, Guimarães JA, Araújo EL, Moreira MAB, Mesquita ALM. 2011. Manejo Integrado de Pragas do Meloeiro. Embrapa Agroindústria Tropical, Fortaleza, Brazil.
- Braga Sobrinho R, Guimarães JA, Freitas J de AD de, Terao D. 2008. Produção Integrada de Melão. Embrapa Agroindústria Tropical, Fortaleza, Brazil.
- CABI (Center for Agriculture and Bioscience International). 2016. Crop Protection Compendium. CAB International Publishing, Wallingford, United Kingdom. http://www.cabi.org/isc/datasheet/8927 (last accessed 1 Jul 2018).
- Chu C, Buckner JS, Karut K, Freeman TP, Nelson DR, Hennebeny TJ. 2003. A survey of sizes and weights of *Bemisia tabaci* (Homoptera: Aleyrodidae) B biotype life stages from field grown cotton and cantaloupes. Insect Science 10: 121–129.
- Costa MG, Barbosa JC, Yamamoto PT, Leal RM. 2010. Spatial distribution of *Di-aphorina citri* Kuwayama (Hemiptera: Psyllidae) in citrus orchards. Scientia Agricola 67: 546–554.
- FAO (Food and Agriculture Organization of the United Nations). 2014. Statistical database 2014. Food and Agriculture Organization of the United Nations, Rome, Italy. http://www.fao.org/faostat/en/#data/QC (last accessed 1 Jul 2018).
- Freitas J de AD de, Sobrinho RB, Andrade APS de. 2009. Produção integrada de melão nos pólos Mossoró-Açu e Baixo Jaguaribe: diagnóstico das conformidades e não conformidades com os requisitos do sistema. Embrapa Agroindústria Tropical, Fortaleza, Brazil.
- Greathead AH. 1986. Host plants, pp. 17–25 *In* Cock MJW [ed.], *Bemisia tabaci* - A Literature Survey on the Cotton Whitefly with Annotated Bibliography. FAO (Food and Agriculture Organization of the United Nations), London, United Kingdom.
- Gusmão MR, Picanço MC, Guedes RNC, Galvan TL, Pereira EJG. 2006. Economic injury level and sequential sampling plan for *Bemisia tabaci* in outdoor tomato. Journal of Applied Entomology 130: 160–166.
- Gusmão MR, Picanço MC, Zanuncio JC, Silva DJH, Barrigossi JAF. 2005. Standardised sampling plan for *Bemisia tabaci* (Homoptera: Aleyrodidae) in outdoor tomatoes. Scientia Horticulturae 103: 403–412.
- Heersink DK, Meyers J, Caley P, Barnett G, Trewin B, Hurst T, Jansen C. 2016. Statistical modeling of a larval mosquito population distribution and abundance in residential Brisbane. Journal of Pest Science 89: 267–279.
- van Lenteren JA, Noldus LPJJ. 1990. Whitefly-plant relationships: behavioral and ecological aspects, pp. 47–89 *In* Gerling D [ed.], Whiteflies: Their Bionomics, Pest Status and Management. Intercept, Andover, United Kingdom.
- Li S-J, Xue X, Ahmed MZ, Ren S-X, Du Y-Z, Wu J-H, Cuthbertson AGS, Qiu B-L. 2011. Host plants and natural enemies of *Bemisia tabaci* (Hemiptera: Aleyrodidae) in China. Insect Science 18: 101–120.
- Maruyama WI, Barbosa JC, Toscano LC. 2006. Distribuição espacial de Oncometopia facialis (Signoret) (Hemiptera: Cicadellidae) em pomar cítrico. Neotropical Entomology 35: 93–100.
- Moura MF, Picanço MC, Guedes RNC, Barros EC, Chediak M, Morais EGF. 2007. Conventional sampling plan for the green leafhopper *Empoasca kraemeri* in common beans. Journal of Applied Entomology 131: 215–220.

- Moura MF de, Picanço MC, Silva ÉM da, Guedes RNC, Pereira JL. 2003. Plano de amostragem do biótipo B de *Bemisia tabaci* na cultura do pepino. Pesquisa Agropecuária Brasileira 38: 1357–1363.
- Pedigo LP, Buntin GD. 1993. Handbook of Sampling Methods for Arthropods in Agriculture. CRC Press, Boca Raton, Florida, USA.
- Pedigo LP, Rice ME. 2014. Entomology and Pest Management, 6th ed. Waveland Press, Long Grove, Illinois, USA.
- Pereira PS, Sarmento RA, Galdino TVS, Lima CHO, dos Santos FA, Silva J, dos Santos GR, Picanço MC. 2017. Economic injury levels and sequential sampling plans for *Frankliniella schultzei* in watermelon crops. Pest Management Science 73: 1438–1445.
- Pessarakli M. 2016. Handbook of Cucurbits: Growth, Cultural Practices, and Physiology. CRC Press, Boca Raton, Florida, USA.
- Picanço MC, Paula SV de, Moraes Junior AR, Oliveira IR de, Semeão AA, Rosado JF. 2004. Impactos financeiros da adoção de manejo integrado de pragas na cultura do tomateiro. Acta Scientiarum 26: 245–252.
- Picanço MC, Pereira EJG, Crespo ALB, Semeão AA, Bacci L. 2002. Manejo integrado das pragas das fruteiras tropicais, pp. 513–578 In Zambolim L [ed.], Manejo Integrado: Fruteiras Tropicais - Doenças e Pragas. Universidade Federal de Viçosa, Viçosa, Minas Gerais, Brazil.
- Pinto CB, Sarmento RA, Galdino TV da S, Pereira PS, Barbosa BG, Lima CHO, Silva NR da, Picanço MC. 2017. Standardized sampling plan for the thrips *Frankliniella schultzei* (Thysanoptera: Thripidae) on watermelon crops. Journal of Economic Entomology 110: 748–754.
- Rijal JP, Brewster CC, Bergh JC. 2014. Spatial distribution of grape root borer (Lepidoptera: Sesiidae) infestations in Virginia vineyards and implications for sampling. Environmental Entomology 43: 716–728.
- Rosado JF, Picanço MC, Sarmento RA, Pereira RM, Pedro-Neto M, Galdino TVS, Saraiva A de S, Erasmo EAL. 2015. Geostatistics as a tool to study mite dis-

persion in physic nut plantations. Bulletin of Entomological Research 105: 381–389.

- Rosado JF, Sarmento RA, Pedro-Neto M, Galdino TVS, Marques RV, Erasmo EAL, Picanço MC. 2014. Sampling plans for pest mites on physic nut. Experimental and Applied Acarology 63: 521–534.
- Southwood TRE. 1978. Ecological Methods with Particular Reference to Study of Insect Populations. Chapman and Hall, New York, New York, USA.
- Souza LA, Barbosa JC, Grigolli JFJ, Fraga DF, Maldonado Jr W, Busoli AC. 2013. Spatial distribution of *Euschistus heros* (F.) (Hemiptera: Pentatomidae) in soybean. Neotropical Entomology 42: 412–418.
- Stansly PA, Naranjo SE [eds.]. 2010. *Bemisia*: Bionomics and Management of a Global Pest. Springer, New York, New York, USA.
- Tonhasca A, Palumbo JC, Byrne DN. 1994. Distribution patterns of *Bemisia taba-ci* (Homoptera: Aleyrodidae) in cantaloupe fields in Arizona. Environmental Entomology 23: 949–954.
- Toscano LC, Boiça Júnior AL, Maruyama WI. 2004. Avaliação de aspectos fisiológicos de três genótipos de tomateiro infestados por *Bemisia tabaci* (Gennadius) biótipo B (Hemiptera: Aleyrodidae). Neotropical Entomology 33: 777–782.
- Varma A, Mandal B, Singh MK. 2011. Global emergence and spread of whitefly (*Bemisia tabaci*) transmitted geminiviruses, pp. 205–292 *In* Thompson WMO [ed.], The Whitefly, *Bemisia tabaci* (Homoptera: Aleyrodidae) Interaction with Geminivirus-infected Host Plants. Springer, Dordrecht, The Netherlands.
- Young L, Young J. 1990. A spatial view of the negative binomial parameter k when describing insect populations. Annual Conference on Applied Statistics in Agriculture. http://newprairiepress.org/cgi/viewcontent.cgi?article= 1428&context=agstatconference (last accessed 1 Jul 2018).
- Young L, Young J. 1998. Statistical Ecology: A Population Perspective. Kluwer Academic Publishers, Boston, Massachusetts, USA.