

## **Examining the Prevalence of Solenopsis Invicta Virus 3 (Solinviviridae: Invictavirus) in Solenopsis invicta (Hymenoptera: Formicidae) Alates Collected in North Florida**

Authors: Arnold, Donna P., Balasubramani, Subramani P., Valles, Steven M., and Hottel, Benjamin A.

Source: Florida Entomologist, 104(1) : 9-12

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.104.0102>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Examining the prevalence of *Solenopsis invicta* virus 3 (Solinviviridae: Invictavirus) in *Solenopsis invicta* (Hymenoptera: Formicidae) alates collected in North Florida

Donna P. Arnold<sup>1</sup>, Subramani P. Balasubramani<sup>2</sup>, Steven M. Valles<sup>3</sup>,  
and Benjamin A. Hottel<sup>1,\*</sup>

---

## Abstract

The red imported fire ant, *Solenopsis invicta* Buren (Hymenoptera: Formicidae), originally is from South America and currently infests over 128 million ha in the US. Its presence has caused significant social, environmental, and economic impacts. Over the decades, chemical insecticides have controlled these pest ants successfully. However, this method is costly and unsustainable because red imported fire ant re-establishes colonies quickly after chemical application. Thus, it is important to develop additional strategies for managing the red imported fire ant in the US. *Solenopsis invicta* virus 3 is a positive-sense, single-stranded RNA virus specific for *S. invicta* that offers promise as a classical biological control agent or biopesticide for control of *S. invicta*. Surveys were conducted to determine the prevalence of *Solenopsis invicta* virus 3 in alates of *S. invicta* collected from 5 urban areas (Tallahassee, Pensacola, Jacksonville, Gainesville, and Panama City) and 5 adjacent rural areas (Quincy, Jay, Macclenny, Lake City, and Blountstown) of North Florida, USA, using the reverse transcription polymerase chain reaction technique. The prevalence of *Solenopsis invicta* virus 3 varied widely from city to city. No statistically significant differences in alate infection rate was found between urban and rural cities sampled. Areas in which no infections of *Solenopsis invicta* virus 3 were detected may be good candidates for the introduction of this virus as a biological control agent to help manage this pest locally.

Key Words: red imported fire ants; biological control; *Solenopsis invicta* virus 3

## Resumen

La hormiga roja de fuego importada, *Solenopsis invicta* Buren (Hymenoptera: Formicidae), es de origen Sudamericano y actualmente infesta más de 128 millones de hectáreas en los Estados Unidos. Su presencia ha provocado importantes impactos sociales, ambientales y económicos. Durante décadas, los insecticidas químicos han controlado estas plagas de hormigas con éxito. Sin embargo, este método es costoso e insostenible porque la hormiga roja de fuego importada restablece las colonias rápidamente después de la aplicación de productos químicos. Por lo tanto, es importante desarrollar estrategias adicionales para el manejo de la hormiga roja de fuego importada en los EE. UU. *Solenopsis invicta* 3 es un virus de ARN monocatenario de sentido positivo específico para *S. invicta* que es prometedor como agente de control biológico clásico o biopesticida para el control de *S. invicta*. Se realizó un sondeo para determinar la prevalencia del virus *Solenopsis invicta* 3 en individuos alados de *S. invicta* recolectados de 5 áreas urbanas (Tallahassee, Pensacola, Jacksonville, Gainesville, y Ciudad de Panamá) y 5 áreas rurales adyacentes (Quincy, Jay, Macclenny, Lake City, y Blountstown) del norte de Florida, EE. UU., utilizando la técnica de reacción en cadena de la polimerasa con transcripción inversa. La prevalencia del virus *Solenopsis invicta* 3 varió ampliamente de una ciudad a otra. No se encontraron diferencias estadísticamente significativas en la tasa de infección de los individuos alados entre las ciudades rurales y urbanas muestreadas. Las áreas en las que no se detectaron infecciones por el virus *Solenopsis invicta* 3 pueden ser buenas candidatas para la introducción de este virus como agente de control biológico para ayudar a manejar esta plaga localmente.

Palabras Clave: hormiga roja de fuego importada; control biológico; virus de *Solenopsis invicta* 3

---

*Solenopsis invicta* Buren (Hymenoptera: Formicidae), commonly referred to as the red imported fire ant, was introduced from South America to the US in the early 1900s at the port of Mobile, Alabama, USA. *Solenopsis invicta* has spread across the US since then (Tschinkel 2006), and currently occupies over 128 million ha from Virginia, USA, to South Florida, and west to California, USA (Williams et al. 2001). *Solenopsis invicta* has become a serious pest in the US (Tschinkel 1998).

These ants thrive in disturbed habitats and may cause serious medical (anaphylaxis from envenomation) and agricultural problems in the US (Porter & Savignano 1990). Costs to manage, prevent, repair damage, and otherwise attempt to mitigate the effects of the pest exceed an estimated \$6 billion annually (Pereira 2003). Insecticides provide effective control; however, this can be costly because continuous applica-

---

<sup>1</sup>Florida A&M University, Center for Biological Control, Tallahassee, Florida 32307, USA; E-mail: donnaarnold16@hotmail.com (D. P. A.), benjamin.hottel@rollins.com (B. A. H.)

<sup>2</sup>Florida A&M University, Center for Viticulture and Small Fruit Research, Tallahassee, Florida 32308, USA; E-mail: balasubra.paranthama@fam.edu (S. P. B.)

<sup>3</sup>USDA-ARS, Center for Medical, Agricultural and Veterinary Entomology, Gainesville, Florida 32608, USA; E-mail: steven.valles@usda.gov (S. M. V.)

\*Corresponding author; E-mail: benjamin.hottel@rollins.com

tion is necessary because fire ants establish new colonies once the use of insecticides is discontinued (Caldera et al. 2008).

Additionally, insecticide use may not be environmentally suitable, nor economically feasible in large areas (i.e., wildlife and agricultural regions) where *S. invicta* has substantial impacts (Drees & Gold 2003). Because of these economic and environmental issues, it is important to explore more sustainable management strategies (Summerlin et al. 1977). *Solenopsis invicta* is not considered a threat in its native range, and this has been attributed to the presence of natural enemies (Porter et al. 1992). A minimum of 30 fire ant natural enemies have been identified in South America, but nearly all of these are absent among US populations (Porter et al. 1997). Thus, permanent sustainable management of *S. invicta* populations in the US likely will depend on self-sustaining biological control agents (Valles 2012). Past research has focused on the discovery and use of biological control agents such as entomopathogenic viruses. In the US, 3 positive-sense single-stranded RNA viruses, *Solenopsis invicta* virus 1 (SINV-1), *Solenopsis invicta* virus 2 (SINV-2), and *Solenopsis invicta* virus 3 (SINV-3), were discovered infecting *S. invicta* using a metagenomic/next generation sequencing approach (Valles et al. 2009; Valles 2012).

Laboratory experiments have been conducted to examine the effects of these viruses on *S. invicta* populations. *Solenopsis invicta* virus 3 was found to cause the highest mortality of all the viruses tested and is the best viral candidate to be used as a biological control agent in the field (Valles et al. 2004). While research has examined the distribution of the virus in Argentina, California, and Florida (Gainesville), no distribution studies have been performed for *Solenopsis invicta* virus 3 on *S. invicta* alates (Hashimoto & Valles 2008). Because the virus is vertically transmitted, alates may serve to disseminate and maintain the *Solenopsis invicta* virus 3 infection. In order to better understand the natural prevalence and potential dissemination of *Solenopsis invicta* virus 3, a survey was conducted to examine the presence or absence of *Solenopsis invicta* virus 3 in *S. invicta* alates collected from urban and rural areas in North Florida.

## Materials and Methods

### FIELD COLLECTION

*Solenopsis invicta* female alates were collected from 5 urban cities (Tallahassee, Pensacola, Jacksonville, Gainesville, and Panama City) and 5 adjacent rural cities (Quincy, Jay, Macclenny, Lake City, and Blountstown) in North Florida from Feb to Oct 2018 (Fig. 1). Cities were classified as rural or urban based on the 2010 US census classification map of an urban area. The rural cities were located within an 80.4 km radius of their paired urban cities with high infestations of *S. invicta*. Female alates were collected from 7 to 10 mounds in each city, with each mound separated from the other by at least 1.6 km. Alates were collected by hand from nests by opening the *S. invicta* nest with a shovel. Alates were placed in 20 mL scintillation vials. Each vial contained a cotton ball soaked with a 10% pure cane sugar solution to keep ants alive for transport. The vials were labelled with the date, time, and location. Forceps were sterilized with a 10% bleach solution before being used at a new mound location. Five randomly selected colonies per city were selected to be screened for *Solenopsis invicta* virus 3 infections.

### VIRUS DETECTION

Collected female *S. invicta* alates were screened for the presence or absence of *Solenopsis invicta* virus 3 using molecular techniques adapted from Valles et al. (2010). RNA was extracted using a Qiagen

RNA easy mini kit (Qiagen Inc., Germantown, Maryland, USA) according to the manufacturer's instructions. For each of the 5 colonies selected per city, total RNA was extracted from female alates from each urban and rural colony sampled. Total RNA was quantified using a Spectrophotometer (Nanodrop 1000; Fisher Scientific, Pittsburgh, Pennsylvania, USA). Samples then were serially diluted with diethyl pyrocarbonate (DEPC)-treated water to achieve a total RNA concentration of 25 to 100 ng per  $\mu$ L. The cDNA was synthesized in a thermal cycler (Eppendorf, Hamburg, Germany) from the extracted RNA template using a reverse transcriptase master mix and a *Solenopsis invicta* virus 3 specific oligonucleotide primer (p812: 5'AATATCAGCATATTGATGATC-CAAAATGCCAA3').

Polymerase chain reaction was performed subsequently on the resulting cDNA. A 10  $\mu$ L polymerase chain reaction master mix was created using 1.0  $\mu$ L of sample buffer, 0.8  $\mu$ L of 25 mM  $MgCl_2$ , 0.2  $\mu$ L of 10  $\mu$ M dNTP mix, 0.5  $\mu$ L Taq DNA polymerase, 1  $\mu$ L of 4  $\mu$ M oligonucleotide primers (p812) and p813 (5'AAGAGAACGTATGCCTACTC-CATCAGAAGCAT3'), 2  $\mu$ L of cDNA and water (variable). Samples were incubated in a thermal cycler for 1.5 h with the following temperature regime: 94 °C for 2 min, then 35 cycles at 94 °C for 15 s, next 60 °C for 15 s, followed by 72 °C for 30 s, and finally 72 °C for 5 min. Polymerase chain reaction products then were separated by electrophoresis in a 2% agarose gel and visualized using SYBR® safe dye. *Solenopsis invicta* virus 3-infected female alates produced an amplicon of 259 base pairs, which indicated a positive response for *Solenopsis invicta* virus 3 viral infection (Valles et al. 2009). A gel documentation system (Bio-Rad, Hercules, California, USA) was used to capture the image of agarose gel for analysis and interpretation. Positive and non-template controls were included in each reverse transcription-polymerase chain reaction.

### STATISTICAL ANALYSIS


Fisher's exact test was used on the count data to determine if there were any statistical differences between the numbers of alates infected with *Solenopsis invicta* virus 3 from colonies collected in rural versus urban cities (Du-Prel et al. 2010). R statistical software version 3.5.1 was used to analyze data (R Foundation for Statistical Computing, Vienna, Austria).

## Results

Reverse transcription-polymerase chain reaction analysis successfully identified *Solenopsis invicta* virus 3 infections in female alates (Fig. 2). *Solenopsis invicta* virus 3 was detected in female alates from *S. invicta* nests from 7 of the 10 collection locations in northern Florida (Fig. 1.). The infection rate ranged from 0 to 100% depending on location. The virus was not detected in alates collected from Jay, Jacksonville, or Pensacola. *Solenopsis invicta* virus 3 was most prevalent in Gainesville with a 100% infection rate per colony (Fig. 2). The overall mean infection rate ( $\pm$  standard error) for North Florida was  $44.0 \pm 11.9\%$ . *Solenopsis invicta* virus 3 infections in rural and urban areas were similar. The mean infection rates were  $40.0 \pm 14.1\%$  and  $48.0 \pm 21.0\%$  for rural and urban areas, respectively. Fisher exact analysis showed no significant difference between the number of colonies infected with *Solenopsis invicta* virus 3 in rural and urban areas ( $P = 0.7761$ ; Odd Ratio 1.38).

## Discussion

The goal of this study was to examine the presence and distribution of *Solenopsis invicta* virus 3 among female alates of *S. invicta* in rural




**Fig. 1.** Map of Florida illustrating the collection locations and distribution of *Solenopsis invicta* virus 3 infection rate of female alates of *Solenopsis invicta* colonies in urban and rural cities of North Florida. Black portions of the pie charts represent the proportion of alates infected with *Solenopsis invicta* virus 3.

and urban areas of North Florida. The *Solenopsis invicta* virus 3 infection rate was higher considerably in female alates compared with earlier studies that examined the infection rate in worker ants (Valles et al. 2009). Female alates were chosen for this study because this life stage had not been surveyed extensively previously, and female alates may portend future establishment of this virus in new areas because the virus is vertically transmitted. Surveys of workers would indicate only the current infection status for an area. The *Solenopsis invicta* virus 3 infection rate was higher than anticipated based on previous surveys. The infection rate of *Solenopsis invicta* virus 3 in *S. invicta* colonies (based on the interrogation of workers) typically is under 10% (Valles et al. 2010). However, pockets of high infection rates (20–50%) have been reported in the US (Valles et al. 2010) and Argentina (Valles et al. 2009). Several factors may contribute to higher *Solenopsis invicta* virus 3 infection rates, including social form and temperature. Higher infection rates typically are observed in polygyne colonies compared with monogyne colonies (Valles et al. 2010). However, we did not examine

the social form of the female alates in this study. *Solenopsis invicta* virus 3 infection exhibits a negative relationship with temperature and is more prevalent during winter months in Florida (Valles et al. 2010).

While several of these cities had high infection rates (Tallahassee, Quincy, and Gainesville), some cities exhibited low infection rates; surprisingly Jacksonville, Pensacola, and Jay had no detectable infection. We fully expected that large cities such as Pensacola and Jacksonville would have *Solenopsis invicta* virus 3 present given the frequent travel of people transporting all sorts of materials between these cities and cities infected with *Solenopsis invicta* virus 3. The absence of *Solenopsis invicta* virus 3 in these cities may be attributed to the small number of colonies sampled. Interestingly, in the paired cities Pensacola (urban) and Jay (rural), *Solenopsis invicta* virus 3 was not detected in female alates, which may represent an actual absence of the virus in these areas.

Knowing the geographic *Solenopsis invicta* virus 3 incidence affords the opportunity to conduct future research comparing the health of



**Fig. 2.** Representative agarose gel depicting detection of *Solenopsis invicta* virus 3 as determined by polymerase chain reaction using cDNA generated from RNA purified from female alates of *Solenopsis invicta*. Arrow represents the 258 bp amplicon from *Solenopsis invicta* virus 3. Lane 1 is a 1 kb DNA ladder and lane 12 is non-template control.

*S. invicta* colonies in cities that have high infection rates versus those that had no infection rate. One could also examine the health of *S. invicta* colonies in cities without the virus such as Jay, Pensacola, or Jacksonville, and then introduce the virus to those cities. One could then compare the before and after health changes of colonies in the field afflicted by *Solenopsis invicta* virus 3. The introductions could follow protocols used by studies that introduced the virus to Gainesville (Valles et al. 2009) and California (Oi et al. 2019).

In conclusion, the results of this study indicate that *Solenopsis invicta* virus 3 is well distributed in female alates in North Florida. Thus, it is likely that *Solenopsis invicta* virus 3 is being disseminated and maintained in *S. invicta* by this life stage. However, female alates from some areas were devoid of *Solenopsis invicta* virus 3, suggesting that they may be good areas for augmentative release of the virus as a natural control agent.

## Acknowledgments

We gratefully acknowledge Worrel Deidrick (Florida A&M University) for assisting with the ant collections, Mehboob Sheikh (Florida A&M University Viticulture Center for Small Fruit Research) for allowing us to use the molecular equipment in his laboratory, Janice Peters for critically reviewing an earlier version of this manuscript and for many helpful suggestions for improvement. We also would like to acknowledge Lambert Kanga (Florida A&M University Center of Biological Control), Oghenekome Onokpise (Florida A&M University), and Jesusa Legaspi (USDA-ARS) for their input over the course of the research project. Partial funding for this study was provided by the USDA, NIFA, CPPM (agreement number: 2017-70006-27290). Mention of a trade or firm name does not constitute an endorsement but is mentioned to benefit the reader.

The dataset may be found here: <https://data.nal.usda.gov/dataset/data-examining-prevalence-solenopsis-invicta-virus-3-solinviviridae-invictavirus-solenopsis-invicta-hymenoptera-formicidae-alates-collected-north-florida> (last accessed 16 Dec 2020).

## References Cited

- Caldera EJ, Ross KC, Detter CJ. 2008. Putative native source of the invasive fire ant *Solenopsis invicta* in the USA. *Biological Invasions* 10: 1457–1479.
- Drees MB, Gold RE. 2003. Development of integrated pest management programs for the red imported fire ant (Hymenoptera: Formicidae). *Journal of Entomological Science* 38: 170–180.
- Du-Prel J-B, Röhrig B, Hommel G, Blettner M. 2010. Choosing statistical tests: part 12 of a series on evaluation of scientific publications. *Deutsches Arzteblatt International* 107: 343–348.
- Hashimoto Y, Valles SM. 2008. Detection and quantitation of *Solenopsis invicta* virus-2 genomic and intermediary replicating viral RNA in fire ant workers and larvae. *Journal of Invertebrate Pathology* 98: 243–245.
- Oi DH, Valles SM, Porter SD, Cavanaugh C, White G, Henke J. 2019. Introduction of fire ant biological controls agents into Coachella Valley of California. *Florida Entomologist* 102: 284–286.
- Pereira RM. 2003. An areawide suppression of fire ant population in pastures: project update. *Journal of Agriculture and Urban Entomology* 20: 123–130.
- Porter SD, Savignano DA. 1990. Invasion of polygyne fire ants decimates native ants and disrupts arthropod community. *Ecological Society of America* 71: 2095–2106.
- Porter SD, Fowler HG, Mackay WP. 1992. A comparison of fire ant densities in North and South America (Hymenoptera: Formicidae). *Journal of Economic Entomology* 85: 1154–1161.
- Porter SD, Williams DF, Patterson RS, Fowler HG. 1997. Intercontinental differences in the abundance of *Solenopsis* fire ants (Hymenoptera: Formicidae): escape from natural enemies. *Environmental Entomology* 26: 373–384.
- Summerlin JW, Hung CF, Vinson SB. 1977. Residues in non-target ants, species simplification and recovery of populations following aerial applications of mirex. *Environmental Entomology* 6: 193–197.
- Tschinkel WR. 1998. The reproductive biology of fire ant societies. *Bioscience* 48: 593–605.
- Tschinkel WR. 2006. *The fire ants*. Belknap/Harvard University Press, Cambridge, Massachusetts, USA.
- Valles SM. 2012. Positive-strand RNA viruses infecting the red imported fire ant, *Solenopsis invicta*. *Psyche: A Journal of Entomology*. Article ID 821591. doi: [10.1155/2012/821591](https://doi.org/10.1155/2012/821591)
- Valles SM, Oi DH, Porter SD. 2010. Seasonal variation and the co-occurrence of four pathogens and a group of parasites among monogyne and polygyne fire ant colonies. *Biological Control* 54: 342–348.
- Valles SM, Varone L, Ramirez L, Briano J. 2009. Multiplex detection of *Solenopsis invicta* viruses-1, -2, and -3. *Journal of Virology* 162: 276–279.
- Valles SM, Strong CA, Dang PM, Hunter WB, Pereira RM, Oi DH, Shapiro AM, Williams DF. 2004. A picorna-like virus from the red imported fire ant, *Solenopsis invicta*: initial discovery, genome sequence and characterization. *Journal of Virology* 328: 151–157.
- Williams DF, Collins HL, Oi DH. 2001. The red imported fire ant (Hymenoptera: Formicidae): an historical perspective of treatment programs and the development of chemical baits for control. *American Entomologist* 47: 146–159.