

Allergy to Crickets: A Review

Author: Pener, Meir Paul

Source: Journal of Orthoptera Research, 25(2) : 91-95

Published By: Orthopterists' Society

URL: <https://doi.org/10.1665/034.025.0208>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Allergy to crickets: a review

MEIR PAUL PENER

Department of Cell and Developmental Biology, The Hebrew University of Jerusalem, Jerusalem, Israel. E-mail: pener@mail.huji.ac.il

Abstract

Cricket allergy is less severe and less common than allergy to locusts and grasshoppers. A partial cross-reactivity exists between cricket and grasshopper allergens. Cricket allergens are proteinaceous compounds, but their nature is insufficiently known; arginine kinase and hexamerin 1B may play a role. Occupational allergy, *i.e.* allergy of personnel working with rearing and breeding of cricket colonies, is the subject of the majority of reports on cricket allergy. Frequent handling of crickets (for example as fish baits) may inflict allergy which may be considered as a kind of occupational allergy. Crickets are edible insects and are widely consumed in many parts of the world. Nevertheless, food allergy to crickets seems to be relatively rare.

Key words

occupational allergy, cricket allergens, hypersensitivity pneumonitis, Gryllidae, *Acheta domesticus*, *Gryllus bimaculatus*, *Gryllus campestris*

Introduction

The present review is a continuation of my former review on allergy to locusts and acridid grasshoppers (Pener 2014). This former review deals with allergy in general and allergy to insects and these subjects are not repeated here. Crickets (Ensifera: Gryllidae) and short-horned grasshoppers or locusts (Caelifera: Acrididae) represent two different suborders in the same insect order (Orthoptera). In this review I compile the relevant literature related to the identity of some cricket allergens, modes of infliction of cricket allergies and management of these allergies.

Allergy to crickets (Gryllidae) is less investigated than allergy to acridids. Nevertheless, cricket allergy is mentioned in many relevant reviews, sometimes including the name of the allergen species (Seward 1999, Barre *et al.* 2014, Dutau & Lavaud 2014, van Huis 2016), but mostly as a general example of allergy to orthopterans (Mathews 1989, Mairesse & Ledent 2002, Micek *et al.* 2014, Stanhope *et al.* 2015), or just to insects/arthropods (Chan-Yeung & Lam 1986, Chan-Yeung 1990, Bellas 1990, van Kampen *et al.* 2000, Barletta & Pini 2003, Jeong *et al.* 2007, Bousquet *et al.* 2008, Yong & Jeong 2009, Moscato *et al.* 2011, and others).

Cricket allergens

Sensitivity to crickets is usually less severe and less common than that to grasshoppers or locusts (Prasad *et al.* 2001, 2009, Patel & Choudhary 2012).

Investigating allergic-asthmatic children in Cincinnati, Ohio,

and using RAST (radioallergosorbent test) of allergen-specific immunoglobulin E (IgE), Lierl *et al.* (1994) revealed a considerable cross-reactivity between cricket and grasshopper allergens. Unfortunately, the authors do not provide the name of the species used in their study, neither for the cricket, nor for the grasshopper.

The cricket-grasshopper cross allergy is not surprising. Cross allergy among different species of insects/arthropods is well known (see for example, Panzani & Ariano 2001). Cross allergy means that the same allergen is present in different taxa and it causes allergy to these taxa. Recently, cross allergy was demonstrated between *Macrobrachium* prawns and the cricket, *Gryllus bimaculatus*; arginine kinase constituted the common allergen (Srinroch *et al.* 2015). This finding contradicts the previous claim that there is no cross allergy between prawns and crickets (Linares *et al.* 2008). However, the cross-reactivity between grasshopper and cricket allergens does not seem to be complete. Comparison of the molecular mass of grasshopper/locust allergens (see Table 1 by Pener 2014) with those of cricket allergens (Bartra *et al.* 2008) reveals partial divergence. On the other hand, similarity (at 65-70 and 130 kDa) exists between grasshopper/locust allergens and hypersensitivity pneumonitis induced antigens, related to arginine kinase, in *Acheta domesticus* (see Park *et al.* 2014).

Srinroch *et al.* (2015) identified hexamerin 1B as a novel and specific allergen in *G. bimaculatus*.

It may be noted that thermal processing may alter allergens. Based on the Bombay locust, *Patanga succincta* (an acridid), Phiriyangkul *et al.* (2015) found differences in the allergens between raw and fried insects. For example, arginine kinase was identified in the former, but not in the latter. Considering that arginine kinase seems to be a common allergen to acridids and gryllids (and other arthropods), it may be assumed that similar changes occur in crickets with thermal procession.

Occupational allergy to crickets

Cazort & Johnston (1955, cited by Bellas 1981) reported sensitivity to crickets in a man raising crickets for fish bait. This was the first report on occupational allergy to crickets.

Two and a half decades later, Bagenstose *et al.* (1980) published two case reports on occupational allergy to crickets. The patients were exposed to crickets while employed at the University of Michigan Amphibian Facility where massive cricket rearing and breeding took place for feeding frogs. Both patients developed allergic rhinitis and bronchial asthma. Temporary cessation of working with crickets resulted in disappearance of these symptoms which reappeared after resumption of the work. The authors prepared an extract of the cricket, *Acheta domesticus*, and tested sensitivity of the patients

by several methods, among them *in vivo* bronchial inhalation challenge and skin prick test, as well as by *in vitro* RAST that revealed specific IgE antibodies in the patients' serum. Interestingly, out of 11 other workers at the same facility, three exhibited positive skin prick test and two out of these three showed positive RAST, without overt symptoms of allergy.

Linares *et al.* (2008) reported a case of a 28-year-old woman with occupational rhinitis and asthma, who was exposed for three years to three species of crickets, *G. campestris* (often connoted as *Acheta campestris*), *G. bimaculatus* and *A. domesticus*, in a farm raising cricket colonies. The authors prepared extracts of each of these three species of crickets with two types of protein extraction. They also made extracts of the food of the crickets. *In vivo* tests included skin prick test, lung function test and bronchial challenge testing. The results of the prick tests were positive for each of the three cricket species, but negative for their food. Specific IgE for each cricket species was determined by *in vitro* enzyme allergosorbent test (EAST), resulting in the values of 2.9, 2.4 and 5.4 kU/L, for *G. campestris*, *G. bimaculatus* and *A. domesticus*, respectively. EAST inhibition assays found a high degree of cross-reactivity among the three species of crickets, up to 86% for *A. domesticus* and *G. campestris*, some cross-reactivity between crickets and the firebug *Pyrrhocoris apterus*, but no cross-reactivity was detected between crickets and prawn, squid, mite or pollen (but see above positive cross-reactivity between *G. bimaculatus* and *Macrobrachium* spp. prawns found by Srinroch *et al.* 2015). Sodium dodecylsulfate – polyacrylamide gel electrophoresis (SDS-PAGE) and IgE immunoblotting showed similarity in IgE binding bands with the three species of whole body cricket extracts. Bands of molecular mass of 64 and 78 kDa were found in non-reducing conditions and bands of 52, 58, 80, 97 and 107 kDa in reducing conditions.

Another case report was published by Bartra *et al.* (2008). A 28-year-old man who had worked in a reptile shop for seven years, where he fed the reptiles with *Acheta campestris*, showed allergy to these crickets. The symptoms of allergy appeared in the fourth year of his work, that is, after a latent period of three years. These symptoms were asthma and rhino-conjunctivitis caused by inhalation and contact urticaria caused by handling the crickets. Skin prick test with a manufactured cricket extract was positive, whereas it was negative for control individuals, for cornmeal (food of the crickets) extracts and for a set of common inhalant allergens. Respiratory function test showed a drop of more than 20% of peak expiratory flow during work period, but it was normal during the weekend when the patient was free from crickets. A specific nasal test with cricket extract yielded positive results whereas the test with the control was negative. Immunoblot experiments of the protein profile of the cricket extract showed several bands with IgE binding, prominently with molecular masses of 17, 32, 47 and 62 kDa.

Investigating the general situation of allergy among pet shop workers, Ranström *et al.* (2011) distributed a questionnaire and received suitable answers from 59 subjects from 24 pet shops around Stockholm, Sweden. The authors tested in details some of the most common allergens like rat and mouse urinary allergens and among insects the darkling beetles (Tenebrionidae), specifically the mealworm beetle (*Tenebrio molitor*) and *Zophobas* droppings and larvae. No special attention was paid to crickets, but the authors mention in a single sentence that "....crickets were reported by several workers to have caused occasional strong allergic reactions" (Ranström *et al.* 2011, p. 1086).

Harris-Roberts *et al.* (2011) made a somewhat similar study, but instead of pet shop workers, they investigated the workers of a specialist insect breeding facility in the UK. Thirty-two persons

answered a questionnaire on work-related respiratory symptoms and underwent lung function test; 18 persons provided serum for specific IgE determination. Sensitivity was found to locusts and mealworm beetles. However, a single asthmatic cricket handling worker was submitted to a more detailed clinical investigation. This person suffered from respiratory problems over the preceding two years with increasing periods of absence from work related to sickness. Expiratory wheeze and reduced lung function were found and the patient's serum revealed IgE to bran, locust and cricket (species not stated). After five months without contact to insects, the work-related respiratory symptoms disappeared.

Mairesse *et al.* (2014) studied allergy by skin prick test in 29 workers in a laboratory of entomology and found two persons showing double sensitivity to the mealworm beetle, *T. molitor*, and the cricket, *A. domesticus*.

Subacute hypersensitivity pneumonitis caused by crickets was recently described by Park *et al.* (2014). This disease is an inflammation of the lungs due to hypersensitivity to inhaled allergens. The patient having the disease was described as a 63-year-old male who previously owned an avian pet shop. Although avian proteins are known allergens, the authors' thorough investigation revealed that in this case the source of the allergen was the house cricket, *A. domesticus*, maintained in colonies by the thousands in boxes. The crickets served for feeding the pets. Cricket proteins were extracted and high titer antibodies were found against the extract with specific antigen-antibody precipitins. Additional investigations identified arginine kinase at 65-70, 130, and 160 kDa mass weights as the major allergen. Other cricket proteins may have acted as minor allergens contributing to the severity of the disease. The patient reduced his exposure to crickets and used a face mask resulting in decline of the disease after six months.

Immediate hypersensitivity to crickets

Immediate type hypersensitivity is characterized by appearance of allergic symptoms within a short while after exposure to the allergen, sometimes just minutes, usually less than one hour, though additional allergic symptoms may appear later.

In some instances these symptoms are inflicted by the classic allergic pathway, that is, IgE mediated. For example, immediate systemic reaction of allergy to bee or wasp stings are mostly (though not necessarily) IgE mediated (Przybilla & Ruëff 2012). In some other instances, the symptoms are of typical allergy, but the underlying mechanisms are not mediated by IgE; Kowalski *et al.* (2011) and Gimenez-Arnau *et al.* (2010) discuss some of such mechanisms.

There is a single case in the literature reporting immediate hypersensitivity to cricket (Harfi 1980). A 16-year-old male on two occasions while fishing and using crickets (species not stated) for baits, developed within 30 minutes lacrimation, angioedema of the eyelids, rhinorrhea and some additional symptoms; cough and wheeze followed in about five hours. Confirmation of the diagnosis was carried out, using cricket whole body extract, by challenge to the right conjunctiva that caused rhinorrhea, by scratch test, and by *in vitro* histamine release (that may, or may not mediated, by the IgE pathway), as well as by other tests.

Allergy associated with crickets in outdoor air

Lierl *et al.* (1994) investigated the effect of outdoor airborne allergens from moth, ant, house fly, grasshopper, cricket (species not stated) and spider. They compared by RAST the effect in "allergic asthmatic" children to that in "nonallergic nonasthmatic" children.

Roughly, the two clusters may be considered as atopic and non-atopic children, respectively. The authors employed several extraction procedures for each group of arthropods and used the most potent extract. Concerning crickets, 8 out of 41 (19%) of allergic-asthmatic children exhibited positive RAST, revealing cricket sensitive IgE in their serum. In contrast, none of the 25 nonallergic-nonasthmatic children showed positive RAST. In this study, cross-reactivity of cricket and grasshopper extracts was revealed, expressed by 59% average RAST inhibition (see also section on cricket allergens, above).

As much as four years before the publication of the article by Lierl *et al.* (1994) an abstract was published by the same authors (Lierl *et al.* 1990) estimating the amount of airborne insect-originating particles in outdoor air. They found 9-300 ng per cubic feet per week for cricket (species not stated).

Food allergy to crickets

A considerable part of the world population, in China, the Far East, and the Middle East, in Africa, Mexico, and South America, as well as indigenous people in Australia, feed on edible insects (Srivastava *et al.* 2009, Ramos-Elorduy 2009, Chen *et al.* 2009, Gahukar 2011, van Huis 2013 and others). Crickets (Gryllidae) and mole crickets (Gryllotalpidae) are important constituents of insect diet (Chen *et al.* 2009, Gahukar 2011, Belluco *et al.* 2013, Barennes *et al.* 2015, Megido *et al.* 2016), having high nutritional value (Belluco *et al.* 2013, Yi *et al.* 2013, van Huis 2013, Barennes *et al.* 2015, Megido *et al.* 2016). Rumpold & Schlüter (2013) published a comprehensive review on nutritional composition and nutritive values of edible insects, often referring to Orthoptera as a single unit (crickets, grasshoppers and locusts together), but presenting some specific details on *A. domesticus* from various sources.

Considering the widespread usage of crickets as human food, cases of food allergy to crickets are relatively rare. Piromrat *et al.* (2008) reported seven cases of anaphylaxis within a two-year period in an emergency department of a hospital in Bangkok, Thailand, caused by consumption of fried grasshoppers and crickets. Unfortunately, the authors did not make distinction between grasshoppers and crickets. Possibly, physicians often do not have the knowledge, or they are not confident, to distinguish among different suborders and families of Orthoptera. Ji *et al.* (2009), summarizing anaphylaxes inflicted by food consumption in China, do not mention crickets, though they report on 27 cases of anaphylaxis caused by eating locusts and another 27 cases due to eating grasshoppers.

Allergy to crickets without defined source

In the studies summarized below, possible former exposure to the allergen was not considered, no species was stated, possible cross allergy was disregarded and no statistical analysis was made. Also, the term grasshopper may mean acridid and/or tettigoniid (see Pener 2014).

Prasad *et al.* (2001) carried out 5,760 intradermal skin tests with 68 different antigens, including of five insects, tested on 108 patients with bronchial asthma who attended a medical college in Lucknow, India. Concerning crickets (species not stated), 31 tests were made and out of these three (9.7%) yielded definitely positive response. For comparison, 31 tests with grasshopper antigen resulted in 11 (35.5%) such response. These findings indicate much higher sensitivity to grasshoppers than to crickets.

Eight years later, Prasad *et al.* (2009) reported a similar study in Lucknow, using skin prick test instead of intradermal injected skin test. Forty-eight patients with nasobronchial allergy were tested.

Eight of these (16.6%) exhibited definitely positive sensitivity to cricket antigen, whereas 10 (20.8%) showed the same to grasshopper antigen.

Another study was conducted by Patel & Choudhary (2012), testing by skin prick various antigens on patients with nasobronchial allergy in Gujarat, India. The authors found two cases (8.3%) of definitely positive response to cricket antigen. Again, for comparison, they found six (25%) cases of such response to grasshopper antigen. Also, sensitivity was higher to many insects, namely locust, moth, butterfly, mosquito, house fly, honey bee, hornet, yellow wasp and ant, than to cricket. These findings are somewhat controversial to Hosen's (1970) older results using a provocative nasal test conducted in 174 patients with perennial respiratory allergy to some insects of Southeast Texas. Out of the 174 patients, only 69 (33.2%) showed sensitivity to the powdered insect antigens tested. Without stating the species, sensitivity to crickets was more common than to red ant, mosquito, cockroach, moth, fire ant and house fly. Only sensitivity to spider (not an insect, but an arachnid) was more common than that to cricket. Hosen (1970) did not test grasshopper/locust antigen.

Management of cricket allergy

Very few publications deal with management/treatment of allergy to crickets. The patient reported by Linares *et al.* (2008) stopped her exposure to crickets and was treated with inhaled bronchodilator and corticosteroid. She became asymptomatic and only occasionally needed inhaled salbutamol. The patient of Harris-Robert *et al.* (2011) was moved to another area of the facility with no direct exposure to insects. An examination five months later revealed that the cricket specific allergy symptoms had disappeared, without change in his inhaled asthma treatment (the patient was diagnosed with asthma 15 years previously). A patient with subacute hypersensitivity pneumonitis caused by house cricket (*A. domesticus*), as reported by Park *et al.* (2014), preferred reducing exposure to crickets and respiratory precaution by face mask instead of taking medicine. Evaluation of the response showed marked improvement of the patient's clinical symptoms.

These examples clearly indicate that the best treatment to cricket allergy is restraining from exposure to crickets. Avoidance of the allergen is a general rule to all kinds of allergy. Sometimes, however, avoidance may be practically difficult. A scientist may not like to change the subject of his/her research. Nevertheless, many entomologists have had to leave their jobs (Stanhope *et al.* 2015 and references therein). A technician may be afraid to lose a job with a reasonable salary.

For personal protection, gloves and dusk mask covering the nose and mouth may be sufficient. A hat is also advisable because dust sticking to hairs may contaminate pillows. Well-fitting laboratory glasses or a whole face mask (instead of a dusk mask that does not cover the eyes) help to avoid allergic conjunctivitis.

In my former article devoted to allergy to locusts and acridid grasshoppers, I described extreme protective measures; complete outfit laboratory coat, trousers and overshoes, as well as specially designed insect room and a changing room (Pener 2014 and references therein). These measures may overshoot the needs of cricket allergy which is less severe and less common than locust/grasshopper allergy. However, they should be implemented partially or fully in a case of a large scale insect rearing and breeding facility.

Allergy to crickets can be treated, like treatments of other allergies, by antihistamines, corticosteroids and bronchodilators. In life threatening cases, like anaphylaxis, intramuscular epinephrine

may be administered (Piromrat *et al.* 2008). It should be stressed that these medicines reduce the symptoms of allergy, but they are not curative. Curative treatments include allergen specific immunotherapy, also termed desensitization, resulting in hyposensitivity to the allergen (see Fujita *et al.* 2012, Berin & Sampson 2013, and in relation to arthropods Schwartz 1990 and Kagen 1990). According to Kagen (1990), Cazort & Johnston (1955) describing a case of inhalant sensitivity to crickets, commented that desensitization was effective.

Additional references regarding control of the environment, prevention, protective measures and therapy are found in several older publications (Bellas 1990, Chan-Yeung 1990, Seward 1999 and reference therein); all these references are not specific to crickets, but related to allergy to insects.

Conclusions

Allergy to crickets is relatively infrequent and it is usually less severe than most common allergies. Nevertheless, crickets are listed as an allergen in many recent reviews on the subject. Cricket allergy is rather mild compared to allergy to locusts and grasshoppers, although there is a partial cross-reactivity between acridid and cricket allergy. Most relevant publications report on occupational allergy to crickets; food allergy seems to be rare despite the fact that crickets are edible insects consumed in many Asian, African and South American countries.

At the species level *Gryllus campestris* (= *Acheta campestris*), *G. bimaculatus* and *Acheta domesticus*, were found to inflict cricket allergy. Although a high level of cross-reactivity was found among these three species, this cross-reactivity was not total. These three are the most commonly maintained colonies of cricket species in the world. To the best of my knowledge, no other cricket species have been investigated as allergens.

Very little information is available on the chemical identity of cricket allergens. It seems that arginine kinase is a considerable cricket allergen, but arginine kinase is a known allergen in many arthropods. Recently, hexamerin B1 was found as a cricket specific allergen, however, hexamerin B1 is known as an allergen also from some non-gryllid insects.

Acknowledgements

The research expenses for this review were covered by a grant to retired faculties of the Hebrew University of Jerusalem. Thanks are due to three anonymous reviewers whose comments improved the manuscript.

References

- Bagenstose III A.H., Mathews K.P., Homburger H.A., Saaveard-Delgado A.P. 1980. Inhalant allergy due to crickets. *Journal of Allergy and Clinical Immunology* 65: 71-74.
- Barennes H., Phimmasane M., Rajaonarivo C. 2015. Insect consumption to address undernutrition, a national survey on the prevalence of insect consumption among adults and vendors in Laos. *Plos One* 10(8): e0136458, doi: 10.1371/journal.pone.0136458.
- Barletta B., Pini C. 2003. Does occupational exposure to insects lead to species-specific sensitization? *Allergy* 58: 868-870.
- Barre A., Caze-Subra S., Gironde C., Bienvenu F., Bienvenu J., Rougé P. 2014. Entomophagie et risque allergique. *Revue Française d'Allergologie* 54: 315-321.
- Bartra J., Carnés J., Muñoz-Cano R., Bissinger I., Picado C., Valero A.L. 2008. Occupational asthma and rhinoconjunctivitis caused by cricket allergy. *Journal of Investigational Allergology and Clinical Immunology* 18: 141-142.
- Bellas T.E. 1981. Insects as a Cause of Inhalant Allergies - A Bibliography. Commonwealth Scientific and Industrial Research Organization, Australia, Division of Entomology Report No. 25.
- Bellas T.E. 1990. Occupational inhalant allergy to arthropods. *Clinical Reviews in Allergy* 8: 15-29.
- Belluco S., Losasso C., Maggioletti M., Alonzi C.C., Paoletti M.G., Ricci A. 2013. *Comprehensive Reviews in Food Science and Food Safety* 12: 296-313.
- Berin M.C., Sampson H.A. 2013. Food allergy: an enigmatic epidemic. *Trends in Immunology* 34: 290-397.
- Bousquet J., Khaltayev N., Cruz A.A., Denburg J., Fokkens W.J., Togias A., Zuberbier T., Baena-Cagnani C.E., Canonica G.W., van Weel C. and 86 more authors. 2008. Allergic rhinitis and its impact on asthma (ARIA) 2008. *Allergy* 63 (Suppl. 86): 8-160.
- Cazort A., Johnston T.G. 1955. Sensitivity to crickets. *Letters to the International Correspondence Society of Allergists, Series XVIII*: 55. (Not seen in original; as cited by Bellas 1981 and Kagen 1990).
- Chan-Yeung M. 1990. Occupational asthma. *Chest* 98 (Suppl.): 148S-161S.
- Chan-Yeung M., Lam S. 1986. Occupational asthma. *American Review of Respiratory Disease* 133: 686-703.
- Chen X., Feng Y., Chen Z. 2009. Common edible insects and their utilization in China. *Entomological Research* 39: 299-303.
- Dutau G., Lavaud F. 2014. Des allergènes émergents: le fénugrec, *Tenebrio molitor*, la méthylisothiazolinone, le macrogol, etc. Et les grillons? *Revue Française d'Allergologie* 54: 287-290.
- Fujita H., Soyka M.B., Akdis M., Akdis C.A. 2012. Mechanisms of allergen-specific immunotherapy. *Clinical and Translational Allergy* 2: article 2.
- Gahukar R.T. 2011. Entomophagy and human food security. *International Journal of Tropical Insect Science* 31: 129-144.
- Gimenez-Arnau A., Maurer M., De La Cuadra J., Maibach H. 2010. Immediate contact skin reactions, an update of contact urticaria, contact urticaria syndrome and protein contact dermatitis - "a never ending story". *European Journal of Dermatology* 20: 552-562.
- Harfi H.A. 1980. Immediate hypersensitivity to cricket. *Annals of Allergy* 44: 162-163.
- Harris-Roberts J., Fishwick D., Tate P., Rawbone R., Stagg S., Barber C.M., Adishes A. 2011. Respiratory symptoms in insect breeders. *Occupational Medicine* 61: 370-373.
- Hosen H. 1970. Insects as a cause of inhalant respiratory allergy. *Annals of Allergy* 28: 596-599.
- Jeong K.Y., Hong C.-S., Yong T.-S. 2007. Domestic arthropods and their allergens. *Protein and Peptide Letters* 14: 934-942.
- Ji K., Chen J., Li M., Liu Z., Wang C., Zhan Z., Wu X., Xia Q. 2009. Anaphylactic shock and lethal anaphylaxis caused by food consumption in China. *Trends in Food Science & Technology* 20: 227-231.
- Kagen S.L. 1990. Inhalant allergy to arthropods. *Clinical Reviews in Allergy* 8: 99-125.
- Kowalski M.L., Makowska J.S., Blanca M., Bavbek S., Bochenek G., Bousquet J., Bousquet P., Celik G., Demoly P., Gomes E.R., Nizankowska-Mogilnicka E., Romano A., Sanchez-Borges M., Sanz M., Torres M.J., De Weck A., Szczeklik A., Brockow K. 2011. Hypersensitivity to nonsteroidal anti-inflammatory drugs (NSAIDs) - classification, diagnosis and management: review of the EAACI/ENDA* and GA2LEN/HANNA*. *Allergy* 66: 818-829.
- Lierl M.B., Riordan M.M., Fischer T.J. 1990. Concentration of airborne insect-derived particles in outdoor air. *Journal of Allergy and Clinical Immunology* 85 (issue 1, part 2): 246 (abstract no. 412).
- Lierl M.B., Riordan M.M., Fischer T.J. 1994. Prevalence of insect allergen-specific IgE in allergic asthmatic children in Cincinnati, Ohio. *Annals of Allergy* 72: 45-50.
- Linares T., Hernandez D., Bartolome B. 2008. Occupational rhinitis and asthma due to crickets. *Annals of Allergy, Asthma & Immunology* 100: 566-569.

- Mairesse M., Ledent C. 2002. Allergie et activités halieutiques. *Allergie et Immunologie* 34: 245-247.
- Mairesse M., Debaugnies F., Doyen V., Ledent C., Michel O., Corazza F., Francis F. 2014. Risque allergique des insectes en alimentation humaine. *Revue Française d'Allergologie* 54: 228 (abstract no. Ali-3).
- Mathews K.P. 1989. Inhalant insect-derived allergens. *Immunology and Allergy Clinics of North America* 9: 321-338.
- Megido R.C., Alabi T., Nieuw C., Blecker C., Danthine S., Bogaert J., Haubruge É., Francis F. 2016. Optimisation of a cheap and residential small-scale production of edible crickets with local by-products as an alternative protein-rich human food source in Ratanakiri Province, Cambodia. *Journal of the Science of Food and Agriculture* 96: 627-632. doi: 10.1002/jsfa.7133
- Micek J., Rop O., Borkovcova M., Bednarova M. 2014. A comprehensive look at the possibilities of edible insects as food in Europe – a Review. *Polish Journal of Food and Nutritional Science* 64: 147-157.
- Moscato G., Rolla G., Siracusa A. 2011. Occupational rhinitis: consensus on diagnosis and medicolegal implications. *Current Opinion in Otolaryngology & Head and Neck Surgery* 19: 36-42.
- Panzani R.C., Ariano R. 2001. Arthropods and invertebrates allergy (with the exclusion of mites): the concept of panallergy. *Allergy* 56, Suppl. 69: 1-22.
- Park M., Boys E.L., Yan M., Bryant K., Cameron B., Desai A., Thomas P.S., Tedia N.T. 2014. Hypersensitivity pneumonitis caused by house cricket, *Acheta domesticus*. *Journal of Clinical & Cellular Immunology* 5: article 248.
- Patel A., Choudhary S. 2012. Prevalence of allergen sensitivity in nasobronchial allergy in Gujarat, India. 2012. *National Journal of Medical Research* 2: 431-434.
- Pener M.P. 2014. Allergy to locusts and acridid grasshoppers: a review. *Journal of Orthoptera Research* 23: 59-67.
- Phiriyangkul P., Srinroch C., Srisomsap C., Chokchaichamnankit D., Punyarit P. 2015. Effect of food thermal processing on allergenicity proteins in Bombay locust (*Patanga succincta*). *International Journal of Food Engineering* 1: 23-28.
- Piromrat K., Chinratanapisit S., Trathong S. 2008. Anaphylaxis in an emergency department: A 2-year study in a tertiary-care hospital. *Asian Pacific Journal of Allergy and Immunology* 26: 121-128.
- Prasad R., Kumar R., Verma S.K., Pandey U.S. 2001. A study of skin sensitivity to various allergens by intradermal test in patients of bronchial asthma. *Indian Journal of Allergy, Asthma and Immunology* 15: 17-21.
- Prasad R., Verma S.K., Dua R., Kant S., Kushwaha R.A.S., Agarwal S.P. 2009. A study of skin sensitivity to various allergens by skin prick test in patients of nasobronchial allergy. *Lung India* 26: 70-73.
- Przybilla B., Ruëff F. 2012. Insect stings. *Deutsches Ärzteblatt International* 109: 238-248.
- Ramos-Elorduy J. 2009. Anthro-entomophagy: Cultures, evolution and sustainability. *Entomological Research* 39: 271-288.
- Ranström A., Olsson M., Hedrén M., Johansson S.G.O., van Hage M. 2011. Pet shop workers: exposure, sensitization and work-related symptoms. *Allergy* 66: 1081-1087.
- Rumpold B.A., Schlüter O.K. 2013. Nutritional composition and safety aspects of edible insects. *Molecular Nutrition and Food Research* 57: 802-823.
- Schwartz H.J. 1990. Inhalant allergy to arthropods. *Clinical Reviews in Allergy* 8: 3-13.
- Seward J.P. 1999. Occupational allergy to animals. *Occupational Medicine: State of the Art Reviews* 14: 285-303.
- Srinroch C., Srisomsap C., Chokchaichamnankit D., Punyarit P., Phiriyangkul P. 2015. Identification of novel allergen in edible insect, *Gryllus bimaculatus* and its cross-reactivity with *Macrobrachium* spp. allergens. *Food Chemistry* 184: 160-166.
- Srivastava S.K., Babu N., Pandey H. 2009. Traditional insect bioprospecting – As human food and medicine. *Indian Journal of Traditional Knowledge* 8: 485-494.
- Stanhope J., Carver S., Weinstein P. 2015. The risky business of being an entomologist: A systematic review. *Environmental Research* 140: 619-633.
- van Huis A. 2013. Potential of insects as food and feed in assuring food security. *Annual Review of Entomology* 58: 563-583.
- van Huis A. 2016. Edible insects are the future? *Proceedings of the Nutrition Society*, 12 pp. Available on CJO2016. doi: 10.1017/S0029665116000069.
- van Kampen V., Merget R., Baur X. 2000. Occupational airway sensitizers: An overview on the respective literature. *American Journal of Industrial Medicine* 38: 164-218.
- Yi L., Lakemond C.M.M., Sagis L.M.C., Eisner-Schadler V., van Huis A., van Boekel M.A.J.S. 2013. Extraction and characterisation of protein fractions from five insect species. *Food Chemistry* 141: 3341-3348.
- Yong T.-S., Jeong K.Y. 2009. Household arthropod allergens in Korea. *Korean Journal of Parasitology* 47, Suppl: S143-S153.

