

Errata: Tettigoniidae (Orthoptera) from Jordan with description of new species and redescription of less known species. (J. Orthoptera Res. 10(1): 25–37)

Authors: Bader, Ahmad Katbeh, and Massa, Bruno

Source: Journal of Orthoptera Research, 10(2) : 376-377

Published By: Orthopterists' Society

URL: [https://doi.org/10.1665/1082-6467\(2001\)010\[0376:ETOFJW\]2.0.CO;2](https://doi.org/10.1665/1082-6467(2001)010[0376:ETOFJW]2.0.CO;2)

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Errata: Tettigoniidae (Orthoptera) from Jordan with description of new species and redescription of less known species. (J. Orthoptera Res. 10(1): 25-37)

AHMAD KATBEH BADER AND BRUNO MASSA

Diogena fausta (Burmeister). This species was previously recorded by Ayal *et al.* (1999) from Israel, thus its actual middle eastern distribution covers Israel and Jordan.

Tylopsis lilifolia (Fabricius). A more detailed examination of Jordanian material revealed that specimens collected at Al Jubayhah, As Salt, Irbid, Jarash and Umm Qays belong to typical *T. lilifolia*, while those collected at Amman and Jordan Valley match the characteristics of *T. peneri*, a specific name proposed by Ragge (1974) for southern Israel specimens, previously cited by himself (Ragge 1964) as a "deserticolous form". This form was also reported by Popov (1981) from Saudi Arabia.

Scotodrymadusa philbyi (Uvarov). Though Uvarov (1939) wrote that the female was unknown, in reality he had already described it (Uvarov 1933). Thus, our description of the female is not new. Additionally, the status of *Paradrymadusa philbyi* and *P. syriaca* Pictet was already changed by Ramme (1939) into, respectively, *Scotodrymadusa philbyi* and *S. syriaca*. These changes are overlooked by Otte (1997) and Naskrecki & Otte (1999), but recorded by Ayal *et al.* (1999).

Parapholidoptera Maran and *Uvarovistia* Maran. When Ramme (1951) described these genera, he did not establish the type species for them. In accordance with the International Code of Zoological Nomenclature, and because it was Maran (1953) who selected the type species, Maran must be the recognized authority of the genera (Koçak 1981). Recently Çiplak (2000) revised the genus *Parapholidoptera*, providing a key to species, on the basis of which the species most related to the Jordanian *P. willemsei* Katbeh Bader & Massa is *P. castaneoviridis* (Brunner) from Turkey.

We wish to thank Prof. Meir Paul Pener who pointed out some of the above listed mistakes.

Literature cited

- Ayal Y., Broza M., Pener M.P. 1999. Geographical distribution and habitat segregation of bushcrickets (Orthoptera: Tettigoniidae) in Israel. *Israel Journal of Zoology* 45: 49-64.
- Çiplak B. 2000. Systematics and phylogeny of *Parapholidoptera* (Orthoptera: Tettigoniidae: Tettigoniinae). *Systematic Entomology* 25: 411-436.
- Koçak A. Ö. 1981. On the nomenclature of some genera of Orthoptera. *Priamus* 1: 126-128.
- Maran J. 1953. Contribution to the knowledge of the genus *Pholidoptera* Wesm. *Acta Entomologica Musei Nationalis Pragae* 27: 209-221.
- Naskrecki P., Otte D. 1999. An Illustrated Catalog of Orthoptera. Vol. I. Tettigoniioidea. (CDROM). The Orthopterists' Society at the Academy of Natural Sciences of Philadelphia.
- Otte D. 1997. Orthoptera Species File. 7. Tettigoniioidea. The Orthopterists' Society at the Academy of Natural Sciences of Philadelphia.
- Popov GB. 1981. Insects of Saudi Arabia. Orthoptera: Fam. Tettigoniidae (Bush Crickets). Pp. 114-148. In: Wittmer W., Buttiker W. (Eds) *Fauna of Saudi Arabia*, vol. 3. Ciba-Geigy Ltd., Basle.
- Ragge D.R. 1964. A revision of the genus *Tylopsis* Fieber (Orthoptera: Tettigoniidae). *Bulletin of the British Museum (Natural History), Entomology* 15: 297-322.
- Ragge D.R. 1974. A new species of *Tylopsis* Fieber from Southern Israel (Orthoptera: Tettigoniidae). *Israel Journal of Zoology* 22: 63-66.
- Ramme W. 1939. Beiträge zur Kenntnis der palaearktischen Orthopterenfauna (Tettig. u. Acrid.). III. Mitteilungen aus dem Zoologischen Museum in Berlin 24: 41- 150.

-
- Ramme W. 1951. Zur Systematik Faunistik und Biologie der Orthopteren von südost-Europa und Vorderasien. *Mitteilungen aus dem Zoologischen Museum in Berlin* 27: 1-432.
- Uvarov B.P. 1933. Notes on new and little-known Orthoptera from Palestine. *Annals and Magazine of Natural History ser. 10, 11*: 663-672.
- Uvarov B.P. 1939. New and less-known Palestinian Orthoptera. *Annals and Magazine of Natural History, Ser. 11, 4*: 216-227.