

## **X1X2O (male) – X1X1X2X2 (female) chromosomal sex determining mechanism in the cricket *Cicloptylodes americanus* (Orthoptera, Grylloidea, Mogoplistidae)**

Authors: Mesa, A., García-Novo, P., and dos Santos, D.

Source: Journal of Orthoptera Research, 11(1) : 87-90

Published By: Orthopterists' Society

URL: [https://doi.org/10.1665/1082-6467\(2002\)011\[0087:XXOMXX\]2.0.CO;2](https://doi.org/10.1665/1082-6467(2002)011[0087:XXOMXX]2.0.CO;2)

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# $X_1X_2O$ (male) – $X_1X_1X_2X_2$ (female) chromosomal sex determining mechanism in the cricket *Ciclotyloides americanus* (Orthoptera, Grylloidea, Mogoplistidae)

A. MESA, P. GARCÍA-NOVO AND D. DOS SANTOS

Departamento de Biologia, Instituto de Biociências, UNESP, Campus de Rio Claro, Av. 24A, nº 1515, 13506-900 Rio Claro, SP, Brazil.  
E-mail: alejomesa@linkway.com.br.

## Abstract

The species of cricket *Ciclotyloides americanus* has males with  $2n = 14$  and females with  $2n = 16$  chromosomes. The autosomes are metacentrics and the sex determining mechanism is of the  $X_1X_2O$  (male) –  $X_1X_1X_2X_2$  (female) type, with  $X_1$  submetacentric and  $X_2$  acrocentric. This mechanism is cited for the first time among species belonging to the order Orthoptera.

## Key words

Scaly cricket, karyotype, sex chromosome, sex determination, meiosis, segregation

## Introduction

*Ciclotyloides americanus* was described by Saussure in 1874 from specimens collected in Cuba, but the species was later found in North America (T. J. Walker, pers. com.) and Brazil, always associated with human dwellings. Due to its small size (5 to 7 mm) and furtive habits it is rarely seen during the day when it remains in hiding. It only moves to feed at night, at which time the very low intensity male calling song is heard.

The basic chromosomal mechanism of sex determination in crickets, as well as in all other Orthoptera, is XO (male) – XX (female), though departures from this primitive scheme are frequent in Grylloidea. *C. americanus* exhibits two X's and the absence of Y, a mechanism which, though common in other groups of invertebrates, e.g., arachnids, has never been found among orthopteran insects.

## Materials and methods

Since 1980, specimens of *C. americanus* have been collected in Rio Claro (São Paulo State, Brazil) and submitted to karyological studies. The meiotic process was analyzed in several males and observations of embryo neuroblasts of both sexes were also carried out in search of mitotic divisions, after 2h pretreatment with 0.05% colchicine in eggs. In both meiotic and mitotic cells, Carnoy I fixative was used, the tissues squashed in lacto-acetic orcein 0.5% and after a few minutes, washed in acetic acid (45%).

## Results

*C. americanus* (Fig. 1a, b) has a diploid number of 14 (male) – 16 (female) (Fig. 2a, b), comprising in males six pairs of metacentric autosomes and two X chromosomes ( $X_1$ ,  $X_2$ ) and in females four

X chromosomes ( $X_1$ ,  $X_1$ ,  $X_2$ ,  $X_2$ ). The autosomes can be grouped in pairs: two large, two medium sized and two small. Within each group it is difficult to identify each pair based on size and morphology. The number of chiasmata from bivalents is two or three and apparently not localized. The  $X_1$  is submetacentric ( $r = 5, 7$ ) and  $X_2$  acrocentric ( $r = 3, 1$ ), near the limit with submetacentric. The  $X_2$  chromosome shows a median light-colored segment in the long arm.

During prophase I,  $X_1$  and  $X_2$  remain together in a single heterochromatic mass until diakinesis (Fig. 3a, b). At metaphase I,  $X_1$  and  $X_2$  remain either connected (Fig. 3f, g, h, i) or independent (Fig. 3c, d, e); but in all cases both chromosomes go to the same pole during anaphase I (Fig. 3j, k). Two kinds of other metaphases are observed, one of them with six autosomes (Fig. 3m), the other with the six autosomes plus the two X's (Fig. 3l).

## Discussion

The derived chromosomal mechanisms of sex determination in crickets are apparently more diverse than in grasshoppers, where only species with neo XY and  $X_1X_2Y$  males were detected. In addition to these mechanisms, species with  $X_1X_2X_3X_4Y$  (Mesa, unpub.) and  $X_1X_2O$  (this paper) males have been discovered in crickets.

Since the basic kind of sex determination in orthopteroid insects is that of XO (males) – XX (females), with the X metacentric, the presence of two Xs in *C. americanus* males could be explained by the dissociation of the original X into two sex chromosomes. But as  $X_2$  is submetacentric and  $X_1$  acrocentric with a relatively conspicuous small arm, further pericentric inversion or adding of new chromatin would need to have occurred after the original X splitting.

The presence of two X chromosomes that, in most of the nuclei, behave independently during first meiotic division, presents us with the unsolved question of how  $X_1$  and  $X_2$  always go to the same pole, in disagreement with the second Mendelian law.

Similar paradoxical chromosome behaviors were detected in males with  $X_1X_2Y$  chromosomes in the crickets *Eneoptera surinamensis* De Geer and *Euscirtus* (probably *hemelytrus* De Haan) (Raychaudhuri & Manna 1950, Smith 1953, Ohmachi & Ueshima 1957) as well as in several other  $X_1X_2Y$  and  $X_1X_2X_3X_4Y$  (male) crickets (Mesa et al. unpub.), species where no prophasic pairing and metaphasic contact took place between the sex chromosomes.

At the chromosomal level, the essence of Mendel's second law consists of the fact that centromeres of maternal and paternal origins of each autosomal bivalent, orientate freely toward opposite




Fig. 1. *Cicloptylodes americanus* a) male, b) female.




Fig. 2. Metaphase chromosomes from c-treated neuroblasts in embryos of *C. americanus*: a) male with  $X_1$  and  $X_2$  and six pairs of autosomes; b) female with  $X_1X_1X_2X_2$  and six pairs of autosomes. Scale = 10mm.

poles during first metaphase, being the neutral poles in this kind of orientation. As a consequence, each bivalent co-orientates its centromeres independently in opposite directions, while the chiasmate momentarily keeps the homologous chromosomes together in the metaphase plate. The same seems to happen in simple or multiple sex chromosomes where a prophase pairing takes place. In those multiple mechanisms with several X and Y chromosomes, these are arranged during metaphase I in a zigzag line, with Xs and Ys alternating positions as in the buprestid *Euchroma gigantea* L. (Mesa & Fontanetti 1984) or in the termite *Neotermes fulvescens* (Silvestri) (Martins & Mesa 1995). When prophase pairing fails in autosomal bivalents, each univalent behaves freely, going in approximately half of the nuclei to the same pole, while in the other half going to the opposite poles.

Slightly different are the XO (male) mechanisms where the X has no chiasmatic link, remaining free and performing several pole to pole movements during first metaphase, until its final transference to one of the poles.

In some species of crickets, in contrast to *C. americanus*, the sex chromosomes do not appear connected by chiasmata or by simple heterochromatin affinities during first metaphase in all or in some nuclei. In other species, neither first prophase pairing takes place. These situations are clearly abnormal, but still the distribution of the sex chromosomes is correct even in the absence of an appropriated metaphase arrangement. For this to happen a temporal structural change in the centrioles, or in the centromeres of the sex chromosomes, or in both of them, is plausible. This must occur in such a way that the direction toward the poles of each sex component is predetermined and in consequence the second Mendelian law is not operative.

## Acknowledgements

The authors thank Dr. T.J. Walker for the identification of the crickets studied here.

## Literature cited

- Claus G. 1956. La formule chromosomique du Gryllodea *Eneoptera surinamensis* De Geer et le comportement des chromosomes sexuels de cette espèce au cours de la Spermatogenèse. *Annales Sc. Nat. Paris Zool.*, 11 (Ser.) 18: 63-106.
- Martins G.A., Mesa A. 1995. Two permanent linear chains of sex chromosomes in *Neotermes fulvescens* and karyotypes of two other neotropical Kalotermitidae species (Insecta, Isoptera). *Genome* 38: 958-967.
- Mesa A., Bran E. J. 1964. Acerca de los cromosomas de *Eneoptera surinamensis* Ann. II. Congr. Latinoam. de Zool. 1: 9-16.
- Mesa A., Fontanetti C.S. 1984. Multiple sex chromosomes, autosomal polymorphism and a high number of S chromosomes in *Euchroma gigantea* L. 1735 (Coleoptera, Buprestidae). *Revista Brasileira de Genética* 7: 629-637.
- Ohmachi F., Ueshima N. 1957. Variations of chromosome complements in *Euscyrus hemelytrus* De Haan (Orthoptera: Grylloidea). *Bull. Fac. Mie Univ.* 15: 1-13.
- Piza S. de T. Jr 1946. Uma nova modalidade de sexo-determinação no grillo Sul-Americano *Eneoptera surinamensis*. *Ann. Esc. Sup. Agríc. "Luiz de Queiroz"* Piracicaba, Brasil, 3: 69-88.
- Raychaudhuri S.P., Manna G.K. 1950. Evidence of a multiple sex chromosome mechanism in a gryllid. *Journal of Heredity*. 42: 277-280.
- Saussure H. de. 1874. Mission scientifique au Mexique et dans L'Amérique Centrale. 6<sup>e</sup> Partie: études sur les Myriapodes et les Insectes. Imprimerie impériale. Paris 531 pp.
- Smith S.G. 1953. A pseudomultiple sex chromosome mechanism in an Indian gryllid. *Chromosoma (Berlin)* 5: 555-573.




Fig. 3. Male meiosis in *C. americanus* a) pachytene; b) diakinesis; c, d, e) first metaphase with  $X_1$  and  $X_2$  unconnected, going to the same pole; f, g, h, i) the same with  $X_1$ ,  $X_2$  connected, going to the same pole; j, k) first metaphases with  $X_1$  and  $X_2$  in contact going to the same pole; l) second metaphase, including unconnected  $X_1$  and  $X_2$ ; m) second metaphase without sex chromosomes. Scale = 10mm.