



## **Radiation Effects on Late-life Neurocognitive Function in Childhood Atomic Bomb Survivors: A Radiation Effects Research Foundation Adult Health Study**

Authors: Ishihara, Kayoko, Kato, Naohiro, Misumi, Munechika, Kitamura, Hiroko, Hida, Ayumi, et al.

Source: Radiation Research, 197(4) : 403-407

Published By: Radiation Research Society

URL: <https://doi.org/10.1667/RADE-21-00122.1>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## SHORT COMMUNICATION

# Radiation Effects on Late-life Neurocognitive Function in Childhood Atomic Bomb Survivors: A Radiation Effects Research Foundation Adult Health Study

Kayoko Ishihara,<sup>a,1</sup> Naohiro Kato,<sup>b</sup> Munechika Misumi,<sup>b</sup> Hiroko Kitamura,<sup>a,c</sup> Ayumi Hida<sup>a</sup> and Michiko Yamada<sup>a</sup>

Departments of <sup>a</sup> Clinical Studies and <sup>b</sup> Statistics, Radiation Effects Research Foundation, Hiroshima, Japan; <sup>c</sup> Occupational Health Training Center, University of Occupational and Environmental Health, Fukuoka, Japan

---

Ishihara K, Kato N, Misumi M, Kitamura H, Hida A, Yamada M. Radiation Effects on Late-life Neurocognitive Function in Childhood Atomic Bomb Survivors: A Radiation Effects Research Foundation Adult Health Study. *Radiat Res.* 197, 403–407 (2022).

High-dose radiation in childhood such as is used in radiation therapy causes cognitive decline, but there is insufficient research on the cognitive effects of low- to medium-dose radiation. We aimed to reveal the association between atomic bomb radiation exposure in childhood and late-life neurocognitive function. In 2011 and 2013, we mailed the Neurocognitive Questionnaire (NCQ) to subjects who were 12 years old or younger at the time of the atomic bombing. We converted the four NCQ subscales (metacognition, emotional regulation, motivation/organization, and processing speed) to T scores and defined the highest 10% of the controls (exposure dose < 5 mGy) as impaired. We used a generalized linear mixed model to evaluate the effect of radiation exposure on T scores and percentage impaired. We enrolled 859 participants. At the time of the bombing, the mean (SD) age was 6.7 (3.8) years for the control (N = 390) and 6.1 (3.8) years for the exposed (N = 469). At the time of replying to the questionnaire, the mean age of all the participants was 73.7 (3.8) years of age. After adjusting for cofactors, older age was related to the decline of all neurocognitive subscales. Sex and education level had relevance to some of the subscales. For neurocognitive function, exposure dose was not related except to percentage impaired, motivation/organization. Late-life neurocognitive function in atomic bomb survivors exposed as children was associated with age, but not clearly with radiation dose. More studies are needed to evaluate the effect of low-dose radiation during childhood on long-term neurocognitive function. © 2022 by Radiation Research Society

## INTRODUCTION

While some studies show that childhood exposure to high-dose radiation such as that administered in cranial radiation therapy (CRT) for leukemia and brain tumors leads to neuropsychological decline for up to about 35 years (1–3), other studies differ, including two Swedish studies on the effect of low-dose radiation used to treat children aged under 18 months of age for cutaneous hemangioma (4, 5). Hall et al. showed a negative dose-response relationship for the cognitive tests of learning ability and logical reasoning in the 1930–1960 Stockholm cohort (4) while in the Stockholm and Gothenburg study, the 1950–1960 cohort showed no clinically relevant effect on global cognition in 18-year-olds (5).

In studies dealing with the cognitive effects of nontherapeutic radiation exposures, Sibley et al. showed that the maximum annual and total lifetime radiation doses were related to death from dementia in female nuclear weapons workers (6) while Yamada et al. reported that cognitive function among total-body-irradiated atomic bomb survivors after adolescence was not significantly associated with radiation dose (7).

Here we aimed to reveal the association between late-life neurocognitive function and radiation exposure among childhood atomic bomb survivors in the Adult Health Study (AHS) of the Radiation Effects Research Foundation (RERF). We used the Childhood Cancer Survivor Study Neurocognitive Questionnaire (NCQ), a subjective self-reported measure of neurocognitive function developed to investigate the neurocognitive effects of therapy for the Childhood Cancer Survivor Study (8).

## MATERIALS AND METHODS

Since 1958, to investigate the health effects of exposure to ionizing radiation, the RERF Adult Health Study (AHS) has conducted biennial health check-ups of people who were in Hiroshima or Nagasaki at the time of the atomic bombings. The cohort was

<sup>1</sup> Address for correspondence: Department of Clinical Studies, Radiation Effects Research Foundation, 5-2 Hijiyama Park, Minami-ku, Hiroshima 732-0815, Japan; email: kishihara-kawase@umin.ac.jp.




FIG. 1. Flowchart showing study participants

expanded in 1977 (9), and since 1995, the AHS has also conducted semiannual mail surveys to obtain additional health information for members of the cohort who had expressed willingness to participate in such a survey. In 2011 and 2013, to examine late-life neurocognitive functions, we mailed the NCQ to everyone who might be eligible for the mail survey with written instructions (10). The RERF Institutional Review Board approved the study. Among members of the mail survey, 1,253 subjects were exposed at 12 years of age or younger (Fig. 1). The NCQ is a self-administered screening tool including 25 subjective neurocognitive complaints over the past six months (8) with options of the following responses: 1 = “Never a Problem”, 2 = “Sometimes a Problem”, and 3 = “Often a Problem”. Using exploratory factor analysis among the cohort control participants in a previous study (10), we identified 4 factors representing 20 of the 25 NCQ items. We labelled them metacognition (Factor 1, 9 items), emotional regulation (Factor 2, 5 items), motivation/organization (Factor 3, 4 items), and processing speed (Factor 4, 2 items) (10). Supplementary Table S1 (<https://doi.org/10.1667/RADE-21-00122.1.S1>) shows items included in each factor. We calculated raw scores for each factor as the sum of items.

Figure 1 is a flowchart of the study participants; 508 were control subjects (brain dose estimated by DS02R1 (11) < 5 mGy), 641 were exposed ( $\geq 5$  mGy), and 104 were unknown-dose subjects. The estimated median brain dose of the exposed was 515 mGy (interquartile range, 177–1,043). We used all subjects who provided at least one factor score for either test. Among mail subjects, that included 80.0% of control subjects and 76.4% of exposed subjects.

In the analyses, we used both NCQ scores in two surveys for those who answered both questionnaires and applied random intercept models of linear and generalized linear mixed effects models for continuous and binary responses to express the within-participant correlation induced by the measurements obtained from the same individuals, respectively (see Supplementary Data Appendix, <https://doi.org/10.1667/RADE-21-00122.1.S1>). In the analyses, we converted factor raw scores to T scores, whose mean (SD) of the control was 50 (10). Higher scores indicated more problems in neurocognitive function. We assumed that the participant was “impaired” for a corresponding group when a T score was in the upper 10th percentile of the control (i.e. T score  $\geq 63$ ). This cutoff definition was used in the

standardization manual for the Brief Symptom Inventory-18 (12) and in the studies of the Childhood Cancer Survivor Study (3, 13).

We performed a separate analysis to estimate the effects of radiation exposures (which were treated as continuous values) to the T scores of each factor after adjusting demographic covariates. The demographic covariates included age at the test (the age was centered and divided by 10), sex, city, and education (primary school, high school, and university). We also estimated the odds ratio of prevalence of “impaired” in the above definition (percentage impaired) using the logistic models.

We estimated the parameters of random intercept models using Bayesian inference. We used OpenBUGS version 3.2.3 (<http://www.openbugs.net>); MCMC software for Bayesian inference and R version 3.5.3 for other statistical analyses.

## RESULTS

The study included 859 participants (390 control and 469 exposed). Five-hundred-and-eighty participants (270 control and 310 exposed) provided at least one factor score for both NCQ. Their mean (SD) ages at the time of the bombing were 6.4 (3.8) years (6.7 (3.8) for the control, 6.1 (3.8) for the exposed,  $p = 0.026$ ) and 73.7 (3.8) years (74.0 (3.8) for the control, 73.5 (3.8) for the exposed,  $p = 0.035$ ) at the time of replying to the questionnaire.

Table 1 shows the sex, city of exposure, and education level of the participants. The control and the exposed did not differ significantly in sex or education level, but the exposed included significantly more Hiroshima survivors.

Table 2 shows the association of those features with radiation exposure, T score, and percentage impaired for each NCQ factor. We showed estimates for T score and odds ratios for percentage impaired with 95% Bayesian confidence intervals (CIs) in the models for 4 NCQ factors. Both T scores and percentage impaired for all factors tended

**TABLE 1**  
**Demographic Features of Control and Exposed Subjects**

Feature, n (%)	Total N = 859	Control N = 390	Exposed N = 469	P
Male	342 (39.8%)	160 (41.0%)	182 (38.8%)	0.554
Hiroshima	525 (61.1%)	212 (54.4%)	313 (66.7%)	<0.001
Education level				0.836
Primary school	229 (26.7%)	102 (26.2%)	127 (27.1%)	
High school	485 (56.5%)	219 (56.2%)	266 (56.7%)	
University	145 (16.9%)	69 (17.7%)	76 (16.2%)	

to increase with age. We could not find a city difference in any endpoints. Females showed more deterioration in Factors 3 and 4 of T score and Factor 4 of percentage impaired. Higher education tended to be associated with lower T scores in Factors 1 and 4 and lower percentage impaired in Factors 1 and 2. We could not find a radiation effect associated with any T score of the four factors or with the percentage impaired except for Factor 3. For Factor 3, percentage impaired increased significantly with radiation dose [Odds ratio per 1 Gy: 2.77, 95% CI (1.35, 6.48)] without significant increase of mean T score.

## DISCUSSION

We have shown here that subjective cognition in late life of participants who were exposed to radiation in childhood was associated with age and partially with sex and education. This result is similar to our previous report for non-exposed subjects, including those aged 66 to 89 years (7).

Few reports can be found on the association of demographic features with subjective cognition. Age is the most significant risk factor for dementia as shown by the Adult Changes in Thought (ACT) cohort study (14), the European Studies of Dementia (EURODEM) study (15), and the Framingham study (16). Lower education levels were associated with Alzheimer's disease, but not with non-Alzheimer's disease dementia in the ACT and EURODEM studies (14, 15), while the ACT study showed no association of sex with Alzheimer's disease (14). Significant deterioration of processing speed (Factor 4) in females in this study is compatible with the results of objective cognition on reaction time (a measure of processing speed) reported that males have faster and less variable reaction times than females (17).

In this study of adults exposed to atomic bomb radiation in childhood, we found no radiation effect in 4 factors of subjective cognition other than for the percentage impaired for motivation/organization. Although dose responses to CRT have been reported in children treated for central nervous system tumors (18), adults who as children received 24 Gy CRT, but not 18 Gy CRT, showed impairment in immediate or delayed memory (19). Children with cutaneous hemangioma after radiation therapy of brain doses up to 250 mGy had no clinically relevant effect on cognition (logical, spatial, technical, verbal or global test scores) at 18 years of age, but, only in the verbal test, with the highest dose group (>250 mGy, median 680 mGy) showing a decreasing score (5). In our study, the median estimated brain dose was 515 mGy, which appears to be below the dose causing cognitive decline.

**TABLE 2**  
**Association of Demographic Factors and Radiation Exposure with Estimates for T Score and Odds Ratios for Percentage Impaired for 4 NCQ Factors**

	Male (vs. Female)	Age (/10 years)	Nagasaki (vs. Hiroshima)	Education (vs. primary school)		Exposure dose (/Gy)
				High school	University	
T score (95% CI)						
Factor 1	-0.51	<b>3.31</b>	0.90	<b>-1.66</b>	<b>-2.42</b>	0.20
Metacognition	(-1.87, 0.89)	<b>(1.80, 4.86)</b>	(-0.52, 2.26)	<b>(-3.17, -0.08)</b>	<b>(-4.46, -0.21)</b>	(-0.78, 1.21)
Factor 2	-1.06	<b>2.09</b>	0.11	-0.96	-1.59	-0.26
Emotional regulation	(-2.40, 0.29)	<b>(0.56, 3.69)</b>	(-1.26, 1.45)	(-2.60, 0.62)	(-3.59, 0.47)	(-1.24, 0.71)
Factor 3	<b>-1.64</b>	<b>3.63</b>	-0.47	-0.66	-0.88	0.85
Motivation/Organization	<b>(-3.01, -0.28)</b>	<b>(2.05, 5.20)</b>	(-1.88, 0.87)	(-2.20, 0.86)	(-2.94, 1.17)	(-0.16, 1.81)
Factor 4	<b>-1.61</b>	<b>4.33</b>	-0.74	<b>-2.20</b>	<b>-2.13</b>	0.47
Processing speed	<b>(-2.88, -0.31)</b>	<b>(2.83, 5.83)</b>	(-2.03, 0.58)	<b>(-3.71, -0.71)</b>	<b>(-4.14, -0.18)</b>	(-0.51, 1.45)
Odds ratios for percentage impaired (95% CI)						
Factor 1	0.63	<b>12.17</b>	1.58	0.49	<b>0.16</b>	0.97
Metacognition	(0.19, 1.82)	<b>(3.34, 58.33)</b>	(0.54, 4.81)	(0.14, 1.58)	<b>(0.02, 0.82)</b>	(0.42, 2.23)
Factor 2	0.41	<b>4.02</b>	1.28	0.45	<b>0.22</b>	0.89
Emotional regulation	(0.14, 1.06)	<b>(1.32, 13.56)</b>	(0.49, 3.39)	(0.15, 1.25)	<b>(0.04, 0.96)</b>	(0.41, 1.78)
Factor 3	0.40	<b>17.29</b>	0.87	0.65	0.43	<b>2.77</b>
Motivation/Organization	(0.12, 1.14)	<b>(4.64, 90.83)</b>	(0.28, 2.55)	(0.18, 2.20)	(0.07, 2.23)	<b>(1.35, 6.48)</b>
Factor 4	<b>0.37</b>	<b>16.07</b>	0.75	<b>0.29</b>	0.32	1.14
Processing speed	<b>(0.14, 0.82)</b>	<b>(5.83, 53.52)</b>	(0.32, 1.72)	<b>(0.10, 0.71)</b>	(0.08, 1.10)	(0.60, 2.15)

Notes. Sample sizes were 820, 816, 838, and 844 for Factor 1, Factor 2, Factor 3, and Factor 4, respectively. Scores were adjusted by age at the test, sex, city, and education. Items with significant difference were expressed in bold font. NCQ: Neurocognitive Questionnaire

Some studies evaluated radiation effects using a subjective cognitive scale. A study after the Chernobyl accident showed the memory problems assessed by a self-reported questionnaire and university attendance of the infant evacuees did not differ from those of the control (20). In adult survivors who had received radiation for CNS tumors or leukemia at adolescent and early young adult, the NCQ score was high for those who had received CRT but not noncranial radiation therapy (21).

In non-CNS cancer survivors, radiation exposure was associated with impairment in task efficiency, memory, and emotional regulation among those diagnosed at less than 6 years of age (13). Young acute lymphoblastic lymphoma or low-grade glioma patients who received CRT showed abnormalities on brain parts related to cognition, such as global cortical atrophy, many white matter hyperintensities, low temporal lobe or hippocampal volume, and thin parietal and frontal cortices (19, 22), even among subjects exposed to a low-fractional dose ( $\leq 2$  Gy). Our study that included people exposed to atomic bombings at less than 13 years of age did not reveal a relationship between radiation and cognition (7), nor did our previous study on objective cognition for people exposed at or over 13 years of age (7).

The factors we selected from our baseline study (7) included 20 out of 25 NCQ items. The limitations of this study are that non-respondents (225, Fig. 1) might have had poorer cognitive function, and that 97% (970/998, Fig. 1) of the respondents addressed at least one factor, but not necessarily all the selected NCQ items. Completeness of each factor was related to sex, age, and education. However, we could not find the relationship between the completeness and radiation dose. Another limitation is that this study was evaluated by subjective complaints. Although such complaints are related to objective cognition (23, 24) or to a hippocampal volume decrease in the elderly (23), additional study on objective cognition is called for. Nevertheless, our survey was taken 67 years postirradiation, so our results may contribute to the long-term effects on cognition of low- to medium-radiation exposure in the young.

#### SUPPLEMENTARY DATA

Appendix: Mathematical information.

Supplementary Table S1: The List of NCQ Items for the Four Factors.

#### ACKNOWLEDGMENTS

The Radiation Effects Research Foundation (RERF), Hiroshima and Nagasaki, Japan, is a public interest foundation funded by the Japanese Ministry of Health, Labour and Welfare (MHLW) and the U.S. Department of Energy (DOE). This publication was supported by RERF Research Protocol 3-11. The views of the authors do not necessarily reflect those of the two governments. We acknowledge Dr. Hirofumi Maruyama (Hiroshima University), Dr. Reid D. Landes (University of Arkansas for Medical Sciences), and Dr. Kevin R. Krull (St. Jude Children's Research

Hospital) for their cooperation. Dr. Miriam Bloom (SciWrite Biomedical Writing & Editing Services) provided English-language support.

Received: June 21, 2021; accepted: November 29, 2021; published online: January 18, 2022

#### REFERENCES

1. Harila MJ, Winqvist S, Lanning M, Bloigu R, Harila-Saari AH. Progressive neurocognitive impairment in young adult survivors of childhood acute lymphoblastic leukemia. *Pediatr Blood Cancer* 2009; 53:156-61.
2. Jereb B, Korenjak R, Krzisnik C, Petric-Grabnar G, Zdravec-Zaletel L, Anzic J, et al. Late sequelae in children treated for brain tumors and leukemia. *Acta Oncol* 1994; 33:159-64.
3. Ellenberg L, Liu Q, Gioia G, Yasui Y, Packer RJ, Mertens A, et al. Neurocognitive status in long-term survivors of childhood CNS malignancies: a report from the Childhood Cancer Survivor Study. *Neuropsychology* 2009; 23:705-17.
4. Hall P, Adami HO, Trichopoulos D, Pedersen NL, Lagiou P, Ekblom A, et al. Effect of low doses of ionising radiation in infancy on cognitive function in adulthood: Swedish population based cohort study. *BMJ* 2004; 328:19.
5. Blomstrand M, Holmberg E, Aberg MA, Lundell M, Björk-Eriksson T, Karlsson P, et al. No clinically relevant effect on cognitive outcomes after low-dose radiation to the infant brain: a population-based cohort study in Sweden. *Acta Oncol* 2014; 53:1143-50.
6. Sibley RF, Moscato BS, Wilkinson GS, Natarajan N. Nested case-control study of external ionizing radiation dose and mortality from dementia within a pooled cohort of female nuclear weapons workers. *Am J Ind Med* 2003; 44:351-8.
7. Yamada M, Landes RD, Mimori Y, Nagano Y, Sasaki H. Radiation effects on cognitive function among atomic bomb survivors exposed at or after adolescence. *Am J Med* 2016; 129:586-91.
8. Krull KR, Gioia G, Ness KK, Ellenberg L, Recklitis C, Leisenring W, et al. Reliability and validity of the Childhood Cancer Survivor Study Neurocognitive Questionnaire. *Cancer* 2008; 113:2188-97.
9. Ozasa K, Cullings HM, Ohishi W, Hida A, Grant EJ. Epidemiological studies of atomic bomb radiation at the Radiation Effects Research Foundation. *Int J Radiat Biol* 2019; 95:879-91.
10. Yamada M, Landes RD, Hida A, Ishihara K, Krull KR. Effects of Demographic Variables on Subjective Neurocognitive Complaints Using the Neurocognitive Questionnaire (NCQ) in an Aged Japanese Population. *Int J Environ Res Public Health* 2019; 16:421.
11. Cullings HM, Grant EJ, Egbert SD, Watanabe T, Oda T, Nakamura F, et al. DS02R1: Improvements to Atomic Bomb Survivors' Input Data and Implementation of Dosimetry System 2002 (DS02) and Resulting Changes in Estimated Doses. *Health Phys* 2017; 112:56-97.
12. Derogatis LR. BSI 18, Brief Symptom Inventory 18: Administration, scoring and procedures manual. Minneapolis: NCS Pearson, Incorporated; 2001.
13. Kadan-Lottick NS, Zeltzer LK, Liu Q, Yasui Y, Ellenberg L, Gioia G, et al. Neurocognitive functioning in adult survivors of childhood non-central nervous system cancers. *J Natl Cancer Inst* 2010; 102:881-93.
14. Kukull WA, Higdon R, Bowen JD, McCormick WC, Teri L, Schellenberg GD, et al. Dementia and Alzheimer disease incidence: a prospective cohort study. *Arch Neurol* 2002; 59:1737-46.
15. Letenneur L, Launer LJ, Andersen K, Dewey ME, Ott A, Copeland JR, et al. Education and the risk for Alzheimer's disease: sex makes a difference. EURODEM pooled analyses. EURODEM Incidence Research Group. *Am J Epidemiol* 2000; 151:1064-71.

16. Bachman DL, Wolf PA, Linn RT, Knoefel JE, Cobb JL, Belanger AJ, et al. Incidence of dementia and probable Alzheimer's disease in a general population: the Framingham Study. *Neurology* 1993; 43:515-9.
17. Dykiert D, Der G, Starr JM, Deary IJ. Sex differences in reaction time mean and intraindividual variability across the life span. *Dev Psychol.* 2012; 48:1262-1276.
18. Armstrong GT, Jain N, Liu W, Merchant TE, Stovall M, Srivastava DK, et al. Region-specific radiotherapy and neuropsychological outcomes in adult survivors of childhood CNS malignancies. *Neuro Oncol* 2010; 12:1173-86.
19. Armstrong GT, Reddick WE, Petersen RC, Santucci A, Zhang N, Srivastava D, et al. Evaluation of memory impairment in aging adult survivors of childhood acute lymphoblastic leukemia treated with cranial radiotherapy. *J Natl Cancer Inst* 2013; 105:899-907.
20. Taormina DP, Rozenblatt S, Guey LT, Gluzman SF, Carlson GA, Havenaar JM, et al. The Chernobyl accident and cognitive functioning: a follow-up study of infant evacuees at age 19 years. *Psychol Med* 2008; 38:489-97.
21. Prasad PK, Hardy KK, Zhang N, Edelstein K, Srivastava D, Zeltzer L, et al. Psychosocial and Neurocognitive Outcomes in Adult Survivors of Adolescent and Early Young Adult Cancer: A Report From the Childhood Cancer Survivor Study. *J Clin Oncol* 2015; 33:2545-52.
22. Douw L, Klein M, Fagel SS, van den Heuvel J, Taphoorn MJ, Aaronson NK, et al. Cognitive and radiological effects of radiotherapy in patients with low-grade glioma: long-term follow-up. *Lancet Neurol* 2009; 8:810-8.
23. Stewart R, Godin O, Crivello F, Maillard P, Mazoyer B, Tzourio C, et al. Longitudinal neuroimaging correlates of subjective memory impairment: 4-year prospective community study. *Br J Psychiatry* 2011; 198:199-205.
24. Amariglio RE, Townsend MK, Grodstein F, Sperling RA, Rentz DM. Specific subjective memory complaints in older persons may indicate poor cognitive function. *J Am Geriatr Soc* 2011; 59:1612-7.