

Cloning of the Heat Shock Protein 60 Gene from the Stem Borer, *Chilo suppressalis*, and Analysis of Expression Characteristics Under Heat Stress

Authors: Cui, Ya-Dong, Du, Yu-Zhou, Lu, Ming-Xing, and Qiang, Cheng-Kui

Source: Journal of Insect Science, 10(100) : 1-13

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.010.10001>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Cloning of the heat shock protein 60 gene from the stem borer, *Chilo suppressalis*, and analysis of expression characteristics under heat stress

Ya-Dong Cui^{1,2}, Yu-Zhou Du^{1a}, Ming-Xing Lu¹, and Cheng-Kui Qiang³

¹Institute of Applied Entomology, Yangzhou University, Yangzhou, 225009, China

²Department of Life Science, Fuyang Teachers College, Fuyang, 236032, China

³Department of Agriculture and Landscape Engineering, Xuzhou Higher Vocational School of Bioengineering, Xuzhou, 221006, China

Abstract

Heat shock protein 60 is an important chaperonin. In this paper, *hsp60* of the stem borer, *Chilo suppressalis* (Walker) (Lepidoptera: Pyralidae), was cloned by RT-PCR and rapid amplification of cDNA end (RACE) reactions. The full length cDNA of *hsp60* consisted of 2142 bp, with an ORF of 1719 bp, encoding 572 amino acid residues, with a 5'UTR of 158 bp and a 3'UTR of 265 bp. Cluster analysis confirmed that the deduced amino acid sequence shared high identity with the reported sequences from other insects (77%-86%). To investigate whether *hsp60* in *C. suppressalis* responds to thermal stress, the expression levels of *hsp60* mRNA in larval haemocytes across temperature gradients from 31 to 39°C were analysed by real-time quantitative PCR. There was no significant difference for *hsp60* expression from 28 to 31°C. The temperatures for maximal induction of *hsp60* expression in haemocytes was close to 36°C. Hsp60 expression was observed by using flow cytometry. These results revealed that thermal stress significantly induced *hsp60* expression and Hsp60 synthesis in larval haemocytes, and the expression profiles of Hsp60 at the mRNA and protein levels were in high agreement with each other from 33 to 39°C.

Keywords: Lepidoptera; Hsp60; RACE; haemocytes; temperature

Abbreviations: Hsp, heat shock protein

Correspondence: yzdu@yzu.edu.cn

Associate Editor: Megha Parajulee was editor of this paper.

Received: 19 January 2010, **Accepted:** 13 March 2010

Copyright : This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 10, Number 100

Cite this paper as:

Cui YD, Du YZ, Lu MX, Qiang CK. 2010. Cloning of the heat shock protein 60 gene from the stem borer, *Chilo suppressalis*, and analysis of expression characteristics under heat stress. *Journal of Insect Science* 10:100 available online: insectscience.org/10.100

Introduction

Heat shock proteins (Hsp) are a family of proteins that help organisms to modulate stress response and protect organisms from environmentally induced cellular damage. They usually act as molecular chaperones, promoting correct refolding and preventing aggregation of denatured proteins (Johnston et al. 1998; Feder and Hofmann 1999). On the basis of molecular weight and homology of amino acid sequences, Hsp can be divided into several families including Hsp90, 70, 60, 40 and small Hsp (Feder and Hofmann 1999; Sørensen et al. 2003).

Hsp60 is mostly located in mitochondria of eukaryotic cells (Gatenby et al. 1991). Under normal conditions, Hsp60 operates the bending and assembling of enzymes and other protein complexes related to energy metabolism; under adverse environmental conditions the synthesis of Hsp60 increases and the protein then renatures damaged proteins to restore their biological activity (Buchner et al. 1991; Cheng et al. 1989; Ostermann et al. 1989; Martin et al. 1991). As a molecular chaperone, Hsp60 helps protect against protein aggregation (Sanders et al. 1992), and in the transport of proteins from cytoplasm to organelles (Fink 1999).

Molecular analysis of thermal stress has been extensively studied in *Drosophila melanogaster* and shows various responses of each Hsp at the transcriptional and translational level (Hoffman et al. 2003; Sørensen et al. 2007). Recently, research has been extended to several other insect species that are important in agricultural, medical and industrial fields (Yocum 2001; Chen et al. 2005, 2006; Sonoda et al. 2006; Huang and Kang 2007; Wang et al. 2007; Kim et

al.2008). The stem borer, *Chilo suppressalis* (Walker) (Lepidoptera: Pyralidae) is one of the most serious pests of rice. This pest has been widely distributed in all rice fields of China, and is constantly adapting to its environment. In the present study, the full-length cDNA of *C. suppressalis hsp60* was cloned. In addition, the expression profiles of Hsp60 in larvae haemocytes from *C. suppressalis* at mRNA and protein levels were analysed by using real-time quantitative PCR and flow cytometry across temperature gradients from 31 to 39°C.

Materials and Methods

Insect rearing and Isolation of haemocytes

The larvae of *C. suppressalis* were initially collected from the paddy fields in suburbs of Yangzhou City, China. The rice stem borers were reared using a method described by Shang et al. (1979) at $28 \pm 1^\circ\text{C}$, 16:8 L:D and RH >80%. Hatched larvae were fed on rice seedlings until larvae reached the 5th stadium.

Larvae from each experimental group were washed two times with distilled water and were anaesthetised by chilling on ice. The proleg was then cut off and haemolymph was collected directly into 1ml Eppendorf tubes containing 200 µl cold PBS. The haemocytes were separated from the haemolymph by centrifugation for 5 min at 800 g at 4°C. Sedimented haemocytes were washed twice with PBS and used for RNA extraction.

RNA extraction and cDNA synthesis

Total RNA from the haemocytes was isolated using TRIzol reagent (Sangon, www.sangon.com). Single-stranded cDNA was synthesized from 1 µg of RNA with the MBI RevertAid First Strand cDNA Synthesis Kit (MBI Fermentas, www.fermentas.com)

according to the manufacturer’s instructions. Single-stranded cDNA for 3’-Rapid amplification of cDNA ends (3’-RACE) and 5’-RACE experiments was synthesized from 1 µg of RNA using the TaKaRa RACE cDNA Amplification Kit (TaKaRa, www.takara-bio.co.jp) according to the manufacturer’s instructions.

Degenerate PCR for isolation of *hsp60* fragments

The partial clones of *hsp60* from *C. suppressalis* were amplified by PCR using primer sets: P1, P2 (Table 1). Pairs of primers were designed using consensus amino acid of insect Hsp60. PCR was carried out using 2 µl cDNA, 15 pmol of each primer, 10 nmol of each deoxynucleoside triphosphate (dNTP), and 1 unit of *Taq* DNA polymerase (TaKaRa) in the supplied buffer giving a final concentration of 2.0 mM MgCl₂ in 25 µl. Cycle conditions were as follows: initial denaturation at 94°C for 4 min; 3°Cycles of 94°C for 40 s, 55°C for 40 s, and 72°C for 1 min; final extension at 72°C for 10 min. Amplification products were purified from 1% agarose gels using a gel extraction kit (BioTeke, www.biotek.com). The purified fragment was cloned into the pMD-18T vector

(TaKaRa) and sequenced.

3’-RACE PC

Two semi-nested 3’-RACE reactions were conducted on 2 µl of the *C. suppressalis* 3’-RACE-ready cDNA. In the first reaction, a sense gene-specific primer designed from P1 and P2 PCR product sequence (*hsp60* 3’-RACE outer primer; Table 1) and an antisense 3’-RACE outer primer (Table 1; TaKaRa) were used. The 3’-RACE inner PCR was performed using the *hsp60* 3’-RACE inner primer (Table 1) and the antisense 3’-RACE inner primer (Table 1; TaKaRa). The 50-µl amplification mix was prepared according to the TaKaRa cDNA protocol using the *La Taq* polymerase mix (TaKaRa). The outer and inner PCR were performed using the following conditions: 3 min at 94°C, followed by 25 cycles of 40 s at 94°C, 40 s at 55°C, and 2 min at 72°C and finishing with chain extension at 72°C for 10 min.

5’-RACE PCR

Two semi-nested 5’-RACE reactions were conducted on 2 µl of *C. suppressalis* 5’-RACE-ready cDNA. In the first reaction, a sense gene-specific primer designed from the sequences obtained following 3’-RACE outer

Table 1. Primer sequences used in the cDNA cloning and real-time PCR

Primer type	Primer sequence(5’-3’)
cDNA isolation (RT-PCR)	
P1	GCKGGDGAYGGNACNACNWC
P2	TCDC CRAADCCRGNGCTTTKA
5’- and 3’-cDNA end isolation(RACE)	
<i>Hsp60</i> 5’RACE Outer primer	GAGATGGTTGCTACTTGTGCG
<i>Hsp60</i> 5’RACE Inner primer	CTTTGGCAACTGTAACTCCGTC
5’RACE Outer primer	CATGGCTACATGCTGACAGCCTA
5’RACE Inner primer	CGCGGATCCACAGCCTACTGATGATCAGTCGATG
<i>Hsp60</i> 3’RACE Outer primer	GTAGCAACCATCTCTGCCAAC
<i>Hsp60</i> 3’RACE Inner primer	CTCCAAGTGGCTGCTGTAAAAG
3’RACE Outer primer	TACCGTCGTTCCACTAGTGATTT
3’RACE Inner primer	CGCGGATCCTCCACTAGTGATTTCACTATAGG
Amplification of <i>Hsp60</i> Complete cDNA	
P3	ACTTCTTGGTTGTTGGTATTTTCG
P4	CAGACTCATCATTGAACAGAAAACACTAC
Primers used in quantitative PCR	
<i>Hsp60</i> sense	ATCCGAAGGGGTGTCATGATGG
<i>Hsp60</i> antisense	GCAGAGATGGTTGCTACTTGTG
18sRNA sense	TCGAGCCGCACGAGATTGAGCA
18sRNA antisense	CAAAGGGCAGGGACGTAATCAAC

and inner PCR amplifications of *hsp60* (*hsp60* 5'-RACE outer primer; Table 1) and an antisense 5'-RACE outer primer (Table 1; TaKaRa) were used. The 5'-RACE inner PCR was performed using the *hsp60* 5'-RACE inner primer (Table 1) and the antisense 5'-RACE inner primer (Table 1; TaKaRa). PCRs were performed as described for the 3'-RACE PCR.

Cloning and sequencing

After gel extraction, the 3'- and 5'-RACE fragments were cloned into a pMD-18T vector (TaKaRa). Recombinant plasmids were isolated using the Plasmid Mini kit (BioTeke), and sequenced. The full-length of the sequence assembled by RACE was verified by sequencing the fragment amplified from the primers P3, P4 (located at 5' UTR and 3'UTR) (Table 1) and subjected to homology analysis.

Amino acid sequence comparisons and phylogenetic analysis

Database searches were performed with the BLAST program (NCBI-BLAST network server). The open reading frame was identified using ORF Finder (<http://www.ncbi.nlm.nih.gov/gorf/gorf.html>) and the amino acid molecular weight was calculated by the SWISS-PROT (ExPASy server) program 'Compute pI/Mw' (<http://au.expasy.org/>). Sequence alignment and homology analysis was performed using Clustal X. A phylogenetic tree (neighbor-joining method) was then constructed with 1000 bootstrap replicates using MEGA version 3.1 based on the deduced amino acid sequence of *C. suppressalis* Hsp60 as well as the known sequences of fourteen other insect species.

Real-time quantitative PCR

Fifteen fifth-instar larvae in three replicates were randomly selected from each

experimental group and exposed to 31, 33, 36 or 39°C for 2 h with 28°C as control. Each treatment was repeated three times.

The haemocytes from control and treated groups were collected as above. The collected haemocytes were immediately used for RNA extraction, and cDNA was synthesized according to the methods described above. 18S rRNA gene in *C. suppressalis* was cloned (GQ265912) and used as the housekeeping gene. Based on the cDNA sequences of *C. suppressalis hsp60* and the 18S rRNA gene sequences, primer pairs (*hsp60* sense / *hsp60* antisense and 18S sense / 18S antisense) for real-time PCR were designed (Table 1). The PCR reactions were performed in a 20 µl total reaction volume including 10 µl of 2 × SYBR® Premix EX Taq™ master mix (TaKaRa), 5 µM each of primers (Table 1) and 1 µl cDNA templates. They were carried out on the ABI Prism 7000 Sequence Detection System (Applied Biosystems, www.appliedbiosystems.com). The thermal cycler parameters were as follows: 10 s at 95°C, then 4°Cycles of 5 s at 95°C, 20 s at 55°C and 20 s at 72°C.

For each amplification, the reaction was carried out in 3 replicates, from which mean threshold cycle (CT) values plus standard deviations were calculated. Relative transcripts of *C. suppressalis hsp60* cDNA amounts were calculated applying the $2^{-\Delta\Delta C^T}$ method (Livak and Schmittgen 2001).

Flow cytometric determination of Hsp60 levels

The larval treatment was carried out as described above. Each temperature treatment was repeated three times. The haemocytes were harvested as described above. The extent of Hsp60 levels in larvae haemocytes was

```

1      cgccgatccacagcctactgatgatcagtcgatggaaaattttataaacctataaccggca
61     gaagaagatcacgttttgcgatcccgcccgccgaatcttcgagaaacagaggtctgaaactt
121    ctgtgttgttggtattttcgattaatttttatcacaaa
159    atgttgcgtttgcctcgtgttgttcgccaaagctgtttcattgcacaaatcacaccaattt
1      M L R L P R V V R Q A V S L H K S H Q F
219    tcaagattttacgccaaagatgtaagggttggtgctgatgtaagagctctcatgcttcaa
21     S R F Y A K D V R F G A D V R A L M L Q
279    ggtgtcgtatgctcctagccgatgcagtagctgttacaatgggcccccaaggcagaaatgtg
41     G V D V L A D A V A V T M G P K G R N V
339    atattggaacaatcttgggttctccgaaaaataacaaaagacggagttacagttgccaaa
61     I L E Q S W G S P K I T K D G V T V A K
399    ggagtgagctcaagacaaattccagaacattggtgccaaagcttgtacaaaatgtggcg
81     G V E L K D K F Q N I G A K L V Q N V A
459    aacagcacaaacgaagaagccggagacggtaccacacagccacagtcctcgcaagagct
101    N S T N E E A G D G T T T A T V L A R A
519    atcgccaaagaaggtttcgaaaaaatctcaaaggggctaacccttgcagatccgaagg
121    I A K E G F E K I S K G A N P I E I R R
579    ggtgtcatgatggcagtagacgccgtgaaagacaaattaaaaagtatgtcgaaaccgtg
141    G V M M A V D A V K D K L K S M S K P V
639    accacaccggaagaaatcgacaaagtagcaaccatctctgccaacggcgacacagcaatc
161    T T P E E I A Q V A T I S A N G D T A I
699    gggaaactgatcgcggacgccatgaagagggtcgccgggacggcgttatcacggtgaag
181    G K L I A D A M K R V G R D G V I T V K
759    gacggcaagacgctcactgacgaactcgaagtattgaaggtatgaaattcgatagagga
201    D G K T L T D E L E V I E G M K F D R G
819    tatatttcaccatatttcatcaactcatccaaggagccaaagtgaattccaagatgct
221    Y I S P Y F I N S S K G A K V E F Q D A
879    ttagttttgtttctgagaaaaaataagcaatgtacagacaatcatccctgctttggag
241    L V L F S E K K I S N V Q T I I P A L E
939    ctagctaatacagcagaggaacaccttagtcatagttgctgaagatgtcgtggggaagca
261    L A N Q Q R K P L V I V A E D V D G E A
999    ctgtcaactttggtgtgaacagactaaaaattggcctccaagtggtgctgtgtaaagca
281    L S T L V V N R L K I G L Q V A A V K A
1059   cctggatttgagaaaaaccgtaagtctacattgagtgatattggccattgctactggtgga
301    P G F G E N R K S T L S D M A I A T G G
1119   gttgtgtttggagatgatcgaatctgatcaagctggaggatgtacaaccatcagatctc
321    V V F G D D A N L I K L E D V Q P S D L
1179   ggccaggtaggtgaggtgtgtattacaaaggatgacacactcttcttgaaggaaagggt
341    G Q V G E V V I T K D D T L F L K G K G
1239   aaaaagacagacattgacaggagggtgaacagattcgatgataaattcaggagactact
361    K K T D I D R R A E Q I R D Q I Q E T T
1299   tcagaatatgagaaggagaagcctcaagaacgtttggctagattggcatcaggtgtggct
381    S E Y E K E K L Q E R L A R L A S G V A
1359   gtattacatgtaggtggatccagtgaaagtgaagtcaacgagaaaaaggacagagtaaat
401    V L H V G G S S E V E V N E K K D R V N
1419   gatgccttgaaatgctacacgtgctgccgttgaagaaggtatagtgccctggagggtctc
421    D A L N A T R A A V E E G I V P G G G S
1479   gctctcctccgatgcattcctgttcttgagcaactctccacctcaaatgctgaccaggca
441    A L L R C I P V L E Q L S T S N A D Q A
1539   actggagttgaaattatcaagaaggccctgcgaatgccatgcatgactattgcacgaat
461    T G V E I I K K A L R M P C M T I A R N
1599   gctggcattgacggctctgtgttgttgcataaagttgaagatttaggacctgagttgga
481    A G I D G S V V V A K V E D L G P E F G
1659   tatgatgctttgaataatgaatatgtaaacatgattgaaaaggcattatgatccacc
501    Y D A L N N E Y V N M I E K G I I D P T
1719   aaagttgtgcgaacagcattacagatgccagtgaggttgcttattgttaacaacagcc
521    K V V R T A L T D A S G V A S L L T T A
1779   gaagcagttatttgtgaaattcctcaagagaaggagcctaattccatggcaggcatggga
541    E A V I C E I P Q E K E P N P M A G M G
1839   ggaatgggtggcatgggtggaatgggtggtatgataa
561    G M G G M G G M G G M M *
1878   gatacaataactaaagacaatcattgcatttaagtgtaaaaataatgggactattattaa
1938   ttagttacatcacctgtgctggtatagctgtaattggaaatgggtatgtagaatttg
1998   tggagtgatgtctttagtatttctgttcaatgatgagctgattgtgttacaaatgta
2058   gtagatttatttagtttaagtctgcagctgcaagcttctgttattcctaataataaaa
2118   taatattttatactaaaaaa

```

Figure 1. The complete cDNA sequence and predicted amino acid sequence of *Hsp60* of *Chilo suppressalis*. A classical mitochondrial *Hsp60* signature is shown in box, and a typical GGM repeat motif at the C terminus is underlined. The stop codon is marked with an asterisk. High quality figures are available online.

investigated following the method of Shen and Zhou (2002).

The collected haemocytes were fixed in 4% paraformaldehyde used for analysis of Hsp60 levels. Each sample was divided into two groups, one group for positive treatment, another as a negative baseline control. Positive treatment was performed as follows: the fixed haemocytes were centrifuged for 5 min at 800 g and washed twice with PBS. Subsequently, the haemocytes were permeabilized in PBS with Triton X-100 and incubated with the rabbit anti-Hsp60 polyclonal antibody (Boster, www.immunoleader.com/) (1:200) for 30 min at 18°C, and then centrifuged for 5 min at 800g and washed twice with PBS. Following treatment with the primary antibody, the haemocytes were incubated for 30 min at 18°C with FITC conjugated goat anti-rabbit secondary antibody (Boster) (1:300) and then centrifuged for 5 min at 800g and washed twice with PBS. Finally the haemocytes were resuspended in PBS for analysis. Negative baseline control procedure was similar to that of positive treatment except that the primary antibody incubation was eliminated.

Cellular fluorescence, reflecting cellular Hsp60 levels, was determined at 525 nm using flow cytometry (Becton Dickinson, www.bd.com). For each group, approximately 1000 cells were analysed. Subsequently, the average fluorescence values of the positive treatment were divided by those of the negative baseline control, and the folds were used as the relative levels of Hsp60 from each sample.

Statistical analysis

Data were expressed as mean values $\pm$ S.D. based on three separate experiments. Statistical analysis was carried out by

Student's t-test. The asterisks denote statistical significance when compared with control: * $p < 0.05$; ** $p < 0.01$.

Results

Sequence analysis

Degenerative primers based on a conserved region of *hsp60* were used to amplify the cDNAs derived from the haemocytes of *C. suppressalis*. A PCR product of 596 bp was obtained. The 596 bp amplified fragment was highly homologous to *hsp60*, and used to obtain 5'- and 3' flanking sequences using RACE 5'- and 3' RACE of *hsp60* produced fragments of 456 and 1142 bp. After assembly of the 2 sequences, a 2142 bp full-length cDNA sequence was obtained (Figure 1). This sequence contained a 158 bp 5'UTR, a 265 bp 3'UTR with a canonical polyadenylation signal sequence (AATAAA), and a poly (A) tail, as well as a single 1,719 bp open reading frame (ORF) encoding a polypeptide comprised of 572 amino acids with a molecular mass of 61014 Da and a pI of 5.69 (GenBank accession number: GQ265913).


The deduced amino acid sequence of Hsp60 of *C. suppressalis* was highly similar to that of other insects including: *Culicoides variipennis* (86%), *Pteromalus puparum* and *Nasonia vitripennis* (85%), *Anopheles gambiae*, *Apis mellifera* and *Culex quinquefasciatus* (83%), *Drosophila melanogaster*, *Liriomyza sativae*, *Lucilia cuprina* and *Aedes aegypti* (82%), *Liriomyza huidobrensis* (81%), *Tribolium castaneum* (79%), *Myzus persicae* (78%), *Acyrtosiphon pisum* (77%). Based on the amino acid sequences of Hsp60, a phylogenetic tree was constructed using the programs of CLUSTALX and MEGA2.1 (Figure 2).

Real-time quantitative analysis of *hsp60* expression

To determine whether *hsp60* responds to heat treatments, fifth-instar larvae of *C. suppressalis* were kept at a target temperature (31, 33, 36 or 39°C) for 2h and expression levels were compared with levels observed in non heat-treated individuals at 28°C. As shown in Figure 3 *hsp60* mRNA was expressed at extremely low levels in the untreated groups (28°C). Under thermal stress, the baseline levels of *hsp60* mRNA were found to vary. A significant induction of *hsp60* was shown at 33, 36 and 39°C, reaching 5.61, 9.13 and 8.67 fold versus the control, respectively, and there was no significant difference for *hsp60* expression from 28 to 31°C. Interestingly, the degree of increase in the level of *hsp60* reached a maximum at 36°C and then dropped at 39°C with increasing temperatures.

Hsp60 verification at protein levels

To study the heat induction of Hsp60 at the translational levels, the expression of Hsp60 was determined by using a flow cytometer. Figure 4 shows that thermal stress significantly elevated the level of Hsp60 synthesis in larvae haemocytes. Compared with the control (28°C), the relative levels of Hsp60 increased to 1.40, 1.47, 1.88 and 1.74 fold at 31, 33, 36, and 39°C, respectively. These results revealed that the expression profiles of Hsp60 at the mRNA and protein levels are in high agreement with each other from 33 to 39°C.






Figure 3. Expression analysis of *hsp60* mRNA of haemocytes from *Chilo suppressalis* under different heat-shock temperature by Real-time PCR. Values are means $\pm$ S.D. of three independent experiments in triplicate. * $p < 0.05$ vs. control, ** $p < 0.01$ vs. control. High quality figures are available online.




Figure 4 Expression levels of Hsp60 of haemocytes from *Chilo suppressalis* under different heat-shock temperature by using Flow cytometry. Values are means $\pm$ S.D. of three independent experiments in triplicate. * $p < 0.05$ vs. control, ** $p < 0.01$ vs. control. High quality figures are available online.

Discussion

Using a combination of RT-PCR and RACE techniques, the full-length *hsp60* cDNA was cloned from haemocytes of *C. suppressalis*. The C-terminal repeats (GGM)_n, which are a characteristic of mitochondrial Hsp60 (Tsugekils et al. 1992), are present in *C. suppressalis* Hsp60, indicating that the isolated gene is a mitochondrial *hsp60*. In addition, the lengths of the cDNA and the ORF, as well as the predicted protein size are similar to those of other Hsp60s. An ATP binding motif, which is highly conserved (Wong et al. 2004), was found in the deduced Hsp60 amino acid sequence. The great similarity in this region may indicate that among Hsp60s, the mechanism of coupling ATP hydrolysis to the substrate-refolding process is similar. Using the BLAST X programme of the NCBI website, the deduced amino acid sequence of Hsp60 of *C. suppressalis* showed high identity and similarity with known Hsp60s of other insect species (more than 77% similarity in all the matches). A phylogenetic tree was constructed based on the full amino acid sequences of Hsp60 of the fifteen insect species in this study.

To find the temperatures for maximal induction of *hsp60* expression, the relative mRNA levels of *hsp60* in larvae haemocytes of *C. suppressalis* were quantified by real-time quantitative PCR at temperatures from 31 to 39°C. It was found that *hsp60* mRNA in larval haemocytes was expressed at extremely low levels in the control groups (under normal conditions). The baseline levels of *hsp60* mRNA were found to vary under heat stress. The results revealed that *hsp60* gene in larvae haemocytes was significantly upregulated with increasing temperatures, reached a maximum at 2h of exposure to a 36°C heat

shock and then dropped at 39°C. However, after exposure at 39°C for 2h, *hsp60* mRNA levels exhibited a reduction in expression, indicating that transcription had decreased. Huang and Kang (2007) observed a similar response in the induction of heat shock in *L. sativae* *hsp60* using real-time quantitative PCR methods, and the expression of *L. sativae* *hsp60* was inhibited when temperatures were higher than 42.5°C for 1 h, which exceeds the tolerance limit of *L. sativae*. However, the majority of previous studies of inducible *hsp60s* expression was virtually undetectable by stress factors. For instance, the expression levels of mitochondrial *hsp60* are not influenced by heat or cold in *Trichinella spiralis* (Wong et al. 2004). Furthermore, the expression of *hsp60* does not respond to various types of stresses, such as H₂O₂ (Martinez et al. 2002), acidic and oxidative stress (Wong et al. 2004). In those studies, mRNA expression was monitored using Northern Blotting or semi-quantitative RT-PCR, which is less sensitive than real-time quantitative PCR.

Hsp60 is known to function as a molecular chaperone in many species and is absolutely essential for the proper functioning of cells under normal and stress conditions (Lindquist 1986; Hemmingsen et al. 1988; Goloubinoff et al. 1989; Hartl 1996). In this study, we detected up-regulation of Hsp60 in haemocytes of *C. suppressalis* in adaptation to thermal stress. Hsp60 in haemocytes was found to be increased at 31°C while *hsp60* was not increased. The results indicate that the thermal responses of Hsp60 in the haemocytes at the mRNA and protein levels are high in agreement with each other from 33 to 39°C. Our findings also agree with those of Wheelock et al. (1999) for *B. plicatilis* in which Hsp60 response increased up to 3–4 fold when heat exposure occurred. Rios-Arana

et al. (2005) also reported that Hsp60 was induced 2–4 fold in *P. patulus* exposed to heat. Other arthropod studies have compared expression levels at gene and protein levels. For example, protein levels of Hsp70 followed thermotolerance and reached the highest levels 49 h after heat hardening in adult *Orchesella cincta*, and the expression of *hsp70* messenger RNA reached a peak within the first 2 h and then sharply decreased after 6 h (Bahrndorff et al. 2009).

Mitochondria are essential eukaryotic organelles that serve as a site for many vital metabolic pathways and supply the cell with oxidative energy. Hsp60 plays a central role in the folding of newly imported and stress-denatured proteins (Martin et al. 1992; Martinus et al. 1995). As so, it was demonstrated that yeast containing mutated mt-Hsp60 do not grow at elevated temperatures (Cheng et al. 1989; Dubaquié et al. 1998) and show irreversible aggregation of a large number of newly imported proteins (Dubaquié et al. 1998). The higher level of Hsp60 expression induced by heat stress strengthens the idea that this protein has a significant role in the adaptation of various environmental conditions.

Some studies indicated that the induction of Hsp60 expression is tissue-specific. Lakhota and Singh (1996) reported Hsp60 heat-induced expression in *D. melanogaster* larval Malpighian tubules following heat shock. A tissue-specific variation in the heat-induced expression of Hsp60 was also reported in grasshopper (*Spathosternum prasiniferum*), cockroach (*Periplaneta americana*) and gram pest (*Heliothis armigera*) (Singh and Lakhota 2000). The level of Hsp60 in *L. cuprina* was significantly enhanced upon heat shock in some tissues (Sunita et al. 2006). Moreover, over-expression of Hsp may cause some

negative effects on growth, development, survival and fecundity (Krebs and Feder 1997; Krebs and Feder 1998; Huang et al. 2007), suggesting that the expression of Hsp may relate to physiological processes (Huang et al. 2007). Interestingly, the expression levels of Hsp60 in the haemocytes of *C. suppressalis* reached a maximum at 36°C and then declined at 39°C with increasing temperatures in the present study. Such a drop of Hsp60 may be due to approach of a threshold limit in the cell. A similar scenario was observed in an earlier study in which Hsp60 synthesis in *L. huidobrensis* were found to increase to a maximum at 42.5°C and then dropped when heat stress was enhanced (Huang and Kang 2007). Kristensen et al. (2002) also reported that inbred larvae of *Drosophila buzzatii* expressed more Hsp70 at high temperatures except at very high temperatures close to the physiological limit. This further supports the fact that Hsp plays a major protective role against cellular damage with high temperature exposure and yet may not be able to protect the cells beyond the threshold limit.

Acknowledgments

This work was supported by the National Basic Research and Development Program of China (2006CB102002).

References

-
- Bahrndorff S, Marien J, Loeschcke V, Ellers J. 2009. Dynamics of heat-induced thermal stress resistance and Hsp70 expression in the springtail, *Orchesella cincta*. *Functional Ecology* 23: 233-239.
- Buchner J, Schmidt M, Fuchs M, Jaenicke R, Rudolph R, Schmid FX, Kieffhaber T. 1991. GroE facilitates refolding of citrate synthase

by suppressing aggregation. *Biochemistry* 30: 1586–1591.

Chen B, Kayukawa T, Monteiro A, Ishikawa Y. 2005. The expression of the HSP90 gene in response to winter and summer diapauses and thermal-stress in the onion maggot, *Delia antiqua*. *Insect Molecular Biology* 14: 697–702.

Chen B, Kayukawa T, Monteiro A, Ishikawa Y. 2006. Cloning and characterization of the HSP70 gene, and its expression in response to diapauses and thermal stress in the onion maggot, *Delia antiqua*. *Journal of Biochemistry and Molecular Biology* 39: 749–758.

Cheng MY, Hartl FU, Martin J, Pollock RA, Kalousek F, Neupert W, Hallberg EM, Hallberg RL, Horwich AL. 1989. Mitochondrial heat shock protein hsp60 is essential for assembly of proteins imported into yeast mitochondria. *Nature* 337: 620–624.

Dubaquie Y, Looser R, Funfschilling U, Jenö P, Rospert S. 1998. Identification of in vivo substrates of the yeast mitochondrial chaperonins reveals overlapping but non-identical requirement for hsp60 and hsp10. *The EMBO journal*. 17: 5868–5876.

Feder ME, Hofmann GE. 1999. Heat-shock proteins, molecular chaperones, and the stress response: evolutionary and ecological physiology. *Annual Review of Physiology* 61: 243–282.

Fink AL. 1999. Chaperone-mediated protein folding. *Physiological reviews* 79: 425–449.

Gatenby AA, Donaldson GK, Goloubinoff P, LaRossa RA, Lorimer GH, Lubben TH, Van

Dyk TK, Viitanen P. 1991. The cellular functions of chaperonins. In: Schlesinger MJ, Santoro MG., Garaci E, editors. *Stress Proteins, Induction and Function*, pp. 57–70. Springer-Verlag.

Goloubinoff P, Gatenby AA, Lorimer GH. 1989. GroE heat-shock proteins promote assembly of foreign prokaryotic ribulose biphosphate carboxylase oligomers in *Escherichia coli*. *Nature* 337: 44–47.

Hartl FU. 1996. Molecular chaperones in cellular protein folding. *Nature* 381: 571–579.

Hemmingsen SM, Woolford C, Vandervies SM, Tilly K, Dennis DT, Georgopoulos CP, Hendrix RW, Ellis RJ. 1988. Homologous plant and bacterial proteins chaperone oligomeric protein assembly. *Nature* 333: 330–334.

Hoffman AA, Sørensen JG, Loeschcke V. 2003. Adaptation of *Drosophila* to temperature extremes: Bringing together quantitative and molecular approaches. *Journal of Thermal Biology* 28: 175–213.

Huang LH, Chen B, Kang L. 2007. Impact of mild temperature hardening on thermotolerance, fecundity, and Hsp gene expression in *Liriomyza huidobrensis*. *Journal of Insect Physiology*. 53: 1199–1205.

Huang LH, Kang L. 2007. Cloning and interspecific altered expression of heat shock protein genes in two leafminer species in response to thermal stress. *Insect Molecular Biology* 16: 491–500.

Johnston JA, Ward CL, Kopito RR. 1998. Aggresomes: a cellular response to misfolded proteins. *Journal of Cell Biology* 143: 1883–1898.

- Kim BG, Shim JK, Kim DW, Kwon YJ, Lee KY. 2008. Tissue-specific variation of heat shock protein gene expression in relation to diapause in the bumblebee *Bombus terrestris*. *Entomological Research* 38: 10-16.
- Krebs RA, Feder ME. 1997. Deleterious consequences of Hsp70 overexpression in *Drosophila melanogaster* larvae. *Cell Stress and Chaperones* 2: 60-71.
- Krebs RA, Feder ME. 1998. Hsp70 and larval thermotolerance in *Drosophila melanogaster*: how much is enough and when is more too much? *Journal of Insect Physiology* 44: 1091-1101.
- Kristensen TN, Dahlgaard J, Loeschcke V. 2002. Inbreeding affects Hsp70 expression in two species of *Drosophila* even at benign temperatures. *Evolutionary Ecology Research* 4:1209-1216.
- Lakhotia SC, Singh BN. 1996. Synthesis of ubiquitously present new hsp60 family protein is enhanced by heat shock only in the Malpighian tubules of *Drosophila*. *Experientia* 52: 751-756.
- Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta C^T}$ method. *Methods* 25:402-4081
- Lindquist S. 1986. The heat-shock response. *Annual review of Biochemistry* 55: 1151-1191.
- Martin J, Langer T, Boteva R, Schramel A, Horwich AL, Hartl FU. 1991. Chaperonin-mediated protein folding at the surface of groEL through a “molten globule”-like intermediate. *Nature* 352: 36-42.
- Martin J, Horwich AL, Hartl FU. 1992. Prevention of protein denaturation under heat stress by chaperonin60. *Science* 258: 995-998.
- Martinez J, Perez-Serrano J, Bernadina WE, Rodriguez-Caabeiro F. 2002. Oxidative, heat and anthelmintic stress responses in four species of *Trichinella*: comparative study. *Journal of Experimental Zoology* 293: 66-474.
- Martinus RD, Ryan MT, Naylor DJ, Herd SM, Hoogenraad NJ, Hoj PB. 1995. Role of chaperones in the biogenesis and maintenance of the mitochondrion. *The FASEB Journal* 9: 371-378.
- Ostermann J, Horwich AL, Neupert W, Hartl FU. 1989. Protein folding in mitochondria requires complex formation with hsp60 and ATP hydrolysis. *Nature* 341: 125-130.
- Rios-Arana1 JV, Gardea-Torresdey JL, Webb R, Walsh EJ. 2005. Heat shock protein 60 (HSP60) response of *Platynus patulus* (Rotifera: Monogononta) to combined exposures of arsenic and heavy metals. *Hydrobiologia* 546: 577-585.
- Sanders BM, Pascoe VM, Nakagawa PA, Martin LS. 1992. Persistence of the heat-shock response over time in a common *Mytilus* mussel. *Molecular Marine Biology and Biotechnology* 1: 147-154.
- Shang ZZ, Wang YS, Zou YH. 1979. Study on rearing method of rice stem borer *Chilo suppressalis* Walker, *Acta Entomologica Sinica* 2: 164-167.
- Shen GX, Zhou RL. 2002. *Experimental Techniques of Modern Immunology*, 2nd

edition. Wuhan: HuBei Science and Technology Press.

Singh AK, Lakhota SC. 2000. Tissue-specific variations in the induction of Hsp70 and Hsp64 by heat shock in insects. *Cell Stress Chaperones* 5: 90–97.

Sonoda S, Muhammad A, Tsumuki H. 2006. Cloning and nucleotide sequencing of three heat shock protein genes (*hsp90*, *hsc70*, and *hsp19.5*) from the diamondback moth, *Plutella xylostella* (L.) and their expression in relation to developmental stage and temperature. *Archives of Insect Biochemistry and Physiology* 62: 90–80.

Sørensen JG, Kristensen TN, Loeschcke V. 2003. The evolutionary and ecological role of heat shock proteins. *Ecology Letters* 6: 1025–1037.

Sørensen JG Nielsen MM, Loeschcke V. 2007. Gene expression profile analysis of *Drosophila melanogaster* selected for resistance to environmental stressors. *Journal of Evolutionary Biology* 20(4): 1624–36.

Sunita S, Manoj SR, Tiwari PK. 2006. Expression of HSP60 homologue in sheep blowfly *Lucilia cuprina* during development and heat stress. *Journal of Thermal Biology* 31: 546–555.

Tsugekils R, Mori H, Nishimura M. 1992. Purification, cDNA cloning and Northern-blot analysis of mitochondrial chaperonin 60 from pumpkin cotyledons. *European Journal of Biochemistry* 209:453–458.

Wheelock CE, Wolfe MF, Olsen H, Tjeerdema ML, Sowby ML. 1999. Hsp60-induced tolerance in the rotifer *Brachionus plicatilis* exposed to multiple environmental

contaminants. *Archives of Environmental Contamination and Toxicology* 36: 281–287.

Wang HS, Wang XH, Zhou CS, Huang LH, Zhang SF, Guo W, Kang L. 2007. cDNA cloning of heat shock proteins and their expression in the two phases of the migratory locust. *Insect Molecular Biology* 16: 207–219.

Wong CS, Mak CH, Ko RC. 2004. Cloning and characterization of the mitochondrial heat-shock protein 60 gene of *Trichinella spiralis*. *Parasitology Research* 93: 461–467.

Yocum GD. 2001. Differential expression of two HSP70 transcripts in response to cold shock, thermoperiod, and adult diapause in the Colorado potato beetle. *Journal of Insect Physiology* 47: 1139–1145.