

Efficacy of Pitfall Trapping, Winkler and Berlese Extraction Methods for Measuring Ground-Dwelling Arthropods in Moist-Deciduous Forests in the Western Ghats

Authors: Sabu, Thomas K., and Shiju, Raj T.

Source: Journal of Insect Science, 10(98) : 1-17

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.010.9801>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Efficacy of pitfall trapping, Winkler and Berlese extraction methods for measuring ground-dwelling arthropods in moist-deciduous forests in the Western Ghats

Thomas K. Sabu^a, Raj T. Shiju

Litter Entomology Research Unit, Post Graduate & Research Department of Zoology, St. Joseph's College, Devagiri, Calicut, Kerala, India 673008

Abstract

The present study provides data to decide on the most appropriate method for sampling of ground-dwelling arthropods measured in a moist-deciduous forest in the Western Ghats in South India. The abundance of ground-dwelling arthropods was compared among large numbers of samples obtained using pitfall trapping, Berlese and Winkler extraction methods. Highest abundance and frequency of most of the represented taxa indicated pitfall trapping as the ideal method for sampling of ground-dwelling arthropods. However, with possible bias towards surface-active taxa, pitfall-trapping data is inappropriate for quantitative studies, and Berlese extraction is the better alternative. Berlese extraction is the better method for quantitative measurements than the other two methods, whereas pitfall trapping would be appropriate for qualitative measurements. A comparison of the Berlese and Winkler extraction data shows that in a quantitative multigroup approach, Winkler extraction was inferior to Berlese extraction because the total number of arthropods caught was the lowest; and many of the taxa that were caught from an identical sample via Berlese extraction method were not caught. Significantly a greater frequency and higher abundance of arthropods belonging to Orthoptera, Blattaria, and Diptera occurred in pitfall-trapped samples and Psocoptera and Acariformes in Berlese-extracted samples than that were obtained in the other two methods, indicating that both methods are useful, one complementing the other, eliminating a chance for possible under-representation of taxa in quantitative studies.

Keywords: forest floor arthropods, soil-litter arthropods, quantitative and qualitative sampling, Acariformes, Araneae, Blattaria, Chalcididae, Chilopoda, Coleoptera, Dermaptera, Diplopoda, Diptera, Formicidae, Hemiptera, Homoptera, Isopoda, Isoptera, Larva, Lepidoptera, Mantoidea, Orthoptera, Hymenoptera, Parasitiformes, Phasmida, Pseudoscorpionida, Psocoptera, Thysanoptera

Correspondence: ^a sabukthomas@gmail.com

Associate Editor: T.X. Liu was editor of this paper.

Received: 11 July 2008, **Accepted:** 15 January 2009

Copyright : This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 10, Number 98

Cite this paper as:

Sabu TK, Shiju RT. 2010. Efficacy of pitfall trapping, Winkler and Berlese extraction methods for measuring ground-dwelling arthropods in moist-deciduous forests in the Western Ghats. *Journal of Insect Science* 10:98 available online: insectscience.org/10.98

Introduction

Inclusion of ground-dwelling arthropods in biodiversity inventories and environmental assessment surveys has increased in the recent past (Oliver and Beattie 1996). Since most ground-dwelling arthropods are minute and numerous, and usually not easy to detect by an unaided eye, assessment of populations of these organisms is hard and labor-intensive. Specific methods that are effective in extracting a high proportion of these taxa are usually necessary. Three methods employed widely to survey ground-dwelling arthropods are pitfall trapping (Holland and Reynolds 2005; King and Porter 2005; Ward et al. 2001; Standen 2000; Brennan et al. 1999; Holland and Smith 1999; Mommertz et al. 1996; Mesibov et al. 1995; Olson 1991; Adis 1979), Berlese extraction (Anu et al. 2009; Palacios-Vargas et al. 2007; Anu and Thomas 2006; King and Porter 2005; Richardson et al. 2005; Edwards 1991; Frith and Frith 1990), and Winkler extraction (Anto and Thomas 2007; Philpott et al. 2007; Robertson 2007; Vineesh et al. 2007; Krell et al. 2005; Leponce et al. 2004; Fisher and Robertson 2002; Longino et al. 2002; Chung et al. 2000; Fisher 1998; Belshaw and Bolton 1994).

Pitfall trapping, the simplest and cheapest method among the three, is efficient in capturing those arthropod taxa that are nocturnally active on the surface, but is inefficient in capturing either the bottom dwellers or those that disseminate by flying (Hansen and New 2005; Leather and Watt 2005; Woodcock 2005; Work et al. 2002; Ward et al. 2001; Standen 2000; Mommertz et al. 1996; Mesibov et al. 1995; Spence and Niemela 1994; Topping and Sunderland 1992; Adis 1979; Greenslade 1964). Pitfall trapping is most effective in open habitats, such as

grasslands and scrub vegetation because the capture values can be affected by vegetation complexity (Melbourne 1999; Majer 1997; Greenslade 1964). Berlese extraction necessitates the use of expensive and unwieldy apparatuses and electricity, which may not be available in remote study sites (Krell et al. 2005; Lasebikan et al. 1978); moreover, in the Berlese extraction method, separation of soil particles and debris that drop into the collection solution along with fauna makes sampling more time-consuming and labour intensive than pitfall trapping (Robertson 2007; Edwards 1991). Winkler extraction is suitable for the extraction of litter-inhabiting, rapidly mobile Formicidae (Delabie et al. 2007; Underwood and Fisher 2006; Longino et al. 2002; Parr and Chown 2001; Bestelmeyer et al. 2000; Delabie et al. 2000; Olson 1991; Nadkarni and Longino 1990) and for the extraction of forest-litter inhabiting Coleoptera (Didham et al. 1998). However, Winkler extraction is less suitable for the extraction of all ground-dwelling arthropod taxa because chances for escape of the larger and more agile taxa are high; moreover, chances for the death of small taxa, with a narrow ecological tolerance, before they drop into the collection jars is also equally high (Didham et al. 1998; Besuchet et al. 1987). Winkler extraction is a relatively slow process in moist and humid environments, and when taxa of Formicidae are abundant in the sample, they may consume other fauna during extraction (Schillhammer 2001; Wheeler and McHugh 1987). Even for the litter-inhabiting Formicidae, for which the Winkler extraction method has proved suitable, it can extract a greater variety of taxa in warm-weather regions than it can in cold-weather regions (Leponce et al. 2004). Therefore, the relative efficiency of Winkler extraction in capturing

ground-dwelling arthropods compared with the other sampling methods needs establishment.

In summary, the effectiveness of these three widely used methods to extract arthropods from soil substrates is being debated. Only a few replicated field studies have attempted to evaluate critically and compare quantitatively the extraction efficiency of the three sampling methods. Most of the earlier efforts evaluated the sampling efficiency of either the Berlese or the Winkler extraction method against pitfall trapping by sampling of a few specific arthropod taxa (e.g., pitfall trapping and Berlese extraction for Carabidae, Spence and Niemela 1994; Coleoptera, Formicidae and Araneae, Oliver and Beatie 1996; litter-inhabiting Formicidae, King and Porter 2005; Diplopoda, Snyder et al. 2006; Pitfall trapping and Winkler extraction for litter Formicidae, Fisher and Robertson 2002, Parr and Chown 2001, Delabie et al. 2000, Fisher 1999, Olson 1991.)

Hence, a considered opinion on the ideal sampling method for the extraction of the whole suite of ground-dwelling arthropods among the three remains to be established. Nonetheless, because of the easy manipulability in terms of time and cost effectiveness, pitfall trapping and Winkler extraction methods are being preferred over the Berlese extraction method in ecological surveys of soil arthropods (Krell et al. 2005; Chung et al. 2000; Didham et al. 1998; Belshaw and Bolton 1994; Spence and Niemela 1994; Hammond 1990; Ward 1987). These conclusions have been arrived at without evaluating the extraction efficiencies of the three methods although it is essential that any chosen method should address minimizing problems associated with complex statistical analysis, which could be

compounded further by low numbers of taxa (Prasifka et al. 2007; Parr and Chown 2001).

In the present paper, the trapping efficiencies of the three widely used ground-dwelling arthropod trapping methods (pitfall, Berlese and Winkler extraction methods) were compared and evaluated using a field trial done in the moist deciduous forests in the Western Ghats in South India. The objective was to determine whether the three methods are equally effective in the separation and to determine which method achieves the best overall population numbers of as many taxa as possible.

Materials and Methods

Study area

The study was carried out in the moist deciduous forests of Sholayar (220 MASL, 20.55 km²) (10° 17'-10° 19' N; 76° 39'-76° 44' E), situated close to the Athirapally–Vazhachal waterfalls, 60 km south of the town of Trichur (Trichur District, Kerala State, India) located in the South-Western Ghats of moist deciduous ecoregion (Wikramanayake et al. 2002). Annual temperature 24-32° C; 40-80% RH; average rainfall 3,000-3,250 mm/year, which occurs mostly in June–November; June, July, and August receive the most rain.

Sampling

Sampling was done in the first week of February 2006. Although a thorough population assessment of any group of invertebrates necessitates sampling at different times in the year (Edwards 1991), the survey period and intensity, although apparently inadequate for definitive inventory, served the purpose of comparing between trap designs, efficiency or capture of trappable fauna. Three parallel line transects, one

dedicated for each extraction type (Berlese extraction method, pitfall trapping, and the Winkler methods) separated by 25 m inter-transect distances, were constructed north-southerly. The 25 m inter-transect distance between two consecutively set trap-transects was meant to minimize possible depletion effects, which can be caused by pitfall trapping (Digweed et al. 1995). Forty pitfall trapped samples and 40 litter samples each for Winkler and Berlese extraction were obtained. All samples were obtained on the same day between 08:00 and 09:30 h.

Each litter sample for Winkler and Berlese extraction was obtained by placing a 50 x 50 cm wooden frame on the forest floor and collecting the leaves, litter and loose humus from within the frame area into a large polythene bag (Frith and Frith 1990). Samples were obtained taking care to prevent possible escape of any invertebrate. The litter thus collected refers to the upper organic litter layer plus the loose humus layer. No underlying compact soil was obtained. Litter samples for Berlese and Winkler extractions were sieved in a 1.5 cm mesh wire sieve to separate larger materials of litter and transported to the laboratory in individual polythene bags.

Fauna was extracted with Berlese extraction apparatus (30.5 cm diameter, 35.6 cm height, 4-6 mm mesh screen, 25 w tungsten-filament lamp) over five days in 70% alcohol. Litter samples for Winkler extraction of the fauna (Besuchet et al. 1987) were placed in coarse-mesh bags, which were suspended inside a large sealed cloth bag suspended over a collecting bottle containing 75% ethanol. The litter and soil were left to dry at room temperature for five days. The litter material was gently mixed every day to ensure that the fauna remained active and to improve their

chances of dropping into the collection cup (Parr and Chown 2001; Besuchet et al. 1987).

Each pitfall trap consisted of a black plastic bowl (210 mm diameter, 150 mm depth), buried to its rim in soil and partly filled with 30 mm of nontoxic, propylene glycol. Each trap was topped with a dark-plastic tray supported on iron bars to prevent either desiccation or flooding; such a system operated for 24 h continuously to avoid bias in catches arising from diurnal activity variation of fauna (Mommertz et al. 1996). Trapped fauna were separated, identified, counted, and the abundance and frequency of occurrence of taxa at each site was recorded.

The fauna obtained from the 40 litter samples from each method were available for data analysis. Taxa with > 30% frequency of occurrence in any of one of the sampling method was categorized as major, and the others as minor taxa. The sampling method, which trapped > 40% frequency of occurrence of a particular taxon, even if that taxon was trapped with < 40% frequency of occurrence in the other two methods was deemed to be 'reasonably effective' in sampling that particular taxon.

Data analysis


Significant differences in the frequency of collection among sampling methods (abundance data with median and with low abundance and total absence of some taxa) made comparisons using common parametric statistics inappropriate. In the data analysis, emphasis was placed on seeking differences in the frequency of occurrence of arthropod types and less on testing for differences in the mean number of arthropod types (Prasifka 2007). Higher frequency of taxa obtained more frequently through a particular method than by the other two methods rendered that

method more reliable. However, to summarize arthropod captures by trap type, the means and standard errors derived from individual trap were calculated for each arthropod group. To test for differences in the frequency with which particular arthropod taxa were collected by the three trap types, 2×3 contingency tables categorized each trap as either successful (one or more individuals collected) or unsuccessful (zero individuals collected); the differences were assessed with chi-square tests. Significant chi-squared values indicated an effect of trap type on the proportion of samples containing one or more individuals of an arthropod taxon (Prasifka et al. 2007). Trap-wise differences in the catch efficiency of individual taxa among the three trap types were assessed with a two sample z-test. Univariate comparisons through Kruskal-Wallis H tests were used to evaluate the significance level of trap-wise difference in faunal abundance. When significant differences were found, a Mann-Whitney U test was applied to determine which pairs of

methods were different significantly (Weiss 2007). All the analyses were done using MegaStat Version 10.0 (Orris 2005).

Results

From the three methods tested, 12,257 individuals belonging to 25 arthropod taxa were collected (Figure 1, Table 1), which could be broadly divided into (i) a major group of 14 arthropod taxa and one minor taxon, which showed significant differences in capture among the tested trapping methods and (ii) a minor group of 10 arthropod taxa with no difference in capture among the tested trapping methods (Tables 2 & 3). From among the 25 taxa, 18 occurred in all the three tested methods, whereas taxa belonging to Phasmida, Isoptera, Lepidoptera, Chilopoda occurred only in Berlese extraction and pitfall trapping methods, taxa belonging to Dermaptera and Mantoidea only occurred in pitfall trapping, and one taxon belonging to Pseudoscorpionida occurred only in Berlese



extraction methods (Figure 1, Table 1). Based on the frequency of occurrence of fauna, the methods effectively trapped 13 major taxa in pitfall traps, nine major taxa and one minor taxa in the Berlese extraction method, and five major taxa in the Winkler extraction method (Figure 2, Table 1).

In the Winkler extraction method, 18 taxa were obtained. The proportionate distribution of dominant taxa in the collection was in the following order: Acariformes (85%) > insect larvae (75%) > Coleoptera (60%) > Araneae (48%) > Formicidae (40%) (Table 1). For the major taxa belonging to Coleoptera, Orthoptera, Blattaria, Hemiptera, Diptera, other Hymenoptera (except Formicidae and Chalcidae), Araneae, and Chalcidae, the Winkler extraction method separated the same frequency of occurrence as that of the Berlese

extraction method, and for Psocoptera and Parasitiformes, the Winkler method obtained the same frequency of occurrence as that obtained in the pitfall trapping method (Table 2). Though the frequency of occurrence was lower than in other methods, 40% of occurrence (value set to consider any method as 'reasonably effective') of five taxa viz., Coleoptera, larval forms of diverse insect orders, Araneae, Acariformes, and Formicidae was obtained in the Winkler extraction method.

In the Berlese extraction method, 23 taxa were obtained (Figure 2, Table 1). The dominance pattern of the major taxa was Acariformes (98%) > insect larvae (93%) > Formicidae (68%) > Coleoptera (63%) > Araneae (60%) > Collembola (45%) > Psocoptera (35%). The Berlese extraction method recorded the

Table 1. Frequency (%) and abundance (median + inter quartiles) of ground dwelling arthropods collected from pitfall trapping (PIT), the Berlese extraction method (BEM) and the Winkler extraction method (WEM).

Taxa/Groups	Pitfall Trapping				Berlese extraction				Winkler extraction			
	Frequency (%)	Abundance			Frequency (%)	Abundance			Frequency (%)	Abundance		
		Median	Q1	Q3		Median	Q1	Q3		Median	Q1	Q3
Collembola	90	3	1	12	45	0	0	2	13	0	0	0
Phasmida	5	0	0	0	3	0	0	0	0	0	0	0
Orthoptera	100	5	2	11	3	0	0	0	5	0	0	0
Dermaptera	5	0	0	0	0	0	0	0	0	0	0	0
Isoptera	8	0	0	0	13	0	0	0	0	0	0	0
Blattaria	90	5	2	12	15	0	0	0	5	0	0	0
Mantoidea	3	0	0	0	0	0	0	0	0	0	0	0
Psocoptera	8	0	0	0	35	0	0	1	13	0	0	0
Thysanoptera	53	1	0	2	30	0	0	1	5	0	0	0
Hemiptera	100	7	3.5	10.5	28	0	0	1	15	0	0	0
Homoptera	10	0	0	0	8	0	0	0	5	0	0	0
Coleoptera	88	10	4	19.5	63	2	0	8	60	1	0	2
Diptera	100	20.5	8	48	20	0	0	0	23	0	0	0
Lepidoptera	10	0	0	0	13	0	0	0	0	0	0	0
Other Hymenoptera	38	0	0	2	5	0	0	0	8	0	0	0
Chalcidae	78	2	1	3	23	0	0	0	18	0	0	0
Formicidae	100	75	36.5	103	68	2	0	8.5	40	0	0	1
Larva	93	7	3	14	93	6	3	10.5	75	2	0.5	3
Araneae	100	14.5	10	17	60	1	0	2	48	0	0	1
Diplopoda	8	0	0	0	15	0	0	0	5	0	0	0
Acariformes	100	13	8	34.5	98	14.5	5	55.5	85	3	1	7
Parasitiformes	5	0	0	0	23	0	0	0	5	0	0	0
Pseud-oscorpionida	0	0	0	0	3	0	0	0	0	0	0	0

highest frequency of occurrence for major taxa belonging to Collembola, Thysanoptera, larval forms of diverse insect orders, Araneae, Formicidae, Psocoptera, and Acariformes and for one minor taxon, Parasitiformes, and an equivalent level of frequency of occurrence as Winkler extraction for 8 of the 14 taxa belonging to Orthoptera, Blattaria, Hemiptera, Coleoptera, Diptera, other Hymenoptera, Araneae, and Chalcidae (Table 2). A higher representation (> 40% of occurrence) of five taxa viz., Collembola, Coleoptera, larval forms of diverse insect orders, Araneae, and Formicidae was obtained in the Berlese extraction method.

In the pitfall trapping method, except Pseudoscorpionida, the remaining 24 taxa were recorded (Figure 2, Table 1).

Proportionate occurrence of the major taxa was in the order of: Acariformes = Formicidae = Orthoptera = Hemiptera = Diptera = Araneae > larval forms of diverse insect orders (93%) > Collembola (90%) > Blattaria (90%) > Coleoptera (88%) > Chalcidae (78%) > Thysanoptera (53%) (Figure 2, Table 1). For 11 out of the 14 taxa belonging to the Collembola, Orthoptera, Formicidae, Blattaria, Hemiptera, Coleoptera, Diptera, other Hymenoptera, Thysanoptera, Araneae, Chalcidae, Formicidae, the pitfall trapping method yielded the highest frequency of occurrence (Table 2). For two taxa (insect larvae and Acariformes) both pitfall trapping and Berlese extraction methods recorded same level of frequency of occurrence. Except for Psocoptera and Acariformes, the pitfall tapping method captured the highest frequency of occurrence for 13 out of the 14 major taxa (Table 2) and the same level of frequency of occurrence for 10 minor taxa.




Table 2. Results of Chi-squared test and two sample z- tests on the variation in the frequency of collection for ground dwelling arthropods using pitfall trapping (PIT), the Berlese extraction method (BEM) and the Winkler extraction method (WEM).

Taxa/Groups	Trapping methods	z-test		Chi-squared test	
		z	P-value	c ²	P-value
Collembola	PIT/BEM	4.9	0.000	48.48	0.000
	PIT/WEM	10.98	0.000		
	BEM/WEM	3.44	0.001		
Orthoptera	PIT/BEM	39.50	0.000	107.496	0.000
	PIT/WEM	27.57	0.000		
	BEM/WEM	-0.59	0.555		
Blattaria	PIT/BEM	10.17	0.000	74.354	0.000
	PIT/WEM	14.50	0.000		
	BEM/WEM	1.51	0.131		
Psocoptera	PIT/BEM	-3.19	0.001	11.466	0.003
	PIT/WEM	-0.75	0.454		
	BEM/WEM	2.45	0.014		
Thysanoptera	PIT/BEM	2.10	0.036	21.862	0.000
	PIT/WEM	5.51	0.000		
	BEM/WEM	3.12	0.002		
Hemiptera	PIT/BEM	10.27	0.000	67.569	0.000
	PIT/WEM	15.06	0.000		
	BEM/WEM	1.38	0.167		
Coleoptera	PIT/BEM	2.70	0.007	8.810	0.012
	PIT/WEM	2.94	0.003		
	BEM/WEM	0.23	0.818		
Diptera	PIT/BEM	12.65	0.000	66.366	0.000
	PIT/WEM	11.74	0.000		
	BEM/WEM	-0.27	0.785		
Hymenoptera	PIT/BEM	3.87	0.000	18.84	0.000
	PIT/WEM	3.44	0.001		
	BEM/WEM	-0.46	0.644		
Chalcidae	PIT/BEM	5.89	0.000	37.214	0.000
	PIT/WEM	6.72	0.000		
	BEM/WEM	0.56	0.575		
Formicidae	PIT/BEM	4.39	0.000	33.839	0.000
	PIT/WEM	7.75	0.000		
	BEM/WEM	2.57	0.010		
Larvae	PIT/BEM	0.00	1.000	7.067	0.029
	PIT/WEM	2.18	0.029		
	BEM/WEM	2.18	0.029		
Araneae	PIT/BEM	5.16	0.000	28.212	0.000
	PIT/WEM	6.65	0.000		
	BEM/WEM	1.13	0.258		
Acariformes	PIT/BEM	1.01	0.311	9.406	0.009
	PIT/WEM	2.66	0.008		
	BEM/WEM	2.03	0.042		
Parasitiformes	PIT/BEM	-2.35	0.019	8.454	0.015
	PIT/WEM	0.00	1.000		
	BEM/WEM	2.35	0.019		

In summary, the pitfall trapping method yielded highest frequency of occurrence for 13 out of the 14 major taxa, Berlese extraction method for 3 out of the 14 taxa and also for one minor taxon, and the Winkler extraction method for none of the major taxa (Table 2). For the 10 minor taxa, all the three methods yielded the same level of frequency of occurrence. The arthropod abundance (median + inter quartiles) data for the ground-dwelling arthropod taxa in the three sampling methods is summarized in Table 1. Pitfall trapping recorded the highest abundance of 19 of the 25 arthropod taxa except for six taxa belonging to Psocoptera, Acariformes, Parasitiformes, Pseudoscorpionida, Chilopoda, and Diplopoda (Table 3). Abundance of all taxa was the lowest in Winkler extracted samples. High abundances of Blattaria, Diptera, and Orthoptera in the pitfall trapping and low abundances of these taxa in both Berlese and Winkler methods occurred.

Discussion

Results of this study lead to recommendations on the relative usefulness of pitfall trapping, and Berlese and Winkler extraction methods for sampling ground-dwelling arthropods in subtropical, moist deciduous forests. A pronounced difference was found in the capture of major ground-dwelling arthropod taxa among the three tested sampling methods. Pitfall trapping yielded the maximal capture (both frequency and numbers) of 87% of the major taxa, followed by the Berlese extraction method (27%). The Winkler method proved ineffective for any of the major taxa in the moist, subtropical South Indian forest conditions. These percentages indicated that pitfall trapping was a useful standard arthropod collection method for ecological studies of ground-surface-dwelling

arthropods compared with the Berlese and Winkler extraction methods. Irrespective of the method tested, few individuals of the minor taxa were captured and such an effect could be attributed to the low population densities of these taxa in the moist forests of the Western Ghats (Anu et al. 2009; Vineesh 2007; Anu 2006). Non-significant differences in the capture of minor taxa among the different trap types are difficult to interpret because of their low frequency of occurrence and abundance.

However, a bias was apparent in the samples obtained via pitfall trapping in comparison with the other two methods. Pitfall trapping captured high numbers of taxa active at the ground level, viz., taxa belonging to Orthoptera, Blattaria, Diptera, Araneae, Formicidae, Collembola, Hemiptera, Coleoptera, and other Hymenoptera (Prasifka et al. 2007; Leather and Watt 2005; Woodcock 2005; Bicknell et al. 2000), with 95-100% frequency of capture when compared to their relatively low frequency of capture in the other two methods. On the contrary, with the Berlese sampling method, high capture was recorded of the less-active taxa associated with moisture and sheltered areas, viz., Acariformes, larvae of diverse insect orders, Psocoptera and Parasitiformes. Such variations and differences in the capture of taxa in relation to their surface activity is an established weakness of pitfall-trapping method, leading to the generalizations that (i) pitfall trap collections of arthropods are measures of activity rather than density estimates, (ii) pitfall trap extractions do not consider the resting and evasive behavior of many taxa thus leading to an under-representation of such taxa, and (iii) a comparison of data from pitfall traps with other methods is impossible in quantitative studies (Woodcock 2005; Oliver and Beattie

Table 3. Results of Kruskal Wallis and Mann-Whitney tests on the variation in the abundance of collection for ground dwelling arthropods using pitfall trapping (PIT), the Berlese extraction method (BEM) and the Winkler extraction method (WEM).

Taxa/Groups	Mann-Whitney		Kruskal-Wallis	
	Trapping methods	P-value	H	P-value
Collembola	PIT/BEM	0.000	50.62	0.000
	PIT/WEM	0.000		
	BEM/WEM	0.001		
Orthoptera	PIT/BEM	0.000	106.56	0.000
	PIT/WEM	0.000		
	BEM/WEM	0.569		
Blattaria	PIT/BEM	0.000	84.25	0.000
	PIT/WEM	0.000		
	BEM/WEM	0.141		
Psocoptera	PIT/BEM	0.002	12.75	0.002
	PIT/WEM	0.448		
	BEM/WEM	0.012		
Thysanoptera	PIT/BEM	0.076	21.67	0.000
	PIT/WEM	0.000		
	BEM/WEM	0.003		
Hemiptera	PIT/BEM	0.000	86.32	0.000
	PIT/WEM	0.000		
	BEM/WEM	0.197		
Coleoptera	PIT/BEM	0.000	29.77	0.000
	PIT/WEM	0.000		
	BEM/WEM	0.077		
Diptera	PIT/BEM	0.000	88.92	0.000
	PIT/WEM	0.000		
	BEM/WEM	0.962		
Other Hymenoptera	PIT/BEM	0.001	19.71	0.000
	PIT/WEM	0.001		
	BEM/WEM	0.671		
Chalcidae	PIT/BEM	0.000	45.23	0.000
	PIT/WEM	0.000		
	BEM/WEM	0.557		
Formicidae	PIT/BEM	0.000	70.91	0.000
	PIT/WEM	0.000		
	BEM/WEM	0.005		
Larvae	PIT/BEM	0.461	27.06	0.000
	PIT/WEM	0.000		
	BEM/WEM	0.000		
Araneae	PIT/BEM	0.000	81.88	0.000
	PIT/WEM	0.000		
	BEM/WEM	0.054		
Acariformes	PIT/BEM	0.736	34.45	0.000
	PIT/WEM	0.000		
	BEM/WEM	0.000		
Parasitiformes	PIT/BEM	0.021	8.84	0.012
	PIT/WEM	1.000		
	BEM/WEM	0.021		

1996; Topping and Sunderland 1992; Adis 1979; Southwood 1978; Greenslade 1964). Moreover, unlike the other two methods, the pitfall trapping method captures arthropod taxa from an unknown area and calculations of absolute density of populations of taxa that is the number of individuals per unit of habitat is nearly impossible (Woodcock 2005; Work et al. 2002; Standen 2000; Spence and Niemela 1994; Halsall and Wratten 1988; Greenslade 1964). Calibrations and adjustments needed for the removal of these complex effects and standardization of data would diminish the time and labour efficiency advantages of using pitfall trapping (Stoyan and Kuschka 2001; Mommertz et al. 1996; Spence and Niemela 1994; Topping and Sunderland 1992). The results support the earlier findings and since the data from pitfall trapping are not useful for estimating absolute abundance (populations per unit area or volume) in multigroup ecological approach involving ground surface dwelling arthropods, it would be prudent to limit their use for production of qualitative data. These setbacks (viz., sampling biases and interpretational difficulties) make density-based estimates from quadrat sampling methods (Berlese or Winkler extraction), which measure populations in numbers of animals/unit area as a better alternative for quantitative multitaxa ecological studies of ground-dwelling arthropods.

Unlike the Berlese or Winkler litter extraction methods, pitfall trapping requires two trips to interior forests: one to set the traps and the second to retrieve traps and samples. A second field trip to retrieve the traps leads to additional expenditure, loss of time and practical difficulties in protected forests with restricted access. Moreover, multiple chances of wildlife disturbing the field-placed traps and inclement weather affecting the sampling

effort in subtropical forests also exist. This leaves the researcher apprehensive about the success of his collection efforts until the second trip and makes pitfall trapping more laborious and chance oriented than the other two methods.

Recent studies have highlighted the Winkler extraction method as a less expensive, more convenient, and a more efficient alternative method for exhaustive extraction of soil macro-invertebrates (Krell et al. 2005; Didham et al. 1998; Nadkarni and Longino 1990). Hence, at the beginning of this study, it was expected that the Winkler extraction method would prove better than the pitfall trapping and Berlese extraction methods by obtaining greater numbers and frequency of different arthropod taxa. However, an entirely different outcome became evident. Firstly, the Winkler extraction method underestimated the abundance and frequency of the major taxa (in South Indian moist deciduous forest conditions), except taxa belonging to Coleoptera, larvae of multiple insect orders, Araneae, Acariformes, and Formicidae; secondly, most of the minor taxa belonging to Dermaptera, Mantoidea, Pseudoscorpionida, Phasmida, Isoptera, Lepidoptera, and Chilopoda were missed out completely. These limitations highlight the less emphatic observations of Besuchet et al. (1987) that Winkler extraction method is less suitable for the extraction of all taxa, and there is possibility of death of taxa with a narrow ecological tolerance before dropping into the collection jars. Although Winkler extractions are cost effective and convenient with limited time needed to sort fauna (Krell et al. 2005) and effective in studies of litter Formicidae and Coleoptera (Fisher 2004; Fisher and Robertson 2002; Didham et al. 1998; Olson 1991; Nadkarni and Longino 1990; Besuchet et al. 1987), the limited volume of quantitative

information they generate for the majority of ground surface dwelling arthropod taxa weaken their value as an effective individual sampling method for intensive ecological studies of ground-dwelling arthropod fauna. Despite that, the Berlese extraction method will be costly and laborious because it requires more time to sort out the fauna from fallen debris and soil in the laboratory (Snyder et al. 2006). Berlese extraction recorded higher capture of taxa per unit volume than the Winkler extraction method. Greater efficiency and thoroughness of the sampling effort makes the Berlese extraction method a better choice than the Winkler extraction method as an individual quantitative sampling method for the ground surface dwelling fauna in a subtropical moist deciduous forest.

Group and trap specific differences noted in the present study supports the findings by Edwards (1991) and Standen (2000) that no single extraction method is the best for all taxa of ground-dwelling arthropods, and it may be necessary to use more than one method based on the aim of the study. The selection of sampling methods for ground-dwelling arthropods should be made based on the data — quantitative or qualitative — required for the study. High trapping efficiency of a majority of the taxa makes pitfall trapping the best method in qualitative inventory studies of ground-dwelling arthropods, but not for quantitative studies because of the above-cited setbacks. Trapping success of pitfall traps conforms to the findings (Spence and Niemelä 1994) that pitfall trapping remains the only realistic way to survey large acreages where qualitative inventory and a comparison of species assemblages of ground-active arthropods is required. However, for quantitative studies of ground-dwelling arthropods, the Berlese extraction method is the best option. Very

high abundance and frequency of occurrence of Blattaria, Diptera and Orthoptera in pitfall trapping and very low trapping with the Berlese extraction method suggest that a combination of pitfall trapping and Berlese extraction and standardization of pitfall trapping data (Stoyan and Kuschka 2001) is more feasible for exhaustive quantitative studies of surface dwelling arthropods.

Conclusions

The relative abundance and frequency of occurrence of fauna was different with the three sampling methods. When cost and time constraints dictate the limiting of ground-dwelling arthropod sampling to one method, the Berlese extraction method is ideal for quantitative estimates, and the pitfall trapping method is ideal for qualitative estimates.

Since the three taxa (Orthoptera, Diptera and Blattaria) with a low catch probability with Berlese extraction were caught efficiently with pitfall trapping, inclusion of pitfall trapping with appropriate adjustments would be the method for comprehensive quantitative surveys of ground-dwelling arthropods. Although pitfall trapping samples tend to include more ground-active species, its efficiency indicates that pitfall trapping is certainly the method of choice for an individual quantitative sampling method for most major taxa except the Psocoptera and insect larval forms, for which the Berlese extraction method is a better option.

As a cost effective, individual quantitative sampling method, Winkler extraction is suitable for obtaining Coleoptera and Acariformes in addition to litter Formicidae, for which it is an established method (Delabie et al. 2007; Underwood and Fisher 2006; Longino et al. 2002; Bestelmeyer et al. 2000;

Delabie et al. 2000), but not for ecological studies involving multiple arthropod groups.

Acknowledgements

We thank the Department of Science and Technology (DST), Government of India for financial assistance, the Kerala Forest and Wildlife Department for logistical support and permissions, and K.T. Thomachan (Devagiri College, Calicut) for the support in statistical analysis. We are grateful to A. Raman (Charles Sturt University, Australia) for critical comments. We thank T.N. Ananthkrishnan (Emeritus Scientist, Chennai), John M. Holland (Game Conservancy Trust, UK), Jerrad Prasifka (USDA - Agricultural Research Service, USA), Tana Wood (Berkely University, USA) for providing relevant literature, P.J. Vineesh, A.K. Aneesh and K.V. Vinod for field collections and A. Anisha for sample sorting.

References

- Adis J. 1979. Problems of interpreting arthropod sampling with pitfall traps. *Zoologischer Anzeiger* 202: 177-184.
- Anto A, Thomas SK. 2007. Biodiversity analysis of forest litter ant assemblages in Wynad region of Western Ghats with taxonomic and conventional diversity measures. *Journal of Insect Science* 6:60, available online at <http://www.insectscience.org/6.60/>
- Anu A. 2006. *Entomofaunal dynamics and biochemistry of litter decomposition in a natural forest with special reference to the systematics of dung Coleoptera (Coleoptera: Scarabaeinae)*. Ph.D. dissertation. University of Calicut, Kerala.
- Anu A, Thomas SK. 2006. Litter arthropod diversity and community structure in an evergreen forest in the Wayanad region of Western Ghats. *Entomon* 31: 181-190.
- Anu A, Sabu TK, Vineesh PJ. 2009. Seasonality of litter insects and relationship with rainfall in a wet evergreen forest in south Western Ghats. *Journal of Insect Science*. 9:46, available online: <http://www.insectscience.org/9.46/>
- Belshaw R, Bolton B. 1994. A survey of the leaf litter ant fauna in Ghana, West Africa (Hymenoptera: Formicidae). *Journal of Hymenoptera Research* 3: 5-16.
- Bestelmeyer BT, Agosti D, Alonso LE, Brandaö CRF, Brown Jr WL, Delabie JHC, Silvestre R. 2000. Field methods for the study of ground-dwelling ants. In: Agosti D, Majer JD, Alonso LE, Shultz TR, editors. *Ants: Standard methods for measuring and monitoring biodiversity*. pp. 122-144. Smithsonian Institution Press.
- Besuchet C, Burckhardt DH, Lobl I. 1987. The "Winkler/Moczarski" elector as an efficient extractor for fungus and litter Coleoptera. *Coleopterists Bulletin* 41: 392-394.
- Bignell DE, Widodo E, Susilo FX, Suryo H. 2000. Soil macrofauna: Ground-dwelling ants, termites, other macroarthropods and earthworms. In: Gillison AN, editor. *Above-ground Biodiversity Assessment Working Group summary report 1996-99: Impact of different land uses on biodiversity*. pp 91-127. International Centre for Research in Agroforestry (ICRAF), Nairobi, Kenya. Available at: <http://www.asb.cgiar.org/pdfwebdocs/Biodiv%20Study%20WG%20reports/C-Sec7.pdf>

Brennan KEC, Majer JD, Reygart N. 1999. Determination of an optimal pitfall trap size for sampling spiders in a Western Australian Jarrah forest. *Journal of Insect Conservation* 3: 297-307.

Chung AYC, Eggleton P, Speight MR, Hammond PM, Chey VK. 2000. The diversity of beetle assemblages in different habitat types in Sabah, Malaysia. *Bulletin of Entomological Research* 90: 475-496.

Delabie JHC, Fisher BL, Majer JD, Wright IW. 2000. Sampling effort and choice of methods. In: Agosti D, Majer JD, Alonso LE, Shultz TR, editors. *Ants: standard methods for measuring and monitoring biodiversity*. pp 145–154. Smithsonian Institution Press.

Delabie JHC, Jahyny B, Nascimento IC, Mariano CSF, Lacau S, Campiolo S, Philpott SM, Leponce M. 2007. Contribution of cocoa plantations to the conservation of native ants (Insecta: Hymenoptera: Formicidae) with a special emphasis on the Atlantic Forest fauna of southern Bahia. *Brazil Biodiversity and Conservation* 16: 359-2384.

Didham RK, Lawton JH, Hammond PM, Eggleton P. 1998. Trophic structure stability and extinction dynamics of Coleoptera (Coleoptera) in tropical forest fragments. *Philosophical Transactions of the Royal Society, Series B* 353: 437-451.

Digweed SC, Currie CR, Carcamo HA, Spence JR. 1995. Digging out the ‘digging-in effect’ of pitfall traps: Influences of depletion and disturbance on catches of ground Coleoptera (Coleoptera: Carabidae). *Pedobiologia* 39: 61-576.

Edwards CA. 1991. The assessment of populations of soil inhabiting invertebrates.

Agriculture Ecosystem and Environment 34: 45-176.

Fisher BL. 2004. Diversity patterns of ants (Hymenoptera: Formicidae) along an elevational gradient on Monts Doudou in Southwestern Gabon. *California Academy of Sciences Memoir* 28: 69-286.

Fisher BL. 1999. Improving inventory efficiency: A case study of leaf-litter ant diversity in Madagascar. *Ecological Applications* 9: 14-731.

Fisher BL. 1998. Ant diversity patterns along an elevational gradient in the Réserve Spéciale d’Anjanaharibe-Sud and on the western Masoala Peninsula, Madagascar. *Fieldiana: Zoology* 90: 9-67.

Fisher BL, Robertson HG. 2002. Comparison and origin of forest and grassland ant assemblages in the high plateau of Madagascar (Hymenoptera: Formicidae). *Biotropica* 34: 155-167.

Frith D, Frith C. 1990. Seasonality of litter invertebrate populations in an Australian upland tropical rain forest. *Biotropica* 22: 181-190.

Greenslade PJM. 1964. Pitfall trapping as a method for studying populations of Carabidae (Coleoptera). *Journal of Animal Ecology* 33: 301-310.

Halsall NB, Wratten SD. 1988. The efficiency of pitfall trapping for polyphagous predatory Carabidae. *Ecological Entomology* 13: 293-299.

Hammond P. 1990. Insect abundance and diversity in the Dumoga-Bone National Park, N. Sulawesi, with special reference to the beetle fauna of lowland rain forest in the Toraut region. In: Knight WJ, Holloway JD,

editors. *Insects and the Rain Forests of South East Asia (Wallacea)*. pp 197-254. Royal Entomological Society of London.

Hansen JE, New TR. 2005. Use of barrier pitfall traps to enhance inventory surveys of epigeic Coleoptera. *Journal of Insect Conservation* 9: 131-136.

Holland JM, Reynolds CJM. 2005. The influence of emptying frequency of pitfall traps on the capture of epigeal invertebrates, especially *Pterostichus madidus* (Coleoptera: Carabidae). *British Journal of Entomology and Natural History* 18:259-263.

Holland JM, Smith S. 1999. Sampling epigeal arthropods: An evaluation of fenced pitfall traps using mark-release-recapture and comparisons to unfenced pitfall traps in arable crops. *Entomologia Experimentalis et Applicata* 91: 347-357.

King JR, Porter SD. 2005. Evaluation of sampling methods and species richness estimators for ants in upland ecosystems in Florida. *Environmental Entomology* 34: 1566-1578.

Krell FT, Chungb AYC, DeBoisea E, Eggletona P, Giustia A, Inwarda K, Krell-Westerwalbesloha S. 2005. Quantitative extraction of macro-invertebrates from temperate and tropical leaf litter and soil: Efficiency and time-dependent taxonomic biases of the Winkler extraction. *Pedobiologia* 49: 175-186.

Lasebikan BA, Belfield W, Gibson NHE. 1978. Comparison of relative efficiency of methods for the extraction of soil microarthropods. *Revue d'Ecologie et de Biologie du Sol* 15: 39-65.

Leather SR, Watt AD. 2005. Sampling theory and practice. In: Leather R, editor. *Insect*

sampling in forest ecosystems. pp 1-15. Blackwell publishing.

Leponce M, Theunis L, Delabie JHC, Roisin Y. 2004. Scale dependence of diversity measures in a leaf-litter ant assemblage. *Ecography* 27: 253-267.

Longino JT, Coddington J, Colwell RK. 2002. The ant fauna of a tropical rain forest: Estimating species richness three different ways. *Ecology* 83: 689-702.

Majer JD. 1997. The use of pitfall traps for sampling ants. *Na critique Mem Mus Victoria* 56: 323-329.

Melbourne BA. 1999. Bias in the effect of habitat structure on pitfall traps: An experimental evaluation. *Australian Journal of Ecology* 24: 228-239.

Mesibov R, Taylor RJ, Brereton RN. 1995. Relative efficiency of pitfall trapping and hand-collecting from plots for sampling of millipedes. *Biodiversity and Conservation* 4: 429-439.

Mommertz S, Schauer C, Kösters N, Lang A, Filser J. 1996. A comparison of D-vac suction, fenced and unfenced pitfall trap sampling of epigeal arthropods in agro-ecosystems. *Annales Zoologici Fennici*. 33: 117-124

Nadkarni NM, Longino JT. 1990. Invertebrates in canopy and ground organic matter in a neotropical montane forest, Costa Rica. *Biotropica* 22: 286-289.

Oliver I, Beattie AJ. 1996. Designing a cost-effective invertebrate survey: A test of methods for rapid assessment of biodiversity. *Ecological Applications* 6: 594-607.

Olson DM. 1991. A comparison of the efficacy of litter sifting and pitfall traps for sampling leaf litter ants (Hymenoptera, Formicidae) in a tropical wet forest, Costa Rica. *Biotropica* 23: 166-172.

Orris JB. 2005. MegaStat Version 10.0. <http://www.mhhe.com/support>. Distributed by Mc Graw-Hill.

Palacios-Vargas JG, Castañõ-Meneses G, Gómez-Anaya JA, Martínez-Yrizar A, Mejía-Recamier BE, Martínez-Sánchez J. 2007. Litter and soil arthropods diversity and density in a tropical dry forest ecosystem in Western Mexico. *Biodiversity and Conservation* 16: 3703-3717.

Parr CL, Chown SL. 2001. Inventory and bioindicator sampling: Testing pitfall and Winkler methods with ants in a South African savanna. *Journal of Insect Conservation* 5: 27-36.

Philpott SM, Bichier P, Rice RA, Greenberg R. 2007. Biodiversity conservation, yield, and alternative products in coffee agroecosystems in Sumatra, Indonesia. *Conservation Biology* 21(4): 975-985.

Prasifka1 JR, Lopez1 MD, Hellmich1 RL, Lewis LC, Dively GP. 2007. Comparison of pitfall traps and litter bags for sampling ground-dwelling arthropods. *Journal of Applied Entomology* 131:115-120.

Richardson BA, Richardson MJ, Soto-adames FN. 2005. Separating the effects of forest type and elevation on the diversity of litter invertebrate communities in a humid tropical forest in Puerto Rico. *Journal of Animal Ecology* 74: 926-936.

Robertson HG. 2007. Comparison of leaf litter ant communities in woodlands, lowland forests and montane forests of north-eastern

Tanzania. *Biodiversity and Conservation* 11: 1637-1652.

Schillhammer H. 2001. Instruction manual on collecting, preserving and preparing Coleoptera (Coleoptera). Natural History Museum, Vienna, Australia.

Snyder BA, Draney ML, Sierwald P. 2006. Development of an optimal sampling protocol for millipedes (Diplopoda). *Journal of Insect Conservation* 10: 277-288.

Southwood TRE. 1978. *Ecological methods, with particular reference to the study of insect populations, 2nd edition*. Chapman and Hall.

Spence JR, Niemela JK. 1994. Sampling carabids assemblages with pitfall traps: The madness and the method. *Canadian Entomologist* 126: 881-894.

Standen V. 2000. The adequacy of collecting methods for estimating species richness of grassland invertebrates. *Journal of Applied Ecology* 37: 884-893.

Stoyan D, Kuschka V. 2001. On animal abundance by estimation based on pitfall traps. *Biometrical Journal* 43: 45-52.

Topping CJ, Sunderland KD. 1992. Limitations to the use of pitfall traps in ecological studies exemplified by a study of spiders in a field of winter wheat. *Journal of Applied Ecology* 29: 485-491.

Underwood EC, Fisher BL. 2006. The role of ants in conservation monitoring: If, when, and how. *Biological Conservation* 132: 166-182.

Vineesh PJ. 2007. *Ecology and diversity of Entomofauna in the litter stands of monoculture and natural forests in Kannur district*. Ph.D. dissertation. University of Calicut, Kerala

Vineesh PJ, Thomas SK, Karmaly KA. 2007. Community structure and functional group classification of litter ants in the montane evergreen and deciduous forests of Wayanad region of Western Ghats, Southern India. *Oriental Insects* 41: 185-200.

Ward PS. 1987. Distribution of the introduced Argentine ant (*Iridomyrmex humilis*) in natural habitats of the lower Sacramento Valley and its effects on the indigenous ant fauna. *Hilgardia* 55: 1–16.

Ward DF, New TR, Yen AL. 2001. Effects of pitfall trap spacing on the abundance, richness and composition of invertebrate catches. *Journal of Insect Conservation* 5: 47-53.

Weiss NA. 2007. Dorling Kindersley (India) Pvt. Ltd., licensees of Pearson Education in South Asia.

Wheeler QD, McHugh JV. 1987. A portable and onvertiblez "Moczarski/Tullgren" extractor for fungus and litter Coleoptera. *Coleopterists Bulletin* 41: 9-12.

Wikramanayake E, Dinerstein E, Loucks CJ, Olson DM, Morrison J, Lamoreux J, McKnight M, Hedao P. 2002. *Terrestrial ecoregions of the Indo-Pacific: A conservation assessment*. Island Press.

Woodcock BA. 2005. Sampling theory and practice. In: Leather SR, editor. *Insect sampling in forest ecosystems*. pp. 37–57. Blackwell.

Work TT, Buddle CM, Korinus LM, Spence JR. 2002. Pitfall trap size and capture of three taxa of litter-dwelling arthropods: Implications for biodiversity studies. *Environmental Entomology* 31: 438-448.