



Molecular Phylogenetic Analysis of the *Stegana ornatipes* Species Group (Diptera: Drosophilidae) in China, with Description of a New Species

Authors: Lu, Jin-Ming, Li, Tong, and Chen, Hong-Wei

Source: Journal of Insect Science, 11(20) : 1-12

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.011.0120>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Molecular phylogenetic analysis of the *Stegana ornatipes* species group (Diptera: Drosophilidae) in China, with description of a new species

Jin-Ming Lu^{1a}, Tong Li^{2,b} and Hong-Wei Chen^{1c*}

¹Department of Entomology, South China Agricultural University, Tianhe, Guangzhou, 510642 China

²Key Laboratory of Zoological Systematics and Evolution, Institute of Zoology, Chinese Academy of Sciences, Datunlu, Chaoyang, Beijing, 100101 China

Abstract

A new species of the *Stegana (Steganina) ornatipes* species group (Diptera: Drosophilidae) is described from Hainan, China, *S. (S.) xipengi* sp. nov. Based on the mitochondrial *ND2* and *COI* gene sequences, the relationships among eight species from mainland China of the *ornatipes* group, and their relationships to the *undulata*, *nigrolimbata* and *shirozui* species groups of the same subgenus, are investigated, using two species of the subgenus *Stegana*, *S. emeiensis* and *S. quadrata*, as outgroups. The result shows that *S. (S.) mengla* is debarred from the *ornatipes* group.

Keywords: Drosophilidae, molecular phylogeny, mitochondrial DNA, *Stegana ornatipes* group, Oriental region
Correspondence: ^a fabregas_l@hotmail.com, ^b holy518125@126.com, ^{c*} hongweic@scau.edu.cn, * Corresponding author

Editor: Zhijian Jake Tu was editor of this paper

Received: 17 December 2009, **Accepted:** 28 September 2010

Copyright : This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 11, Number 20

Cite this paper as:

Lu JM, Li T, Chen HW. 2011. Molecular phylogenetic analysis of the *Stegana ornatipes* species group (Diptera: Drosophilidae) in China, with description of a new species. *Journal of Insect Science* 11:20 available online: insectscience.org/11.20

Introduction

So far five species groups have been identified in the subgenus *Steganina* Wheeler (Diptera: Drosophilidae) of the genus *Stegana* Meigen: *coleoprata* group (Laštovka and Máca 1982; Chen and Chen 2008), *nigrolimbata* group (Sidorenko 2002; Cao and Chen 2008), *shirozui* group (Chen et al. 2009), *undulata* group (Sidorenko 2002) and *ornatipes* group (Cheng et al. 2009), and they included 51 species; most of them were from the Oriental region except for some species of the *coleoprata* group from the Palearctic region. The *ornatipes* group includes ten species from the Oriental region: *S. (S.) vietnamensis* Sidorenko, 1997 from Vietnam; *S. (S.) albiventralis* Cheng, Gao et Chen, 2009; *S. (S.) angusigena* Cheng, Gao et Chen, 2009; *S. (S.) chitouensis* Sidorenko, 1998; *S. (S.) lingnanensis* Cheng, Gao et Chen, 2009; *S. (S.) mengla* Cheng, Gao et Chen, 2009; *S. (S.) nulliseta* Cheng, Gao et Chen, 2009; *S. (S.) ornatipes* Wheeler et Takada, 1964; *S. (S.) pilosella* Cheng, Gao et Chen, 2009 and *S. (S.) zhaofengi* Cheng, Gao et Chen, 2009 from China. This group is supported by the following morphological characters as the diagnosis: surstylus large, with a strong prensiseta apically and several thin, long setae; 10th sternite mostly narrowed, nearly arcuate, with a pair of projections posterolaterally; gonopods with a pair of projections sublaterally. On the other hand, this group is similar to the *nigrolimbata* group in sharing the following morphological characters: palpus mostly black, sometimes yellow basally; gena yellow to brown, narrow ($ch/o \leq 0.10$); aedeagus basally contiguous to aedeagal apodeme; which shows the both are more closely related to each other than other members of the subgenus *Steganina*.

Recently, some studies of molecular phylogeny were appeared to the subfamily Steganinae (Otranto et al. 2008; He et al. 2009a, b; Zhao et al. 2009; Li et al. 2010). Otranto et al. (2008) reconstructed the phylogenetic relationships among 13 species of 8 genera of Steganinae based on the DNA sequences of the cytochrome oxidase subunit I (*COI*) gene, however, in their phylogenetic analysis, only two *Stegana* species were sampled as the representative. Li et al. (2010) investigated the phylogenetic relationships among seven of the Chinese species of the subgenus *Stegana* (s.s.) based on the DNA sequences of the NADH dehydrogenase subunit 2 (*ND2*) gene, using two species of the subgenus *Steganina* (*S. nigrolimbata* Duda, 1924 and *S. ctenaria* Nishiharu, 1979) as outgroup taxa.

In the present study, we described a new species of the *ornatipes* group from Hainan, China. We also constructed the molecular phylogeny based on the mtDNA sequences of *ND2* and *COI* genes. To investigate the relationships in *ornatipes* group and with the other species groups of subgenus *Steganina*, we employed the additional seven species from mainland China of this group and *S. nigrolimbata* Duda, *S. xiaoleiae* Cao and Chen, *S. ctenaria* Nishiharu, and *S. undulata* de Meijere which belong to *nigrolimbata*, *shirozui* and *undulata* species groups of the subgenus *Steganina* as ingroup taxa. Two species from subgenus *Stegana*, *S. emeiensis* Sidorenko and *S. quadrata* Cao and Chen were chosen as outgroup taxa.

Materials and Methods

All materials were collected on tussock and tree trunks along streams in forest, preserved in 75% ethanol immediately and identified

(Table 1). A small piece of tissue was removed from the fly abdomen and used for the DNA extraction; then, the body and terminalia parts were dried and deposited in the Department of Entomology, South China Agricultural University, Guangzhou, China (SCAU). McAlpine (1981) was followed for morphological terminology and Zhang and Toda (1992), and Chen and Toda (2001) for the definitions of measurements, indices and abbreviations.

DNA extraction and sequencing

The total DNA was extracted using the DNA extraction Kit (TIANGEN®) according to the manufacture's protocol. The *ND2* gene and the 5' end of *COI* gene were amplified. Primers used were given in table 2. The PCR cycle program comprised an initial 3 min of predenaturation at 94 °C, 35 cycles of amplification (50 s of denaturation at 94 °C; 1 min of annealing at 53 °C for *ND2*, 49 °C for *COI*; 1 min of extension at 72 °C), and a final elongation for 5 min at 72 °C. When possible, purified amplified products were directly run on an ABI 3730 sequencer for sequencing, otherwise they were cloned into the pMD18-T plasmid vector (TAKARA®), and then sequenced. The related *ND2* sequences of *S. emeiensis*, *S. quadrata*, *S. ctenaria* and *S. nigrolimbata* were retrieved from the

National Center for Biotechnology Information (NCBI); the related *COI* sequences of *emeiensis*, *S. nigrolimbata* and *S. undulata* were also retrieved from the NCBI.

Phylogenetic analyses

The sequences were aligned by the Clustal W (Thompson et al. 1994) method implemented in program MEGA 4.0 (Tamura et al. 2007) with default options. A partition homogeneity test (PHT) between the *ND2* and *COI* sequences was performed with PAUP 4.0b10* (Swofford 2002). The program DAMBE 5.0.80 (Xia and Xie 2001) was used to measure the nucleotide substitution saturation using the method of Xia et al. (2003) as the substitution saturation masked the phylogenetic signal (Lopez et al. 1999; Philippe and Froterre 1999). Base compositions were investigated by means of the software PAUP 4.0b10* (Swofford 2002), and a χ^2 test was also used to test the nucleotide composition homogeneity. Uncorrected pairwise divergence was estimated by program MEGA 4.0 (Tamura et al. 2007).

Phylogenetic trees were constructed by using the maximum parsimony (MP) and maximum likelihood (ML) in PAUP 4.0b10* (Swofford 2002), the Bayesian inferring (BI) method

Table 1. Collection data of samples for DNA sequencing, and accession numbers of the *ND2* and *COI* sequences.

Subgenera	Groups	Species	Species numbers in SCAU	Localities	Latitude and longitude	Accession numbers of <i>ND2</i>	Accession numbers of <i>COI</i>
Stegana		<i>emeiensis</i>	I20268	Menglu, Yunnan, China	21°41'N, 101°25'E	EU805515	HM636455
		<i>quadrata</i>	I20084	Mengla, Yunnan, China	21°28'N, 101°38'E	EU805516	HQ270147
Steganina	<i>nigrolimbata</i>	<i>nigrolimbata</i>	I20390	Guangzhou, Guangdong, China	23°10'N, 112°34'E	EU805513	HM636458
		<i>xiaoleiae</i>	I20453	Maoershan, Guangxi, China	25°51'N, 110°27'E	GQ259982	HQ260633
	<i>shirozui</i>	<i>ctenaria</i>	I20321	Kumamoto, Japan	32°37'N, 130°51'E	EU805514	HQ270148
	<i>undulata</i>	<i>undulata</i>	I20175	Mengla, Yunnan, China	Ditto	GQ249196	HM646466
	<i>ornatipes</i>	<i>albiventralis</i>	I20610	Jingdong, Yunnan, China	24°32'N, 101°01'E	GQ259983	HQ260634
		<i>angustigena</i> -GX	I20627	Nonggang, Guangxi, China	22°22'N, 106°51'E	GQ259984	HQ260637
		<i>angustigena</i> -HN	I20626	Jianfengling, Hainan, China	18°41'N, 108°52'E	GQ259985	HQ260635
		<i>angustigena</i> -YN	I20634	Mengla, Yunnan, China	Ditto	GQ259986	HQ260636
		<i>lingnanensis</i>	I20636	Guangzhou, Guangdong, China	Ditto	GQ259987	HQ260638
		<i>pilosella</i>	I20642	Mengla, Yunnan, China	Ditto	GQ259988	HQ260640
		<i>nulliseta</i>	I20656	Mengla, Yunnan, China	Ditto	GQ259991	HQ260639
		<i>xipengi</i> sp. nov.	I20691	Jianfengling, Hainan, China	Ditto	GQ259992	HQ260641
		<i>zhaofengi</i> -YN1	I20459	Mengla, Yunnan, China	Ditto	GQ259993	HQ260643
		<i>zhaofengi</i> -YN2	I20464	Mengla, Yunnan, China	Ditto	GQ259994	HQ260642
		<i>zhaofengi</i> -YN3	I20665	Mengla, Yunnan, China	Ditto	GQ259995	HQ260644
	?	<i>mengla</i> -YN1	I20653	Mengla, Yunnan, China	Ditto	GQ259989	HQ260645
		<i>mengla</i> -YN2	I20652	Mengla, Yunnan, China	Ditto	GQ259990	HQ260646

Table 2. Primers used for PCR and sequencing.

Target gene	Primer	Primer sequence (5'-3')	Reference
ND2	ND2-H	AAGCTACTGGGTTTCATACC	Park 1999
	ND2-T3	AGGCGATAGATTGTAAATC	Li 2010
COI	COI-F1	CGCCTAAACTTCAGCCACTT	He et al. 2009a
	LCO1490	GGTCAACAAATCATAAAGATATTGG	Folmer et al. 1994
	HCO2198	TAAACTTCAGGGTGACCAAAAAATCA	Folmer et al. 1994

performed in MrBayes 3.2.1 (Huelsenbeck and Ronquist 2001; Ronquist and Huelsenbeck 2003). The MP and ML trees were searched by the heuristic method, with initial trees obtained by randomly adding taxa, and the TBR algorithm was used in branching swapping. Branch support for each node in the MP and ML trees was assessed by 1000 bootstrap replicates. The nucleotide substitution models of ML and BI analyses were selected by MrModeltest 2.3 (Nylander 2004) using the hierarchical likelihood ratio test (hLRT) criterion (Posada and Crandall 1998). In the BI analyses, the site-specific models were assigned to dataset partitioned by locus (2 data partitions) and by codon positions (6 data partitions). Two independent runs with 2,000,000 generations were implemented in parallel, sampling frequency of every 100 generations was employed. When the average deviation of split frequencies fell well below 0.01, the two runs were stopped. For each running, the 5,000 early-phase samples were burn-in, the rest samples were used in summarizing and a majority rule tree showing all the compatible partitions was obtained.

Partition Bremer support (PBS) was used to show the contribution of each gene partition to the Bremer support of the simultaneous analysis (Baker and DeSalle 1997). Values can be positive, negative or zero and sum of all the partitioned Bremer support values at a node will equal the Bremer support value for that node. A positive PBS value suggests support for the node by that gene, whereas a negative PBS value indicates that the partition

lends conflict to a given node, and zero indicate that the partition lends neither support nor conflict to a given node. The partitioned Bremer support values were calculated using the partitioned constraint file in TreeRot v3 (Sorenson 1999).

Results


Stegana (Steganina) xipengi sp. nov.
(Figures 1, 2)

Diagnosis

This species is related to *S. (S.) albiventralis* from Yunnan in having the entirety white katepisternum, but clearly distinguishable from it by the palpus yellow basally, black distally, the mesonotum brown, without stripe (in *albiventralis*: palpus entirely yellow; mesonotum brown, with yellow stripe medially).

Description

Male: Frons and face not rectangular in profile. Eyes red. Ocellar triangle black, with 1 pair of small setae above ocellar setae.



Figures 1-2. *Stegana (Steganina) xipengi* sp. nov., ♂: 1. Epandrium, cercus and surstylus (lateral view); 2. surstylus (frontal view). Scale bars = 0.1 mm. High quality figures are available online.

Postvertical setae slightly behind vertex ridge. Frons shiny, brown, with sporadic, minute setulae submedially, and a black, transverse band above ptilinal fissure. Proclinate orbital setae slightly nearer to ptilinal fissure than to inner vertical setae. Pedicel brown; first flagellomere yellow only basally, mostly black. Face black with yellow, transverse band medially, broadened ventrally; facial carina absent. Clypeus black medially, yellow laterally. Palpus yellow basally, black distally, with 1-2 longer setae distally and several shorter setae basally. Gena yellow, narrow. Vibrissa prominent; other orals small. Occiput glossy, yellow, but black around occipital foramen. Mesonotum brown. Mesopleuron with a black longitudinal stripe above (running from propleuron to base of halter). Postpronotal lobe brown on upper part, white on lower part, with 1 long and a few short setae. Acrostichal setulae approximately in 10 irregular rows. Prescutellar setae 1 pair. Katepisternum entirely white. Scutellum brown; basal setae divergent; apical setae crossing with each other. Wing dark brown anteriorly, pale posteriorly, curved downward on distal part. Basal medial-cubital crossvein present. C_1 with 2 isometric setae. Costal vein with 9 minute spinules on ventral surface between veins R_{2+3} and R_{4+5} . Vein R_{2+3} obviously curved to costa at tip; Veins R_{4+5} and M_1 convergent distally. Halteres white basally, greyish brown distally. Legs whitish yellow, brown on apical part of fore femur, and fore and hind tarsomeres, dark brown to black on medially on mid and hind femora, with 2 dark brown rings on fore and mid tibiae. Fore femur with 3-4 setae on distal part of ventral surface. Apical seta present on mid tibia. Preapical dorsal setae present on all tibiae. Mid tibia (misused to mid tarsus in Cao and Chen 2008; Cheng et al. 2009) with 5 strong setae on basal part of dorsal surface. Mid and hind tarsomeres with 2 and 1 row(s)

of minute cuneiform setulae on ventral surface, respectively; fore and hind 1st tarsomeres slightly shorter than the rest combined; mid 1st tarsomere longer than the rest combined. Abdominal all tergites dark brown. Sternites brown; 3rd to 5th broadened; 6th covered with 5th. Epandrium pubescent except for anteroventral margins, with approximately 21 setae near posterior margin on each side (Figure 1). Cercus separated from epandrium, setigerous, lacking pubescence (Figure 1). Surstylus separated from epandrium, with several thin, long setae on inner margin and surface (Figure 2), apically strongly curved and with 1 strong prensiseta (Figure 2). The hypandrium, gonopods, aedeagus and aedeagal apodeme were lost when clearing them in KOH solution.

Measurements

BL = 2.76 mm in holotype; ThL = 1.32 mm; WL = 2.58 mm; WW = 1.12 mm. Indices: arb = 8/7, avd = 0.83, adf = 1.20, flw = 1.80, FW/HW = 0.36, ch/o = 0.08, prorlb = 1.16, rcorb = 0.82, vb = 0.30, dcl = 0.40, presctl = 0.60, sctl = 1.80, sterno = 0.90, orbito = 2.20, dcp = 0.20, setlp = 1.00, C = 1.86, 4c = 1.22, 4v = 1.74, 5x = 1.40, ac = 9.33, M = 0.61, C3F = 0.66.

Type

Holotype: ♂ (SCAU, No. 120589), CHINA: Jianfeng, Ledong, Hainan, 18°41'N, 108°52'E, alt. 750 m, 14.iv.2008, *ex* tussock, X.P. Chen.

Etymology

Patronym of the collector Xipeng Chen (SCAU).

Distribution

China (Hainan).

Table 3. Results of model selection, composition homogeneity test and test of substitution saturation.

	ND2				COI			
	1st-CP*	2nd-CP	3rd-CP	All	1st-CP	2nd-CP	3rd-CP	All
Model selection								
Model selected	GTR+G	HKY+G	HKY+G	GTR+I+G	SYM+G	F81	HKY+I+G	GTR+I+G
Proportion of invariable sites (I)	0	0	0	0.297	0	0	0.113	0.6104
Gamma distribution shape parameter	0.2678	0.1055	0.5701	0.4971	0.1581	-	0.534	1.359
Composition homogeneity test								
df	54	54	54	54	54	54	54	54
χ^2	13.00	2.67	44.86	20.45	3.67	0.32	31.94	7.36
P	1.0000	1.0000	0.8075	1.0000	1.0000	1.0000	0.9927	1.0000
Test of substitution saturation								
<i>I_{ss}</i>	0.1484	0.0783	0.3847	0.1742	0.0585	0.0093	0.4286	0.3728
<i>I_{ss.c}</i> (for an asymmetrical tree)	0.6935	0.6935	0.6935	0.7605	0.6793	0.6793	0.6793	0.7347
P	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
<i>I_{ss.c}</i> (for an extreme asymmetrical tree)	0.4575	0.4575	0.4575	0.5479	0.4390	0.4390	0.4390	0.5123
P	0.0000	0.0000	0.0031	0.0000	0.0000	0.0000	0.7226	0.0000

* CP = codon position;

Molecular analysis

Data set analysis

The alignment was 1739 nucleotide positions (1029 for *ND2* and 710 for *COI*, respectively) in length. There were end gaps in the *ND2* sequence of *S. undulata* (sites 1-22) and in the *COI* sequence of *S. xiaoleiae* (sites 1-33). The base composition of *ND2* and *COI* were generally AT rich with a mean of 83% and 69%, respectively. It contained high AT contents in the 3rd (94.2% and 94.2%, respectively) codon positions. Performance of the Chi-square test was showed in table 3. It yielded a homogeneous base composition in the *ND2*-alignments and *COI*-alignments or in the separate condon positions of the two mitochondrial genes.

The test of substitution saturation showed that the observed index of substitution saturation (*I_{ss}*) for *ND2*-alignments or for *COI*-alignments was significantly lower than the corresponding critical index substitution saturation (*I_{ss.c}*), indicating that there was little saturation in our sequences. However, when considering partitions separated by codon, we identified substitution saturation in the third codon position of the *COI*-

alignments [*I_{ss}* = 0.4286 < *I_{ss.c}* = 0.4390 (for an extreme asymmetrical tree, $p = 0.72$)] (Table 3). Since none of the resulted trees of the present study are extremely asymmetric, there should be little substitution saturation in our sequence.

Table 4 shows the uncorrected pairwise p-distances for the *ND2* and *COI* sequences. The genetic divergence of *ND2* sequences of species within the *ornatipes* group ranged from 5.28% to 14.24%, and genetic divergence of *COI* sequences ranged from 4.28% to 10.49%, however, when we took no account of the *S. mengla*, the upper limits would declined to 8.37% and 8.42% for *ND2* and *COI*, respectively. Within the *ornatipes* group, divergences between *S. mengla* and other species ranged from 9.36% to 15.64% for *ND2* and from 8.57% to 10.93% for *COI*, whereas the genetic variance among groups ranged from 11.65% to 14.34% for *ND2* and from 8.57% to 11.23% for *COI*.

Phylogenetic analysis

The PHT resulted in a p value of 0.062, indicating that no significant incongruence was found between the *ND2* and *COI* data sets. The best-fit models selected for the ML

Table 4. Uncorrected pairwise p-distance among the *ND2* and *COI* sequences of the *ornatipes* species group.


		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19
1	<i>S. emeiensis</i>		0.0798	0.1241	0.1270	0.1064	0.1196	0.1167	0.1182	0.1182	0.1182	0.1152	0.1137	0.1329	0.1137	0.1137	0.1137	0.1152	0.1241	0.1241
2	<i>S. quadrata</i>	0.0747		0.1108	0.1137	0.1064	0.1196	0.1211	0.1270	0.1241	0.1241	0.1196	0.1226	0.1241	0.1211	0.1285	0.1285	0.1329	0.1078	0.1078
3	<i>S. nigrolimbata</i>	0.1464	0.1365		0.0768	0.1064	0.0931	0.0857	0.0916	0.0886	0.0886	0.0857	0.0916	0.0871	0.0916	0.0916	0.0916	0.0945	0.1108	0.1108
4	<i>S. xiaoleiae</i>	0.1554	0.1404	0.0697		0.1123	0.1108	0.1004	0.0975	0.0975	0.0975	0.0931	0.0975	0.0901	0.0901	0.0990	0.0990	0.1019	0.1064	0.1064
5	<i>S. ctenaria</i>	0.1434	0.1255	0.1245	0.1345		0.0960	0.1064	0.1049	0.1019	0.1019	0.1004	0.1093	0.1078	0.0975	0.1064	0.1064	0.1078	0.1049	0.1049
6	<i>S. undulata</i>	0.1663	0.1514	0.1404	0.1434	0.1165		0.0975	0.1004	0.1004	0.1004	0.0916	0.0990	0.0975	0.0931	0.1019	0.1019	0.1049	0.1093	0.1093
7	<i>S. albiventralis</i>	0.1534	0.1375	0.1076	0.1106	0.1255	0.1534		0.0458	0.0428	0.0428	0.7390	0.0591	0.0561	0.0635	0.0576	0.0576	0.0606	0.1049	0.1049
8	<i>S. angustigena</i> -HN	0.1564	0.1384	0.1006	0.0996	0.1325	0.1414	0.0677		0.0044	0.0044	0.0665	0.0620	0.0502	0.0473	0.0650	0.0650	0.0694	0.1049	0.1049
9	<i>S. angustigena</i> -YN	0.1573	0.1394	0.1056	0.1006	0.1345	0.1424	0.0677	0.0120		0.0000	0.0694	0.0620	0.0532	0.0502	0.0635	0.0635	0.0679	0.1034	0.1034
10	<i>S. angustigena</i> -GX	0.1584	0.1404	0.1066	0.1016	0.1355	0.1434	0.0687	0.0129	0.0010		0.0694	0.0620	0.0532	0.0502	0.0635	0.0635	0.0679	0.1034	0.1034
11	<i>S. lingnanensis</i>	0.1484	0.1295	0.1046	0.1036	0.1235	0.1504	0.0697	0.0568	0.0608	0.0618		0.0827	0.0753	0.0606	0.0857	0.0857	0.0871	0.0945	0.0945
12	<i>S. nulliseta</i>	0.1514	0.1315	0.1096	0.1096	0.1335	0.1544	0.0837	0.0767	0.0807	0.0817	0.0777		0.0724	0.0709	0.0620	0.0620	0.0679	0.0975	0.0975
13	<i>S. pilosella</i>	0.1534	0.1394	0.1036	0.1125	0.1315	0.1544	0.0807	0.0578	0.0598	0.0608	0.0667	0.0807		0.0591	0.0679	0.0679	0.0739	0.1049	0.1049
14	<i>S. xipengi</i>	0.1554	0.1325	0.1026	0.1076	0.1265	0.1504	0.0677	0.0767	0.0548	0.0538	0.0588	0.0817	0.0598		0.0798	0.0798	0.0842	0.1034	0.1034
15	<i>S. zhaofengi</i> -YN1	0.1414	0.1285	0.0946	0.1066	0.1245	0.1544	0.0737	0.0687	0.0697	0.0707	0.0647	0.0677	0.0717	0.0697		0.0000	0.0059	0.1049	0.1034
16	<i>S. zhaofengi</i> -YN2	0.1444	0.1295	0.0936	0.1076	0.1255	0.1564	0.0747	0.0528	0.0707	0.0717	0.0657	0.0687	0.0737	0.0707	0.0050		0.0059	0.1049	0.1034
17	<i>S. zhaofengi</i> -YN3	0.1404	0.1275	0.0936	0.1056	0.1235	0.1544	0.0727	0.0667	0.0687	0.0697	0.0637	0.0667	0.0707	0.0687	0.0040	0.0010		0.1049	0.1034
18	<i>S. mengla</i> -YN1	0.1454	0.1245	0.1355	0.1365	0.1175	0.1375	0.1404	0.1384	0.1384	0.1394	0.1375	0.1434	0.1424	0.1315	0.1345	0.1355	0.1345		0.0015
19	<i>S. mengla</i> -YN2	0.1454	0.1245	0.1355	0.1365	0.1175	0.1375	0.1404	0.1384	0.1384	0.1394	0.1375	0.1434	0.1424	0.1315	0.1345	0.1355	0.1345	0.0000	

The matrix in the lower left was the uncorrected pairwise p-distance among the *ND2* sequences; the matrix in the upper right was the uncorrected pairwise p-distance among the *COI* sequences.

reconstruction and Bayesian inference were listed in table 3.

The relationships within the *ornatipes* group were not stable revealed by different tree-building methods as the low supports for the basal nodes (Figures 3, 4 and 5), but it was surprising that *S. mengla* was debarred from

the *ornatipes* group in all trees, and it was placed at the most basal clade of subgenus *Steganina* receiving great supports (MP BP, or bootstrap percentages of the MP analysis = 100; ML BP = 100; PP or posterior probability of the 2-/6-partition Bayesian inferring = 1.00/1.00). The remaining species of the *ornatipes* group were recovered as a




monophyletic group with robust supports in all trees (MP BP = 100; ML BP = 100; PP = 1.00/1.00 in the 2-/6-partition Bayesian analyses, respectively). The *nigrolimbata* group appeared to be the closest relative to this monophyletic group with well supports (MP BP = 100; MLBP = 100; PP = 1.00/1.00) (Figures 3, 4 and 5). The Bayesian analysis yielded a general topology (Figure 3), which was mostly congruent with the result of the ML analysis (Figure 4). The monophyletic group diverged into two branches. One consist of *S. zhaofengi* triple and *S. nulliseta*, and the other further diverged into *S. albiventralls*, *S. lingnanensis*, *S. pilosella*, *S. xipengi* and *S. angustigenai* triple orderly in the Bayesian tree, whereas *S. pilosella* diverged prior to *S. lingnanensis* (ML BP = 41), leaving *S. lingnanensis* and *S. xipengi* as sister group in the ML reconstruction, but with a low support (ML BP = 32). The MP tree (Figure 5) differed from the ML and Bayesian tree at several points. It suggested a very basal position for *S. lingnanensis* in the *ornatipes*


group and *S. xipengi* clustered with *S. pilosella* which was consistent with the Bayesian analysis. The Yunnan (-YN), Guangxi (-GX) and Hainan (-HN) samples of *S. angustigena* clustered together with well support (MP BP = 100; ML BP = 100; PP = 1.00/1.00), and so did in the YN1, YN2 and YN3 samples of *S. zhaofengi* (Figures 3, 4 and 5).

To determine the relative contributions of the two data partition to the combined analysis tree, partition Bremer supports were calculated and given in table 5. Support for combined analysis phylogeny came from *ND2* was a little bit more than that from *COI*. Nodes 1, 3, 6, 7 and 12 showed a mixture of positive and negative PBS scores.

Discussion

The phylogenetic trees showed that the *ornatipes* group clearly appeared to be paraphyletic. To eliminate the effect of





individual difference, another sample of *S. mengla* was included in the analysis, but the situation did not change. In morphological viewpoints, the *S. mengla* holds the same diagnostic characters of the *ornatipes* group, which contradict with our molecular phylogeny as it formed a separate branch in the phylogenetic tree. The amount of genetic divergences between *S. mengla* and other species within the *ornatipes* group were high and overlapped to some extent with the divergence between species groups. Although speculative, the morphological convergence should be the reason for this situation. The convergent morphological evolution seems to be common in the subfamily *Steganinae* (Otranto et al. 2008), which is similar to the suggestion made in this research concerning convergent morphological evolution in *S. mengla*. Considering the closer relationship of *S. mengla* with the outgroup *S. emiensis* respect to the other species showed in the

phylogenetic tree, it is possible that *S. mengla* is the interim species of the divergent between subgenus *Steganina* and subgenus *Stegana*. Of course, this hypothesis should be proved with analysis of suitable species of both the subgenus *Steganina* and subgenus *Stegana*. Except the *S. mengla*, the branch consist of the rest species of the *ornatipes* group showed the closer relationship with the *nigrolimbata*

Table 5. Partition Bremer support for all nodes in the MP tree (Fig. 5).

Node	Bremer support	Individual gene partitions	
		<i>ND2</i>	<i>COI</i>
1	8	8.50	-0.50
2	22	15.00	7.00
3	1	1.50	-0.50
4	1	1.00	0.00
5	40	23.50	16.50
6	4	-1.90	5.90
7	1	-4.17	5.17
8	13	7.00	6.00
9	19	13.50	5.50
10	17	13.50	3.50
11	9	3.83	5.17
12	4	-2.50	6.50
13	70	43.17	26.83
Total PBS	209	121.93	87.07

group than other species groups of subgenus *Steganina* was consistent with the morphological affinity in the two groups (Cao and Chen 2008).

In general, the NADH dehydrogenase subunit genes are rapidly evolving, but the cytochrome oxidase subunit is more slowly evolving (Simon 1994). It was supposed that the *ND2* gene was better than the *COI* gene suited for species-level analysis, but the PBS analysis indicating that the contribution to the MP reconstruction in this research of the *ND2* gene was nearly the same as the *COI* gene. Our PBS analysis had implications for the conflicts of the two genes at some nodes (e.g., nodes 1, 3, 6, 7 and 12), suggesting that the two partitions data (*ND2* and *COI*) may be favoring an alternative tree topology. The relationships of these nodes should be viewed cautiously.

The genetic distances between Yunnan, Guangxi and Hainan samples of *S. angustigena* [p-distance of *ND2* = 0.0129 (-GX vs. -HN), 0.0120 (-HN vs. -YN), 0.001 (-GX vs. -YN); p-distance of *COI* = 0.0044 (-GX vs. -HN), 0.0044 (-HN vs. -YN), 0.0000 (-GX vs. -YN)] were among the mean intraspecific variability of Meier et al. 2008 ($1.3 \pm 1.6\%$) for Diptera. In addition, no essential morphological character was found to distinguish the specimens of these three samples, indicating that they should be taken as conspecific ones. It was the same as the case of the YN1, YN2 and YN3 samples of *S. zhaofengi*. The genetic data [p-distance of *ND2* = 0.0050 (-YN1 vs. -YN2), 0.0040 (-YN2 vs. -YN3), 0.0010 (-YN1 vs. -YN3); p-distance of *COI* = 0.0000 (-YN1 vs. -YN2), 0.0059 (-YN2 vs. -YN3), 0.0059 (-YN1 vs. -YN3)] also indicated the conspecific status of the three samples of *S. zhaofengi*.

Some relationships within the *ornatipes* group were not well resolved, especially the alternative placement of *S. lingnanensis*, *S. xipengi* and *S. pilosella*. Therefore, it may be worthy to increase either the genetic markers (such as nuclear markers) or the number of samples in the future phylogenetic analysis of the *ornatipes* group.

Acknowledgements

We thank Dr. JJ Gao (Yunnan University, China) for helping in fieldwork; Ms. Y Cheng for providing the figures 1-2. This work was supported by the National Natural Science Foundation of China (No. 30970396).

Editor's note: Paper copies of this article will be deposited in the following libraries. The date of publication is given in 'About the Journal' on the JIS website.

Universitaetsbibliothek Johann Christian Senckenberg, Frankfurt Germany; National Museum of Natural History, Paris, France; Field Museum of Natural History, Chicago, Illinois USA; University of Wisconsin, Madison, USA; University of Arizona, Tucson, Arizona USA; Smithsonian Institution Libraries, Washington D.C. USA; The Linnean Society, London, England.

References

- Baker RH and DeSalle R. 1997. Multiple sources of character information and the phylogeny of Hawaiian drosophilids. *Systematic Biology* 46: 654-673.
- Cao HZ and Chen HW. 2008. Revision of the *Stegana* (*Steganina*) *nigrolimbata* species group from the Oriental region (Diptera, Drosophilidae). *Zootaxa* 1848: 27-36.

- Chen XP and Chen HW. 2008. The *Stegana coleoptrata* species group (Diptera, Drosophilidae) from mainland China. *Zootaxa* 1891: 55-65.
- Chen XP, Gao JJ and Chen HW. 2009. The *Stegana shirozui* species group (Diptera, Drosophilidae). *Journal of Natural History* 43: 1909-1927.
- Chen HW and Toda MJ. 2001. A revision of the Asian and European species in the subgenus *Amiota* Loew (Diptera, Drosophilidae) and establishment of species-groups based on phylogenetic analysis. *Journal of Natural History* 35: 1517-1563.
- Cheng Y, Gao JJ and Chen HW. 2009: The *Stegana ornatipes* species group from the Oriental Region (Diptera, Drosophilidae). *Zootaxa* 2216: 37-48.
- Folmer O, Black M, Hoeh W, Lutz R and Vrijenhoek R. 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3: 294-299.
- He XF, Gao JJ, Cao HZ, Zhang XL and Chen HW. 2009a. Taxonomy and molecular phylogeny of the *Phortica hani* species complex (Diptera: Drosophilidae). *Zoological Journal of the Linnean Society* 157: 359-372.
- He XF, Jiang JJ, Cao HZ and Chen HW. 2009b. Taxonomy and molecular phylogeny of the *Amiota nagatai* species group (Diptera, Drosophilidae). *Zootaxa* 2193: 53-61.
- Huelsenbeck JP, Ronquist FR. 2001. MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* 17: 754-755.
- Laštovka P and Máca J. 1982. European and North American species of the genus *Stegana* (Diptera, Drosophilidae). *Annotations Zoologicae et Botanicae* 149: 1-38.
- Li T, Cao HZ, Gao JJ and Chen HW. 2010. A revision of the subgenus *Stegana* (s. str.) (Diptera, Drosophilidae) from mainland China. *Zoological Journal of the Linnean Society* 158: 726-739.
- Lopez P, Forterre P and Philippe H. 1999. The root of the tree of life in the light of the covarion model. *Journal Molecular Evolution* 49: 496-508.
- McAlpine JF. 1981: Morphology and terminology – adults. In J. F. McAlpine (ed.), *Manual of Nearctic Diptera*, 1: 9-64. Research Branch Agriculture Canada Monograph, 27. Research Branch, Agriculture Canada, Ottawa.
- Meier R, Zhang G and Ali F. 2008. The use of mean instead of smallest interspecific distances exaggerates the size of the “Barcoding Gap” and leads to misidentification. *Systematic Biology* 57: 809-813.
- Nylander J A A. 2004. MrModeltest v2. Program distributed by the author. Evolutionary Biology Centre, Uppsala University.
- Otranto D, Stevens JR, Testini G, Cantacessi C and Máca J. 2008. Molecular characterization and phylogenesis of Steganinae (Diptera, Drosophilidae) inferred by the mitochondrial cytochrome c oxidase subunit 1. *Medical and Veterinary Entomology* 22: 37-47.

- Park J. 1999. Molecular phylogenetic studies of the *Drosophila (Drosophila) virilis* section (Diptera, Drosophilidae). PhD. Thesis, Tokyo Metropolitan University.
- Philippe H and Forterre P. 1999. The rooting of the universal tree of life is not reliable. *Journal Molecular Evolution* 49: 509-523.
- Posada D and Crandall KP. 1998. Modeltest: testing the model of DNA substitution. *Bioinformatics* 14: 817-818.
- Ronquist F, Huelsenbeck JP. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572-1574.
- Sidorenko VS. 2002. Phylogeny of the tribe Steganini Hendel and some related taxa (Diptera, Drosophilidae). *Far Eastern Entomologist* 111: 1-20.
- Simon C, Frati F, Beckenbach A, Crespi B, Liu H and Flook P. 1994. Evolution, weighting and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved PCR primers. *Annals of the Entomological society of America* 87: 651-710.
- Sorenson MD. 1999. TreeRot, vesion 2. Boston University, Boston, Massachusetts.
- Swofford DL. 2002. PAUP*: Phylogenetic analysis using parsimony (*and other methods), Version 4. Sinauer Associates, Sunderland, MA.
- Tamura K, Dudley J, Nei M and Kumar S. 2007. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software28 version 4.0. *Molecular Biology and Evolution* 24: 1596-1599.
- Thompson JD, Higgins DG and Gibson TJ. 1994. Clustal W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position specific gap penalties and weight matrix choice. *Nucleic Acids Research* 22: 4673-4680.
- Xia XH and Xie ZH. 2001. DAMBE: Data analysis in molecular biology and evolution. *Journal of Heredity* 92: 371-373.
- Xia XH, Xie ZH, Salemi M, Chen L and Wang Y. 2003. An index of substitution saturation and its application. *Molecular Phylogenetics and Evolution* 26: 1-7.
- Zhang WX and Toda MJ. 1992. A new species-subgroup of the *Drosophila immigrans* species-group (Diptera, Drosophilidae), with description of two new species from China and Revision of Taxonomic Terminology. *Japan Journal Entomology* 60: 839-850.
- Zhao F, Gao JJ and Chen HW. 2009. Taxonomy and molecular phylogeny of the Asian Paraleucophenga Hendel (Diptera, Drosophilidae). *Zoological Journal of the Linnean Society* 155: 615-629.