

### The Semiaquatic Nematoceran Fly Assemblages of Three Wetland Habitats and Concordance with Plant Species Composition, a Case Study from Subalpine Fennoscandia

Author: Salmela, Jukka

Source: Journal of Insect Science, 11(35): 1-28

Published By: Entomological Society of America

URL: https://doi.org/10.1673/031.011.0135

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.





### The semiaquatic nematoceran fly assemblages of three wetland habitats and concordance with plant species composition, a case study from subalpine Fennoscandia

Jukka Salmela

Department of Biology, Zoological Museum, FI-20014 University of Turku, Finland

### Abstract

Semiaquatic flies (Diptera, Nematocera) are an ecologically important and species rich group of insects within the boreal and arctic biomes. Community structure, species richness and abundance of semiaquatic flies were studied in three habitat types (aapa mires, springs and headwater streams), totaling 19 study sites, within the subalpine ecoregion of northern boreal Finland. Concordance of semiaquatic fly species composition with plant assemblages (higher plants and mosses), and geographical and environmental distance matrices were also studied. The collected insect material consisted of 94 species and 9038 specimens. According to non-metric multidimensional scaling ordination (visual inspection), multi-response permutation procedure and analysis of similarity tests, fly assemblages of aapa mires were clearly different from those of springs and headwater streams, but no differences were found between spring and headwater stream assemblages. The cumulative number of species was highest in headwater streams. Alpha diversity varied within the habitat types but was generally highest among headwater streams. Semiaquatic fly communities of headwater streams were the most abundant (number of specimens) and their rank-abundance distributions were relatively skewed; assemblages of aapa mires were less abundant and rather even. Community composition of combined plant material (219 taxa), higher plants (116 taxa) and mosses (103 taxa) were all in concordance with the flies; the strongest matrix correlation was found between higher plants and flies (Mantel test). The influence of geographical distance of the study sites to species composition was statistically significant but rather weak; instead, much stronger concordance was noted with environmental variables (Mantel test). Plants, especially higher plants, may be potential surrogates for semiaquatic fly assemblage composition. However, more studies of community concordance in a larger geographic area and within one habitat type are needed.

Keywords: Lapland, Finland, Tarvantovaara wilderness area, aapa mires, springs, headwater streams, dipteran diversity Abbreviations: ANOSIM, analysis of similarity test; MRPP, multi-response permutation procedure; NMS, nonmetric multidimensional scaling Correspondence: jueesal@utu.fi Received: I February 2010, Accepted: 8 June 2010 Copyright : This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed. ISSN: 1536-2442 | Vol. 11, Number 35 Cite this paper as: Salmela J. 2011. The semiaquatic nematoceran fly assemblages of three wetland habitats and concordance with plant species composition, a case study from subalpine Fennoscandia. *Journal of Insect Science* 11:35 available online: insectscience.org/11.35

Journal of Insect Science | www.insectscience.org

#### Introduction

of Species composition terrestrial communities is a consequence of several hierarchical and interacting factors. The regional species pool is a result of climate, geology, historical events and evolutionary processes. Further, dispersal ability and habitat selection of the species affect the identity of the potential colonizers and interspecific interactions facilitate or inhibit the co-occurrence of the species in a given locality (Morin 1999). Assemblages (sensu Fauth et al. 1996) are phylogenetically related members of the biological community. Most often taxonomically related assemblages rather than whole communities are surveyed, because inventories of small bodied and species rich taxa need huge scientific effort (Lawton et al. 1998) and are beyond possibility in most cases. Ecological investigations of lesser known groups (such as many insects, nematodes and fungi) are needed, because in most ecosystems diversity patterns and composition of two or more taxonomic groups are seldom or only weakly correlated (Gaston and Spicer 2004; Cox and Moore 2005).

Positive cross-taxon concordance in either alpha or beta diversity refers to a situation in which one (or more) taxonomic group could be used as a surrogate for other groups; congruence in assemblage structure is a highly interesting aspect in applied ecology, such as biomonitoring (e.g. Carlisle et al. 2008; Mykrä et al. 2008) and nature conservation (e.g. Su et al. 2004; Bilton et al. 2006). There are, however, rather conflicting results about community concordance in different habitats or regions. For example, in two rather large studies of terrestrial habitats in the southern hemisphere, no support was found for the use of a reliable surrogate group (van Jaarsveld et al. 1998; Lawton et al. 1998). Instead, in a study of meadow communities in North America. positive concordance а in assemblage structure, but not in species richness, was reported (Su et al. 2004). In the boreal zone, biotic groups of streams and rivers showed concordant patterns only if the spatial gradients were long enough (Paavola et al. 2006). To clarify, different lotic groups differently to environmental responded variables within a scale of a catchment, but between-ecoregion concordance was probably caused by biogeographic factors (post-glacial history, latitudinal distribution patterns) (Paavola et al. 2006). Concordant patterns in assemblage structure have most often been studied with relatively well known taxonomic groups. such as fishes. benthic macroinvertebrates, diatoms and macrophytes in freshwater ecosystems (e.g. Paavola et al. 2006; Grenouillet et al. 2008) and plants, butterflies, beetles and birds in terrestrial ones (Blake et al. 2003; Su et al. 2004; Similä et al. 2006; Davis et al. 2008). Notable exceptions are e.g. studies by Sääksjärvi et al. (2006) on parasitoid wasps and tropical plants and Murray et al. (2006) on curculionid beetles, lichens, bryophytes and higher plants. Thus, there is need for concordance studies gathering poorly known taxonomic groups and habitats.

In terms of species richness, habitat associations and functional feeding groups, semiaquatic flies (Diptera, Nematocera) are a diverse group of insects, of which most species are dwellers of wetlands and small water bodies (Salmela 2004; Salmela et al. 2007; de Jong et al. 2008; Wagner et al. 2008). Many semiaquatic fly species are stenotopic, that is, their life cycles are bound to rather narrow habitat niches, such as

springs, rich fens, headwater streams, fruiting bodies of fungi or decaying wood (e.g. Alexander 1920; Brindle 1967; Salmela et al. Hancock et al. 2009). Hence, 2007: semiaquatic flies are considered to have great potential for biomonitoring, conservation and assessment of freshwater habitats or other wetlands (Chadd and Extence 2004; Salmela 2008). Semiaquatic flies belong to 12 families and 424 semiaquatic fly species are currently known from Finland (Salmela 2010). Within the semiaquatic flies, craneflies sensu lato (Tipulidae, Limoniidae. Pediciidae. Cylindrotomidae) are the most species rich group (330 spp), followed by moth flies (Psychodidae, 57 spp); other families are relatively species poor, each of them having (Ptychopteridae, 1-15 species Dixidae. Thaumaleidae. Synneuridae, Canthyloscelidae, Pachyneuridae, Pleciidae). Semiaquatic flies have been rather neglected and poorly known in Finland up till the beginning of 2000's; studies dealing with the ecology of the species and community parameters such as diversity, abundance and species turn-over, have been very few in number (see Salmela 2008 for a short review about northern Fennoscandia).

Within the northern boreal vegetation zone in Finland, the local richness (alpha diversity) of semiaquatic flies does not decrease with increasing latitude, but a remarkable change occurs in the regional species pool at the border of coniferous forests and subalpine ecoregion (Salmela 2008). On a large scale, within the extent of northern boreal zone, species composition of semiaquatic flies are influenced by biogeographical factors and local conditions, especially factors which are connected to hydrogeology (e.g. ground water influence, mineral composition of the bedrock) and to the dominating habitat type (Salmela 2008). The variation of species richness and community composition on a smaller spatial scale (i.e. within a drainage area or ecoregion) and the factors influencing them, are insufficiently known. In addition, the community concordance of semiaquatic flies with other biotic groups, like plants, is not well known.

The main goal of the present study is to investigate the variation in assemblage species richness composition and of semiaquatic flies in the subalpine ecoregion, northernmost Finland. Specifically, this study is trying to seek out patterns in the following issues: (i) what is the amount of variation in the species richness and assemblage composition in three habitat types (aapa mires, springs, headwater streams) and (ii) to study the correlation of semiaquatic flies with plant species composition (higher plants and mosses), environmental variables and geographical distance (spatial autocorrelation). It has recently been shown that terrestrial arthropod assemblages are well predicted by plant species composition; in fact, plants proved to be better determinants of arthropod species composition than environmental variables physical or vegetation structure (Schaffers et al. 2008). It is thus highly interesting to compare the performance of environmental variables and plant species composition as surrogates for semiaquatic fly assemblage composition. This study is part of the national survey of semiaquatic flies which aims to assess the redlist status and distribution patterns of the species, and develop tools for the conservation and assessment semiaquatic fly communities (J. Salmela, unpublished data).

#### **Materials and Methods**

#### Study area

The study area is located in northernmost Finland 68°35'N 22°40'E). (ca. biogeographical province of Lapponia enontekiensis, belonging to the northern boreal vegetation zone and to its subzone, the subalpine ecoregion (Lindholm and Heikkilä 2006). In some publications this subzone is also called the "Arctic-alpine" ecoregion (e.g. Soininen et al. 2004). The subalpine ecoregion is characterized by virtual lack of coniferous forests; the tree line is formed by mountain birch, Betula pubescens ssp. czerepanovii (Orlova) Hämet-Ahti (Fagales: Betulaceae) and large districts are treeless fell areas. The study area is part of the Tarvantovaara wilderness area (67030 ha), its landscape is dominated by mountain birch forests, mires and fells. This district is practically lacking any year-round settlements or public roads. The altitude in the area ranges between 360 to 630 m a.s.l. and the bedrock is composed of variety of rocks, including granodiorite, quartzite, metavolcanites and gabbro (Table 1). The climate is continental, that is, relatively low annual precipitation (ca. 450 mm) and cold winters (mean temperature of January ca. -16° C) are prevailing (Piirainen 2002; Tikkanen 2006). Evaporation does not exceed precipitation during the thermal growing season (105-115 days  $\geq$  5° C, Piirainen 2002), and thus, runoff from the drainage areas is relatively large despite of low amount of precipitation. This phenomenon is strikingly seen in the occurrence of numerous sloping fens, which indicate high moisture level.

# Selection of study sites and measurement of environmental variables

A total of 19 study sites were selected; eight of these were aapa mires, four were springs and seven were headwater streams (Table 1, Figure 1). *Aapa mires* in the subalpine ecoregion are minerotrophic fens, characterized by flarks (i.e. inundated, mudbottom or carpet level vegetation, mainly inhabited by horizontally growing bryophytes, see Rydin and Jeglum 2006 for terminology) surrounded by irregular pattern of strings, (hummock-level vegetation, dominated by dwarf shrubs). Most of the studied aapa mires were poor and intermediate rich in their floristic composition, that is, they were dominated by Sphagnum sp. and Warnstorfia procera (Renauld and Arnell) Tuomikoski (Hypnales: Amblystegiaceae) mosses. Some aapa mires were rich fens, i.e. they were characterized by Scorpidium mosses. Springs are points of emerging groundwater, usually stable in terms of water temperature and discharge. One of the studied springs had a minerogenous bottom of the spring brook while in other sites the outflow took place over soft bottom or moss vegetation (e.g. Warnstorfia exannulata (W.Gümbel) Loeske, Philonotis fontana (Hedwig) Bridel). Springs were quite sharply separated from the surrounding biotopes (birch forests, alpine heaths and mires). Unfortunately, only four springs were studied. The original plan was to include more springs, but suitable sites were either not available in certain areas or not accessable due to difficult terrain. Headwater streams are small lotic waters located in the upper reaches of the catchment. In relation to springs, headwater streams are much more variable in their discharge and water temperature. The headwater streams of this study were first or second order streams, characterized by small drainage areas (92-1004 ha) and minerogenous bottoms (sand, gravel, stones and boulders); mosses such as Fontinalis dalecarlica Schimper and Hygrohypnum alpestre (Sw. Ex Hedwig) Loeske were common inhabitants on the submerged and emergent stony substrates of the streams. Riparian zones of headwater streams were rich in plant species, which

study site	N	E	habitat type	birch z <sup>2</sup>	prevailing bedrock type <sup>3</sup>
Lavivaara N	7613921	3318967	Aapamire; intermediate flark fen	I.	Quartz monzodiorite
Guhkesgielas N	7621821	3327056	Aapamire; rich flark fen	Ĩ	Gabbro
Syvävuoma	7618817	3323684	Aapamire; rich flark fen	L	Quartz, arkosite, mica shcist
Tuulivaara N	7621820	3328829	Aapamire; intermediate flark fen	Ľ	Mafic and felsic metavolcanite
Tuolvanenä SW	7626529	3332154	Aapamire; intermediate flark fen	0	Quartz, arkosite, mica shcist
Stuorrahanoaivi S	7627823	3328685	Aapamire; intermediate flark fen	0	Quartz monzodiorite
Tomuttirova I	7623848	3319419	Aapamire; intermediate flark fen	1	Mafic and felsic metavolcanite
Tomuttirova II	7623638	3318845	Aapamire; intermediate swamp fen	I	Mafic and felsic metavolcanite
Marttavaara	7616276	3321834	Spring; helocrene, mesotrophic	I.	Quartz, arkosite, mica shcist
Veitikielas	7627041	3325798	Spring; limnocrene, mesoeutrophic	0	Tonalite and trondhjemite gneiss
Hietakero	7625969	3323074	Spring; helocrene, mesotrophic	0	Gabbro
Teräväpäänharjut	7620432	3317637	Spring; rheocrene, mesoeutrophic	L	Quartz, arkosite, mica shcist
Pulkanoja	7622864	3328802	Hw stream, cathment area 866 ha	T	Mafic and felsic metavolcanite
Stuorrahanoaivi	7627522	3328421	Hw stream, cathment area 92 ha	0	Quartz monzodiorite
Ravaltolommol SE	7626054	3326760	Hw stream, cathment area 151 ha	0	Quartz monzodiorite
Ravaltojärvi	7628764	3324787	Hw stream, cathment area 424 ha	I	Tonalite and trondhjemite gneiss
Saappajärvi	7625346	3323536	Hw stream, cathment area 63 ha	0	Gabbro
Pahtavaara SE	7625151	3321084	Hw stream, cathment area 113 ha	Ī	Mafic and felsic metavolcanite
Tomuttioja	7622477	3318433	Hw stream, cathment area 1004 ha	1	Mafic and felsic metavolcanite

<sup>1</sup>N=north coordinate, E=east coordinate, uniform grid coordinate system (27°E), <sup>2</sup>location in the birch forest zone=1, location in the tree-less fell area=0, <sup>3</sup>prevailing bedrock type, modified from the Sutigis-database of Metsähallitus.

included species typical for mires, herb-rich forests and alpine meadows.

The study sites were chosen beforehand to cover gradients in the bedrock composition (siliceous -base rich stones) and altitude (birch forest - fell areas). The shortest distance between the study sites was 0.4 km and the longest distance 16.9 km, average distance between the sites was 7.4 km; the study sites were located in an area ca. 225 km<sup>2</sup>. Malaise traps (see below) were set during the first visit in the beginning of June (8-12 June 2009). In aapa mires, the traps were placed in the immediate vicinity of flarks. In springs and headwater streams, the traps were set over the flowing water or in the bank area as close as possible to the spring pool or flowing water. Each trap formed the centre of the study site which was a  $10 \times 30$  m rectangle. The following environmental variables were measured from the study sites: tree basal area (i.e. the cross-sectional area over the at breast height,  $m^2/ha$ ), coverage (%) of different moisture levels and soil types (water, flark, intermediate, hummock, total peatland area and minerogenous substrates),

water temperature (GEFU digital thermometer, precision +/- 0.1) and pH (pHep-portable instrument, precision +/- 0.1); water temperature was measured three times (June, July, end of August) and pH in July. Finally, altitude was measured from topographic maps and exact location with a GPS-navigator (Garmin etrex, precision +/- 8 m).

## Sampling of adult insects and identification of plant species composition

One Malaise trap was placed in each study site. Malaise (length 110, height 140, width 70 cm) is a trap model made of cloth (black sides, white cover) and is suitable for collecting low-flying insects, especially dipterans. efficient for hymenopterans, trichopterans and plecopterans. Based on experience of ten years and >400 trapping sites, no protected invertebrates or any vertebrates have been caught by the traps. The traps were set in the beginning of June (8-12 June 2009), collecting jars were emptied in the middle of July (18-24 July) and in the end of August (23-27 August); traps also were removed from the field during this last

occasion. A solution of 50 % ethylene glycol + few drops of detergent was used as a preservative in the traps. The collected material was stored in 80 % ethanol. The semiaquatic fly specimens were sorted from the material in the laboratory and were identified to species level. During the last field trip in August sweep net samples (ca. 15 min collecting effort in each site) were taken and this material was later combined with the Malaise trap material (i.e. final species × sample units matrix). Literature for the identification of semiaguatic flies comes from hundreds of sources, too numerous to be referred here. Labeled museum samples (in 80 % ethanol, 2 ml plastic tubes) of most species are deposited in the Zoological Museum, University of Turku, Finland (ZMUT).

Higher plants (Tracheophyta) and bryophytes (mosses and liverworts) were identified during the field trip in July; the aim was to detect all plant taxa present, to species or genus level. For each site the plant species composition was inventoried (presence/absence) from the  $10 \times 30$  m study plots. In the relatively monotonous aapa mires the inventories were performed within 30-45 minutes and in the structurally more complex lotic sites ca. 60-90 minutes. A hand lens  $(30\times)$  was used in the field and samples were taken for later identification in the laboratory. Nomenclature of the higher plants and bryophytes follows Hämet-Ahti et al. (1998) and Ulvinen and Syrjänen (2009), respectively (Appendix 2).

#### Data analysis

One-way ANOVA was used to test for differences between the mean values of the variables (species richness, abundance etc.) from the studied habitats (aapa mires, springs, headwater streams). In pair-wise *post hoc* comparisons, Tukey's HSD test was used. In order to validate the use of parametric method, the homogeneity of the variances and normal distribution of the variables were assessed with the tests of Levene and Shapiro-Wilk, respectively. If these assumptions were not satisfied, the non-parametric Kruskal-Wallis test was used with the Mann-Withney U-test in pairwise comparisons. Correlations between the species richness of semiaquatic flies. plant species richness and environmental variables were studied by using non-Spearman rank correlation parametric coefficient. The chi square test ( $\gamma^2$ ) was used to examine whether there were differences in the number of occurrences of noteworthy species (red-listed and national responsibility species) in the studied habitats.

Semiaquatic fly community composition of the study sites were examined and viewed with a number of multivariate methods. NMS (non-metric multidimensional scaling) is an ordination method, in which the original ranked distances (based on distance measure) of the sample units in the *p* dimensional species space are forced to a reduced, kdimensional ordination (Legendre and Legendre 1998; McCune and Grace 2002). Spearman correlation coefficients were calculated between the ordination's coordinates of the sampling units and environmental variables. Significant correlations are instrumental in the assessment of which variables are associated with the position of study sites on different dimensions of the ordination. McCune and Grace (2002, pp. 107-108) questioned whether it is appropriate to present *p*-values in this connection because coordinate points of the sampling units along the dimensions are not independent variables. However, it is possible to describe and interpret the main directions of variation along the dimensions of the ordination.

MRPP (multi-response permutation procedure) is a non-parametric method for testing the null hypothesis no difference in assemblage composition of two or more a priori defined groups (McCune and Mefford 1999). The within-group homogeneity of each group (observed  $\delta$ ) is compared to a random arrangement of the sampling units (expected  $\delta$ ) and this difference is the effect size A (chance-corrected within-group agreement). A=1, if all sampling units within a group are identical and A=0, if the within group variation in the species composition equals expectations by chance. The statistical inference (*p*-value) is based on a comparison of frequency distribution of randomized values and observed  $\delta$ . ANOSIM (analysis of similarity) is a non-parametric method for testing whether there are differences in the assemblage composition of two or more a priori defined groups (Legendre and Legendre 1998). Similarity (or distance) is calculated between all sample units within each group and afterwards these similarities are ranktransformed (i.e. sample units, which have highest resemblance, rank-order is 1). The same procedure is performed with betweengroup similarities. Test value *R* is based on the remainder of the between-group and withingroup mean rank-order similarities (see Legendre and Legendre 1998 for details) and p-value is based on permutation (10000 permutation were used). R values may range between 1 and 0: the higher the value, the more differentiated assemblages between the groups.

In the above mentioned multivariate methods (NMS, MRPP, ANOSIM) log (x+1) transformed data matrices of semiaquatic flies and Bray-Curtis metric were used. Logarithmic transformation was seen as necessary, because the abundances (number of specimens) were quite variable between the

study sites. This transformation reduces the importance of the most numerous species and thus, gives more weight to less numerous species.

The Mantel test was used to examine concordance of semiaquatic flies with (i) higher plants, (ii) bryophytes, (iii) combined plant material (higher plants + bryophytes), (iv) geographical distance of the study sites and (v) environmental variables. The Mantel test is used to test the null hypothesis of no relationship between two distance matrices. i.e. the test evaluates linear correlation between two distance matrices. Each matrix is calculated from a different set of variables, measured for the same sample units (study sites) (Legendre and Legendre 1998; McCune and Grace 2002). The test value  $r_M$  is the Pearson correlation analogous to coefficient (range -1 and 1). Statistical significance is calculated by permutation (9000 permutations were used). Presence/absence data matrix of semiaquatic flies was used because plant species composition was measured to the accuracy of presence/absence; it is recommended to use the same matrix transformation in the comparisons of two distance matrices (Heino 2008). geographical Because and environmental distance matrices (Euclidean distance in both) showed positive Mantel correlations with semiaquatic fly distance matrix, partial Mantel tests were also used in the semiaquatic  $fly \times plant$  matrices correlations. Partial Mantel test can be used to examine the relationship between two resemblance matrices while eliminating the linear effect of third matrix (Legendre and Legendre 1998).

The investigation of species accumulation in the habitat types was based on the assessment of the number of samples (=number of study

sites per habitat type) and number of individuals. Because there was only one trap per study site, it was not possible to use means and SDs to study the effect of different number of traps on the species accumulation within a single site. Instead, the cumulative number of species was calculated for the combined material for each habitat type (Mao-Tau method, see Colwell 2009). The rarefaction method was used for the calculation of individual based accumulation curves. Rarefaction is instrumental if the sampling efficiency between the study sites has varied and the samples should be standardized to a given number of specimens (Krebs 1998).

Non-parametric species richness estimators (Jackknife<sub>1</sub>, Jackknife<sub>2</sub>, Chao<sub>1</sub>, Chao<sub>2</sub>) were used to evaluate the total number of species (observed + unseen species) in the combined material for all study sites (local species pool) and for habitat types (local species pool within each habitat type). These estimators are based on the assumption that the observed species richness is lower than the true richness of the study sites (e.g. Colwell and Coddington 1994; McCune and Grace 2002; Magurran 2004; Colwell 2009). These estimators are best suited for an analysis of such communities, in which a given study site has several plots, quadrats, traps or other sample units of similar size (Krebs 1998; Magurran 2004). In this study these estimators were seen as appropriate because at each site the sampling intensity was of the same magnitude and the combined material from the habitat types most probably covered >50 % of the estimated number of species (Ulrich and Ollik 2005).

Mean values of the Simpson diversity index (1-D) were calculated for the habitat types. This index is calculated as

$$1 - \sum (p_i)^2$$
 (1)

where  $p_i$  is the proportion of the species *i* of the total number of specimens in a community (Krebs 1998). The diversity index is linked to the probability that two randomly chosen specimens belong to different species. This index was used together with the rankabundance distributions (Whittaker plots) and the percentages of the most abundant species, to evaluate how skewed the species' abundances are among the habitat types.

MRPP test was calculated by using PC-ORD 5.0 (McCune and Mefford 1999), species accumulation curves and non-parametric species richness estimators by using EstimateS 8.2.0 (Colwell 2009) and Mantel tests by using Fstat 2.9.3.2 (Goudet 1995) and other analyses by using PAST 1.94b software (Hammer et al. 2001).

#### Results

#### Assemblage composition of semiaquatic flies, relationships with plant species composition, environmental variables and geographical distance

The collected material of adult semiaquatic flies was composed of 9038 specimens of 94 species (<u>Appendix 1</u>). A total of 1592 specimens (exx.) and 62 species (spp.) were

n (+standard doviation) values in the habi

	Aapamires (n=8)	Springs (n=4)	Hw streams (n=7)
T C°'	12.0±1.3	3.4±1.3	10.7±1.4
pН	5.3±0.4	6.4±0.2	6.8±0.6
peatland area % <sup>2</sup>	100±0	95±6.6	74±24.9
flark area % <sup>3</sup>	45±22.6	18±10	0
lawn-carpet area %	36±23.5	13±6.5	2±3.8
hummock area %	20±15.2	64±11.1	72±23.3
mineral soil area %	0	0	11±24.1
water surface area %	0	6±6.7	15±5.4
tree basal area /ha	0.3±0.7	3±2.4	10±11.2
altitude (m a.s.l.)	441±26.4	466±58.1	471±42

Table 2. Environmental variables measured from the study sites and

<sup>1</sup>Water temperature, mean value of three measurements, <sup>2</sup>Total area % of mire vegetation (=peat forming plant assemblages) in the study area, <sup>3</sup>Flark, lawn-carpet and hummock indicate the moisture levels of mire vegetation, flark being the most wet and hummock the most dry.

**Table 3.** Correlation cofficients (Spearman) and the associated p-<br/>values between the NMS ordination scores of semiaquatic flies<br/>(dimensions 1-3) and environmental variables.

	I. dimen.	2. dimen.	3. dimen.
pН	0.80***	0.22	0.31
peatland	-0.87	-0.04	-0.51*
flark	-0.75***	-0.03	-0.63**
lawn-carpet	-0.77***	0.17	-0.43
hummock	0.64**	-0.25	0.45
mineral soil	0.49*	-0.12	0.34
water surface	0.85***	-0.001	0.45
tree basal area	0.81*	0.37	0.46*
water T C°	-0.43	-0.05	0.15
altitude	0.04	-0.46*	0.12
tatistically significa	ant correlation (*p	o<0.05, **p<0.01	, ***p<0.001).

found from aapa mires, the corresponding figures for the spring and headwater streams are 1288 (exx.), 49 (spp.) and 6158 (exx.), 82 (spp.), respectively. According to NMS ordination (Figure 2), headwater streams are clearly separated from the aapa mires, springs are located closer to headwater streams but do not form their own separable group. The distribution of the study sites along the first dimension correlates significantly with several environmental variables which are related to the conditions typical for mires (positive correlation) and lotic waters (negative correlation); the second dimension is correlated only with altitude (Table 3). The third dimension correlates weakly with the same parameters as the first dimension, and this dimension is practically redundant with the first dimension (Table 3). In other words, the arrangement of the study sites along the first dimension is mirroring a gradient of hydrological conditions and the second gradient discriminates the sites according to an altitudinal gradient.

According to MRPP test, the assemblages of the habitat types were significantly different (A=0.326, p<0.001). In pair-wise comparisons aapa mires differed from springs (A=0.218, p=0.002) and headwater streams (A=0.341, p < 0.001), but no difference was found between the assemblages of springs and headwater streams (A=0.076, p=0.11). The result of the ANOSIM test was in accordance with the MRPP, since the *a priori* classification of the assemblages was highly significant (R=0.532, p<0.001). Pair-wise comparison revealed that aapa mires differed from springs (p=0.021) and headwater streams (p < 0.001), but no difference was found between springs and headwater streams (p=0.313).

According to Mantel tests, the community concordance of semiaquatic flies with higher plants ( $r_M=0.636$ ), bryophytes ( $r_M=0.451$ ) and combined plant material (r<sub>M</sub>=0.612) were highly significant (Table 4, Figure 3); in other words, distance matrices of semiaquatic flies and plants were positively correlated. Partial Mantel statistics (i.e. the controlling effect of environmental variables and geographical proximity) had an only minor effect on the matrix correlations (Table 4). The association semiaquatic fly assemblages of and geographical distance between the study sites was positive but rather weak ( $r_M=0.221$ ).

**Table 4.** Mantel  $r_M$  test values, *p*-values and coefficient of determination (R<sup>2</sup> %, variance explained) for two-matrix correlations of semiaquatic flies (SF) between geographic distance, environmental distance and between distance matrices of higher plants, bryophytes and combined plant material. Partial Mantel test values (i.e. two assemblage matrices controlled against the effect of environmental and geographical distance matrices) are given.

	Geographical	Environment	Higher plants	Bryophytes	<b>Combined plant material</b>
Semiaquatic flies $r_{M}$	0.182	0.395	0.636	0.451	0.612
partial (environment)			0.546	0.351	0.518
partial (geographic)			0.641	0.447	0.614
Semiaquatic flies p	0.02	<0.001	<0.001	<0.001	<0.001
Semiaquatic flies R <sup>2</sup>	3.3	15.6	40.4	20.3	37.5

Journal of Insect Science | www.insectscience.org

**Table 5.** Mean  $(\pm SD)$  species richness of higher plants, bryophytes and semiaquatic flies (SF) in the habitat types. Mean values are also presented for diversity and abundance variables of semiaquatic flies.

	Aapamires (n=8)	Springs (n=4)	Hw streams (n=7)
higher plant species	15.1±4.2	33.3±7.4	44.9±7.3
bryophyte species	12.8±3.8	22.3±4.2	24.9±10.2
SF species richness	25.7±7.3	27.8±7.8	34.9±7.4
SF dominance %'	29.1±7.0	41.7±11.0	52.3±12.9
SF Simpson I-D	0.86±0.03	0.78±0.07	0.68±0.13
SF abundance <sup>2</sup>	199±123.9	322±160.9	880±662.9

<sup>1</sup>Proportion (%) of the most abundant species (1. rank-order), <sup>2</sup>Total number of specimens.

Instead, correlation between environmental variables and semiaquatic flies was higher ( $r_M$ =0.452) (Table 4, Figure 4). Matrix correlation between geographical distance and environmental variables of the study sites was non-significant (Mantel test:  $r_M$ =0.063, p=0.405). However, higher plant species composition (R<sup>2</sup>=40.4 %) was superior as an explanatory variable of semiaquatic fly species composition over bryophytes (R<sup>2</sup>=20.3 %), environmental variables (R<sup>2</sup>=40.4 %) and geographical distances (R<sup>2</sup>=4.9 %) (Table 4, Figures 3 and 4).

#### Patterns in species richness and abundance of semiaquatic fly communities, rankabundance distributions and estimation of species richness

The species richness of semiaguatic flies varied in the habitat types and was generally highest amongst headwater streams (Table 5, Figure 5a), near-significantly (ANOVA F=2.9, df=2, p=0.08). The habitat types studied differed in the proportion of their most abundant species (ANOVA F=10.3, df=2, *p*=0.0013, square-root transformed), pair-wise post hoc comparisons indicate a significant difference between aapa mires (lowest dominance of the most numerous species) and headwater streams (highest dominance of the most numerous species) (p=0.0009) (Table 5, Figure 5b). Similar results were obtained by using a diversity index: mean values of the Simpson (1-D) index were different between the sites (Kruskal-Wallis H=11.2, p=0.004) (Table 5, Figure 5c). Pair-wise comparisons show that the index value was higher among than headwater mires streams aapa (Bonferroni corrected *p*=0.006). Finally, mean values of the raw-abundance (number of specimens) differed between the habitats (ANOVA F=8.7, df=2, p=0.0028, log transformed) (Table 5, Figure 5d); according to pair-wise comparisons the average number of specimens in aapa mires was lower than that headwater of streams (p=0.02).Correlations rank-correlation (Spearman coefficient) of the semiaquatic fly species richness with the combined plant material (r=0.4, p=0.09) and higher plant species richness (r=0.4, p=0.08) were positive but not significant: correlation with bryophyte richness was also positive but also not significant k (r=0.35, p=0.15).

The rank-abundance distributions of the semiaquatic fly communities among the habitats were quite similar, resembling truncated log-normal type distribution (Figure 6a-c). A typical feature of the assemblages is the numerical dominance of a few species, most members of the community being low in numbers.

Accumulation of species richness (or cumulative number of species) was highest in the headwater streams (Figure 7). The cumulative number of species in springs would probably have risen, had there been greater sampling effort (only four sites studied). According to rarefaction, in all habitat types the number of observed species rises rapidly in the beginning, but levels-off after ca. 50 % of the species are captured (Figure 8a-c). For example, a standardized sample size of 800 specimens would have captured 53 (85 % of the observed number of species), 44 (90 %) and 48 (59 %) species for

**Table 6.** Observed  $(S_{obs})$  and estimated species richness of semiaquatic flies in the combined material (n=19) and amongst the habitat types. Estimated values are based on non-parametric estimators (Jackknife<sub>1</sub>, Jackknife<sub>2</sub>, Chao<sub>1</sub>, Chao<sub>2</sub>), percentual values are the additions of estimators to the observed values.

	combined	aapamires	springs	hw streams
S <sub>obs</sub>	94	62	49	82
l.site <sup>1</sup>	18	21	17	32
2. site <sup>2</sup>	10	8	12	8
singletons <sup>3</sup>	9	15	13	21
doubletons⁴	6	3	I	6
Jackk	.  (+ 8 %)	80.4 (+30 %)	61.7 (+26 %)	109.4 (+33 %)
Jackk <sub>2</sub>	118.7 (+26 %)	91 (+47 %)	66.2 (+35 %)	127.5 (+55 %)
Chaoı	99.1 (+5.4 %)	99.5 (+60 %)	133.5 (+172 %)	118.6 (+45 %)
Chao <sub>2</sub>	107.2 (+14 %)	89.6 (+45 %)	61.4 (+25 %)	146 (+78 %)
Spacias prasa	nt in a single s	tudy site 25pec	ies present in tw	o study sites

<sup>&</sup>lt;sup>1</sup>Species present in a single study site, <sup>2</sup>Species present in two study sites <sup>3</sup>One observed specimen of a species, <sup>4</sup>Two observed specimens of a species.

aapa mires, springs and headwater streams, respectively.

According to Jackknife and Chao estimators, the observed species richnesses among the habitat types were 25-172 % lower than estimated number of species (Table 6). For the combined material of the study area (19 study sites, 94 spp), the observed number of species was 5.4-26 % lower than estimated number of species.

#### Frequency and abundance of the most common semiaquatic flies, rare species and preliminary assessment of the conservation value of the habitat types

The most common (frequency >75 % across all study sites) species of the studied material were *Tipula excisa* Schummel (100 %), *Tipula* subnodicornis Zetterstedt (100 %), *Tricyphona immaculata* (Meigen) (95 %), *Phylidorea squalens* (Zetterstedt) (84 %), *Dicranomyia distendens* Lundström (89 %), *Ptychoptera minuta* Tonnnoir (84 %), *Tipula limbata* Zetterstedt (84 %), *Pedicia rivosa* (L.) (79 %) and *Idioptera pulchella* (Meigen) (78 %); only one out of nine species was not a cranefly. As one can expect on the basis of community analyses, the abundances and frequencies of the species were highly variable between different habitat types. For example. Idioptera linnei Oosterbroek, Prionocera subserricornis Zetterstedt and Pneumia ussurica Wagner were typical inhabitants of aapa mires (Figure 9). No such abundant species (total number of specimens >30) displayed fidelity to, or were found exclusively from springs. However, Euphylidorea meigenii (Verrall) and Dicranota guerini Zetterstedt were more numerous in springs than either in aapa mires or headwater streams (Figure 9). Headwater streams harbored several species that were not present elsewhere (e.g. Dicranophragma separatum [Walker]) or the species occurred abundantly in springs and headwater streams Molophilus flavus Goetghebuer, (e.g. Parabazarella subnegleta [Tonnoir], Figure 9). Tipula excisa and Tricyphona immaculata, among the most eurytopic species, were abundant in the all habitat types (Figure 9).

A total 13 semiaquatic fly species were classified as noteworthy, i.e. the species are considered to indicate conservation value of their habitats (Appendix 1). These species are either red-listed, that is, threatened or near threatened in the Finnish Red-Data book or listed as national responsibility species of Finland (see e.g. Salmela 2009, Penttinen et al. 2010). [Additional information dealing with these species is available from the author]. Most of the occurrences of these species were encountered in aapa mires (36 records), followed by headwater streams (24) and springs (19); no statistical difference between the habitat types was found ( $\chi^2$ =5.88, df=2, p=0.053). There was no difference in the mean number of noteworthy species in the habitat types (aapa mires 4.5, springs 3.4 and headwater streams 4.8, ANOVA F=0.6, df=2, p=0.56).

#### Discussion

#### Community structure, classification of habitat types and assemblage concordance of semiaquatic flies and plants

The semiaquatic fly assemblages of aapa mires were clearly differentiated from those of springs and headwater streams, but no difference was found between the two latter habitat types. This result is rather surprising, since springs deviate from other wetland habitats due to their constant temperature and discharge of water. There are several crenobiontic (spring dependent) and crenophilous (favoring springs) semiaquatic fly species, especially in southern boreal Finland (e.g. Salmela et al. 2007) and Central Europe (e.g. Reusch and Hohmann 2009). In the subalpine fell areas of Fennoscandia, headwater streams may often be spring-fed, but the ones studied here displayed higher summer water temperatures than the springs (see Table 2) and received their water from up-stream lakes or mires. Thus, it can be safely concluded that the studied headwater streams were not appreciably influenced by the groundwater. In a study conducted in the northern boreal Finnish zone, spring semiaquatic fly communities did not form their own, plain cluster group or assemblage type, but showed compositional similarity to rich fens and headwater streams (Salmela 2008). It is very likely that the semiaquatic fly fauna of northern boreal region is characterized by low number of crenobiontic species. Such almost strictly spring-dwelling species are Rhabdomastix parva (Siebke) Tipula fendleri Mannheims and Pneumia pilularia (Tonnoir) (Salmela 2008, J. Salmela, Crenophilous unpubl.). species (e.g. Dicranomyia caledonica Edwards, D. stylifera Lackschewitz and Dicranota guerini) are usually abundant or occur frequently in springs, but are also common in lotic waters or rich fens. Hence, in this light, it is not so puzzling that semiaquatic fly assemblages of springs in the subalpine ecoregion are not so distinctive.

In this study, only two taxa were exclusively present in springs (Dicranota bimaculata [L.] and Dicranomyia (Melanolimonia) sp, female, most likely D. caledonica or D. stylifera). Abundant and/or frequent members of the spring assemblages were also abundant in headwater streams (e.g. Molophilus flavus, Dicranota exclusa, Parabazarella subnegleta). Quite evidently, several springdwelling semiaquatic flies are lotic species that are able to complete their life cycles in thermally constant springs. Based on the author's personal assessment, a part of the northern boreal spring-dwelling semiaquatic flies are calciphilous, in other words, prefer sites with high pH value and specific conductivity. The springs of the study area were not calcareous (assessed from the plant composition, pН species and bedrock composition), and thus, the lack of suitable habitat conditions may explain the absence of such species in this study. Calciphilous northern species are at least D. caledonica, D. stvlifera. Pneumia pilularia and Rhabdomastix parva; the last mentioned species is perhaps restricted to the Caledonian mountains in Finland (Salmela 2008). Based on the results of this study and previous studies, the semiaquatic fly community composition of springs in the northern boreal zone is mostly "nested" (see e.g. Cutler 1998, Ulrich and Gotelli 2007) with surface fed streams and the species richness of springs is usually lower compared to that of headwater streams. On the other hand, influence of upwelling groundwater usually is in connection with high species richness of semiaquatic flies in small lotic waters and mires (Salmela 2008).

In the present study the comparison of semiaquatic fly faunas between ecologically rather different wetland habitats was seen meaningful, because the variation in species composition and richness in such a small spatial scale (ca. 225 km<sup>2</sup>) has not been investigated before in Fennoscandia. It is quite clear that the habitat type can be used as a rough predictor of the semiaquatic fly assemblage, although there is more or less between-habitat overlap in species composition. The spatial structure, i.e. geographical distance between the study sites was not strongly associated with their faunistic resemblance. Instead, much stronger association was noted with the environmental distance and species composition. Despite the several distinguishing characteristics, and the differences between the aapa mires, springs and headwater streams, the studied habitats and their semiaquatic fly assemblages do form wetland gradient, which is mainly а influenced by conditions and resources of peatlands, lotic waters and their riparian zones; altitude is part of this multidimensional habitat space. In addition to environmental factors, biotic interactions (competition, predation. parasitism) are also central determinants of community composition (Morin 1999). It is unfortunate that knowledge about these biotic factors and their importance for community level organization among semiaquatic flies is almost negligible. According to an elegant study by Freeman (1967), physical properties of environment (e.g. substrate, moisture) are likely to be the most important drivers of the soil-dwelling *Tipula* communities, although (intraspecific) larval competition for space may occur and cause high mortality. Freeman (1967) stresses that the coexistence of several *Tipula* species within small spatial scale (some hundreds of square meters) is explainable by the fine-scale niche separation between the species, i.e. differential use of microhabitats. Further, temporal separation (phenology) between closely related species perhaps lessens the probability of interspecific competition (Freeman 1967).

According to Mantel tests, plant species composition explained very well the variation resemblance in of semiaquatic flv assemblages. Mantel test values and associated coefficients of determination showed high matrix concordance between semiaquatic flies and higher plants and between semiaquatic flies and combined plant material; the concordance between bryophytes was much weaker. Indeed, higher plant and combined plant species composition  $(R^2)$ : 37.5-40.4 %) explained much better the variation in semiaquatic flv species composition than either geographical distance  $(R^2=3.3 \%)$  or environmental variables  $(R^2=15.6 \%)$ . Thus, the results of this study are in accordance with Schaffers et al. (2008), who claimed that arthropod assemblages are best predicted by plant species composition (their study encompassed spiders and several terrestrial insect groups in meadow habitats). It has traditionally been considered that insect communities are determined by environmental conditions and (physical) vegetation structure. However, plant species composition is largely shaped by environmental conditions and summarizes these conditions that may fluctuate over time, and plants also modify their own environment (Schaffers et al. 2008). The integrating nature (referring to conditions, physical structure and microhabitats) of plant communities is apparently important for arthropods in all trophic levels both directly (obligate herbivores) and indirectly (e.g. soil dwellers and epigean predators). To conclude,

in the subalpine ecoregion the nematoceran plant assemblages are co-varying and according to the same wetland gradients. As pointed out by Schaffers et al. (2008), this covariation is partly indirect, probably due to a similar pattern of response to underlying environmental variables, and partly direct, due to the obligate associations between biotic groups. For example, in the present study the relatively common Phalacrocera replicata (Cylindrotomidae) eats and dwells among aquatic bryophytes (Peus 1952), larvae of the peatland species *Idioptera linnei* (Limoniidae) inhabit Sphagnum mosses (Boardman 2004) and adult flies may feed on liquids secreted by woody plants (Stubbs 2005) or flower nectars (Chernov and Lantsov 1992).

Community concordance between different taxonomic groups may be rather weak (Similä et al. 2006; Carlisle et al. 2008; Davis et al. 2008; Mykrä et al. 2008) and significant only after the environmental gradients (e.g. location of study sites in multiple ecoregions) are rather long (Paavola et al. 2006). The relatively low concordance may reflect divergent responses of different taxonomic groups to the prevailing environmental factors (Virtanen et al. 2009). On the other hand, the positive cross-taxon congruence may be relatively high, although the same groups would not display correlation in species richness (Su et al. 2004). One should bear in mind that, despite the close geographical proximity of the study sites, the environmental gradients in this study were long, because a selection of such varied habitat types was included. Usually community concordance has been studied within a single habitat, such as streams (Paavola et al. 2006), ponds (Bilton et al. 2006) and springs (Virtanen et al. 2009). As noted above, plant species composition, especially that of higher plants, seems to be the best predictor of semiaquatic fly assemblage This result structure. has importance in nature conservation and monitoring, whether assemblages of a single taxonomic group are to be used as a surrogate for other groups. However, more studies are needed (i) in a larger spatial scale and (ii) within а habitat type before sound recommendations about the potential congruence between higher plants and semiaquatic flies can be made.

Although semiaquatic fly assemblages of small lotic waters (springs and headwater streams) were clearly distinguishable from those of aapa mires, there was a notable overlap in species composition between the habitat types. For example, several peatlanddwelling species were present in headwater Phylidorea streams (e.g. squalens, Alexander. Dicranomvia terraenovae Prionocera ringdahli Tjeder); these species have most probably spent their immature stages in mire vegetation in the vicinity of the traps. Further, eurytopic generalist species (e.g. Tipula excisa, Tricyphona immaculata), which are common in a multitude of moist habitats, were encountered in all habitat types. Hence, the observed species overlap between the habitat types is most likely explainable by two factors: (i) small lotic waters are not extremely clear-cut but part of the wetland gradient and (ii) high frequency of occurrence of generalist species increase the between site similarity. The trapping method used in this study (Malaise) is non-selective; the trap collects insects from a wider area than emergence traps, for example. Nevertheless, the Malaise trap is passive, it does not attract insects in similar manner as the light trap. Based on the author's experience from >400Malaise trapping sites, it can be estimated that the vast majority of the collected specimens have spent their immature life cycle within or in the near vicinity of the  $10 \times 30$  m study

plots. Larger adult flies, especially craneflies, probably have good potential for dispersal, but are usually found in close vicinity to their larval habitat (Freeman 1968).

## Rank-abundance distributions and patterns in species richness

The rank-abundance distribution patterns of semiaquatic fly assemblages in the different habitat types could perhaps be classified as truncated log-normal type (Magurran 2004; Ulrich and Ollik 2005), although no tests of goodness-of-fit to any statistical distribution model was performed. This kind of rankabundance distribution is commonly recorded across different biomes and taxonomic groups (Ulrich and Ollik 2005). Log-normal distribution is characterized by the dominance of one or a few species, most of the species being relatively scarce, i.e. in low numbers. In spite of the seemingly similar rank-abundance patterns in this study, the habitat types were differentiated by the proportion of the most numerous species. The mean proportion of the most abundant species was lowest in aapa mires, higher in springs and headwater streams, in ascending order. In the same vein, the Simpson diversity index values were highest in aapa mires, lower in the springs and headwater streams. In other words. semiaquatic fly assemblages of aapa mires were relatively even, and species abundances were more skewed amongst springs and Nevertheless, headwater streams. rankabundance distribution in a pristine aapamire can be heavily skewed. A sample of >500 specimens belonging to seven species were collected from a carpet-lawn level, poor fen in the northern boreal zone, Kittilä (Finland): tipulid species Tipula subnodicornis accounted for 91 % of the total number of specimens (Salmela 2008). In contrast, intermediate rich and rich fens (as the aapa mires of this study) are more species rich than poor fens (Salmela 2004, 2008) and possibly also display higher evenness.

Cumulative species richness was clearly highest in headwater streams, in other words its species pool was richer than in aapa mires or springs. As there were only four springs studied, the cumulative species richness would probably had been higher had more sites been sampled. Actually, none of the species accumulation curves did not reach an asymptote, and it is likely that species belonging to the local species pool were unrecorded (see below). Individual based species accumulation curves (rarefaction) indicate that observed species richness rose rapidly at first but leveled-off after ca. 50 % of the species were captured. It is hard to give any recommendations about the representative sampling effort (number of traps/study sites or collected specimens). One should consider what is adequate to fulfill the aims of the study: is it sufficient to collect the most common or abundant species, which would allow beta diversity comparison, or is the aim to collect also the rarest members of the assemblage? In species rich taxonomic groups, the probability of catching a rare species is higher the larger the collected sample is (Martikainen and Kouki 2003). Furthermore, the accumulation of rare species is lower than that of common species. A study performed in only one field season does not possibly record all low abundance species (Martikainen and Kaila 2004).

According to non-parametric species richness estimators, there were unseen species in all studied habitat types, perhaps more so in headwater streams than aapa mires or springs. These estimates are, of course, just indications, not absolute facts, because the number of study sites for each habitat type was rather low. Perhaps the best estimate is

obtained by combining the material gathered from 19 sites (94 observed species), which indicates that 5.4-26 % of the species in the study area were not collected. A total of 192 semiaquatic fly species are recorded from the biogeographical province of *Lapponia enontekiensis* (J. Salmela, unpublished data), and thus there is a high probability of encountering more species from the study area than were sampled in the present study.

#### General notes on the observed semiaquatic fly fauna and conservation value assessment

The most frequently encountered semiaquatic fly species of the studied material are widespread and common inhabitants of various wetlands in Finland. The only exception is *Tipula excisa*, a northern species, which is not present south of the middle zone (J. Salmela, unpublished boreal observation). Almost 20 % of the semiaquatic fly species were rather rare, since they were collected from only one site. As already from rank-abundance observed the distributions, there were rather few very abundant species: seven of the most abundant species (>350 specimens) accounted for 65 % of the total number of collected specimens. Most of the abundant or relatively numerous species showed rather high fidelity for a certain habitat type (see Figure 10); in the light of community analyses, this is obvious.

The most remarkable faunistic record of this study was a limoniid, *Dicranomyia intricata* Alexander, a species which have hitherto been found from the northern Baltic in Finland, some 600 km south from the study area (J. Salmela, unpublished observation). There is one old record from North Sweden, Abisko, a collection of a holotype male of *D. suecica* (Nielsen 1953), which is a synonym of *D. intricata*. In the Palaearctic region, the species

is not known to occur outside Finland and Sweden (Oosterbroek 2010). The species was Canada; described from the Nearctic, apparently the species occurs there in boreal (Alexander 1927). Dicranomvia mires intricata was quite numerous in two close lying aapa mires (Tomuttirova I and II), in three other sites only singletons were present (aapamire, spring and headwater stream). Other rare or otherwise notable species were e.g. Dicranomyia lulensis (Tjeder) (endemic to Fennoscandia), Prionocera abscondita Lackschewitz (internationally rare, arctic species), Prionocera woodorum Brodo (rare and poorly known, northern species) and Tipula laccata Lundström and Frev (rare. northern).

Thirteen species, 14 % of the total number of were observed species. classified as noteworthy. These species will be threatened or near threatened in the Red-Data book of species are National Finland. or the Responsibility Species of Finland (Penttinen et al. 2010, J. Salmela, unpublished data). statistically Although no significant differences of the occurrences of such species between habitat types were noted, aapa mires harbored more occurrences (36) than headwater streams (24) or springs (19). Most of these noteworthy species are principally mire species (8 spp), four species could be classified inhabitants of headwater streams or alpine wetlands and only species is crenophilous. Thus, it is not surprising that aapa mires display high conservation value as habitats for semiaquatic flies. From an international view point, aapa mires are important because many mire species having viable populations with no current threat in Finland (e.g. Priocera chosenicola Alexander, P. pubescens Loew, Idioptera linnei) are extremely rare or threatened in the other parts

of Europe (Boyce 2004; Martinovský and Barták 2005).

#### Acknowledgements

I want to show my special thanks to Hannu Jurkkala and Aapo Kahilainen for their vigorous participation in the hard field work in Tarvantovaara wilderness area. Katriina Peltonen helped me in the plant inventories in July and Riikka Juutinen was helpful in numerous ways (higher plant samples, verification of some bryophyte identifications, and comments to the earlier draft of the manuscript). Two anonymous referees, and John Kramer, Jari Ilmonen and Jouni Penttinen gave constructive comments on the manuscript, AK and JI provided good consultation with the Mantel test and other analyses. statistical Field work and identification of the collected material was financially supported by Societas Entomologica Fennica. Vuokon Luonnonsuojelusäätiö Metsähallitus and (Lapin luontopalvelut).

#### References

Alexander CP. 1920. The crane-flies of New York. Part II. Biology and phylogeny. *Memoirs, Cornell University Agricultural Experiment Station* 38: 691-1133.

Alexander CP. 1927. Records and descriptions of crane-flies from Alberta (Tipulidae, Diptera). I. *Canadian Entomologist* 59: 214-225.

Bilton DT, McAbendroth L, Bedford A, Ramsay PM. 2006. Applied Issues: How wide to cast the net? Cross-taxon congruence of species richness, community similarity and indicator taxa in ponds. *Freshwater Biology* 51: 578-590. Blake S, McCracken DI, Eyre MD, Garside A, Foster GN. 2003. The relationship between the classification of Scottish groud beetle assemblages (Coleoptera, Carabidae) and the National Vegetation Classification of British plant communities. *Ecography* 26: 602-616.

Boardman P. 2004. Notes on the autecology of the cranefly Idioptera linnei Oosterbroek, 1992 (Diptera, Limoniidae). *Dipterists Digest* 11: 167-170.

Boyce DC. 2004. A review of the invertebrate assemblage of acid mires. *English Nature Research Reports* 592: 1-110.

Brindle A. 1967. The larvae and pupae of the British Cylindrotominae and Limoniinae. *Transactions of the Society for British Entomology* 17: 151-216.

Carlisle DM, Hawkins CP, Meador MR, Potapova M, Falcone J. 2008. Biological assessments of Appalachian steams based on predictive models for fish, macroinvertebrate, and diatom assemblages. *Journal of North American Benthological Society* 27: 16-37.

Chadd R, Extence C. 2004. The conservation of freshwater macroinvertebrate populations: a community-based classification scheme. *Aquatic Conservation: Marine and Freshwater Ecosystems* 14: 597-624.

Chernov II, Lantsov VI. 1992. Why do Tipulomorpha (Diptera, Insecta) succeed in the arctic conditions? *Acta Zoologica Cracoviensia* 35: 193-197.

Colwell RK. 2009. Statistical estimation of species richness and shared species from samples. Version 8.2.0. URL http://viceroy.eeb.uconn.edu/estimates

Colwell RK, Coddington JA. 1994. Estimating terrestrial biodiversity through extrapolation. *Philosophical Transactions of the Royal Society, London. Ser B* 345: 101-118.

Cox BC, Moore JA. 2005. *Biogeography. An Ecological and Evolutionary Approach*. Blackwell Publishing, 7<sup>th</sup> Edition.

Cutler AH. 1998. Nested patterns of species distribution: processes and implications. In: McKinney ML and Drake JA, editors. *Biodiversity Dynamics. Turnover of populations, Taxa, and Communities*, pp. 212-231. Columbia University Press.

Davis JD, Hendrix SD, Debinski DM, Hemsley CJ. 2008. Butterfly, bee and forb community composition and cross-taxon incongruence in tallgrass prairie fragments. *Journal of Insect Conservation* 12: 69-79.

Fauth JE, Bernardo J, Camara M, Resetarits Jr, WJ, van Buskirk J, McCollum SA. 1996. Simplifying the jargon of community ecology: a conceptual approach. *The American Naturalist* 147: 282-286.

Freeman BE. 1967. Studies on the ecology of larval Tipulinae (Diptera, Tipulidae). *Journal of Animal Ecology* 36: 123-146.

Freeman BE. 1968. Studies on the Ecology of Adult Tipulidae (Diptera) in Southern England. *Journal of Animal Ecology* 37: 339-362.

Gaston KJ, Spicer JI. 2004. Biodiversity. An Introduction. Second Edition, Blackwell Publishing. Goudet J. 1995. FSTAT V1.2: a computer program to calculate F-statistics. *Journal of Heredity* 86: 485-486

Grenouillet G, Brosse S, Tudesque L, Lek S, Baraille Y, Loot G. 2008. Concordance among stream assemblages and spatial autocorrelation along fragmented gradient. *Diversity and Distributions* 14: 592-603.

Hammer Ø, Harper DAT, Ryan PD. 2001. PAST: Paleontological Statistics Software Package for Education and Data Analysis. *Palaeontologia Electronica* 4(1): 9pp.

Hancock GE, Hewitt SM, Godfrey A, Mullin M. 2009. Thoracic spiracular gill structure of *Lipsothrix* (Diptera, Limoniidae) in Britain described from scanning electron micrographs. *Zoosymposia* 3: 77-87.

Heino J. 2008. Influence of taxonomic resolution and data transformation on biotic matrix concordance and assemblage – environment relationships in stream macroinvertebrates. *Boreal Environment Research* 13: 359-369.

Hämet-Ahti L, Suominen J, Ulvinen T, Uotila P. 1998. editors: *Retkeilykasvio*. Luonnontieteellinen keskusmuseo, Kasvimuseo.

van Jaarsveld AS, Freitag S, Chown SL, Muller C, Koch S, Hull H, Bellamy C, Krüger M, Endrödy-Younga S, Mansell MW, Scholtz CH. 1998. Biodiversity assessment and conservation strategies. *Science* 279: 2106-2108.

de Jong H, Oosterbroek P, Gelhaus J, Reusch H, Young C. 2008. Global diversity of craneflies (Insecta, Diptera: Tipulidea or

Tipulidae sensu lato) in freshwater. *Hydrobiologia* 595: 457-467.

Krebs CJ. 1998. *Ecological methodology*, 2<sup>nd</sup> edition. Addison Wesley Longman.

Lawton JH, Bignell DE, Bolton B, Bloemers GF, Eggleton P, Hammond PM, Hodda M, Holt RD, Larsen TB, Mawdsley NA, Stork NE, Srivastava DS, Watt AD. 1998. Biodiversity inventories, indicator taxa and effects of habitat modification in tropical forest. *Nature* 391: 72-76.

Legendre P, Legendre L. 1998. *Numerical Ecology*. Developments in Environmental Modelling 20. Second English Edition, Elsevier.

Lindholm T, Heikkilä R. 2006. Geobotany of Finnish forests and mires: the Finnish approach. In: Lindholm T and Heikkilä R, editors. *Finland – land of mires*. pp. 95–104. *The Finnish Environment* 23.

Magurran AE. 2004. *Measuring biological diversity*. Blackwell Publishing.

Martikainen P, Kouki J. 2003. Sampling the rarest: threatened beetles in boreal forest biodiversity inventories. *Biodiversity and Conservation* 12: 1815-1831.

Martikainen P, Kaila L. 2004. Sampling saproxylic beetles: lessons from a 10-year monitoring study. *Biological Conservation* 120: 171-181.

Martinovský J, Barták M. 2005. Tipulidae (tiplicovití). In: Farkač J, Král D, Škorpík M, editors. *Red list of threatened species in the Czech Republic. Invertebrates.* 245-246. Agentura ochrany přídory a krajiny CŘ, Praha. McCune B, Mefford M J. 1999. PC-ORD for Windows. Multivariate Analysis of Ecological Data. Version 4.0. MjM Software, Gleneden Beach, Oregon, USA.

McCune B, Grace J B. 2002. *Analysis of ecological Communities. With a contribution from Dean L. Urban.* MjM Software Design, Gleneden Beach.

Morin PJ. 1999. *Community Ecology*. Blackwell Science.

Murray TJ, Dickinson KJM, Barrat BIP. 2006. Associations between weevils (Coleoptera: Curculionidea) and plants, and conservation values in two tussock grasslands, Otago, New Zealand. *Biodiversity and Conservation* 15: 123-137.

Mykrä H, Aroviita J, Hämäläinen H, Kotanen J, Vuori K-M, Muotka T. 2008. Assessing stream condition using macroinvertebrates and macrophytes: concordance of community responses to human impact. *Archiv für Hydrobiologie* 172: 191-203.

Nielsen P. 1953. Diagnosen über fünf neue europäische Limoniinae (Dipt. Tipulidae). Zeitschrift der Wiener Entomologischen Gesellschaft 38: 34-36.

Paavola R, Muotka T, Virtanen R, Heino J, Jackson D, Mäki-Petäys A. 2006. Spatial scale affects community concordance among fishes, benthic macroinvertebrates and bryophytes in streams. *Ecological Applications* 16: 368-379.

Penttinen J, Ilmonen J, Jakovlev J, Salmela J, Kuusela K, Paasivirta L. 2010. Sääsket -Thread-horned flies, Diptera: Nematocera. In: Rassi P, Hyvärinen E, Juslén A, Mannerkoski I, editors. Suomen lajien uhanalaisuus –

Punainen kirja 2010. The 2010 Red List of Finnish Species. Ympäristöministeriö& Suomen ympäristökeskus, Helsinki. pp. 477-489

Peus F. 1952. 17. Cylindrotomidae. In: Lindner E, editor. *Die Fliegen der palaearktischen Region* 3(5)3 Lief. 169: 1-80.

Piirainen M. 2002. 5 Kasvillisuus. In: Kajala L. editor. Tarvantovaaran erämaa-alueen ja Lätäsenon-Hietajoen soidensuojelualueen luonto ja käyttö. *Metsähallituksen luonnonsuojelujulkaisuja Sarja A* 140: 42-68.

Reusch H, Hohmann M. 2009. Stelzmücken (Diptera: Limoniidae et Pediciidae) aus Emergenzfallen im "Nationalpark Harz" (Sachsen-Anhalt). *Lauterbornia* 68: 127-134.

Rydin H, Jeglum J. 2006. *The Biology of Peatlands*. Oxford University Press.

Salmela J. 2004. Semiaquatic flies (Diptera, Nematocera) of three mires in southern boreal zone, Finland. *Memoranda Societatis Pro Fauna Flora Fennica* 80: 1-10.

Salmela J. 2008. Semiaquatic fly (Diptera, Nematocera) fauna of fens, springs, headwater streams and alpine wetlands in the northern boreal ecoregion, Finland. *w-album* 6: 3-63.

Salmela J. 2009. The subgenus *Tipula* (*Pterelachisus*) in Finland (Diptera, Tipulidae) – species and biogeographic analysis. *Zoosymposia* 3: 245-261.

Salmela J. 2010. Craneflies of Finland. http://sites.google.com/site/cranefliesoffinland

Salmela J, Autio O, Ilmonen J. 2007. A survey on the nematoceran (Diptera)

communities of southern Finnish wetlands. Memoranda Societatis Pro Fauna Flora Fennica 83: 33-47.

Schaffers AP, Raemakers IP, Sýkora KV, Ter Braak CJF. 2008. Arthopod assemblages are best predicted by plant species composition. *Ecology* 89: 782-794.

Similä M, Kouki J, Mönkkönen M, Sippola A-L, Huhta E. 2006. Co-variation and indicators of species diversity: Can richness of forest-dwelling species be predicted in northern boreal forests? *Ecological Indicators* 6: 686-700.

Soininen J, Paavola R, and Muotka T. 2004. Benthic diatom communities in boreal streams: community structure in relation to environmental and spatial gradients. *Ecography* 27: 330-342.

Stubbs AE. 2005. Observations of craneflies (Diptera, Cylindrotomidae and Limoniidae) feeding of leaf-surfaces. *Dipterists Digest* 12: 134.

Su JC, Debinski DM, Jakubauskas ME, Kindscher K. 2004. Beyond species richness: Community similarity as a measure of crosstaxon congruence for coarse-filter conservation. *Conservation Biology* 18: 167-173.

Sääksjärvi IE, Ruokolainen K, Tuomisto H, Haataja S, Fine PVA, Cardenas G, Mesones I, Vargas V. 2006. Comparing composition and diversity of parasitoid wasps and plants in an Amazonian rain-forest mosaic. *Journal of Tropical Ecology* 22: 167-176.

Tikkanen M. 2006. Unsettled weather and climate of Finland. In: Lindholm T, Heikkilä

R, editors. *Finland – land of mires*, pp 7-16. *The Finnish Environment* 23:

Ulrich W, Gotelli N. 2007. Disentangling community patterns of nestedness and species co-occurrence. *Oikos* 116: 2053-2061.

Ulrich W, Ollik M. 2005. Limits to the estimation of species richness: The use of abundance distributions. *Diversity and Distributions* 11: 265-273.

Ulvinen T, Syrjänen K. 2009. Suomen sammalten levinneisyys eliömaakunnissa. In: Laaka-Lindberg S, Anttila S, Syrjänen K. editors. *Suomen uhanalaiset sammalet*. 309-342. Suomen ympäristökeskus, Helsinki. Ympäristöopas.

Virtanen R, Ilmonen J, Paasivirta L, Muotka T. 2009. Community concordance between bryophyte and insect assemblages in boreal springs: a broad-scale study in isolated habitats. *Freshwater Biology* 54: 1651-1662.

Wagner R, Barták M, Borkent A, Courtney G, Goddeeris B, Haenni J-P, Knutson L, Pont A, Rotheray GE, Rozkošný R, Sinclair B, Woodley N, Zatwarnicki T, Zwick P. 2008. Global diversity of dipteran families in freshwater (excluding Simuliidae, Culiciidae, Chironomidae, Tipulidae and Tabanidae). *Hydrobiologia* 595: 489-519.



Salmela

**Figure 1.** Location of the study area in Finland, subalpine ecoregion (above). Graphical representation of the study sites (below). Vertical and horizontal axes are north and east coordinates (Finnish uniform grid 27° E coordinate system, coordinates are shown in 5 km intervals). Circle=aapamire, triangle=spring and diamond=headwater stream. Filled symbols are located between altitudes of 410-450 m a.s.l. (birch zone) and open symbols are located in the tree-less fell area (460-540 m a.s.l.). Lavivaara, Hietakero and Tarvantovaara are fells, Tomuttirova is a glacifluvial formation (esker) and Ravaltojärvi is a lake. High quality figures are available online.







**Figure 3.** Correlations between assemblage dissimilarities (Bray-Curtis) of a) semiaquatic flies and higher plants, b) semiaquatic flies and bryophytes and c) semiaquatic flies and combined plant material (higher plants + bryophytes). Coefficients of determination (R<sup>2</sup>) are given in the Table 4. High quality figures are available online.





**Figure 4.** Correlation between a) semiaquatic fly assemblage dissimilarity (Bray-Curtis) and geographical distance (km), b) semiaquatic fly dissimilarity and environmental distance (dissimilarity, Euclidean) and c) geographical distance and environmental distance. Coefficients of determination (R<sup>2</sup>) are given in the Table 4 (except for Figure 4C, which is 0.014). High quality figures are available online.











eurytopic craneflies, were rather abundant in all habitat types. High quality

figures are available online.



Salmela

**Appendix 1.** Semiaquatic flies and their abundances (total number of specimens) and frequencies (%) among the studied habitat types. \* indicates noteworthy species (red-listed in 2010 or national responsibility species). Aapamires n=8, springs n=4, headwater streams n=7.

		1 1	MUMMAN ST	1	0.270705
	exx.	freq.%	exx.	freq.%	exx.
Limoniidae					
Austrolimnophila (Archilimnophila) harperi (Alexander, 1926)					2
Dicranophragma (Brachylimnophila) separatum (Walker, 1848)	2000				79
Eloeophila trimaculata (Zetterstedt, 1838)	1	-	-	-	29
Euphylidorea meigenii (Verrall, 1886)	1	13	69	50	23
ldioptera linnei Oosterbroek, 1992	174	88	4	75	1
dioptera pulchella (Meigen, 1830)	19	75	4	75	13
Phylidered (Phylidered) longicernis (Schummel 1829)	17	12	1	25	73
Dhulidanaa (Dhulidanaa) oongloonnis (Schuhnmen, 1829)	11	13	1	25	/3
Phylidorea (Phylidorea) squalens (Zettersteat, 1838)	60	100	9	50	17
Pilaria meridiana (Staeger, 1840)	4	38	-	-	
Arctoconopa zonata (Zetterstedt, 1851)	8 <b>0</b> 7	121		12	4
Cheilotrichia (Empeda) areolata (Lundström, 1912)	16	63	26	50	11
Erioconopa diuturna (Walker, 1848)	9	63	1	25	1
Erioptera (Erioptera) beckeri Kuntze, 1914*	3	25	-	1.0	~
Erioptera (Erioptera) flavata (Westhoff, 1882)	7	50			1
Sonomvia (Gonomvia) stackelbergi Lackschewitz 1935	2	13	1	25	i
Molobbilus (Molobbilus) ater (Moison 1904)	2 F	25		25	401
Malashika (Malashika) dara Carashika (200	5	25	1	25	491
molophilus (Molophilus) flavus Goetghebuer, 1920	-	-	265	75	1728
Molophilus (Molophilus) propinquus (Egger, 1863)	1	13		1.0	37
Ormosia (Ormosia) ruficauda (Zetterstedt, 1838)	27	50	22	50	152
Phyllolabis macroura (Siebke, 1863)	3 <b>-</b> 2		11	25	2
Scleroprocta sororcula (Zetterstedt, 1851)					4
Symplecta (Psiloconoba) meigeni (Zetterstedt 1838)	с. г	12			2
Dieranomuja (Dieranomuja) didama (Maiana 1004)	3	13		-	4
Dicranomyia (Dicranomyia) didyma (Meigen, 1804)	-		•	-	11
Dicranomyia (Dicranomyia) distendens Lundström, 1912	11	88	19	75	33
Dicranomyia (Dicranomyia) cf. halterata f	0.5		-		1
Dicranomyia (Dicranomyia) hyalinata (Zetterstedt, 1851)*	19	50	17	50	3
Dicranomyia (Dicranomyia) patens Lundström, 1907	12	13	-		3
Dicranomyia (Dicranomyia) terraenovae Alexander, 1920	1	13	-		6
Dicranomvia (Dicranomvia) ventralis (Schummel, 1829)	1	13	627	1.2	
Disranomyla (Idiobuga) bateralla Edwards 1921		13	14	75	
Dicraholnyid (lalopyga) haitereid Edwards, 1921	1	13	16	/5	80
Dicranomyia (lalopyga) intricata Alexander, 1927*	88	38	1	25	1
Dicranomyia (Idiopyga) Iulensis (Tjeder, 1969)*	52	50	1	25	3
Dicranomyia (Idiopyga) ponojensis Lundström, 1912*	109	88	6	100	13
Dicranomyia (Idiopyga) stigmatica (Meigen, 1830)	5	50	12	50	4
Dicranomyia (Melanolimonia) rufiventris (Strobl, 1900)	3	38	9	100	54
Dicranomyia (Melanolimonia) sp f	1		1	25	-
Limonia sylvicola (Schummel, 1829)				-	10
Tipulidae	0.54		0.50		10
Angestibule tumidicersis (Lundetröm 1907)*	24	(2	2	75	
	26	63	3	75	-
Dictenidia bimaculata (Linnaeus, 1760)		-	-		1
Prionocera abscondita Lackschewitz, 1933*	4	25	1	1.41	
Prionocera chosenicola Alexander, 1945*	•	-			2
Prionocera pubescens Loew, 1844	28	63	1	25	1
Prionocera turcica (Fabricius, 1787)	7	50	-	-	1
Prionocera recta Tieder, 1948	3	25	-	-	1
Prionocera ringdahli Tieder, 1948	2	25			1
Deionocora corricornie (Zerocorade 1020)	4	25	-	-	
rionocera serricornis (Zettersteat, 1838)	12	38		-	
Prionocera subserricornis (Zetterstedt, 1851)	73	63	2	25	1
Prionocera woodorum Brodo, 1987	1	13	0.50	1.2	1
Tanyptera (Tanyptera) atrata (Linnaeus, 1758)	1	13			3
Tanyptera (Tanyptera) nigricornis (Meigen, 1818)	1	13	-		1
Tipula (Lunatipula) trispinosa Lundström, 1907	7(#7		-		12
Tibula (Platytibula) luteibennis Meigen 1830	17	50			
Tibula (Platitibula) malanacaras Schummel 1933	1/2			-	
The (Providential metanoceros schummer, 1855	112	88	-		4
npula (rterelachisus) mutila VVahlgren, 1905		13	. 7		-
Tipula (Savtshenkia) gimmerthali Lackschewitz, 1925	5	13	23	75	35
Tipula (Savtshenkia) grisescens Zetterstedt, 1851	9	25	17	75	7
Tipula (Savtshenkia) invenusta Riedel, 1919	19	63	4	50	19
Tipula (Savtshenkia) limbata Zetterstedt, 1838	14	75	11	100	27
Tibula (Saytshenkia) subnodicornis Zetterstedt 1838	45	100	59	100	64
Titula (Schummelia) uzviisemie Schummel 1022	03	100	30	100	04
npula (schummella) variicornis schummel, 1833	-	•	•	-	20
Iipula (Vestiplex) excisa Schummel, 1833	243	100	243	100	471
Tipula (Vestiplex) laccata Lundstrom & Frey, 1916*	1.0	0.00	1.00	(	8
Tipula (Vestiplex) montana verberneae Mannheims & Theowald, 1959	9	25	4	50	14
Tipula (Vestiplex) nubeculosa Meigen, 1804		-	-		1
	_		0.24		2
Tibula (Vestiblex) tchukchi Alexander 1934	1020	1.1.1.1.1			
Tipula (Vestiplex) tchukchi Alexander, 1934	12	-	-	-	

Sal	me	la
-----	----	----

Pediciidae						
Dicranota (Dicranota) bimaculata (Schummel, 1829)	-	-	52	25	-	-
Dicranota (Dicranota) guerini Zetterstedt, 1838	-	-	59	75	14	57
Dicranota (Paradicranota) gracilipes Wahlgren, 1905	1	13	-	-	46	43
Dicranota (Rhaphidolabis) exclusa (Walker, 1848)			36	50	- 11	57
Pedicia (Pedicia) rivosa (Linnaeus, 1758)	13	50	16	100	39	10
Tricyphona (Tricyphona) immaculata (Meigen, 1804)	87	88	133	100	443	10
Cylindrotomidae						
Cylindrotoma distinctissima (Meigen, 1818)	-	-	-	-	9	57
Phalacrocera replicata (Linnaeus, 1758)	17	50	3	50	1	14
Ptychopteridae						
Ptychoptera hugoi Tjeder, 1968	1	13	-	-	I	14
Ptychoptera minuta Tonnoir, 1919	52	88	9	100	17	71
Psychodidae						
Berdeniella freyi (Berdén, 1954)	-	-	-	-	351	57
Parabazarella subnegleta (Tonnoir, 1922)	-	-	41	50	880	7
Pericoma formosa Nielsen, 1964	-	-	-	-	25	14
Pneumia borealis (Berdén, 1954)*	1	13	24	100	87	71
Pericoma rivularis Berdén, 1954	-	-	25	100	123	43
Pneumia stammeri (Jung, 1954)	-	-	-	-	458	71
Pneumia ussurica (Wagner, 1994)*	83	63	1	25	1	14
Chodopsycha lobata (Tonnoir, 1940)	-	-	1	25	-	-
Logima satchelli (Quate, 1955)	6	25	6	50	19	7
Psycha grisescens (Tonnoir, 1922)	1	13	-	-	-	
Psychoda phalaenoides (Linne, 1758)	1	13	-	-	6	5
Psychoda spl	-	-	1	25	12	29
Dixidae						
Dixella borealis (Martini, 1929)	-	-	-	-	I	14
Dixella laeta (Loew, 1849)	13	63	1	25	2	14
Dixella naevia (Peus, 1934)	1	13	1	25	1	14
Dixella obscura (Loew, 1849)	2	13	3	25	1	14
Synneuridae						
Synneuron annulipes Lundström, 1910	-		-		2	14

**Appendix 2.** Higher plants and bryophytes and their frequencies (%) amongst the studied habitat types. Species are in alphabetical order. Aapamires n=8, springs n=4, headwater streams n=7.

Higher plants	aapamires	springs	hw streams
Agrostis mertensii	-	-	43
Alchemilla sp			57
Andromeda polifolia	88	50	29
Antennaria dioica	-	-	14
Anthoxanthum odoratum ssp. alpinum	-		86
Astragalus alpinus ssp. arcticus	-	1.00	14
Bartsia alpina		25	57
Betula nana	88	100	71
Betula pubescens	25	75	71
Bistorta vivipara	-	75	86
Calmagrostis canescens	-	: <b>-</b> :	14
Calmagrostis lapponica	5	50	29
Calamagrostis purpurea ssp. phragmitoides	-	50	71
Calmagrostis stricta	-	50	14
Caltha palustris	-		5/
Cardamine pratensis	-	-	14
Carex aquatilis	25	25	57
Carex brunnescens		-	14
		- 100	04
Carex constant	25	25	29
Carex chardorrhiza	25	25	27
Carex dioica	-	50	14
Carex lasiocarpa	38	-	14
Carex limosa	50	-	-
Carex livida	13		
Carex magellanica ssp. irrigua	38	50	43
Carex nigra ssp. juncella	-	50	43
Carex nigra ssp. nigra	13	50	43
Carex pauciflora	13	25	-
Carex rariflora	13	-	
Carex rostrata	75	50	29
Carex rotundata	63	-	-
Carex vaginata	13	529	57
Cerastium fontanum	-	25	14
Chrysosplenium tetrandrum	-	25	-
Cirsium helenioides	-	50	71
Cornus suecica	-	25	43
Deschampsia flexuosa	-	25	86
Empetrum nigrum	50	100	71
Ebilobium angusujonum	-	- 75	27
Epilobium homemannii	- 13	100	57
Friothorum angustifolium	88	75	57
Friophorum vaginatum	38	50	14
Envisetum arvense		75	29
Eauisetum fluviatile	25	25	-
Equisetum palustre	-	-	29
Equisetum pratense		-	43
Equisetum sylvaticum	13	75	57
Euphrasia sp	-	-	43
Festuca rubra	-		14
Filipendula ulmaria		1.00	57
Galium palustre	13	-	29
Galium uliginosum	-	25	43
Geranium sylvaticum	-	25	86
Gnaphalium norvegicum	-		43
Gymnocrpium dryopteris	-	1	14
Hieracium sect. Alpina	-	-	43
Hierochloe hirta	-	<u> </u>	14
Juncus filiformis	-		71
Juniperus communis		25	100
Ledum palustre	50	50	-
Linnea borealis	-	25	14
	-	25	43
Luzula pilosa	-		43
lycobodium annotinum	-	25	29
Melambyrum bratense	-	- 25	27
Melica nutans			14
Menvanthes trifoliata	50	820	
Nardus stricta			43
Parnassia balustris	-	50	71
Pedicularis lapponum	13	50	14
Petasites frigidus	-	25	-
Phegopteris connectilis	-	-	29
Phleum alpinum	-	-	14
Phyllodoce caerulea	-	25	71
Pinguicula vulgaris	-		57

Poa sp	13	100	100
Polemonium acutiflorum	-	-	14
Potentilla palustris	38	100	57
Prunus þadus	-	-	14
Pyrola minor	-	50	71
Pyrola rotundifolia	-	25	14
Ranunculus hyperboreus	-	25	-
Ribes spicatum	-	-	29
Rubus arcticus	-	25	86
Rubus chamaemorus	63	100	29
Rubus saxatilis	-	-	14
Salix glauca	13	50	86
Salix lapponum	75	50	71
Salix myrsinifolia ssp. myrsinifolia	-	-	29
Salix myrsinifolia ssp. borealis		25	-
Salix myrtilloides	13	-	-
Salix phylicifolia	13	100	71
Salix sp		-	14
Saussurea albina		50	71
Solosinella colosinoidea		25	57
Selaginena selaginolaes		23	3/
Solidago virgaurea	-	/5	86
Sorbus aucuparia	-	-	14
Sparganium sp	-	-	14
Stellaria borealis	-	25	14
Taraxacum sp	-	-	71
Tofieldia pusilla	-	25	-
Trichophorum cespitosum	63	-	14
Trientalis europaea	13	75	100
Trollius europaeus	-	-	86
Vaccinium microcarþum	75	75	29
Vaccinium myrtillus	50	75	57
Vaccinium oxycoccos	38	-	-
Vaccinium uliginosum	38	100	71
Vaccinium vitis-idaea	38	100	86
Viola hiflora			14
Viola epitosila	25	50	86
Bryonbytes	25	50	
Aulasomnium balustro	20	75	42
Rindia acuta	50	75	43
Bindia acuta	· ·	-	14
Brachythecium rivulare	-	25	-
Brachythecium sp		25	14
Bryum sp	-	-	29
Bryum pseudoriquetrum	-	25	57
Bryum weigelii	-	75	14
Calliergon cordifolium	13	-	14
Calliergon giganteum	13	-	-
Campylium stellatum	-	-	29
Cinclidium subrotundum	13	-	-
Climacium dendroides	-	-	43
Dichelyma falcatum	-	-	29
Dichodontium palustre	-	-	43
Dicranum sp	25	100	86
Fissidens osmundoides	13	-	29
Fontinalis antipyretica	1 -	- 1	43
Fontinglis dalecarlica	1.	I	71
Grimmia sh		-	14
Helodium blandowii	+	50	
Hugenhubnum albertre	+		
Hygronyphum alpestre	+ -	· ·	43
riygronypnum duriusculum		-	43
Hygronypnum ochraceum		25	14
Hylocomnium splendens	13	50	14
Loeskypnum badium	13	-	-
Meesia triquetra	-	-	4
Mnium stellare	-	-	14
Oncophorus wahlenbergii	-	-	43
Paludella squarrosa	25	100	14
Philonotis fontana	-	25	-
Philonotis seriata	-	25	-
Philonotis tomentella	-	50	14
Plagiomnium ellipticum	-	75	43
Plagiomnium medium	-	-	14
Pleurozium schreberi	63	100	71
	1 12		
Pohlia nutans	1 14		
Pohlia nutans Pohlia wahlenbergii	13	100	
Pohlia nutans Pohlia wahlenbergii Pohlia ob	-	100	14
Pohlia nutans Pohlia wahlenbergii Pohlia sp	-	100 25	86
Pohlia nutans Pohlia wahlenbergii Pohlia sp Polytrichastrum sp	-	100 25 -	14 86 14
Pohla nutans Pohlia wahlenbergii Pohla sp Polytrichastrum sp Polytrichum commune	-	100 25 - -	14 86 14 86
Pohlia nutans Pohlia syn Pohlia sp Polytrichastrum sp Polytrichum commune Polytrichum strictum	- - - - - 50	100 25 - - 100	14 86 14 86 14
Pohlia nutans Pohlia wahlenbergii Pohlia sp Polytrichastrum sp Polytrichum commune Polytrichum strictum Polytrichum schwartzii	- - - - 50 13	100 25 - 100 -	14 86 14 86 14 -
Pohlia nutans Pohlia wahlenbergii Polytichastrum sp Polytrichastrum sp Polytrichum commune Polytrichum strictum Polytrichum schwartzii Polytrichum sp	- - - - 50 13	100 25 - 100 - -	14 86 14 86 14 - 14
Pohla nutans Pohla wahlenbergii Pohla sp Polytrichastrum sp Polytrichum strictum Polytrichum strictum Polytrichum strictum Polytrichum sp Polytrichum sp Pseudobryum cinclidioides	- - - 50 13 - 13	100 25 - 100 - - 50	14 86 14 86 14 - 14 14 57

Rhodobryum roseum			14
Sanionia uncipata	-	_	57
Schistidium sb		-	29
Scarbidium revolvons	25	-	14
Scorpidium revolvens	25	-	14
Scorpiolum scorpioldes	25	-	14
	-	25	-
Spragnum annulatum coll.	13	-	-
Sphagnum balticum	25	-	-
Sphagnum capilifolium		25	-
Sphagnum compactum	25	-	-
Sphagnum fallax	13	-	-
Sphagnum fimbriatum	-	-	57
Sphagnum fuscum	50	25	-
Sphagnum lindbergii	63	25	-
Sphagnum magellanicum	13	-	14
Sphagnum majus	25	-	-
Sphagnum palillosum	-	-	14
Sphagnum platyphyllum	-	-	14
Sphagnum riparium	63	50	29
Sphagnum rubellum	13	-	-
Sphagnum russowii	75	75	71
Sphagnum sauarrosum		-	43
Sphagnum subfulvum	25		
Sobaanum subsecundum	38		14
Sphagnum toros	13	-	29
Spridghum teres	13	- 75	29
Spragnum warnstorn	23	/5	14
Straminergon stramineum	88	100	57
layloria lingulata	-	-	14
Tomentypnum nitens	-	50	14
Warnstorfia exannulata	-	100	43
Warnstorfia fluitans	13	25	-
Warnstorfia procera	100	-	-
Warnstorfia sarmentosa	38	-	14
Warnstorfia trichophylla	13	-	29
Barbilophozia kunzeana	25	-	-
Barbilophozia sp	-	25	29
Cephalozia sp	25	25	29
Chiloscyphos polyanthus	-	-	14
Gymnocolea inflata	25	-	-
Harpanthus flotovianus	· ·	50	-
lungermannia exsertifolia		-	29
lungermannia obovata		-	43
lophozia sp	25	50	57
Marchantia polymorpha		100	14
Marsubella emarginata			29
Marsubella		-	14
Malisupena	-	-	14
Mylia anomala	25	-	-
rema sp		-	100
Pullalum ciliare		25	29
Riccardia sp	13	-	4
Scapania hyperborea	13	-	-
Scapania irrigua	13	25	86
Scapania paludicola	25	-	-
Scapania paludosa	-	25	-
Scapania subalpina	-	-	14
Scapania undulata	-	-	43
Tritomaria polita	· ·	-	29