

Quantitative Characterization of Cantharidin in the False Blister Beetle, *Oedemera podagrariae ventralis*, of the Southern Slopes of Mount Elborz, Iran

Authors: Abtahi, S.M., Nikbakhtzadeh, M.R., Vatandoost, H., Mehdinia, A., Rahimi-Foroshani, A., et al.

Source: Journal of Insect Science, 12(152) : 1-5

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.012.15201>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Quantitative characterization of cantharidin in the false blister beetle, *Oedemera podagrariae ventralis*, of the southern slopes of Mount Elborz, Iran

S.M. Abtahi,^{1a} M.R. Nikbakhtzadeh,^{2b} H. Vatandoost,^{1c} A. Mehdinia,^{3d} A. Rahimi-Foroshani,^{4e} M. Shayeghi^{1f*}

¹Department of Medical Entomology and Vector Control, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

²Department of EEOB, 300 Aronoff Laboratory, 318 W 12th Avenue, Ohio State University, Columbus, OH 43210, USA

³Department of Marine Living Resources, Iranian National Institute for Oceanography, Tehran, Iran

⁴Department of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

Abstract

Cantharidin, a potent vesicant and antifeedant agent, is produced by two families of beetles, Meloidae and Oedemeridae (Coleoptera). In this study, the cantharidin content of oedemerid beetles of central Iran was investigated using the GC-MS method. Cantharidin in both sexes of *Oedemera podagrariae ventralis* Meïneitrieās (Oedemeridae) was found in an average of 3.89 µg/beetle in males and 21.68 µg/beetle in females, which are amounts sufficient to irritate human skin. The average of cantharidin in virgin and coupled beetles was 1.35 and 1.62 (µg cantharidin/mg of beetle) respectively. Females had five to six times more cantharidin in their bodies than males, but there was no significant difference between the amount of cantharidin in virgin and coupled females. The results of this study revealed the production of cantharidin in both sexes of beetle.

Keywords: Oedemeridae

Correspondence: ^a abtahi.mohamad@yahoo.com, ^b nik.nikbakht@gmail.com, ^c hvatandoost1@yahoo.com, ^d mehdinia@inco.ac.ir, ^e rahimifo@tums.ac.ir, ^f mansorehshayeghi@yahoo.com, *Corresponding author

Received: 20 February 2011 **Accepted:** 18 August 2011

Copyright : This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 12, Number 152

Cite this paper as:

Abtahi SM, Nikbakhtzadeh MR, Vatandoost H, Mehdinia A, Rahimi-Foroshani A, Shayeghi M. 2012. Quantitative characterization of cantharidin in the false blister beetle, *Oedemera podagrariae ventralis*, of the southern slopes of Mount Elborz, Iran. *Journal of Insect Science* 12:152. Available online: <http://www.insectscience.org/12.152>

Introduction

Cantharidin, $C_{10}H_{12}O_4$ (Figure 1), has an important role in the ecology of some insects that use or produce it as a defense mechanism (Carrel and Eisner 1974). Only two families of beetles, Meloidae and Oedemeridae (Coleoptera), are recognized as cantharidin producers in the animal kingdom. The main function of cantharidin in these beetles is to preserve their eggs from predators (Capinera 2008; Mullen and Durden 2009). Cantharidin can cause severe skin blisters, especially when the insects discharge it from their junctions as a defense system or get crushed on a human body (Mizota 2001; Arnett and Thomas et al. 2002; Nikbakhtzadeh and Targari 2008).

The Oedemeridae family consists of about 100 genera and 1500 species. The species are distributed worldwide, and so are meloid beetles (Vazquez 2002). There are a few reports about the existence of some oedemerid species in Iran (Svihla 1999, 2006, 2007), but there is no information about their cantharidin content. In this study, the presence of cantharidin was recorded and quantified in the dominant species in the southern slopes of the Elborz Mountains, part of the central region of Iran.

Materials and Methods

Beetles

Oedemerids are pollen-feeding insects (Vazquez 2002); therefore, they were collected by hand on flowers during May–June 2009. Beetles were transported alive to the laboratory, and 30 adult beetles were placed in two groups based on sex by investigation of terminal genitalia (15 male and 15 female). The virginity of beetles was

determined by investigation of the internal reproductive system.

Chemical analysis

Extraction. All samples were put into small, fused, test tubes separately and freeze-dried for 24 hr at -50°C and 9×10^{-2} mbar pressure using a LYOTRAP PLUS freeze-dryer (Scientific Laboratory Supplies Ltd., <http://www.scientificlabs.co.uk/>) to determine the dry weight. Whole, dried bodies were hydrolyzed in fused test tubes with 200 μl 6 N hydrochloric acid (Technical HCl, 37%, MERCK, <http://www.merck.com/>) at 120°C for 4 hr to dissolve all body structures and to free bound cantharidines. Subsequently, an equal volume of chloroform was added, each vial was vigorously shaken on a Vortex mixer for 60 sec, and samples were centrifuged (ROTO FIX 32A centrifuge, Hettich, <http://www.hettichlab.com/>) at $3000 \times g$ for 5 min. Using a Pasteur pipette, the organic phase was filtered and transferred into vials (Holz and Strell 1994; Mebs and Pogoda et al. 2009). All test tubes and vials used had been already silanized for 24 hr by dimethyldichlorosilane solution $(\text{CH}_3)_2\text{SiCl}_2$, (MERCK).

GC-MS. The samples were injected in splitless mode via a 1 μl Hamilton syringe into a 6890N (Agilent, <http://www.agilent.com/>) gas chromatograph equipped with a HP-5 (5% phenyl, 95% methylpolysiloxane, non-polar) fused silica capillary column (30 m, ID 0.25 mm ID, 0.33 μm film thickness) that in turn connected to a 5973 mass selective detector (Agilent). The chromatographic conditions were as follows: initial temperature 60°C , temperature increased to 275°C at $10^{\circ}\text{C}/\text{min}$, an isotherm period of 10 min in 275°C , and then cooled down to 60°C . Helium was used as a carrier

gas at a constant flow of 2 mL/min. The electron Impact Ionization (EI 70 ev) provided mass spectra with a characteristic fragmentation of cantharidin: two base peaks of fragments were m/z 128 and 96 (M^+ : 196). The total mass Spectra analyzed by MSD chemstation software and base peaks were compared by the Wiley 275 registry of mass spectral data bank. The quantization of cantharidin was performed using a standard regression curve, drawn by injections of ascending concentrations of authentic cantharidin (Aktin Chemical, Inc., <http://www.aktinchemicals.com/>) from 0.01 to 250 ppm with $y = 61507x - 3E + 06$ formula ($R^2 = 0.994$).

Results

All the collected samples were *Oedemera podagrariae ventralis* Meïneitrieāš (Coleoptera: Oedemeridae) based on a related key (Svihla 1999). Cantharidin was detected in both sexes. It had a peak in the chromatogram with the retention time of 12.8 and mass spectra with base peaks at m/z 128 and 96. Despite the obvious sexual dimorphism (Figure 2), there was no significant difference between the weights of both sexes ($p = 0.972$). However, intraspecific comparisons showed that females had five to six times more cantharidin in their bodies than males. Quantitative measurements are presented in Table 1. Three females were virgin, but there was no significant difference between the cantharidin amount in coupled and virgin females ($p = 0.478$) (Table 2).

Table 1. Cantharidin content in studied oedemerid beetles. Values are means \pm SD.

Beetle gender	Body mass (mg)	μ g cantharidin / beetle	μ g cantharidin / mg of beetle	Cantharidin / beetle (%)
Male	14.9 \pm 5.63	3.89 \pm 3.74	0.3 \pm 0.29	0.03 \pm 0.03
Female	14.8 \pm 6.69	21.68 \pm 8.45	1.62 \pm 0.73	0.16 \pm 0.07

Table 2. Cantharidin amount in virgin and coupled females.

Females	μ g cantharidin / mg of beetle
Virgin	1.35 \pm 0.26
Coupled	1.62 \pm 0.73

Discussion

The main result of this study was the demonstration of cantharidin in *O. p. ventralis* collected from the southern slopes of Mount Elborz, Iran. This subspecies also occurs in south Azerbaijan and southwest Turkmenistan (Svihla 1999). *O. p. ventralis* were found to have 0.03–0.16% of their weight be cantharidin. This percentage of body weight is lower than that in previously studied blister beetles in Iran (0.2–6% of beetle weight) (Nikbakhtzadeh and Targari 2002), but it is still sufficient to irritate human skin (Carrel 1986).

Previous studies of cantharidin content of three other species of oedemerids are shown in Table 3. This study quantified cantharidin contents in both virgin and mated female *O. p. ventralis*. There was no significant difference between them (Table 2), showing that both sexes can produce cantharidin. This was previously shown in another oedemerid beetle, *Oedemara femorata* (Holz and Strell 1994). Oedemerid beetles are unique in that both sexes are able to produce cantharidin, whereas only male *O. p. ventralis* produce cantharidin, transferring it to females during copulation (Capinera 2008; Mullen and Durden 2009). Moreover, the amount of cantharidin in female *O. p. ventralis*, even in virgin ones, is 5–6 fold more than males. This result supports the hypothesis that females produce cantharidin. The higher content of

Table 3. Cantharidin content in oedemerid beetles in both other studies and the current study.

Species	Sex	Cantharidin content		Reference
		Relative (%)	Total (μ g)	
<i>Oxocopsis thoracica</i>	F	0.34	35.2	Carrel et al. 1986
	M	0.23	15.5	Carrel et al. 1986
<i>Heliocis repanda</i>	F	0.72	7.4	Carrel et al. 1986
	M	0.29	2.1	Carrel et al. 1986
<i>Oedemera femorata</i>	F	No information available	8.4	Holz et al. 1994
	M		3.5	Holz et al. 1994
<i>Oedemera femorata</i>	F	0.15-0.21	21-31	Frenzel et al. 1994
	M	0.03-0.09	2.8-9.6	Frenzel et al. 1994
<i>Oedemera flavipes</i>	F	0.16	22	Frenzel et al. 1994
	M	0.21-0.34	16.5-38.5	Frenzel et al. 1994
<i>Oedemera lurida</i>	F	0.23	12.4	Frenzel et al. 1994
<i>Oedemera podagrariae</i>	F	0.16	21.68	Present study
	M	0.03	3.89	Present study

this chemical has been attributed to achieving a greater degree of protection for survival of their progeny (Capinera 2008; Mullen and Durden 2009).

Acknowledgments

The authors would like to thank Professor Vladimir Svihla (Department of Entomology, National Museum of Prague, Czech Republic), who kindly helped us in species diagnosis, Ms. Atefeh Khazeni (Department of Medical Entomology and Vector Control, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran), who kindly cooperated with us in the field, and the instrumental support of the Iranian National Institute for Oceanography.

References

- Arnett RH, Thomas MC, Skelley PE, Frank JH, Editors. 2002. *American beetles (Polyphaga: Scarabaeoidea through Curculionoidea)*. CRC Press.
- Capinera JL, Editor. 2008. *Encyclopedia of entomology*. Springer Science.
- Carrel JE, Doom JP, McCormick JP. (1986). Identification of cantharidin in false blister beetles (Coleoptera, Oedemeridae) from Florida. *Journal of Chemical Ecology* 12(3): 741-747.
- Carrel JE, Eisner T. 1974. Cantharidin: potent feeding deterrent to insects. *Science* 183: 755-757.
- Frenzel M, Dettner K. 1994. Quantification of cantharidin in canthariphilous Ceratopogonidae (Diptera), Anthomyiidae (Diptera) and cantharidin-producing Oedemeridae (Coleoptera). *Journal of Chemical Ecology* 20(8): 1795-1812.
- Holz C, Streil G, Dettner K. (1994). Intersexual transfer of a toxic terpenoid during copulation and its paternal allocation to developmental stages: Quantification of cantharidin in cantharidin-producing oedemerids (Coleoptera, Oedemeridae) and canthariphilous pyrochroids (Coleoptera, Pyrochroidae). *Zeitschrift fur Naturforschung* 49(C): 856-864.
- Mebs, D., W. Pogoda, et al. (2009). Cantharidin and demethylcantharidin (palasonin) content of blister beetles (Coleoptera: Meloidae) from southern Africa. *Toxicon* 53(4): 466-468.
- Mizota K. 2001. Additional records on dermatitis caused by three oedemerid species (Coleoptera: Oedemeridae). *Medical Entomology and Zoology* 52(1): 63-66.
- Mullen G, Durden L, Editors. 2009. *Medical and Veterinary Entomology*. Academic Press.
- Nikbakhtzadeh MR, Targari S. 2002. Cantharidin component of Iranian blister beetles (Col: Meloidae) and their differences between Iranian and exotic species *Iranian Journal of Public Health* 31(3-4): 113-117.
- Nikbakhtzadeh MR, Targari S. 2008. Medically important beetles (Insecta: Coleoptera) of Iran. *Journal of Venomous Animals and Toxins including Tropical Diseases* 14: 597-618.
- Svihla V. 1999. Revision of the subgenera *Stenaxis* and *Oedemera* s.str. of the genus *Oedemera* (Coleoptera: Oedemeridae). *Folia Heyrovskyana Supplementum* 4: 1-117.

Svihla V. 2006. Revision of *Chitona* species (Coleoptera: Oedemeridae) from the eastern Mediterranean. *Acta Entomologica Musei Nationalis Pragae* 46: 107-121.

Svihla V. 2007. Preliminary revision of the genus *Sparedrus* (Coleoptera: Oedemeridae) from eastern and southeastern Asia. *Acta Entomologica Musei Nationalis Pragae* 47: 153-168.

Vazquez XA. 2002. *European fauna of Oedemeridae (Coleoptera)*. Argania Editio.




Figure 1. Chemical structure of $C_{10}H_{12}O_4$. High quality figures are available online.



Figure 2. Male (Left) and female (right) of *Oedemera podagrariae ventralis* Meïneitrieāš. High quality figures are available online.