

Resistance Mechanisms to Chlorpyrifos and F392W Mutation Frequencies in the Acetylcholine Esterase Ace1 Allele of Field Populations of the Tobacco Whitefly, *Bemisia tabaci* in China

Authors: Zhang, Ning-ning, Liu, Cai-feng, Yang, Fang, Dong, Shuang-lin, and Han, Zhao-jun

Source: Journal of Insect Science, 12(41) : 1-13

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.012.4101>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Resistance mechanisms to chlorpyrifos and F392W mutation frequencies in the acetylcholine esterase ace1 allele of field populations of the tobacco whitefly, *Bemisia tabaci* in China

Ning-ning Zhang^a, Cai-feng Liu^b, Fang Yang^c, Shuang-lin Dong^{d*} and Zhao-jun Han^e

College of Plant Protection, Education Ministry Key Laboratory of Integrated Management of Crop Diseases and Pests, Nanjing Agricultural University, Nanjing 210095, China

Abstract

The tobacco whitefly B-biotype *Bemisia tabaci* Gennadius (Hemiptera: Aleyrodidae) is a worldwide pest of many crops. In China, chlorpyrifos has been used to control this insect for many years and is still being used despite the fact that some resistance has been reported. To combat resistance and maintain good control efficiency of chlorpyrifos, it is essential to understand resistance mechanisms. A chlorpyrifos resistant tobacco whitefly strain (NJ-R) and a susceptible strain (NJ-S) were derived from a field-collected population in Nanjing, China, and the resistance mechanisms were investigated. More than 30-fold resistance was achieved after selected by chlorpyrifos for 13 generations in the laboratory. However, the resistance dropped significantly to about 18-fold in only 4 generations without selection pressure. Biochemical assays indicated that increased esterase activity was responsible for this resistance, while acetylcholine esterase, glutathione S-transferase, and microsomal-O-demethylase played little or no role. F392W mutations in ace1 were prevalent in NJ-S and NJ-R strains and 6 field-collected populations of both B and Q-biotype from locations that cover a wide geographical area of China. These findings provide important information about tobacco whitefly chlorpyrifos resistance mechanisms and guidance to combat resistance and optimize use patterns of chlorpyrifos and other organophosphate and carbamate insecticides.

Keywords: carbamate insecticides, esterase activity, insensitivity, organophosphate insecticides, resistance selection
Abbreviations: **AChE**, acetylcholine esterase; **DEM**, diethyl meleate; **GST**, glutathione S-transferase; **MFO** microsomal-O-demethylase; **NJ-R**, Nanjing Jiangsu resistant *B. tabaci* strain; **NJ-S**, Nanjing Jiangsu susceptible *B. tabaci* strain; **OP**, organophosphorus; **PBO**, piperonyl butoxide; **TPP**, triphenyl phosphate
Correspondence: ^a skyning456@163.com, ^b caifengliu2003@njau.edu.cn, ^c yangfangheliang2008@126.com, ^d sldong@njau.edu.cn, ^e zjhan@njau.edu.cn, * Corresponding author
Editor: Nannan Liu was Editor of this paper.
Received: 30 May 2011, **Accepted:** 19 July 2011
Copyright : This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.
ISSN: 1536-2442 | Vol. 12, Number 41

Cite this paper as:

Zhang N-n, Liu C-f, Yang F, Dong S-l, Han Z-j. 2012. Resistance mechanisms to chlorpyrifos and F392W mutation frequencies in the acetylcholine esterase ace1 allele of field populations of the tobacco whitefly, *Bemisia tabaci* in China. *Journal of Insect Science* 12:41 available online: insectscience.org/12.41

Introduction

The tobacco whitefly *Bemisia tabaci* Gennadius (Hemiptera: Aleyrodidae) is a small pest with great agricultural importance worldwide (Bellows et al. 1994). The insects not only feed on leaves resulting in delayed growth and even death of the plants (Liu et al. 2007), but also deposit honeydew on leaves that often lead to sooty mold and reduction in photosynthesis (Ghanim et al. 1998). Additionally, the insect is also known to transmit various plant viruses. With increasing acreage of Bt-transgenic crops such as cotton, which has no resistance to piercing–sucking insects (Ffrench-Constant et al. 2004), the tobacco whitefly problem becomes more serious due to the absence of co–control effects from insecticides previously used to control Lepidopteran pests.

The control of the tobacco whitefly has depended heavily upon synthetic insecticides for decades. As a result, considerable resistance development to a variety of insecticides is very well documented. Understanding of the resistance mechanisms is essential for combating the resistance and improving control efficacy.

In the early days, studies of whitefly insecticide resistance mechanisms were mainly at biochemical and toxicological levels (Hansen and Hodgson 1971; Gunning et al. 1992; Byrne et al. 1995; Gunning et al. 1996; Valles and Woodson 2002). These studies revealed two main mechanisms: reduced penetration and enhanced metabolism of the involved insecticides, the latter playing a more important role. Three groups of enzymes—esterase, glutathione S-transferase (GST), and microsomal-O-demethylase (MFO)—have been proven to be involved in metabolic

resistance. For example, Mouches et al. (1986) showed that an esterase gene is responsible for resistance to a variety of organophosphate (OP) insecticides in *Culex* mosquitoes. GST has been showed to play a major role in detoxification of insecticides in mosquitoes (Huang et al. 1998; Vontas et al. 2001; Vontas et al. 2002).

Acetylcholine esterase (AChE, EC 3.1.1.7), a key enzyme in neurotransmission, is the target of organophosphate and carbamate insecticides. Studies with many insect species indicate that resistance to these two classes of insecticides was associated with reduced sensitivity of AChE to insecticides (Mutero et al. 1994b; Walsh et al. 2001; Li and Han 2002; Weill et al. 2003). AChE genes have been cloned in insects the orders Diptera, Hemiptera, Lepidoptera, Hymenoptera, and others (Mutero et al. 1994a; Zhu et al. 1996; Walsh et al. 2001; Vontas et al. 2002; Nabeshima et al. 2003; Cassanelli et al. 2006). Some AChE gene mutations have been confirmed to associate with insect resistance to organophosphate and carbamate insecticides in multiple insect species including *B. tabaci* (Mutero et al. 1994a; Mutero et al. 1994b; Zhu et al. 1996; Walsh et al. 2001; Li and Han 2002; Weill et al. 2003; Alon et al. 2008; Jiang et al. 2009).

Chlorpyrifos has been used to control tobacco whitefly and other insect pests for many years (Pasteur and Sinègre 1978; Milio et al. 1987; Rust and Reiersen 1991; Archer 1994; Guides et al. 1996; Liu et al. 2005; Curtis and Pasteur 2009), and is still used to some extent in China and other countries. To combat the resistance problem and prolong the utility of this insecticide, it is essential to understand the resistance mechanisms and resistance levels of field tobacco whitefly populations.

This paper reports the biochemical mechanisms associated with a lab selected chlorpyrifos resistant strain and the frequencies of F392W mutated ace1 allele in six field populations from locations covering a wide geographic area across China.

Materials and Methods

Insects

Tobacco whiteflies were initially collected from cotton fields in the suburb of Nanjing, Jiangsu, China in 2005. They were identified as B-biotype by mtDNA COI sequence analysis (Frohlich et al. 1999). A chlorpyrifos resistant strain was derived from them by exposing part of the population to chlorpyrifos using a spray method at a selection dose around LC_{70} . After 26 generations with 22 generations exposed to chlorpyrifos, a resistant strain (NJ-R) was obtained. In the meantime, the rest of the collected population was maintained with no insecticide pressure for 26 generations to obtain a susceptible strain (NJ-S). All insects were reared on caged non-Bt cotton plants at 26 ± 2 °C, 60% RH, and 16:8 L:D photoperiod in the laboratory.

For the F392W mutated ace1 allele frequencies study, six field populations of tobacco whiteflies were collected from locations in Nanjing, Beijing, Wuhan, Guangdong, Guangxi, and Xijiang provinces in 2010 (Table 1). After being brought back to the lab, live adults were immediately transferred to Eppendorf tubes immersed in liquid nitrogen and stored at -75 °C until used in the experiment. The biotypes of these populations were determined by the same method as above.

Insecticides and chemical reagents

Technical grade chlorpyrifos (purity > 95%) and triphenyl phosphate (TPP, reagent grade)

were purchased from Nantong Insecticide Co., Jiangsu, China; diethyl meleate (DEM, reagent grade), p-Nitro anisole (p-NA), and 1-chloro-2,4-dinitrobenzene (CDNB) from Shanghai Chemical Reagent Co. Ltd. (www.richjoint.com); piperonyl butoxide (PBO, reagent grade), NADPH, glutathione (GSH), editic acid (EDTA), 1,4-dithiothreitol (DTT), phenylmethylsulfonyl fluoride (PMSF) phenylthiourea (PTU), and 5,5-Dithiobis (DTNB) disulfide from Sigma-Aldrich (www.sigmaaldrich.com); 1-Naphthyl acetate and fast blue RR salt from Farco Chemical Supplies; and acetylthiocholine iodide (ATChI), bovine serum albumin (BSA), and Triton X-100 from Fluka (www.sigmaaldrich.com). Other chemicals were of analytical quality and purchased from available commercial suppliers.

Leaf-dip bioassay

The susceptibility of all tobacco whitefly populations used in the experiments was determined using a leaf-dip bioassay adopted from Elbert and Nauen (2000). Briefly, chlorpyrifos stock solutions were prepared in acetone and serially diluted to desired concentrations. Cotton leaf discs (35 mm diameter) were dipped for 10 sec in solutions of insecticides containing 0.2 g L^{-1} Triton X-100 as a non-ionic wetting agent. After the surface was air-dried, a leaf disc was placed onto a bed of agar (1.5 g L^{-1} , 10 mm depth) in a plastic dish (35 mm diameter) with the adaxial surface facing downwards. Adult females collected from rearing cages using a pump-powered aspirator were anesthetized with CO_2 , and 25 insects were placed onto each leaf disc. The dishes were sealed with a ventilated lid and stored upside down. Insect mortality was scored after 48 hours. LC_{50} and 95% FL were calculated by Probit regression. The bioassay was conducted at 26 ± 2 °C, 60% RH, and 16:8 L:D photoperiod.

Synergism assay

Synergism was measured using the above described leaf-dip bioassay. Instead of pure water, 100 mg L⁻¹ synergist solutions (TPP, PBO, or DEM) were used as solvents. Control leaf discs were dipped in 100 mg L⁻¹ synergist solutions. Preliminary experiments indicated that 100 mg L⁻¹ synergist solutions had no direct toxicity against tobacco whitefly adults. Mortality was scored after 48 hours. LC₅₀ values were calculated by Probit regression. Synergism ratio (SR) was calculated as LC₅₀ of insecticide alone/LC₅₀ of insecticide with synergist.

Detoxification enzyme activity assays

Esterase. 40 adults from NJ-S or NJ-R were homogenized in an 800 µL ice-cold sodium phosphate buffer (0.2 M, pH 7.8) at 4 °C. The homogenate was then centrifuged at 10,000 g for 20 min. The supernatant was used as an enzyme source. Esterase activity was measured according to Han et al. (1998) by adding 80 µL enzyme source into 120 µL 0.2 M substrate solution containing Fast Blue RR salt in sodium phosphate buffer (0.2 M, pH 7.8) and 1 mM 1-naphthylacetate. Reactions were read by a Versamax kinetic microplate reader (Molecular Devices, LLC, www.moleculardevices.com) recording at 20 sec intervals for 7 min at 450 nm and 27 °C.

MFO. 20 mg of adult tobacco whitefly (mixed sexes) from NJ-S or NJ-R strains was homogenized with 500 µL ice-cold sodium phosphate buffer (0.2 M, pH 7.8, contained EDTA, DTT, PTU, PMSF, and glycerol). The homogenate was then centrifuged at 13,000 g for 30 min at 4 °C. After filtered with glass wool, the supernatant was re-centrifuged at 13,000 g for 20 min, and the supernatant was used as an enzyme source. Enzyme activity was measured by mixing 100 µL enzyme

source with 20 µL 2 mM substrate solution (p-NA) and 10 µL 9.6 mM NADPH. The reaction was read by the microplate reader at 20 sec intervals for 15 min at 405 nm and 27 °C (Hansen and Hodgson 1971).

GSTs. Enzyme source was prepared from 80 adults (mixed sexes) in the same way as MFO with no filtration process after the first centrifugation. Activity was measured by mixing 100 µL enzyme source with 20 µL 1.2 mM substrate solution CDNB and 100 µL 6 mM GSH. The reaction was read by the microplate reader at 20 sec intervals for 10 min at 340 nm and 27 °C (Oppenoorth et al. 1979).

All measurement was done in five replicates. SOFTmax software was used to fit kinetic plots by linear regression. Enzyme activity (V_{max}) was expressed in mOD/min.

AChE kinetics parameters and Ki assay

60 adults (mixed sexes) from NJ-S or NJ-R strain were homogenized with 500 µL ice-cold sodium phosphate buffer (0.2 M, pH 7.6, contained 0.05% Triton X-100). The homogenate was centrifuged at 13,000 g for 10 min at 4 °C, and the supernatant was centrifuged again at 13,000 g for 20 min. The supernatant that resulted from the second centrifugation served as the enzyme source.

AChE kinetics parameters were measured according to the method by Li and Han (2002). The reaction system in volume of 100 µL contained substrate analogue acetylthiocholine (ATChI) at final concentrations ranging from 31.25 µM to 500 µM and DTNB (in buffer solution) at a final concentration of 450 µM. AChE activity was measured at 30 sec interval for 30 min by the microplate reader at 405 nm and 25 °C.

Double reciprocal method was used to obtain K_m and V_m (expressed in mOD/min).

K_i of AChE was determined according to the method reported by Moores et al. (1996). The reaction solution was prepared by mixing 100 μL of enzyme source and 100 μL chlorpyrifos-methyl (50 ppm) at 25 °C. An aliquot of 20 μL reaction solution was taken out every other 20 sec, mixed with 80 μL 0.02 mol L^{-1} sodium phosphate buffer (pH 7.0), 100 μL DTNB (450 μM), and 100 μL ATChI (1.5 mM). The reaction was subsequently read with the microplate reader at intervals of 30 sec for 30 min at 405 nm and 25 °C. All measurements were done in five replicates. SOFTmax software was used to fit kinetic plots by linear regression.

Total protein content of all used enzyme sources was determined by Coomassie brilliant blue method using bovine serum albumin as a standard (Bradford 1976).

Cloning and analysis of ace1 gene fragments

Total RNA was extracted using TRIzol® reagent (Invitrogen, www.invitrogen.com) from 100 adults according to manufacturer instructions. First-strand cDNA was synthesized from the total RNA using ThermoScript™ reverse transcriptase (Invitrogen). PCR for cloning BT-ace1 fragments was performed by LA Taq polymerase (TaKaRa Co. www.takara-bio.com) and 2 × GC Buffer II (TaKaRa Co.) with following parameters: 94 °C for 2 min followed by 40 cycles at 94 °C for 30 sec, 60 °C for 30 sec, 72 °C for 2 min, and one additional cycle at 72 °C for 10 min. The sense and antisense primers used for this PCR were designed from the reported BT-ace1 sequence (ncbi nucleotide: EF675188.1), which are 5'→3'

ATGGACTTCGATCACCTCCCTCTCA and W5'→3'

CGGTGACGAATGACTGGATAAT, respectively. PCR products were separated by agarose gel electrophoresis, purified with AxyPrep™ DNA Gel Extraction Kit (Axygen Biosciences, www.axxygenbio.com), and then cloned into pGEM-T easy vector (Promega, www.promega.com). The ligation reactions were used for transformations with the DH5a competent cells. Positive clones were screened with blue/white and standard ampicillin selection. Recombinant plasmids were fully sequenced by Invitrogen.

For PCR of individual whiteflies, the same method and protocol were used with the following modifications. All reagents used in RNA extraction were in half amounts. PCR conditions were 94 °C for 2 min followed by 40 cycles at 94 °C for 30 sec, 60 °C for 30 sec, 72 °C for 40 sec, and one additional cycle at 72 °C for 10 min, with two primers of F5'→3' CCTTCCTGGACGAGATGCC and R5'→3' CGCCGCACGATGAAGTTGT.

PCR-RFLP assay

The PCR-RFLP assay was adopted from Tsagkarakou et al. (2009). Genomic DNA (gDNA) was extracted from individual adults by DNeasy Blood and Tissue Kit (Qiagen, Germany). The primer pairs used in the PCR were Test-F (5'-TAGGGATCTGCGACTTCCC-3') and Test-R (5'-GTTCAGCCAGTCCGTGTA-3'), by which a 287 bp fragment was amplified. This fragment was fully digested with the restriction enzyme BsrI (MBI Fermentas, www.fermentas.com). As susceptible ace1 allele contains two sites and resistant ace1 allele contains three sites for restriction endonuclease BsrI, digestion of the PCR product with BsrI yields a restriction pattern of three fragments (201, 79, and 7 bp) for the

susceptible *ace1* allele and four fragments (140, 61, 79, and 7 bp) for the resistant *ace1* allele.

Amplifications were performed with approximately 20 ng gDNA in 10x Ex Taq reaction buffer (TaKaRa) with 4 μ L MgCl₂ at a final concentration of 25 mM, 10 μ M of each primer (Test-F and Test-R), 1 μ L, 2.5 mM dNTP 41 μ L and 1.25 U Ex taq. PCR cycling conditions were 94 °C for 5 min, 35 cycles of 95 °C for 15 sec, 52 °C for 30 sec, 72 °C for 40 sec, followed by 72 °C for 10 min. The amplification product was incubated for three hours in a reaction buffer (TaKaRa) with 5 U BsrI. Digested products were electrophoresed using a 3% (w/v) agarose gel. The 7 bp fragments could not be detected, as they were too small to be visualized by electrophoresis. Sixteen to eighteen individuals were examined for each of the field populations of the tobacco whitefly.

Results

NJ-R strain establishment

To investigate the development process of the tobacco whitefly resistance to chlorpyrifos, a resistant strain (NJ-R strain) was selected from a field population in the laboratory (Figure 1). During the course of resistance selection, the LC₅₀ increased slowly but steadily in 1st to 9th generations (from 143.90 ppm to 1458.30 ppm), and afterwards LC₅₀ increased in a much faster pace to reach 4874.10 ppm at the 13th generation. At this point, the selection was stopped for the following four generations, and as a result, the LC₅₀ declined sharply to about 2500 ppm measured in 17th generation. However, with additional selection the LC₅₀ was recovered to 4818.02 ppm at the 21st generation. Continuing selection in the 21st to 26th generation did not result in increasing LC₅₀,

but maintained a value around 4800 ppm. This field collected tobacco whitefly population had a 33.94-fold chlorpyrifos resistance based on the LC₅₀ ratio after facing selections in 22 of the 26 generations. In the meantime, from the part of the same population used for resistance selection, a relative susceptible strain (NJ-S strain) was obtained by maintaining it without exposure to any insecticide for 26 generations. These selected resistant and susceptible strains were further used to explore the resistance mechanisms.

Metabolic enzyme activity of NJ-R and NJ-S strains

To understand the metabolic resistant mechanism involved in the chlorpyrifos resistance of NJ-R strain, the activities of three major metabolic enzymes were measured and compared between NJ-R and NJ-S strains (Table 2). The results indicated that the esterase activity of NJ-R was significantly higher than that of NJ-S (increased 1.53-fold). However, no significant difference was found in GSTs and MFO activities between NJ-R and NJ-S strains.

Synergism of TPP, DEM and PBO

TPP, DEM, and PBO are specific inhibitors of esterase, GSTs, and MFO, respectively. To confirm the results of enzyme activity measurement, the synergisms of TPP, PBO,




Figure 1. Resistance development of NJ-R *Bemisia tabaci* strain selected with chlorpyrifos in a dose of around LC₇₀ in laboratory. LC₅₀s were examined every two or three generations. The selection by chlorpyrifos was stopped from the 13th to 16th generation, and then restored at the 26th generation. High quality figures are available online.

```

      *           20           *           40           *           60
SUD-S : FLDEMPKSLATKNFKKTNILMGSNTEEGNYFIMYYLTDLFRKEENIHVSRDQFIQAVSELNPNYFIVR : 69
R-CK  : FLDEMPKSLATKNFKKTNILMGSNTEEGNYFIMYYLTDLFRKEENIHVSRDQFIQAVSELNPNYFIVR : 69
S-1   : FLDEMPKSLAAKNFKKTNILMGSNTKEGNYWIMHYLTDLFREEENIHVSRDQFIQAVSELNPNYFIVR : 69
S-2   : FLDEMPKSLATKNFKKTNILMGSNTEEGNYFIMYYLTDLFRKEEDIHVSRDQFIQAVSELNPNYFIVR : 69
S-3   : FLDEMPKSLAAKNFKKTNILMGSNTKEGNYWIMHYLTDLFREEENIHVSRDQFIQAVSELNPNYFIVR : 69
S-4   : FLDEMPKSLAAKNFKKTNILMGSNTKEGNYWIMHYLTDLFREEENIHVSRDQFIQAVSELNPNYFIVR : 69
S-5   : FLDEMPKSLAAKNFKKTNILMGSNTKEGNYWIMHYLTDLFREEENIHVSRDQFIQAVSELNPNYFIVR : 69
S-6   : FLDEMPKSLAAKNFKKTNILMGSNTKEGNYWIMHYLTDLFREEENIHVSRDQFIQAVSELNPNYFIVR : 69
S-7   : FLDEMPKSLAAKNFKKTNILMGSNTKEGNYWIMHYLTDLFREEENIHVSRDQFIQAVSELNPNYFIVR : 69
S-8   : FLDEMPKSLAAKNFKKTNILMGSNTKEGNYWIMHYLTDLFREEENIHVSRDQFIQAVSELNPNYFIVR : 69
S-9   : FLDEMPKSLAAKNFKKTNILMGSNTKEGNYWIMHYLTDLFREEENIHVSRDQFIQAVSELNPNYFIVR : 69
S-10  : FLDEMPKSLAAKNFKKTNILMGSNTKEGNYWIMHYLTDLFREEENIHVSRDQFIQAVSELNPNYFIVR : 69
R-1   : FLDEMPKSLAAKNFKKTNILMGSNTKEGNYWIMHYLTDLFREEENIHVSRDQFIQAVSELNPNYFIVR : 69
R-2   : FLDEMPKSLATKNFKKTNILMGSNTEEGNYWIMYYLTDLFRKEENIHVSRDQFIQAVSELNPNYFIVR : 69
R-3   : FLDEMPKSLATKNFKKTNILMGSNTEEGNYWIMYYLTDLFRKEENIHVSRDQFIQAVSELNPNYFIVR : 69
R-4   : FLDEMPKSLATKNFKKTNILMGSNTEEGNYWIMYYLTDLFRKEENIHVSRDQFIQAVSELNPNYFIVR : 69
R-5   : FLDEMPKSLATKNFKKTNILMGSNTEEGNYWIMYYLTDLFRKEENIHVSRDQFIQAVSELNPNYFIVR : 69
R-6   : FLDEMPKSLAAKNFKKTNILMGSNTKEGNYWIMHYLTDLFRKEENIHVSRDQFIQAVSELNPNYFIVR : 69
R-7   : FLDEMPKSLAAKNFKKTNILMGSNTKEGNYWIMHYLTDLFRKEENIHVSRDQFIQAVSELNPNYFIVR : 69
R-8   : FLDEMPKSLAAKNFKKTNILMGSNTKEGNYWIMHYLTDLFRKEENIHVSRDQFIQAVSELNPNYFIVR : 69
R-9   : FLDEMPKSLATKNFKKTNILMGSNTEEGNYWIMYYLTDLFRKEENIHVSRDQFIQAVSELNPNYFIVR : 69
R-10  : FLDEMPKSLAAKNFKKTNILMGSNTKEGNYWIMHYLTDLFRKEENIHVSRDQFIQAVSELNPNYFIVR : 69

```

Figure 2. Examination of F392W point mutation in 10 NJ-S *Bemisia tabaci* (from S-1 to S-10) and 10 NJ-R tobacco whiteflies (from R-1 to R-10) by cloning a 211 bp-*ace1* gene fragment. SUD-S (ncbi protein: ABV45413.1) and R-CK (ncbi protein: ABV45421.1) served as susceptible and resistant references. All 10 NJ-R individuals and 9 out of 10 NJ-S individuals carried the F392W mutation. High quality figures are available online.

and DEM with GSTs and MFO activities on NJ-R and NJ-S strains were determined. The results revealed that TPP had an obvious synergism to chlorpyrifos on both strains, with the synergism ratios of 4.46 and 2.43 in NJ-R and NJ-S strains, respectively. However, DEM and PBO had no significant synergism to chlorpyrifos in both strains (Table 3). The tests confirmed that enhanced esterase activity is at least partially responsible for the observed chlorpyrifos resistance.

Inhibition kinetics of AChE

To explore the target mechanisms of this chlorpyrifos resistance in NJ-R strain, the K_m , V_m , and K_i of AChE preparations from both NJ-R and NJ-S strains are shown in Table 4. There was no significant difference in K_m , V_m , and K_i between NJ-R and NJ-S strains. The results suggested that AChE was not involved in this chlorpyrifos resistance of the NJ-R strain relative to the NJ-S strain of the tobacco whitefly.

Frequencies of F392W mutation in NJ-R and NJ-S

Since F392W mutation in *ace1* gene was already reported to be associated with OP resistance in the tobacco whitefly (Alon et al. 2008), the F392W mutation frequencies in NJ-R and NJ-S strains were determined. *Ace1* fragments were cloned using cDNA template from a pooled sample of 100 insects of NJ-R and NJ-S strains, respectively. Three clones of PCR products from each strain were randomly chosen to be sequenced. Unexpectedly, the reported F392W mutation was present in all six sequenced clones, and no other mutation was found in the *ace1*-1895 bp fragments.

To confirm this result, a PCR assay of individual tobacco whitefly was conducted using primers that are able to amplify a 221 bp fragment across the F392W mutation site in *ace1* specifically. One out of 10 NJ-S individuals showed the sensitive genotype (F392) and the other nine had the F392W mutation, whereas all 10 individuals from NJ-R strain displayed the F392W mutation

(Figure 2). The results indicated that F392W mutation was not responsible for the chlorpyrifos susceptibility difference between NJ-S and NJ-R stains, and that both NJ-S and NJ-R strains had a similar level of target resistance.

Frequencies of F392W mutated *ace1* allele in field population

As it was reported that the F392W mutation in *ace1* gene resulted in the OP resistance of the target insensitivity, frequencies of F392W mutation in *ace1* gene in 6 geographically distinct populations across China were investigated by the PCR-RFLP assay. High frequencies of the mutation were found in all 6 field populations (Table 5). All 18 tested whiteflies from the Beijing population were mutant homozygotes, and most individuals (> 88%) from the other 5 populations also were mutant homozygotes. In addition, 2 of the 16 whiteflies from Guangdong and Xinjiang population were mutant heterozygotes. Among the tested individuals, only 4 individuals (two from Nanjing population, one from Wuhan population, and one from Guangxi population) were wild homozygotes.

Discussion

This study showed that B-biotype tobacco whitefly can develop chlorpyrifos resistance (NJ-R) under continuous selection pressure. The 34-fold laboratory selected resistance involves metabolic mechanisms that confer resistance to a certain extent as the LC_{50} reached a plateau in the selection process. The substantial drop of LC_{50} from the plateau level (33.9 fold) to the level of 16.9 fold in only 4 generations without selection pressure suggested that such metabolic resistance has a high fitness cost. This high fitness cost of the metabolic resistance to chlorpyrifos can partly explain that chlorpyrifos still retains a

relatively higher control efficacy against tobacco whitefly after decades of use in the field. Tobacco whitefly control in China always employs several insecticides with different modes of action, and chlorpyrifos is rarely applied consecutively more than 5 times. This practice curbs the development of the resistance and should be continued.

Metabolic enzyme activity analysis showed that esterase plays a major role in the resistance as no significant difference in GSTs and MFO activities between NJ-S and NJ-R stains was found. Synergism experiments delivered the same conclusion as only TPP resulted in a higher synergism ratio (SR) for NJ-R. This result agrees with Alon et al. (2008), but differs from abamectin resistance in tobacco whitefly and *T. urticae*, where detoxification of MFO and GSTs was indicated as a key factor (Stumpf and Nauen 2002; Wang and Wu 2007). This is not necessarily unexpected, as insecticides of different action modes often induce resistance with different mechanism even in same insect species.

As the target of OPs and carbamates, insensible OP site mutations of AChE have been identified in insects (Mutero et al. 1994a, 1994b; Zhu et al. 1996; Walsh et al. 2001; Vontas et al. 2002; Nabeshima et al. 2003; Cassanelli et al. 2006). Specific to tobacco whitefly, an F392W mutation was shown to be responsible for OPs resistance (Alon et al. 2008). However, AChE kinetic parameters in our study showed no significant difference between NJ-R and NJ-S strains. This result suggests that resistance in NJ-R was not due to AChE site mutations. However, to our surprise, sequence analysis of *ace1*-1895 bp fragments from pooled samples of NJ-R or NJ-S showed consistent F392W mutations compared to the wild type SUS-S strain (ncbi

protein: ABV45413.1). A further analysis with a single tobacco whitefly showed that all 10 tested NJ-R individuals and nine out of 10 NJ-S individuals possessed the F392W mutation. Therefore, the NJ-S strain used in our study was a 'susceptible' strain already carrying the target site resistance. This is indirectly supported by the fact that chlorpyrifos LC_{50} of SUD-S strain in Alon et al. (2008) was much lower (4.57 ppm) compared to NJ-S strain (137.55 ppm) in our study. The results indicated that resistance can have multiple mechanisms for any given insecticide and insect species. Conducting a complete investigation in understanding resistance mechanisms, as shown in this study, is a good method for future research.

The investigation of mutant ace1 gene frequencies in field populations revealed that high frequencies (88-100%) of F392W mutant ace1 allele were found in all six field populations of different biotypes from a wide geographic area of China, and most individuals (92%) were resistant homozygotes. This result indicated that F392W mutant ace1 in *B. tabaci* associated to OP, and carbamate insecticide resistance is widespread; this should be taken into consideration when designing insecticide rotation programs for whitefly management.

As an invasive pest, B-biotype *B. tabaci* was first introduced into China at the end of 1990s. By 2003, it had rapidly spread into 25 provinces and become the dominating biotype. Fast and strong development of insecticide resistance was one of the key factors contributing to this successful invasion and rapid spreading (Liu et al. 2008). Byrne and Devonshire (1993) reported that a large proportion of B-biotype whiteflies in United Kingdom carried insensitive AChE capable of conferring extremely high resistance to OP

and carbamate insecticides. The fact that a similar high level of ace1 mutation frequencies (88-100%) were detected in all six geographically different populations leads us to speculate that this mutation was already present at the time of invasion, and the insecticide selection pressure after invasion had little effect on the mutation frequency because the insecticide use patterns as well as invasion time were different among the six locations where the tested populations were collected. Unfortunately, no baseline data (at the time of invasion) on the mutated ace1 frequency in tobacco whitefly field populations (B- or Q-biotype) are available in China. This speculation remains a hypothesis waiting to be accepted or rejected.

Acknowledgements

This study was supported by the Doctoral Program of Higher Education of China (20060307010) and the Priority Academic Program, Jiangsu Development of Higher Education Institutions (PAPD).

References

-
- Alon M, Alon F, Nauen R, Morin S. 2008. Organophosphates' resistance in the B-biotype of *Bemisia tabaci* (Hemiptera: Aleyrodidae) is associated with a point mutation in an ace1-type acetylcholinesterase and overexpression of carboxylesterase. *Insect Biochemistry and Molecular Biology* 38(10): 940-949.
- Archer T. 1994. Chlorpyrifos resistance in greenbugs (Homoptera: Aphididae): Cross-resistance and synergism. *Journal of Economic Entomology* 87(6): 1437-1440.
- Bellows T, Perring T, Gill R, Headrick D. 1994. Description of a species of *Bemisia* (Homoptera: Aleyrodidae). *Annals of the*

Entomological Society of America 87(2): 195-206.

Bradford MM. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* 72: 248-254.

Byrne FJ, Cahill M, Denholm I, Devonshire AL. 1995. Biochemical identification of interbreeding between B-type and non B-type strains of the tobacco whitefly *Bemisia tabaci*. *Biochemical Genetics* 33(1-2): 13-23.

Byrne F, Devonshire A. 1993. Insensitive acetylcholinesterase and esterase polymorphism in susceptible and resistant populations of the tobacco whitefly *Bemisia tabaci* (Genn.). *Pesticide Biochemistry and Physiology* 45(1): 34-42.

Cassanelli S, Reyes M, Rault M, Carlo Manicardi G, Sauphanor B. 2006. Acetylcholinesterase mutation in an insecticide-resistant population of the codling moth *Cydia pomonella* (L.). *Insect Biochemistry and Molecular Biology* 36(8): 642-653.

Curtis C, Pasteur N. 2009. Organophosphate resistance in vector populations of the complex of *Culex pipiens* L. (Diptera: Culicidae). *Bulletin of Entomological Research* 71(1): 153-161.

Elbert A, Nauen R. 2000. Resistance of *Bemisia tabaci* (Homoptera: Aleyrodidae) to insecticides in southern Spain with special reference to neonicotinoids. *Pesticide Science* 56(1): 60-64.

Ffrench-Constant R, Daborn P, Goff G. 2004. The genetics and genomics of insecticide

resistance. *TRENDS in Genetics* 20(3): 163-170.

Frohlich DR, Torres-Jerez II, Bedford ID, Markham PG, Brown JK. 1999. A phylogeographical analysis of the *Bemisia tabaci* species complex based on mitochondrial DNA markers. *Molecular Ecology* 8(10): 1683-1691.

Ghanim M, Morin S, Zeidan M, Czosnek H. 1998. Evidence for transovarial transmission of tomato yellow leaf curl virus by its vector, the whitefly *Bemisia tabaci*. *Virology* 240(2): 295-303.

Guides C, Nariko R, Dover B, Kambhampati S. 1996. Resistance to Chlorpyrifos-Methyl, Pirimiphos-Methyl, and Malathion in Brazilian and US Populations of *Rhyzopertha dominica* (Coleoptera: Bostrichidae). *Journal of Economic Entomology* 89(1): 27-32.

Gunning R, Balfe M, Easton C. 1992. Carbamate resistance in *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae) in Australia. *Journal of the Australian Entomological Society* 31: 97-103.

Gunning RV, Moores GD, Devonshire AL. 1996. Insensitive Acetylcholinesterase and Resistance to Thiodicarb in Australian *Helicoverpa armigera* Hubner (Lepidoptera: Noctuidae). *Pesticide Biochemistry and Physiology* 55(1): 21-28.

Han Z, Moores G, Denholm I, Devonshire A. 1998. Association between Biochemical Markers and Insecticide Resistance in the Cotton Aphid, *Aphis gossypii* Glover. *Pesticide Biochemistry and Physiology* 62(3): 164-171.

Hansen LG, and Hodgson E. 1971. Biochemical characteristics of insect microsomes. N- and O-demethylation. *Biochemical Pharmacology* 20(7):1569-1578.

Huang H, Hu N, Yao Y, Wu C, Chiang S, Sun C. 1998. Molecular cloning and heterologous expression of a glutathione S-transferase involved in insecticide resistance from the diamondback moth, *Plutella xylostella*. *Insect Biochemistry and Molecular Biology* 28(9): 651-658.

Jiang X, Qu M, Denholm I, Fang J, Jiang W, Han Z. 2009. Mutation in acetylcholinesterase1 associated with triazophos resistance in rice stem borer, *Chilo suppressalis* (Lepidoptera: Pyralidae). *Biochemical and Biophysical Research Communications* 378(2): 269-272.

Li F, Han Z. 2002. Two different genes encoding acetylcholinesterase existing in cotton aphid (*Aphis gossypii*). *Genome* 45(6): 1134-1141.

Liu H, Xu Q, Zhang L, Liu N. 2005. Chlorpyrifos resistance in mosquito *Culex quinquefasciatus*. *Journal of Medical Entomology* 42(5): 815-820.

Liu J, Zhang G, Wan F, Wang J. 2008. Mechanisms of inter- and intra-specific competitive replacement by the *Bemisia tabaci* B biotype (Homoptera: Aleyrodidae). *Biodiversity Science* 16(3): 214-224.

Liu SS, De Barro PJ, Xu J, Luan JB, Zang LS, Ruan YM, Wan FH. 2007. Asymmetric mating interactions drive widespread invasion and displacement in a whitefly. *Science* 318(5857): 1769-1772.

Milio J, Koehler P, Patterson R. 1987. Evaluation of three methods for detecting chlorpyrifos resistance in German cockroach (Orthoptera: Blattellidae) populations. *Journal of Economic Entomology* 80(1): 44-46.

Mouches C, Pasteur N, Berge JB, Hyrien O, Raymond M, de Saint Vincent BR, de Silvestri M, Georghiou GP. 1986. Amplification of an esterase gene is responsible for insecticide resistance in a California *Culex* mosquito. *Science* 233(4765): 778-780.

Mutero A, Bride JM, Pralavorio M, Fournier D. 1994a. *Drosophila melanogaster* acetylcholinesterase: identification and expression of two mutations responsible for cold- and heat-sensitive phenotypes. *Molecular and General Genetics* 243(6): 699-705.

Mutero A, Pralavorio M, Bride JM, Fournier D. 1994b. Resistance-associated point mutations in insecticide-insensitive acetylcholinesterase. *Proceedings of the National Academy of Sciences USA* 91(13): 5922-5926.

Nabeshima T, Kozaki T, Tomita T, Kono Y. 2003. An amino acid substitution on the second acetylcholinesterase in the pirimicarb-resistant strains of the peach potato aphid, *Myzus persicae*. *Biochemical and Biophysical Research Communications* 307(1): 15-22.

Oppenoorth F, Van der Pas L, Houx N. 1979. Glutathione S-transferase and hydrolytic activity in a tetrachlorvinphos-resistant strain of housefly and their influence on resistance. *Pesticide Biochemistry and Physiology* 11: 176-188.

- Pasteur N, Sinègre G. 1978. Chlorpyrifos (Dursban(R)) resistance in *Culex pipiens pipiens* L. from Southern France: Inheritance and linkage. *Cellular and Molecular Life Sciences* 34(6): 709-711.
- Rust M, Reiersen D. 1991. Chlorpyrifos resistance in German cockroaches (Dictyoptera: Blattellidae) from restaurants. *Journal of Economic Entomology* 84(3): 736-740.
- Stumpf N, Nauen R. 2002. Biochemical markers linked to abamectin resistance in *Tetranychus urticae* (Acari: Tetranychidae). *Pesticide Biochemistry and Physiology* 72(2): 111-121.
- Tsagkarakou A, Nikou D, Roditakis E, Sharvit M, Morin S, Vontas J. 2009. Molecular diagnostics for detecting pyrethroid and organophosphate resistance mutations in the Q biotype of the whitefly *Bemisia tabaci* (Hemiptera: Aleyrodidae). *Pesticide Biochemistry and Physiology* 94(2-3): 49-54.
- Valles SM, Woodson WD. 2002. Insecticide susceptibility and detoxication enzyme activities among *Coptotermes formosanus* Shiraki workers sampled from different locations in New Orleans. *Comparative Biochemistry and Physiology C: Toxicology and Pharmacology* 131(4): 469-476.
- Vontas J, Small G, Hemingway J. 2001. Glutathione S-transferases as antioxidant defence agents confer pyrethroid resistance in *Nilaparvata lugens*. *Biochemical Journal* 357(1): 65-72.
- Vontas J, Small G, Nikou D, Ranson H, and Hemingway J. 2002. Purification, molecular cloning and heterologous expression of a glutathione S-transferase involved in insecticide resistance from the rice brown planthopper, *Nilaparvata lugens*. *Biochemical Journal* 362(2): 329-337.
- Walsh S, Dolden T, Moores G, Kristensen M, Lewis T, Devonshire A, Williamson M. 2001. Identification and characterization of mutations in housefly (*Musca domestica*) acetylcholinesterase involved in insecticide resistance. *Biochemical Journal* 359(1): 175-181.
- Wang L, Wu Y. 2007. Cross-resistance and biochemical mechanisms of abamectin resistance in the B-type *Bemisia tabaci*. *Journal of Applied Entomology* 131(2): 98-103.
- Weill M, Lutfalla G, Mogensen K, Chandre F, Berthomieu A, Berticat C, Pasteur N, Philips A, Fort P, Raymond M. 2003. Comparative genomics: Insecticide resistance in mosquito vectors. *Nature* 423(6936): 136-137.
- Zhu K, Lee S, Clark J. 1996. A point mutation of acetylcholinesterase associated with azinphosmethyl resistance and reduced fitness in Colorado potato beetle. *Pesticide Biochemistry and Physiology* 55(2): 100-108.

Table 1. Collection locations, host plants, and biotypes of *Bemisia tabaci* field populations.

Population	Collection location	Host plant	Biotype
Beijing	Beijing municipality	<i>Solanum lycopersicum</i>	Q
Nanjing	Nanjing in Jiangsu province	<i>Gossypium hirsutum</i>	Q
Wuhan	Wuhan in Hubei province	<i>Gossypium hirsutum</i>	Q
Guangdong	Guangzhou in Guangdong province	<i>Cucumis sativus</i>	B
Guangxi	Nanning in Guangxi Zhuang Autonomous Region	<i>Ipomoea batatas</i>	B
Xinjiang	Urumchi in Xinjiang Uygur Autonomous Region	<i>Cucurbita moschata</i>	B

Table 2. Metabolic enzyme activities of NJ-R and NJ-S strains.

Metabolic enzyme	Strain	Activity \pm SE	Ratio
Esterase	NJ-S	27.96 \pm 3.24 *	1.00
	NJ-R	42.73 \pm 3.75	1.53
Glutathione S-transferase	NJ-S	8.94 \pm 1.56	1.00
	NJ-R	9.89 \pm 0.42	1.11
Microsomal-O-demethylase	NJ-S	0.35 \pm 0.07	1.00
	NJ-R	0.35 \pm 0.09	1.00

Enzyme activity: mOD min^{-1} mg protein $^{-1}$. *The esterase activity between NJ-S and NJ-R is significantly different (*t*-test; *p* < 0.05).

Table 3. Synergism of TPP, PBO, and DEM on chlorpyrifos in NJ-R and NJ-S strains of *Bemisia tabaci*.

Treatment	Strain	LC ₅₀ (ppm) \pm SE	SR
Chlorpyrifos	NJ-S	137.55 \pm 8.05	1
	NJ-R	4 884.44 \pm 481.03	1
Chlorpyrifos + TPP	NJ-S	56.41 \pm 10.33	2.43
	NJ-R	1 093.56 \pm 131.28	4.46
Chlorpyrifos + DEM	NJ-S	142.35 \pm 8.99	0.97
	NJ-R	3 853.27 \pm 205.55	1.27
Chlorpyrifos + PBO	NJ-S	131.60 \pm 8.26	1.04
	NJ-R	3 949.52 \pm 530.49	1.24

SR = LC₅₀ of insecticide alone/LC₅₀ of insecticide after synergist.

Table 4. Kinetic parameters and Ki values of AChE from NJ-R and NJ-S strains of *Bemisia tabaci*.

Strain	Km \pm SE	Ratio	Vm \pm SE	Ratio	Ki \pm SE	Ratio
NJ-S	0.20 \pm 0.03	1	0.44 \pm 0.03	1	0.13 \pm 0.001	1.00
NJ-R	0.15 \pm 0.01	0.75	0.30 \pm 0.01	0.69	0.14 \pm 0.006	1.07

Ratio = value/value of NJ-S strain.

Table 5. Genotypes and F392W mutation frequencies in ace1 of six field *Bemisia tabaci* populations from China, detected by the PCR-RFLP assay.

Field population	Total tested (individuals)	RR (individuals)	RS (individuals)	SS (individuals)	Frequency of mutated ace1 allele (%)
Beijing	18	18	0	0	100.00
Nanjing	17	15	0	2	88.23
Wuhan	16	15	0	1	93.75
Guangdong	16	14	2	0	93.75
Guangxi	18	17	0	1	94.44
Xinjiang	16	14	2	0	93.75

RR (mutated homozygote); RS (mutated heterozygote); SS (wild homozygote).