

### The Effects of Temperature on the Development of the Moth Athetis lepigone, and a Prediction of Field Occurrence

Authors: Li, Li-Tao, Wang, Yu-Qiang, Ma, Ji-Fang, Liu, Lei, Hao, Yan-Tang, et al.

Source: Journal of Insect Science, 13(103): 1-13

Published By: Entomological Society of America

URL: https://doi.org/10.1673/031.013.10301

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



## The effects of temperature on the development of the moth Athetis lepigone, and a prediction of field occurrence

Li-Tao Li<sup>1a#</sup>, Yu-Qiang Wang<sup>1,2b#</sup>, Ji-Fang Ma<sup>1c</sup>, Lei Liu<sup>1d</sup>, Yan-Tang Hao<sup>3e</sup>, Chao Dong<sup>3f</sup>, Yao-Jin Gan<sup>1g</sup>, Zhi-Ping Dong<sup>1h</sup>\*, Qin-Ying Wang<sup>4i</sup>\*

Institute of Millet Crops, Hebei Academy of Agriculture and Forestry Sciences, Shijiazhuang, China

<sup>2</sup>Plant Protection Station of Xing Tai City, Xingtai, China

<sup>3</sup>Plant Protection Station of Guantao County, Handan, China

<sup>4</sup>College of plant protection, Agricultural University of Hebei, Baoding, China

<sup>#</sup>These authors contributed equally to this work

#### Abstract

Athetis lepigone (Möschler) (Lepidoptera: Noctuidae) is an important insect pest of corn crops in China. To determine the effect of temperature on A. lepigone growth, and to provide a forecasting model for this pest, the development and fecundity of A. lepigone under five different temperatures (18, 21, 24, 27, 30° C) was investigated, and an experimental population life table was constructed based on the obtained results. The results showed that the duration of development of A. lepigone decreased as the temperature increased from 18 to  $30^{\circ}$  C. Approximately 95% of mature larvae stopped pupating at 18° C, and about 70% of mature larvae stopped pupating at 21° C. When the growth chamber temperature was above 24° C, no growth arrest was observed. The results indicated that the optimum growth temperature of A. lepigone was about 26.47° C. In this study, the highest survival rate, fecundity per female, and population index trend were observed when the temperature was set at 27° C. The percentages of larvae that could spin cocoons after the 5th or 6th instar differed at the different temperatures. The developmental threshold temperatures for A. lepigone eggs, larvae, pre-pupae, pupae, preoviposition females, and the whole generation (i.e., egg to oviposition) were 11.03, 9.04, 15.08, 11.79, 11.63, and 10.84° C, respectively, and their effective accumulative temperatures were 63.51, 339.42, 30.04, 118.41, 35.06 and 574.08 degree-days, respectively. Based on the effective accumulative temperature law, this pest insect can have four generations in most of the Huang-Huai region of China, and two to three generations annually in some cold regions. Athetis *lepigone* may have four generations in the mid-southern part of Hebei Province. This prediction matches the field survey results.

Keywords: developmental duration, developmental threshold temperature, effective accumulative temperature, generation, life table Correspondence: a wangyill126@126.com, b wangyuqiang0202@163.com, c jifangma@sohu.com, d jiaby808@sina.com, e hbgtzbz@126.com, f gtdongchao@126.com, s zhibaoshi001@163.com, h dzping001@163.com, i wqinying@yahoo.com.cn, \*Corresponding authors Received: 17 August 2012 Accepted: 6 December 2012 Published: 6 October 2013 Copyright: This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed. ISSN: 1536-2442 | Vol. 13, Number 103

Cite this paper as:

Li L-T, Wang Y-Q, Ma J-F, Liu L, Hao Y-T, Dong C, Gan Y-J, Dong Z-P, Wang Q-Y. 2013. The effects of temperature on the development of the moth *Athetis lepigone*, and a prediction of field occurrence. *Journal of Insect Science* 13:103. Available online: <u>http://www.insectscience.org/13.103</u>

#### Introduction

Athetis lepigone (Lepidoptera, Noctuidae) (Möschler 1860) is known to occur in many European countries, east-central parts of Russia, Japan, North Korea, and Mongolia (Nowacki et al. 2001; Nikolaevitch et al. 2003; Lindeborg 2008; Poltavsky et al. 2009). It was first reported in Shenyang, China, in 1999 (Chen 1999; Zhang et al. 2009). Because *A. lepigone* was not known to cause severe damage to field crops, this insect was only briefly described (Szőcs et al. 1981; Nieminen et al. 1998). However, *A. lepigone* has caused severe damage to maize crops in China in recent years, but its growth and life cycle characteristics have not been investigated.

In recent years, leaving plant straws in fields and no-tillage seeding have become common practices in China. This new cultivation system has created a suitable ecological environment and rich food source for A. lepigone. Consequently, since 2005 A. *lepigone* has become a major pest in summer corn fields in the Hebei Province. Athetis *lepigone* larvae drill and eat on the corn stems. resulting in the wilting and later death of the plant. It also chews on the aerial parts of corn roots, causing lodging and severe yield loss of corn crops (Jiang et al. 2008). A recent study by Zhu et al. (2012) indicated that, due to the changes in cultivation practices, the population of this pest insect in several provinces, including Shandong, Shanxi. Henan, Anhui, and Jiangsu, increased to a level of outbreak in 2011. In the fields with a serious occurrence of A. lepigone, the average number of larvae per corn plant ranged from 3 to 5, with maximum numbers of about 20 per plant (Jiang et al. 2008). Field surveys conducted in recent years indicated that A. lepigone had three significant emergence peaks (mid-June, mid to late July, and late August to early September) annually in the Hebei Province. The earliest emergence time for adult A. lepigone was in mid-April (Zhang et al. 2011).

Among the all known environmental factors, temperature is the main factor affecting insect growth and reproduction (Zhang 2002). Many studies were made to determine the relationship between temperature. development, and reproduction of agriculture pest insects. The results of these studies have provided useful information for predicting the occurrence of various pests in specific regions, predicting the number of generations each year, and designing control protocols ( Luo and Li 1993; Infante 2000; Fantinou et al. 2004; Feng et al. 2007; Kang et al. 2009; Ju et al. 2011).To elucidate the biological characteristics of A. lepigone, and to establish a forecasting model for the pest in China, the growth duration, the developmental threshold temperature, and the effective accumulative temperature for A. lepigone was investigated.

An experimental population life table and a forecasting model are presented for this important pest.

#### **Materials and Methods**

#### Source of *Athetis epigone*

Athetis lepigone larvae were taken from corn fields in Shijiazhuang, Hebei Province, China, in July 2011, and kept indoors until emergence. Male and female moths were transferred together into insect cages and fed with 10% honey solution. The cages were covered with a layer of gauze during *A. lepigone* spawning. *A. lepigone* was grown inside a laboratory set at  $26 \pm 1^{\circ}$  C,  $50\% \sim 70\%$ RH, and natural daylight. The progeny of the laboratory-reared *A. lepigone* were used in this experiment.

## Observation of developmental duration and data analysis

Over 150 newly-laid eggs were collected and treated with 5% formaldehyde for disinfection. After air-drying, the eggs were placed in plastic containers and incubated in growth chambers (Model E30B, Percival Scientific, www.percival-scientific.com) set at 18, 21, 24, 27 and  $30 \pm 0.5^{\circ}$  C, respectively, and at  $80 \pm 5\%$  RH and 14:10 L:D photoperiod. The total number of eggs hatched under each temperature was recorded, and the developmental duration of each egg (i.e., incubation period) was recorded.

After hatching, 90 larvae were placed individually in wells  $(3.5 \times 1.5 \text{ cm})$  of 6-well, flat-bottomed, tissue culture plates and were fed with fresh maize leaves until larvae matured. If the newly-emerged larvae had a high mortality rate at a temperature, the 1st instar larvae reared at 27° C were supplied to determine the development and survival of immature stages at this temperature. The head

capsule popping off was used to determine larval molting, and the complete formation of a cocoon was considered to be pre-pupae. In this study, the survival rate and the developmental duration of each growth stage were recorded. The survival rate was calculated as:

$$S = \frac{N1}{N2}$$

S is survival rate, N1 is the number of insects that progressed into the next developmental stage, and N2 is the number of insects in the previous developmental stage.

The eggs, larvae, pre-pupae, pupae, and adults were examined and recorded at three different times (8:30, 14:30, and 20:30) each day for their development progresses. The experiment was repeated three times.

In the adult stage, one newly emerged female and one newly emerged male (less than 24 hr old) were paired in an insect cage at each treatment temperature, fed with 10% honey solution, and this was replicated at least five times. Male and female adults were determined based on the appearance of the hair clusters at the ends of their bodies. The female hair cluster is open, and its gonopore is visible. The pleon end of an adult male is semitriangular, and its hair cluster is closed. The number of eggs laid under each temperature condition by each female was counted, and the pre-oviposition period, oviposition period, and longevity of adults were also recorded.

Temperature requirements and the rates of each developmental stage were calculated by using the results from the different treatments. The least squares method was used to calculate the developmental threshold temperature (C) and the effective

accumulative temperature (K) for instars at various stages as described previously (Zhang, 2002). The following equations were used:



Where T represents the temperature, V is the

developmental rate, r is the correlation coefficient, n is the sample number in the indicated temperature group, Sc is the standard error for the developmental threshold temperature, and Sk is the standard error for the effective accumulative temperature.

Within the tested temperature range, the generation survival rate was parabolic. The relationship between the generation survival (Y) rate and temperature (T) was expressed as:  $Y = aT^3 + bT^2 + cT + d$ .

The experimental population life tables were based on formulated the effects of temperature on A. lepigone survival, growth, development, and reproduction. The index of population trend was calculated by using the data shown in the life tables as described previously (Feng et al. 2007). Means and standard errors were determined by using the DPS version 7.05 software (Tang and Feng 2007), and the significant difference (p <0.05) between different treatments was determined by using the single-factor analysis (one-way ANOVA), Duncan's multiple range test (p < 0.05), and correlation analysis (rvalue).

Developm	antal stage		Dev	velopmental du	ration (days)		
Developin	ental stage	18°C	21°C	24°C	27°C	30°C	R
E	gg	$9.10 \pm 0.05$ a	$6.29 \pm 0.04 \text{ b}$	$5.03 \pm 0.06 \text{ c}$	$3.89 \pm 0.10 \text{ d}$	$3.38 \pm 0.05 \text{ e}$	-0.9577
Larva 1st in	1st instar	$8.45 \pm 0.05$ a	$5.82 \pm 0.04 \text{ b}$	$4.10 \pm 0.02 \text{ c}$	$3.21 \pm 0.03$ d	$3.01 \pm 0.03 e$	-0.9420
Larva formed	2nd instar	$7.00 \pm 0.05$ a	$5.97 \pm 0.05$ b	$4.20 \pm 0.03 \text{ c}$	$3.31 \pm 0.03$ d	$2.99 \pm 0.03 e$	-0.9744
	3rd instar	$7.26 \pm 0.08$ a	$6.40 \pm 0.04 \text{ b}$	$4.31 \pm 0.02 \text{ c}$	$3.90 \pm 0.02 \text{ d}$	$3.70 \pm 0.05 \text{ e}$	-0.9443
cocoon at	4th instar	$7.50 \pm 0.05$ a	$5.80\pm0.09~b$	$4.45\pm0.02$ c	$3.65 \pm 0.02 \text{ d}$	$3.30 \pm 0.03 \text{ e}$	-0.9679
Jui instai	5th instar	$7.03 \pm 0.05$ a	$5.90 \pm 0.05 \text{ b}$	$4.35 \pm 0.03 \text{ c}$	$3.91 \pm 0.02 \text{ d}$	$3.20 \pm 0.03 \text{ e}$	-0.9795
	1st instar	8.50 ± 0.14 a	$4.80 \pm 0.05 \text{ b}$	$4.20 \pm 0.03 \text{ c}$	$3.23 \pm 0.04 \text{ d}$	$2.81 \pm 0.04 \text{ e}$	-0.9061
Larva	2nd instar	$7.00 \pm 0.14$ a	$4.49\pm0.04~b$	$3.95\pm0.04~\text{c}$	$3.36 \pm 0.04 \text{ d}$	$2.70\pm0.06~e$	-0.9324
formed	3rd instar	$6.83 \pm 0.22$ a	$4.51 \pm 0.03 \text{ b}$	$3.85\pm0.05~c$	$2.95 \pm 0.04 \text{ d}$	$2.80 \pm 0.03 \text{ d}$	-0.9320
cocoon at	4th instar	6.42 ± 0.17 a	$4.33\pm0.02~b$	$3.83\pm0.04\ c$	$3.38 \pm 0.06 \text{ d}$	$2.63 \pm 0.04 \text{ e}$	-0.9432
6th instar	5th instar	$6.50 \pm 0.14$ a	$4.71 \pm 0.04 \text{ b}$	$4.20\pm0.05~c$	$3.33 \pm 0.04 \text{ d}$	$2.91 \pm 0.04 \text{ e}$	-0.9638
	6th instar	$6.25 \pm 0.14$ a	$4.59 \pm 0.05 \text{ b}$	$4.10\pm0.02$ c	$3.34 \pm 0.05 \text{ d}$	$2.80\pm0.03~e$	-0.9699
La	rva	$37.45 \pm 0.16$ a	$29.03\pm0.24~b$	$22.50\pm0.26\ c$	$18.73 \pm 0.27 \text{ d}$	$16.30\pm0.26\ e$	-0.9980
Pre-	pupa		$5.40 \pm 0.29$ a	$3.20 \pm 0.17 \text{ b}$	$2.33 \pm 0.13$ c	$2.17 \pm 0.18$ c	-0.8854
Pı	ipa		$13.05 \pm 0.46$ a	$9.68\pm0.15~b$	$7.53 \pm 0.13$ c	$6.65 \pm 0.18 \text{ d}$	-0.9708
Pre-ovi	position		$4.00 \pm 0.14$ a	$2.50 \pm 0.76$ b	$2.31 \pm 0.33$ b	$2.00\pm0.40~b$	-0.8860
Femal	e adult		$17.20 \pm 0.41$ a	$13.30\pm0.37~b$	$11.50 \pm 0.45$ c	$8.70 \pm 0.37 \text{ d}$	-0.9896
Male	adult		$18.50 \pm 0.55$ a	$13.90\pm0.40\ b$	$12.30 \pm 0.37$ c	$9.10 \pm 0.51 \text{ d}$	-0.9830
Generatio	on (egg to sition)		57.76	42.90	34.78	30.50	-0.9797

	•	1 0 0	1.00				
Table	2.	Influence of	different	temperatures	on age and	pupation of	t Athetis ledigone.

Tomporature (9C)	Percentage that formed	Percentage that formed	<b>Proportion of mature larvae</b>
Temperature (°C)	cocoons at 5th instar (%)	cocoons at 6th instar (%)	that stopped pupating (%)
18	95.00 ± 2.89 a	$5.00 \pm 2.89 \text{ d}$	$95.00 \pm 2.89$ a
21	$35.00 \pm 0.96 \text{ d}$	65.00 ± 0.96 a	$70.00 \pm 0.96$ b
24	57.89 ± 1.75 c	42.11 ± 1.75 b	
27	63.16 ± 1.75 c	36.84 ± 1.75 b	
30	72.22 ± 1.07 b	27.78 ± 1.07 c	
			a 1160 - 0.00

The data in the table are mean  $\pm$  SE. Different letters in the same column indicate a significant difference at p < 0.05.

Table 3. Developmental threshold temperature and effective accumulative temperature of Athetis lepigone.

Develop sta	omental Ige	Developmental threshold temperature	Effective accumulative temperature	<b>Regression</b> equation	Sc	Sk	R
E	gg	11.03	63.51	T = 11.03 + 63.51V	0.14	2.06	0.9984
Ŧ	1st instar	11.85	51.56	T = 11.85 + 51.56V	0.37	4.59	0.988
Larva	2nd instar	10.56	56.72	T = 10.56 + 56.72V	0.37	5.00	0.988
ionned	3rd instar	7.97	76.14	T = 7.97 + 76.14V	0.66	12.26	0.963
5th instar	4th instar	9.06	67.41	T = 9.06 + 67.41V	0.21	3.35	0.996
Jui ilistai	5th instar	8.65	69.13	T = 8.65 + 69.13V	0.31	5.09	0.991
	1st instar	11.49	50.88	T = 11.49 + 50.88V	0.35	4.25	0.989
Larva	2nd instar	9.76	55.35	T = 9.76 + 55.35V	0.36	4.80	0.988
formed	3rd instar	9.71	53.99	T = 9.71 + 53.99V	0.42	5.43	0.985
cocoon at	4th instar	9.11	56.22	T = 9.11 + 56.22V	0.46	6.17	0.982
6th instar	5th instar	8.18	63.32	T = 8.18 + 63.32V	0.26	3.91	0.994
	6th instar	8.12	62.09	T = 8.12 + 62.09V	0.28	4.09	0.993
La	rva	9.04	339.42	T = 9.04 + 339.42V	0.36	7.83	0.999
Pre-	pupa	15.08	30.04	T = 15.08 + 30.04V	1.23	3.14	0.971
Pu	ipa	11.79	118.41	T = 11.79 + 118.41V	0.92	7.64	0.994
Pre-ovi	position	11.63	35.06	T = 11.63 + 35.06V	2.90	7.12	0.956
Generatio	on (egg to	10.84	574.08	T = 10.84 + 574.08V	0.97	37.20	0.996

Sc is the standard error of developmental threshold temperature. Sk is the standard error of effective accumulative temperature. R is the related coefficient.

#### **Observation of adult occurrence**

*A. lepigone* adults were trapped by using the automatic pest forecasting lamps (JDAO-III, Hebi Jiaduo Science, <u>www.jiaduo.net</u>). The lamps were set up in fields in Guantao County. The observation period was from 1 April to 31 October 2011. The number of trapped adults was recorded daily, and the annual occurrence figure of *A. lepigone* adults was then prepared.

#### Results

#### Development of *Athetis lepigone* under different temperature conditions

The results of this study showed that temperature had significant effects on the developmental durations of *A. lepigone*. For the same developmental stages, the development duration significantly decreased as the temperature was increased from 18 to 30° C (Table 1) (eggs:  $F_{4, 595} = 8145.2$ , p < 0.0001; for larvae spun cocoon after the 5th instar: 1st instar:  $F_{4, 176} = 3223.9$ , p < 0.0001; 2nd instar:  $F_{4, 176} = 1771.3$ , p < 0.0001; 3rd instar:  $F_{4,176} = 1327.8, p < 0.0001$ ; 4th instar:  $F_{4,176}$ = 1905.6, p < 0.0001; 5th instar:  $F_{4, 176}$  = 1712.8, p < 0.0001; for larvae spun cocoon after the 6th instar: 1st instar:  $F_{4,110} = 654.0$ , p < 0.0001; 2nd instar:  $F_{4, 110} = 341.0, p < 0.0001$ 0.0001; 3rd instar:  $F_{4,110} = 418.1$ , p < 0.0001; 4th instar:  $F_{4,110} = 237.7$ , p < 0.0001; 5th instar:  $F_{4,110} = 302.1$ , p < 0.0001; 6th instar:  $F_{4,10}$  $_{110}$  = 316.3, p < 0.0001; larvae:  $F_{4, 291}$  = 4192.7, p < 0.0001; pre-pupae:  $F_{3,231} = 160.2$ , p < 0.0001; pupae:  $F_{3,223} = 782.5$ , p < 0.0001; pre-oviposition:  $F_{3,16} = 5.3$ , p < 0.01; female adult:  $F_{3,16} = 78.7$ , p < 0.0001; male adult:  $F_{3,16}$  $_{16} = 71.2, p < 0.0001;$ ).

The results shown in Table 2 indicate that the instar number of *A. lepigone* was influenced by different temperatures. It also indicates that the *A. lepigone* larvae could have 5 or 6

instars before pupation, and the percentages of 5 and 6 instar larvae varied under the different temperatures. For example, the highest percentage of larvae that formed cocoons after the 5th instar was observed at 18° C, and the highest percentage of larvae that formed cocoons after the 6th instar was observed at 21° C. The percentage of larvae that formed cocoons after the 5th instar increased when the temperature was increased from 21° C to 30° C. Approximately 95% of the population failed to complete their life cycle at 18° C and most of them stopped pupating in the cocoons. When the temperature was increased to 24° C or above, all larvae developed successfully into the adult stage.

#### Developmental threshold temperature and effective accumulative temperature of different developmental stages

The developmental threshold temperatures of *A. lepigone* eggs, larvae, pre-pupae, pupae, and pre-oviposition ranged from 9.04 to  $15.08^{\circ}$  C, and the effective accumulative temperatures were from 30.04 to 339.42 degree days (Table 3). In the larval stage, the developmental threshold temperature was highest for 1st instar larvae (11.85 and 11.49° C). For all developmental stages, the developmental threshold temperature was highest for prepupae (15.08° C). The developmental threshold temperature for the entire generation was 10.84° C, and the effective accumulative temperature of the entire generation was 574.08 degree days (Table 3).

#### The effect of temperature on *Athetis lepi*gone survival

The effects of temperature on the survival of *A. lepigone* varied at different developmental stages. Larvae at the 1st and last instar stages were more susceptible to the influence of temperature. The survival rate of 1st instar larvae was 20.56, 50.48, and 62.49% at 18,



21, and 30° C, respectively (Figure 1). Survival rates increased to over 95% when the temperatures were 24 and 27° C. At 18° C, mature larvae stopped pupating. Within the temperature range of  $21-30^{\circ}$  C, the relationship between the generation survival (Y) and temperature (T) can be expressed as: Y =- 0.0914T<sup>3</sup> +2.57 T<sup>2</sup> +56.076T-1450.7. When the derivative was 0, 26.47° C was the optimum temperature.

## Experimental population life table of *Athetis lepigone*

The experimental population life tables were produced based on the survival rates of the different developmental stages and the adult fecundity data obtained from the different temperature treatments. The number of live eggs under different temperatures was assumed to be 100 (Table 4). The number of instars going into one stage was calculated according to the actual survival rates. The optimum temperatures for pupae and adult of A. lepigone were between 24 and 27° C. At 18° C, both pupae and adult A. lepigone stopped developing. No significant differences were observed between adult sex ratios under various temperature conditions. Fecundity was highest at 27° C (345.15 eggs per female) and showed approximately 17% reduction when the temperatures were reduced to 24 and 21° C, and 12.5% reduction when the temperature was increased to 30° C. The highest popula-

Developmental	Number	Number of individuals at the beginning of different stages							
stage	18°C	21°C	24°C	27°C	30°C				
Egg	100.00	100.00	100.00	100.00	100.00				
1st instar	100.00	100.00	99.50	98.74	96.50				
2nd instar	20.56	50.48	95.75	95.25	60.30				
3rd instar	20.56	50.48	95.75	95.25	60.00				
4th instar	20.53	49.37	95.55	95.25	60.00				
5th instar	20.35	48.46	95.02	95.25	60.00				
6th instar	1.57	32.56	38.05	20.42	18.25				
Pre-pupa	0.00	37.80	92.15	93.34	57.60				
Pupa	0.00	11.34	90.26	90.25	55.75				
Adult	0.00	10.56	85.26	86.36	54.38				
Female-male ratio		0.88:1	0.86:1	0.87:1	0.89:1				
Number of eggs laid per female		287.36	285.40	345.15	304.24				
Total eggs of next generation expected		1420.41	11250.84	13867.50	7790.83				
Population trend index	0.00	14.20	112.51	138.68	77.91				

Table 4. Experimental population life tables of Athetis lepigone at different temperatures.



tion trend index of *A. lepigone* was observed at 27° C, suggesting that under this condition *A. lepigone* can propagate rapidly and may result in an outbreak if food sources are available.

# Estimated generation numbers of *Athetis lepigone* in different geographical regions each year

The data suggest that *A. lepigone* begins to develop when the temperature is above  $10.84^{\circ}$  C. Assuming the regional effective accumulative temperature is the sum of temperature above  $10.84^{\circ}$  C each month and throughout the year, the total effective

accumulative temperature in Guantao, Hebei Province, in 2011 was 2183.34 degree days. Because *A. lepigone* requires approximately 574.08 degree days to complete a whole generation cycle, it was proposed that *A. lepigone* could occur 3.80 generations in Guantao in 2011. Using the same method, it was propose that the estimated generations per year for Zhengding, Zhengzhou, Jinan, Bozhou, and Xuzhou would be 4.14, 4.01, 4.04, 4.20, and 4.16, respectively. For Shenyang, Chengde, Tangshan, and Jincheng, the estimated generation numbers are 2.67, 2.57, 3.51, and 3.23, respectively (Table 5).

As shown in Figure 2, adult *A. lepigone* (overwintered adults) appeared in early April 2011 in the fields of Guantao, and its population remained small until May. A small emergence peak was observed on 30 April. Three obvious emergence peaks were observed in 2011 in Guantao. During the first emergence peak period (mid-June), the number of moths captured by using the insect trapping lamps ranged from 100 to 523 per day. The second peak period was observed between mid and late July, and the largest number of moths captured was 806 per day.

Ta

Place				Month average temperature (°C)								Total effective accumulative	Concration	
Flace	1	2	3	4	5	6	7	8	9	10	11	12	temperature (degree/day)	Generation
Guantao	-4.3	1.0	8.6	15.1	20.0	26.4	27.7	24.9	18.6	14.5	7.6	—	2183.34	3.8
Zhengding	-3.0	1.2	9.6	16.3	21.2	27.3	28.2	26.0	19.5	15.1	7.5	_	2378.74	4.14
Shenyang	-17.5	-5.7	9.0	9.2	17.3	21.3	25.0	24.1	16.5	10.3	5.0	-	1533.88	2.67
Chengde	-10.9	-3.8	2.7	10.6	17.0	22.2	23.7	23.3	16.1	9.4	1.7	-	1474.48	2.57
Tangshan	-7.4	-1.9	6.0	13.2	19.9	24.7	26.2	25.4	19.1	13.2	5.5	—	2015.94	3.51
Jincheng	-5.1	1.0	6.3	15.1	19.2	24.7	25.1	22.6	16.9	12.8	7.5	—	1851.94	3.23
Xuzhou	-2.0	3.3	9.0	16.4	21.3	26.1	27.5	25.7	20.8	15.8	11.1	-	2386.34	4.16
Zhengzhou	-1.3	3.1	9.8	16.4	20.8	26.9	28.1	24.7	18.5	15.6	9.0	-	2299.44	4.01
Jinan	-3.2	2.7	8.9	15.5	21.0	27.3	27.5	24.9	19.5	16.0	9.0	-	2320.64	4.04
Bozhou	-0.8	4.1	9.8	16.9	21.2	26.6	28.0	25.5	20.0	16.5	10.5	_	2412.44	4.20

<b>Fable 6.</b> The theoretical date, the actual occurrence, and the annual generations of Athetis lepigone adults in Guantao.											
	Pasis	Overwin	iter	1st gener	ation	2st generation		3rd generation		Generation per	
	Dasis	Theoretical	Actual	Theoretical	Actual	Theoretical	Actual	Theoretical	Actual	Theoretical	Actual
	Atmospheric temperature	5/2	4/29	6/18	6/14	7/21	7/22	9/2	9/5	3.88	4

The third peak period occurred in late August until early September, with less than 125 moths captured per day. This observation indicates that *A. lepigone* can have four generations per year in Guantao.

To predict the peak period of adult A. lepigone of each generation, the effective accumulative temperature was calculated using the local average temperature provided by the meteorological department. The laboratory observations indicated that, after overwintering, mature larvae could start their pupation just 1 day after incubation at room temperature (data not published). The meteorological data collected in 2011 showed that the daily mean temperatures of 6 days in March were above the developmental threshold temperature for pre-pupae, and 4 of the 6 days had temperatures above 20° C. If the emergence peak of the winter generation adults was predicted using the pupae developmental threshold temperature and the average daily temperature, the number of theoretical A. lepigone generations in Guantao per year would be 3.88 (Table 6). Using the average monthly temperature provided by the local meteorological department, the theoretical generation number in

Guantao would be 3.80, similar to that predicted using the average daily temperature. The predicted adult emergence peaks, using the local temperature data, were on 2 May, 18 June, 21 July, and 2 September for the overwintering, first, second, and third generations, respectively. Thus, the prediction agrees with the field survey results.

#### Discussion

The results of this study show that the developmental threshold temperature for different developmental stage A. lepigone was above 10° C, with the optimum temperatures ranging from 21 to 30° C. Within this temperature range, the developmental duration of each instar decreased as the temperature was increased, indicating a significant negative correlation between the temperature and the developmental duration. Both high and low temperatures had adverse effects on the survival of A. lepigone larvae and the fertility of adults. In addition to the tested temperatures in this experiment, the effect of high temperature on the mortality of larvae and adults was tested. The results showed that the newly hatched larva and adults had a higher mortality rate when the temperature

reached above 33° C, indicating that *A. lepigone* was sensitive to high temperatures (Ma et al. 2012a). This observation is consistent with previously published results (Infante 2000; Ma et al. 2004; Ju et al. 2011; Li et al. 2011a). For example, *Corythucha ciliate* eggs did not hatch at 16° C, and the survival rate of eggs until the adult stage was lowest at 19° C. In addition, most nymphs did not develop further at 36° C (Ju et al. 2011).

It was reported that A. lepigone larvae had 5 (Li et al. 2011b) or 6 instars (Bai et al., 2012). However, we have shown here that the number of instars of A. lepigone was variable, and both 5 and 6 instars could be observed under the same temperature. The percentages of 5th or 6th instar larvae were influenced by temperature. Fan (2008) and others reported that male Pristophora conjugata larvae had 5 instars and female P. conjugata larvae had 6 instars. Under the same growth conditions, Euproctis pseudoconspersa larvae from different geographical populations had different numbers of larval instars (Yin et al. 2011). In our study, the sex and the source of population did not affect the number of larval instars. The significance of having different numbers of larval instars, other factors that determine A. lepigone growth (e.g., photoperiod and nutrition), and which larval instar is the most susceptible to the external influence require further investigation.

Many insects alter their developmental rate, or even suppress a particular developmental stage, to cope with environmental conditions differing from place to place and/or time to time (Danks 2002). Several studies have examined the developmental responses to environmental conditions, especially photoperiod and temperature (Li et al. 2008; Wang et al. 2009). Our study showed that temperature had a significant influence on the development of *A. lepigone*, causing some larvae to stop pupating in their cocoons during maturation at  $18^{\circ}$  C and  $21^{\circ}$  C. Low temperature was the inducing factor for *A. lepigone* dormancy (or diapause). In addition to the detailed developmental responses to temperature, the influence of photoperiod on *A. lepigone* development is currently under the investigation.

Each year, during the overlapping growing season between the wheat and corn crops in the Huang-Huaihai area of China, large amounts of wheat straw and stubble are left in the fields, and these leftover materials create a favorable environment for A. lepigone. In addition. newly-planted summer corn seedlings provide A. lepigone with sufficient food sources. Under suitable temperature conditions, A. lepigone larvae grow well and mature quickly, and the adults are often more fertile, leading to rapid reproduction and then an outbreak of the pest within a short period of time in June, the month with an average temperature of 26° C, in the south-central part of the Hebei Province (Jiang et al. 2008).

Due to irregular developmental patterns, some mature A. lepigone larvae do not produce cocoons, and some pupae survive during the warm winters in some regions. These surviving mature larvae and pupae often represent part of the first generation for the following year (Ma et al. 2012b). The developmental threshold temperature of prepupae is estimated to be 15.08° C. Based on the daily average temperatures in Guantao, it was estimated that the emergence peak of the overwintered A. lepigone would occur in late May. The actual field survey showed, however, that the overwintered A. lepigone adults first appeared in early April and reached the emergence peak in late April in 2011. The possible explanation for this earlier

field emergence peak is that the accumulative temperature of the overwintered generation met the requirement sooner than had been predicted. The weather information obtained indicated that the daily highest temperatures in late May often exceeded 15.08° C. Consequently, a short period of warmer weather had led A. lepigone larvae to pupate sooner than had been predicted. Because the developmental threshold temperature for pupae is lower than that for the other developmental stages, the adult A. lepigone captured in April might have emerged from the overwintered pupae. Therefore, the prediction of the emergence peak of the overwintered adults, based on the developmental threshold temperature for pupae, agreed with the actual field survey results.

The predicted number of generations per year using the developmental threshold temperature and the effective accumulative temperature was similar to that observed in the field using insect trapping lamps. Although the developmental threshold temperature and the effective accumulative temperature presented in this paper were obtained under constant temperature conditions, and the variable temperatures in the field were reported to be more favorable for insect development and reproduction (Cai et al. 2001; Mironidis and Savopoulou-Soultani 2008), the prediction of A. lepigone emergence peaks are still useful. It is noteworthy A. lepigone often survives inside the leftover plant materials in the field, which provide environmental conditions that are quite different from the conditions reported by meteorological data. In addition, the law of effective temperature emphasizes only the effect of temperature on development. Other factors, including food sources, can also have a significant impact on insect development duration, longevity, and fecundity (Sison and Shanower 1994; Shanower et al. 1997; Jalali et al. 2010; Saeed et al. 2010). Thus, feeding characteristics should also be considered when predicting the emergence peaks of *A. lepigone* using this method (Zhang 2002). Different generations of *A. lepigone* have different food sources. The impacts of different food sources on *A. lepigone* growth remain unknown and require further investigation.

The results showed that the A. lepigone generation number, the emergence peaks in a given year, and the duration of individual developmental stages could be predicted using the developmental threshold temperature, effective accumulative temperature, and temperature data provided by the local meteorological departments. This prediction provides useful information on A. lepigone occurrence dynamics and control strategies. According to the field surveys made in recent occurred vears. A. lepigone in four generations annually in the south-central part of Hebei Province and often caused severe damage to the summer maize crops in late June until July. Overlapping between A. lepigone generations was seen frequently in field. The overwintered A. lepigone larvae could develop irregularly, resulting in small numbers of adult A. lepigone observed in the field in April of the following year (Ma et al. 2012c). The first, second, and third generation adults produced three emergence peaks each year (mid-June, mid to late July, and late August until early September). The number of adults captured during the first and second emergence peaks was significantly higher than that captured during the third peak period, with a daily maximum number of more than 1,000. Our investigation showed that A. *lepigone* was distributed unevenly in the field. For example, adults and larvae often stayed in shade to avoid significant temperature

changes caused by direct sun light (Ma et al. 2012c). Although no report is available on A. lepigone damage to the maize crops in the northern part of Hebei Province, this insect was captured previously in Shenyang area using insect killer traps (Zhang 2009). A. lepigone larvae are known to be omnivorous. Indoor feeding surveys showed that this insect could eat leaves, stems, fruits, and even dry plant materials of more than 30 plant species belonging to 13 different families. These plant species include maize, wheat, soybean, peanut, sweet potato, cabbage, tomato. spinach, amaranth, goosefoot, purslane, and more (Ma et al. 2012a). Field investigations also indicated that A. lepigone larvae were able to feed on maize, wheat, sweet potato, soybean, peanut, and several vegetable species (Ma et al. 2012c). Although no report was available on A. lepigone damage to these field obvious crops. it is that favorable environments for this pest are widely present. Close attention should be paid to the population dynamics of this pest in the field.

#### Acknowledgements

This study was supported by the Project of Hebei Department of Science and Technology (11220301D).

#### References

Bai XF, Li GQ, Liu SY, Wang P, Wang YL. 2012. Preliminary study on the biological characteristics of *Athetis lepigone*. *China Plant Protection* 32(1): 31–32. (in Chinese)

Cai WZ, Pang XF, Hua BZ. 2001. *General entomology*. China Agricultural University Press. (in Chinese)

Chen YX. 1999. Fauna Sinica, Lepidoptera: Noctuidae.Science Press. (in Chinese) Danks HV. 2002. The range of insect dormancy responses. *European Journal of Entomology* 99: 127–142.

Fantinou A, Perdikis D, Zota O. 2004. Reproductive responses to photoperiod and temperature by diapausing and nondiapausing populations of *Sesamia nonagrioides* Lef. (Lepidoptera – Noctuidae). *Physiological Entomology* 29: 169–175.

Fan LQ. 2008. Larval instars and duration of *Pristophora conjugate*. *Chinese Bulletin of Entomology* 45(1): 144–145. (in Chinese)

Feng HZ, Wang LY, Dong HQ. 2007. Life tables of the laboratory population of *Coccinella undecimpunctata* L. at different temperatures. *Acta Entomologica Sinica* 50(3): 259–263. (in Chinese)

Infante F. 2000. Development and population growth rates of *Prorops nasuta* (Hym., Bethylidae) at constant temperatures. *Journal of Applied Entomology* 124: 343–348.

Jalali MA, Tirry L, Arbab A, De Clercq P. 2010. Temperature-dependent development of the two-spotted ladybeetle, *Adalia bipunctata*, on the green peach aphid, *Myzus persicae*, and a factitious food under constant temperatures. *Journal of Insect Science* 10:124. Available online: http://www.insectscience.org/10.124/

Jiang JY, Li XQ, Xu YH, Li ZH, Zhang ZY, Xu H. 2008. Preliminary studies on *Athetis* (*Proxenus*) *lepigone*. *Plant Protection* 34(3): 123–126. (in Chinese)

Ju RT, Wang F, Li B. 2011. Effects of temperature on the development and population growth of the sycamore lace bug, *Corythucha ciliate. Journal of Insect Science* 

11:16. Available online: http://www.insectscience.org/11.16/

Kang L, Chen B, Wei JN, Liu TX. 2009. Roles of thermal adaptation and chemical ecology in *Liriomyza* distribution and control. *Annual Review of Entomology* 54: 127–145.

Li HB, Shi L, Lu MX, Wang JJ, Du YZ. 2011a. Thermal tolerance of Frankliniella occidentalis: Effects of temperature, exposure time, and gender. *Journal of Thermal Biology* 36(7): 437–442.

Li WX, Li JC, Coudron TA, Lu XY, Pan WL, Liu XX, Zhang QW. 2008. Role of photoperiod and temperature in diapause induction of endoparasitoid wasp *Microplitis mediator* (Hymenoptera: Braconidae). *Annals of the Entomological Society of America* 101(3): 613–618.

Li YF, Dang ZH, Zhang LJ, Gao ZL. 2011b. Morphological discrimination and development period of *Athetis lepigone* Möschler. *Journal of Hebei Agricultural Sciences* 15(4): 23–24. (in Chinese)

Lindeborg M. 2008. Remarkable records of Macrolepidoptera in Sweden. *Entomologisk Tidskrift* 129(1): 43–52.

Luo LZ, Li GB. 1993. Partition of generation region and effective accumulated temperature of *Loxostege sticticalis*. *Acta Entomologica Sinica* 36(8): 332–338. (in Chinese)

Ma CS, Hau B, Poehling HM. 2004. The effect of heat stress on the survival of the rose grain aphid, *Metopolophium dirhodum* (Hemiptera: Aphididae). *European Journal of Entomology* 101: 327–331.

Mironidis GK, Savopoulou-Soultani M. 2008. Development, Survivorship, and Reproduction of *Helicoverpa armigera* (Lepidoptera: Noctuidae) Under Constant and Alternating Temperatures. *Environmental Entomology* 37(1): 16–28.

Ma JF, Li LT, Wang XY, Gan YJ, Dong ZP. 2012a. The morphological characteristics, habits and injury loss of *Athetis lepigone* larva. *China Plant Protection* 32(5): 16–19. (in Chinese)

Ma JF, Wang YQ, Li LT, Jiang JY, Gan YJ, Dong ZP. 2012b. The field investigation and the developmental stage of *Athetis lepigone* before winter. *China Plant Protection* 32(1): 28–30. (in Chinese)

Ma JF, Wang XY, Li LT, Gan YJ, Dong ZP. 2012c. The occurrence regularity and the control technology of *Athetis lepigone*. *China Plant Protection* 32(5): 26–29. (in Chinese)

Nieminen M, Hanski I. 1998. Metapopulations of moths on islands: a test of two contrasting models. *Journal of Animal Ecology* 67: 149–160.

Nikolaevitch PA, Vjatcheslavovna IE. 2003. The Noctuidae (Lepidoptera) of the Daghestan Republic (Russia) II. *Phegea* 31(4): 167–181.

Nowacki J, Holowinski M, Palka K. 2001 *Athetis lepigone* (Möschler, 1860) (Lepidoptera. Noctuidae), a noctuid moth new for the Polish fauna. *Polskic Pismo Entomologiczne* 70: 271–275.

Poltavsky AN, Matov AY, Ivlievc PP. 2009. Heteroceran moths (Lepidoptera, Heterocera) of the Don River Delta. *Entomological Review* 89(9): 1072–1081.

Saeed S, Sayyed AH, Ahmad <u>I</u>. 2010. Effect of host plants on life-history traits of *Spodoptera exigua* (Lepidoptera: Noctuidae). *Journal of Pest Science* 83: 165–172.

Shanower TG, Yoshida M, Eter JA. 1997. Survival, Growth, Fecundity, and Behavior of *Helicoverpa armigera* (Lepidoptera: Noctuidae) on Pigeonpea and Two Wild *Cajanus* Species. *Journal of Economic Entomology* 90(3): 837–841.

Sison MLJ, Shanower TG. 1994. Development and Survival of *Helicoverpa armigera* (Lepidoptera: Noctuidae) on Short-Duration Pigeonpea. *Journal of Economic Entomology* 87(6): 1749–1753.

Szőcs G, Tóth M, Novák L. 1981. Sex attractants for eight lepidopterous species. *zeitschrift für angewandte entomologie* 91: 272–280.

Tang QY, Feng MG. 2007. DPS Data Processing System: Experimental Design, Statistical Analysis, and Data Mining. Science Press. (in Chinese)

Wang XP, Yang QS, Zhou XM, Xu S, Lei CL. 2009. Effects of phtoperiod and temperature on diapause induction and termination in the swallowtail, *Sericinus montelus*. *Physiological Entomology* 34(2): 158–162.

Yin KS, Tang MJ, Xiao Q. 2011. The difference on larval instars and duration of *Euproctis pseudoconspersa* Strand from different geographic population. *China Tea* 33(3): 14–16. (in Chinese)

Zhang XL, Zhang YG, Li HQ, Liu L, Dong ZP. 2011. Damage Characteristics, Occurrence Regularity and Control Technology of *Athetis (Proxenus) lepigone*. *Journal of Hebei Agricultural Sciences* 15(12): 1–4. (in Chinese)

Zhang XX. 2002. *Insect Ecology and Forecast.* pp. 205–237. China Agriculture Press. (in Chinese)

Zhu YB, Ma JF, Dong L, Li LT, Jiang JY, Li ZH, Dong ZP, Dong JG, Wang QY. 2012. Analysis of genetic polymorphism of *Athetis lepigone* (Lepidoptera: Noctuidae) populations from China based on mtDNA COI gene sequences. *Acta Entomologica Sinica* 55(4): 457–465. (in Chinese)

Zhang ZL, Zhao Y, Ding XY. 2009. *Shenyang insect illustrated handbook*. Liaoning National Publishing House. (in Chinese)