



Composition and Diversity Analysis of the Gut Bacterial Community of the Oriental Armyworm, *Mythimna separata*, Determined by Culture-Independent and Culture-Dependent Techniques

Authors: He, Cai, Nan, Xiaoning, Zhang, Zhengqing, and Li, Menglou

Source: Journal of Insect Science, 13(165) : 1-11

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.013.16501>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Composition and diversity analysis of the gut bacterial community of the Oriental armyworm, *Mythimna separata*, determined by culture-independent and culture-dependent techniques

Cai He^{1,2a}, Xiaoning Nan^{1b}, Zhengqing Zhang^{1c}, and Menglou Li^{1d*}

¹College of Forestry, Northwest A&F University, Yangling, Shaanxi, 712100, P. R. China

²Wuwei Academy of Forestry Sciences, Wuwei, Gansu, 733000, P.R. China

Abstract

The intestinal bacteria community structure and diversity of the Oriental armyworm, *Mythimna separata* (Walker) (Lepidoptera: Noctuidae), was studied by analysis of a 16S rDNA clone library, denaturing gradient gel electrophoresis, and culture-dependent techniques. The 16S rDNA clone library revealed a bacterial community diversity comprising Cyanobacteria, Firmicutes, Actinobacteria, Gracilicutes and Proteobacteria, among which *Escherichia coli* (Migula) (Enterobacteriales: Enterobacteriaceae) was the dominant bacteria. The intestinal bacteria isolated by PCR-denaturing gradient gel electrophoresis were classified to Firmicutes, Proteobacteria, and Gracilicutes, and *E. coli* was again the dominant bacteria. The culture-dependent technique showed that the intestinal bacteria belonged to Firmicutes and Actinobacteria, and *Staphylococcus* was the dominant bacteria. The intestinal bacteria of *M. separata* were widely distributed among the groups Cyanobacteria, Firmicutes, Actinobacteria, Gracilicutes, Proteobacteria, and Gracilicutes. 16S rDNA clone library, denaturing gradient gel electrophoresis, and culture-dependent techniques should be integrated to obtain precise results in terms of the microbial community and its diversity.

Abbreviations: DGGE, denaturing gradient gel electrophoresis

Keywords: 16S rDNA, intestinal bacteria, PCR-DGGE

Correspondence: ^a brittany0825@hotmail.com, ^b nanningqi2008@yahoo.com.cn, ^c zhangzhengqing0802@163.com, ^d limenglou@126.com, *Corresponding author

Received: 22 May 2012 **Accepted:** 15 July 2012 **Published:** 31 December 2013

Copyright: This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 13, Number 165

Cite this paper as:

He C, Nan X, Zhang Z, Li M. 2013. Composition and diversity analysis of the gut bacterial community of the Oriental armyworm, *Mythimna separata*, determined by culture-independent and culture-dependent techniques. *Journal of Insect Science* 13:165. Available online: <http://www.insectscience.org/13.165>

Introduction

The Oriental armyworm, *Mythimna separata* (Walker) (Lepidoptera: Noctuidae), is a typical seasonal migrating pest in China that causes heavy losses to agricultural production and food security (Chen et al. 1995; Lee et al. 1995). For efficient control of *M. separata*, a lot of research on their biological and ecological habits has been performed, but little has been done relative to its population genetics and molecular ecology (Zhang and Zhai 2008). *Mythimna separata* is widespread in China and other countries and has been used as a test insect to explore new agricultural pesticides. For instance, the bioactivity of anthradin (Zhang et al. 2003), coriaria lactone (Guo et al. 2007), tutin (Zhuang et al. 2007), *Datura stramonium* (Zhou et al. 2008), and celangulin V (Lü et al. 2008) have been measured based on *M. separata*. Recently, molecular biological approaches to the study of insect intestinal bacteria have received a great deal of attention, including studies on *Reticulitermes speratus* (Hongoh et al. 2003), *Ixodes ricinus* (Claudia et al. 2003), *Apriona germari* (Zhang et al. 2004), *Hepialus gonggaensis* (Liu et al. 2001; Zhuo et al. 2005), *Bombyx mori* (Xiang et al. 2007), and *Apis mellifera*, *Vespula vulgaris*, *Vespa crabro*, and *Nauphoeta cinerea oliv* (Mrazek et al. 2008). The insect guts may represent a large source of microbial diversity. These bacteria utilize a wide range of organic polymers and can be involved in methanogenesis and nitrogen fixation (Nardi et al. 2002). The gut microflora also play an important role in pheromone production, pesticide degradation, vitamin synthesis, and pathogen prevention (Reeson et al. 2003). In addition, these bacteria also take part in resisting the invasion and propagation of microbes and strengthening the function of the immune

system (Dillon et al. 1995, 2002a, 2002b; Tokuda et al. 1997). Therefore, analysis of the composition and diversity of the intestinal bacteria of *M. separata* have application value in pest biocontrol and new pesticide exploration.

Bacterial communities in insect guts have been studied mainly by the cultivation-dependent technique, which does not reflect the entire microbial communities (Gilliam et al. 1997). 16S rDNA genes, which exist in all nuclear biological cells, have been widely used to estimate the diversity of the insect gut bacterial microbiota (Reeson et al. 2003; Broderick et al. 2004). However, little information is available on the gut bacterial communities of many insects, including *M. separata*. In our study, a detailed analysis of bacterial diversity and community structure in the gut of the *M. separata* is reported, through the construction of the 16S rDNA clone library, PCR denaturing gradient gel electrophoresis (DGGE), and culture-dependent techniques. The objective of this work was to demonstrate the bacterial community structure and diversity in *M. separata*, which would help to provide a novel method of pest biocontrol and to explore potential pesticides as well.

Materials and Methods

Sample collection

The *M. separata* were purchased from Northwest A&F University biorational pesticide research center, China. Corn leaves were used for food, and the insects were maintained at $22 \pm 1^\circ \text{C}$. The isolated guts of 10 starved *M. separata* were put on ice and gently crushed using a pestle in liquid nitrogen. The entire experiment was repeated three times.

DNA extraction and 16S rDNA amplification

The DNA extraction of culture bacteria and intestinal total bacteria was performed according to the procedure described by Yuan et al. (2009). The quality of genomic DNA was analyzed on 1.0% agarose gel, followed by staining with ethidium bromide.

The universal primers 27mf (5'- AGA GTT TGA TCM TGG CTC AG -3') and 1492r (5'- ATG GGY TAC CTT GTT ACG ACT T -3') (Weisburg et al. 1991) were used to amplify the full-length 16S rDNA gene of the culturable bacteria. 357f-GC (5'- GC-clamp- CCT ACG GGA GGC AGC AG -3) and 517r (5'- ATT ACC GCG GCT GCT GG -3') were used to amplify the 16S rDNA V3 region for DGGE, 341F (5'- CCT ACG CGA GGC AGC AG -3') and 517r (5'- ATT ACC GCG GCT GCT GG -3') (Muyzer et al. 1993) for the construction of a cloning library. The amplification was performed in a 50 μ L reaction mixture consisting of 25 μ L 2x ES Taq MasterMix (CWBio, www.cwbio.com), 2 μ L each primer, 2 μ L genomic DNA, and 19 μ L RNA-free water. The PCR reaction was performed in a DNA Engine Dyad Peltier Thermal Cycler (Bio-Rad, www.bio-rad.com). The PCR amplification conditions were as follows: predenaturing at 94° C for 5 min, followed by 29 cycles of 94° C for 1 min, 55° C for 1 min, 72° C for 1.5 min, and a final extension at 72° C for 10 min. Finally, the amplification product was purified using a universal DNA purification kit (TIANGEN, www.tiangen.com).

Isolation and culture of the intestinal bacteria

Guts of ten healthy *M. separata* larvae were homogenized in 1 mL sterilized distilled water, and then serial 8-fold dilutions were

made, also using sterilized distilled water. Aliquots (0.1 mL) of the 10^{-3} to 10^{-5} dilutions were spread onto conventional agar plates (potato dextrose agar, nutrient agar medium, beef-extract-peptone-dextrose, Miller and Schroth medium, and glucose ammonium salt culture medium). The plates were cultivated for 48 hr at 37° C. Selected different bacteria strains were pure cultured for 48 hr at 37° C, then extracted DNA from pure cultured strains was selected for PCR and sequencing.

Construction of 16S rDNA library

The intestinal total bacteria DNA of *M. separata* was amplified and purified according to the above DNA extraction and amplification process. PCR products were cloned into pUCm-T according to the manufacturer's instructions (Sangon, www.sangon.com) and then transported into competent cells (SK2301, Sangon). Plasmid insertions were screened following the manufacturer's instructions. Segment size was checked by PstI single enzyme cut and agarose gel electrophoresis to confirm whether the target fragment was present or not. The cloning vector was sequenced by the universal primer m13.

DGGE analysis

DGGE was performed with the Dcodek Universal Mutation Detection System (Bio-Rad), according to the manufacturer instructions. DGGE of the amplified 16S rDNA gene was performed with 8% acrylamide gel containing a denaturant gradient of 35–60% (100% defined as urea 7 M, 40% deionized formamide). The total of PCR products was concentrated to a 30 μ L final volume and loaded on gel, which was electrophoresed at a constant voltage of 80 V for 1 hr, 60 V for 16 hr, and 60° C in 1 TAE buffer (40 mM Tris, 20 mM acetate, 1.0 mM Na₂-EDTA).

After electrophoresis, DGGE gel was stained with ethidium bromide and photographed with UV transillumination. Then, characteristic bands were excised from the gel and dissolved with sterilized distilled water at 4° C for a night. The supernatant after centrifugation (10000 rpm, 5 min, 4° C) was used as a DNA template for 16S rDNA-V3 amplification using the same primers without a GC-clamp. The PCR products were analyzed by electrophoresis in 1.5% agarose gel. Finally, the amplification products were purified using a universal DNA purification kit and sequenced.

Phylogenetic sequence analysis

The sequences were compared with known sequences listed in the GenBank nucleotide sequence databases. The BLAST search op-

tion of the National Center for Biotechnology Information (NCBI) (GenBank ID: BA123456) (<http://www.ncbi.nlm.nih.gov>) was used to search for close evolutionary relatives in the GenBank database (Altschul et al. 1997). A neighbor-joining tree of the aligned sequences was constructed using MEGA 5.0 (Kumar et al. 2001).

Results

Cultured bacterial community of *M. separata*

In our study, 30 bacterial strains were isolated from five different culture mediums, and 20 16S rDNA sequences were obtained with the BLAST search from the database of the National Center of Biotechnology Information. Four bacterial strain sequences were the same as the strain B-5, and one was the same as the N-4 strain, so these strains are not seen in Figure 1. The cultured bacterial sequences obtained in our study are available in the GenBank nucleic acid sequence database under the accession numbers JN995574-JN995588. The results of our study were compared the identities of these bacterial strain sequences to the reported sequences in the NCBI database, and they were over 99% similar, except M-3, which only had 97% similarity with *Bacillus pumilus* Meyer and Gottheil (Bacillales: Bacillaceae). Four different phylotypes were isolated. *Staphylococcus* was the dominant component (9/20, 45%). The other strains belonged to the genera *Bacillus*, *Arthrobacter*, and *Enterococcus*. The identities of these strain sequences reported in the NCBI database were over 99%. The neighbor-joining method of MEGA 5 was used to construct a phylogenetic tree (Figure 1). The results indicated that *Staphylococcus saprophyticus* (Fairbrother) (Bacillales: Staphylococcaceae) (7/9, 77.78%), *B. pu-*




Figure 1. The phylogenetic tree of *Mythemna separata* intestinal bacteria obtained using the culture-dependent method. The following reference species were used: *Staphylococcus saprophyticus*: B-5, B-7, B-8; *Staphylococcus equorum*: B-6; *Staphylococcus sciuri*: M-2; *Arthrobacter*: G-2; *Bacillus pumilus*: N-3, N-7, N-8, M-3; *Enterococcus faecium*: B-4; *Enterococcus mundtii*: N-2, N-4, N-6; *Bacillus safensis*: P-1. High quality figures are available online.

Table 1. The intestinal bacteria 16S rDNA sequences identified by the clone library.

Group	Accession No.	Affiliation	Closest sequence	Identity (%)
NC-43	JQ013053	Actinobacteria	<i>Frigoribacterium</i> sp. (GU585010.1)	100
NC-52	JQ013058	Firmicutes	<i>Enterococcus mundtii</i> (GU372705.1)	100
NC-61	JQ013051	Gracilicutes	<i>Pseudomonas aeruginosa</i> (HM149547.1)	100
NC-85	JQ013052	Gracilicutes	<i>Pseudomonas veronii</i> (HQ824976.1)	100
NC-42	JQ013050	Proteobacteria	<i>Beta proteobacterium</i> (JN371609.1)	100
NC-62	JQ013056	Proteobacteria	<i>Beta proteobacterium</i> (GU213427.1)	100
NC-18	JQ013046	Proteobacteria	<i>Escherichia coli</i> (JN578647.1)	100
NC-19	JQ013047	Proteobacteria	<i>Escherichia coli</i> (JN578647.1)	100
NC-22	JQ013049	Proteobacteria	<i>Escherichia coli</i> (JN578647.1)	100
NC-45	JQ013048	Proteobacteria	<i>Escherichia coli</i> (JN578647.1)	100
NC-47	JQ013042	Proteobacteria	<i>Escherichia coli</i> (GU132179.2)	100
NC-48	JQ013043	Proteobacteria	<i>Escherichia coli</i> (JF508253.1)	100
NC-53	JQ013045	Proteobacteria	<i>Escherichia coli</i> (JN578647.1)	100
NC-69	JQ013059	Proteobacteria	<i>Escherichia coli</i> (JN578645.1)	100
NC-71	JQ013044	Proteobacteria	<i>Escherichia coli</i> (JN578664.1)	100
NC-54	JQ013054	Proteobacteria	<i>Ochrobactrum anthropi</i> (U88444.1)	100
NC-78	JQ013057	Proteobacteria	<i>Ralstonia pickettii</i> (FR873796.1)	100
NC-15	JQ013040	Uncultured bacterium	Uncultured bacterium clone CO1July A8 (JN684317.1)	99
NC-56	JQ013055	Uncultured bacterium	Uncultured bacterium clone BULK-16 (JN161843)	100
NC-92	JQ013041	Uncultured bacterium	Uncultured bacterium isolate DGGE band he-24 (FJ235654.1)	100

milus (3/4, 75%) and *Enterococcus mundtii* Collins (Lactobacillales: Enterococcaceae) (4/5, 80%) were the dominant species.

Different bacterial culture strains were isolated from five bacterial culture mediums. The results showed that more bacteria, including *Enterococcus*, *Bacillus*, *Staphylococcus*, were isolated from NA and Miller and Schroth media than other media. *Staphylococcus* and *Enterococcus* were isolated from the beef-extract-peptone-dextrose medium. *Arthrobacter* and *Bacillus* were isolated from both the glucose ammonium salt culture medium and the potato dextrose agar medium.


Construction of 16S rDNA gene clone library

The 16S rDNA gene clone library consisted of 100 clones from *M. separata*. Twenty typical clones were selected to sequence. According to the sequences, intestinal bacteria were classified into four domains: Firmicutes, Actinobacteria, Gracilicutes, and Proteobacteria. Thirteen clones were clustered to Proteobacteria and nine clones of those were closely related to the genus *Escherichia*. Four clones of these had a high degree of similarity (100%) to the previously sequenced organisms *Ralstonia pickettii*


(clone NC-78), *Ochrobactrum anthropi* (clone NC-54), and *Beta-proteobacteria* (clones NC-42 and NC-62). Firmicutes and Actinobacteria had one clone each, *Enterococcus mundtii* and *Frigoribacterium*, respectively. Three clones had high identities with uncultured bacterium sequences from the NCBI database (clones NC-56, NC-15, and NC-92) (Table 1). All the clone sequences obtained in our study have been assigned to the GenBank nucleic acid sequence database, with accession numbers JQ013040- JQ013059.

DGGE pattern and phylogenetic analysis of the intestinal bacteria of *M. separata*

For phylogenetic identification, sequences were compared with 16S rDNA sequence information of known bacteria listed in the GenBank databases. The predominant DGGE bands of *M. separata* intestinal bacteria are shown in Figure 2, and the results of phylogenetic analysis are shown in Figure 3, revealing the presence of a variety of different genera. Eight bacterial sequences were divided into three main groups through neighbor joining analysis: Firmicutes, Proteobacteria, and Gracilicutes. Similarity values were between 99% and 100% to *Enterococcus* (band 3), *Pantoea* (band 8), *Pseudomonas* (band 2), *Escherichia* (band



5), and *Staphylococcus* (band 9). Band 7 was the most closely related (91%) to *Pseudomonas*, bands 4 and 6 had highly similar identities (100%) to uncultured bacterium sequences from the NCBI database (FJ490436.1 and HM115934.1). All the DGGE sequences obtained in this study have been assigned to the GenBank nucleic acid sequence database, with accession numbers JQ039351- JQ039358.



Compared *M. separata* intestinal bacteria gained by different methods

Eleven known bacterial genera were isolated by three methods in this experiment. Table 2 shows that the construction gene library of bacterial species was detected to be the highest percentage, accounting for 72.72% (8/11). DGGE accounted for 45.45% (5/11), and culture-dependent accounted for 36.36% (4/11). The common bacteria determined by construction of the gene library, DGGE separation, and conventional culture was *Enterococcus*. *Bacillus*, *Arthrobacter*, and *Staphylococcus* were only isolated by culture-dependent methods, while *Escherichia*, *Beta proteobacterium*, *Frigoribacterium*, *Ochrobactrum*, *Pantoea*, *Pseudomonas*, and *Ralstonia* were detected by molecular biolo-

Table 2. Comparison of intestinal bacteria gained by different methods.

Genera	Clone	Culture-Dependent	DGGE
<i>Arthrobacter</i>	-	+	-
<i>Bacillus pumilus</i>	-	+	-
<i>Bacillus safensis</i>	-	+	-
<i>Beta proteobacterium</i>	+	-	-
<i>Enterococcus faecium</i>	-	+	-
<i>Enterococcus mundtii</i>	+	+	+
<i>Escherichia sp.</i>	+	-	-
<i>Escherichia coli</i>	+	-	+
<i>Frigoribacterium sp.</i>	+	-	-
<i>Ochrobactrum anthropi</i>	+	-	-
<i>Pantoea ananatis</i>	-	-	+
<i>Pseudomonas aeruginosa</i>	+	-	+
<i>Pseudomonas veronii</i>	+	-	-
<i>Ralstonia pickettii</i>	+	-	-
<i>Staphylococcus equorum</i>	-	+	-
<i>Staphylococcus saprophyticus</i>	-	+	-
<i>Staphylococcus sciuri</i>	-	+	-
<i>Staphylococcus xylosus</i>	-	+	-
<i>Staphylococcus gallinarum</i>	-	-	+

+ = the detected genera; - = the undetected genera

gy methods (PCR-DGGE and construction of the gene library) (Table 2).

Discussion

The culture-independent nucleic acids-based techniques, such as DGGE and clone library techniques, were used to reveal the intestinal bacteria community structure of *M. separata*. Eleven genera of intestinal bacteria in *M. separata* were isolated in this study. Different intestinal bacteria were obtained by three methods. Four genera of intestinal bacteria were gained by culture-dependent techniques, among which *Staphylococcus* was the predominant bacteria. *Escherichia* was the predominant genus among the six genera of intestinal bacteria isolated by constructing the clone library and the five genera of intestinal bacteria isolated by DGGE. The intestinal bacteria of *M. separata* obtained from construction of the gene library were greater in number than the number obtained using the DGGE and culture-dependent methods, indicating that the construction of the gene library revealed higher bacterial diversity than the DGGE and culture-

dependent methods. Thus, the construction of a gene library could identify the most bacteria strains, while DGGE isolated the predominant bacteria, and the culture-dependent method was useful for identifying a few bacteria. Using DGGE or a culture-dependent technique alone reduces the integrity of the experimental results and results in a lower diversity of bacteria. Moreover, the three methods showed different predominant bacteria. The difference in results is because the culture-dependent technique inevitably screens for microorganisms. For instance, some of the bacteria present in low amounts would grow rapidly and become the dominant bacteria in a short time, while some highly abundant bacteria could not be identified, because the artificial separation culture conditions are not suitable for their growth. In addition, most of the highly abundant bacteria gained by molecular biology methods cannot be isolated or cultured by traditional bacteria culture methods, and bacteria isolated by culture-dependent techniques often exhibited low abundance when molecular biology methods are used.

DGGE band 1 was identified as 18S rDNA of *M. separata*. Comparative sequence analysis revealed that the universal primer 517r completely annealed to the *M. separata* 18S rDNA. Although primer 341f is a bacterial-specific 16S rDNA primer, and comparison to the *M. separata* 18S rDNA sequence revealed four unannealed base pairs, 18S rDNA was amplified with the primer set 341f and 517r under the given PCR conditions (Claudia et al. 2003). Nested PCR and touch down PCR amplification could increase the specificity of the 16S rDNA amplification and circumvent this problem.

Most of the bacteria that were isolated in this study have been found in other insects. Spe-

cies of the genus *Staphylococcus*, which are gram-positive bacteria and typically unpathogenic, can often be found in the gut of insects. Eutick et al. (1978) reported the intestinal bacteria of 10 kinds Australia termites and found that *Staphylococcus* was the predominant bacteria. *Staphylococcus* was also isolated from the gut of *Apriona germari* with traditional culture-dependent techniques (Zhang et al. 2004), from *Hepialus gonggaensis* by culture-dependent and culture-independent techniques (Zhuo et al. 2005), and from *Bombyx mori* using library sequence analysis (Xiang et al., 2007). Therefore, *Staphylococcus* is the normal intestinal bacteria of many insects. From the point of view of animal micro-ecology, the nature, amount, and location conditions of the normal gut microflora may have a very close relationship to insect nutrition and digestive physiology (He et al. 1994). Lots of research has been done on the mechanism of action in *Bacillus*, *Pseudomonas*, *Escherichia*, and *Enterococcus*. Mead et al. (1988) studied one kind of grasshopper (*Melanoplus sanguinipe*) intestinal *Enterococcus* and found that many *Enterococcus* species could produce acetate, and speculated that the accumulation of a large amount of acetate reduced the intestinal pH of the grasshopper. Broderick et al. (2004) surveyed the intestinal bacteria of *Lymantria dispar* larvae and found that *Enterococcus* was the dominant bacterium; furthermore, the authors suggested that reducing intestinal pH could protect the host from toxins. Prescott et al. (2002) found that the common characters of *Pseudomonas* and *Escherichia* species were organic nutrition and facultative anaerobes, and that they could degrade carbohydrates via glycolysis and the pentose phosphate pathway.

Yang et al. (2010) found *Bacillus*, *Pseudomonas*, *Pantoea*, and *Proteus* in the gut of *Apis mellifera ligustica*. Both He et al. (2001) and Zhang et al. (2004) found *Staphylococcus*, *Bacillus*, *Escherichia*, and *Proteus* in the gut of *Apriona germari*. Broderick et al. (2004) found *Bacillus*, *Pseudomonas*, *Pantoea*, *Staphylococcus*, and *Enterococcus* in the gut of *Lymantria dispar*. So, we speculated that the gut bacterial community composition of *M. separata* would have many similarities with insects of Hymenoptera, Coleoptera, and Lepidoptera. These insects have different living conditions, habits and food, but *Bacillus*, *Pseudomonas*, *Pantoea*, *Staphylococcus*, *Enterococcus*, *Proteus*, and *Escherichia* were found in *M. separata* guts. This indicated that these intestinal bacteria have no relationship with feeding methods and that they are the intrinsic gut bacteria of insects.

As shown in our study, three methods were used to identify bacterial diversity of *M. separata*. Nevertheless, as with each method, the applied approach is not free from bias. Limitations such as preferential amplification of 16S rDNA of some bacterial taxons (Farrelly et al. 1995; Polz and Cavanaugh 1998) and identical electrophoretic migration of sequences with multiple differences lead to an underestimation of bacterial diversity in DGGE community fingerprints (Jackson et al. 2000). In addition, DGGE bands have short sequences (about 150 bp in the present study, which may have affected the resolution of the taxa on the plots and may limit the available analysis (Asakawa and Kimura 2008)). Clone library analysis can identify more bacterial strains, but are not suitable for handling multiple samples due to heavy workload and high cost. Culture-dependent methods cannot reveal bacterial community effectively,

because most of bacteria have strict living conditions. Therefore, culture-independent and culture-dependent methods should be combined so that the composition of microbial community structure and diversity can be better revealed.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (No. 31170608). We thank professor T.-X. Liu, Northwest A&F University, for assisting with PCR-DGGE techniques.

References

Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research* 25: 3389–3402.

Asakawa S, Kimura M. 2008. Comparison of bacterial community structures at main habitats in paddy field ecosystem based on DGGE analysis. *Soil Biology & Biochemistry* 40:1322–1329.

Broderick NA, Raffa KF, Goodman RM, Handelsman J. 2004. Census of the bacterial community of the gypsy moth larval midgut by using culturing and culture-independent methods. *Applied and Environmental Microbiology* 70(1): 293–300

Chen RL, Sun YJ, Wang SY. 1995. Migration of the oriental armyworm *Mythimna separata* in East Asia in relation to weather and climate. In: Drake VA, Gatehouse AG, Editors. *Insect migration: tracking resources through space and time*. pp. 93–104. Cambridge University Press.

Claudia SG, Werner L, Sabine R. 2003. Application of broad-range 16S rRNA PCR amplification and DGGE fingerprinting for detection of tick-infecting bacteria. *Journal of Microbiological Methods* 52: 251–260.

Dillon RJ, Charnley AK. 2002. Mutualism between the desert locust *Schistocerca gregaria* and its gut microbiota. *Research In Microbiology* 153(8): 503–509.

Dillon RJ, Vennard CT, Charnley AK. 2002. Gut bacteria produce components of a locust cohesion pheromone. *Journal of Applied Microbiology* 92: 759–763.

Dillon RJ, Charnley AK. 1995. Chemical Barriers to Gut Infection in the Desert Locust: In Vivo Production of Antimicrobial Phenols Associated with the Bacterium *Pantoea agglomerans*. *Journal of Invertebrate Pathology* 66: 72–75.

Eutick ML, O'Brien RW, Slaytor M. 1987. Bacteria from the gut of Australian Termites. *Applied and Environmental Microbiology* 35(5): 823–828.

Farrelly V, Rainey FA, Stackebandt E. 1995. Effect of genome size and *rrn* gene copy number on PCR amplification of 16S rRNA genes from a mixture of bacterial species. *Applied and Environmental Microbiology* 61: 2798–2801.

Gilliam M. 1997. Identification and roles of non-pathogenic microflora associated with honey bees. *Fems Microbiology Letters* 155: 1–10.

Guo XR, Zhang YL, Li ML. 2007. Effect of Coriaria Lactone on the Activities of Three Enzymes in *Mythimna separata*. *Acta*

Botanica Boreali-Occidentalia Sinica 27(8): 1656–1660.

He MQ. 1994. *Animal ecology*. China Agriculture Press.

He ZB, Yin YP, Cao YQ, Dong YM, Zhang W. 2001. The bacterial community of the gut of *Apriona germari* larvae. *Acta Microbiologica Sinica* 41(6): 471–473.

Jackson CR, Roden EE, Churchill PF. 2000. Denaturing gradient gel electrophoresis can fail to separate 16S rDNA fragments with multiple base differences. *Journal of Molecular Biology* 1: 49–51.

Kumar S, Tamura K, Jakobson IB, Nei M. 2001. MEGA2: Molecular Evolutionary Genetics Analysis software. *Bioinformatics* 17: 1244–1245.

Lee JH, Uhm KB. 1995. Migration of the oriental armyworm *Mythimna separata* in East Asia in relation to weather and climate.. In: Drake VA, Gatehouse AG, Editors. *Insect migration: tracking resources through space and time*. pp. 105–116. Cambridge University Press.

Liu L, Wang ZK, Yu HW, Chen SH, Yan GF, Xia YY, Yin YP. 2001. Analysis of the bacteria diversity in intestines of *Hepialus gonggaensis* larvae. *Acta Microbiologica Sinica* 48(5): 616–622.

Lü M, Kou J, Liu HX, Wu WJ. 2008. Studies on the induction of celangin in V to cytochrome P450 and glutathione S-transferase of armyworm, *Mythimna separata*. *Journal of Northwest Agricultural and Forestry University (Natural Science Edition)* 36(6): 180–184.

Mead LJ, Khachatourians GG, Jones GA. 1988. Microbial ecology of the gut in laboratory stocks of the migratory grasshopper, *Melanoplus sanguinipes* (Fab) (Orthoptera:Acrididae). *Applied and Environmental Microbiology* 54(5): 1174–1181.

Mrazek J, Strosova L, Fliegerova K, Kott T, Kopečný J. 2008. Diversity of insect intestinal microflora. *Folia Microbiologica* 53 (3): 229–233.

Muyzer G, DeWaal EC, Uitterlinden AG. 1993. Profiling of complex microbial populations by denaturing gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. *Applied and Environmental Microbiology* 59: 695–700.

Nardi JB, Mackie RI, Dawson JO. 2002. Could microbial symbionts of arthropod guts contribute significantly to nitrogen fixation in terrestrial ecosystems? *Journal of Insect Physiology* 48: 751–763.

Polz MF, Cavanaugh CM. 1998. Bias in template-to-product ratios in multitemplate PCR. *Applied and Environmental Microbiology* 64: 3724–3730.

Prescott LM, Harley JP, Donald AK. 2002. *Microbiology*. 5th edition. pp. 498-507. Higher Education Press/McGraw- Hill Companies.

Reeson AF, Jankovic T, Kasper ML, Rogers S, Austin AD. 2003. Application of 16S rDNA- DGGE to examine the microbial ecology associated with a social wasp *Vespa germanica*. *Insect Molecular Biology* 12: 85–91.

Tokuda G, Watanabe H, Matsumoto T. 1997. Cellulose digestion in the wood-eating higher termite *Nsutiterme takasafoensis* (Shiraki): distribution of cellulase and properties of Endo- β -1,4-glucanase. *Zoological Science* 14: 83–97.

Xiang H, Li MW, Zhao Y, Zhao LP, Zhang YH, Huang YP. 2007. Bacterial community in midguts of the silkworm larvae estimated by PCR-DGGE and 16S rDNA gene library analysis. *Acta Entomologica Sinica* 50(3): 22–233.

Yuan XJ, Li HP, Huang DZ, Huang QX, Su XY, Hou XJ. 2009. Genomic DNA extraction and optimization of PCR reaction system of *Apriona germari* intestinal bacteria. *Science of Sericulture* 35(2): 379–383.

Zhang W, He ZB, Deng XP, Yin YP. 2004. Variety and distribution of intestinal flora of *Apriona germari* larvae. *Journal Southwest Agricultural University (Natural Science Edition)* 26(2): 169–172.

Zhang GY, Zhai BP. 2008. Construction and identification of enriched microsatellite library from *Pseudaletia separata* genome. *Acta Microbiologica Sinica* 28(8): 3860–3867.

Zhang YL, Zhuo Y, Zhang ZY. 2003. Effect of cantharidin on the midgut of the orient armyworm (*Mythimna separata*) and diamondback moth (*Plutella xylostella*). *Acta Entomologica Sinica* 46(3): 272–276.

Zhou J, Hu GF, Liu MY, Li YQ, Li JJ, Yu HT. 2008. Contact and antifeeding effects on four species of the genus *Datura* against *Mythimna separata* and *Pieris rapae*.

Journal of Gansu Agricultural University 3: 102–106.

Zhuang SH, Zong N, Li ML. 2007. Tutin's effect on poisoned larvae's malpighian tubules and midgut tissue. *Journal of Northwest A & F University (Natural Science Edition)* 35(7): 127–130

Zhuo FP, Chen SJ, Yin YP, Wang ZK, Xia YX. 2005. The analysis of intestinal bacteria community of *Hepialus gonggaensis* larvae. *Acta Microbiologica Sinica* 27(11): 26–29.

Weisburg WG, Barns SM, Pelletier DA, Lane DJ. 1991. 16S ribosomal DNA amplification for phylogenetic study. *Journal of Bacteriology* 173(2): 697–703.