



The Community of Hymenoptera Parasitizing Necrophagous Diptera in an Urban Biotope

Authors: Frederickx, Christine, Dekeirsschieter, Jessica, Verheggen, François J., and Haubruge, Eric

Source: Journal of Insect Science, 13(32) : 1-14

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.013.3201>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



The community of Hymenoptera parasitizing necrophagous Diptera in an urban biotope

Christine Frederickx^{a*}, Jessica Dekeirsschieter^b, François J. Verheggen^c, Eric Haubruge^d

Department of Functional and Evolutionary Entomology, Gembloux Agro-Bio Tech, University of Liege, Passage des Déportés 2, 5030 Gembloux, Belgium

Abstract

Most reports published in the field of forensic entomology are focused on Diptera and neglect the Hymenoptera community. However, Hymenoptera are part of the entomofaunal colonization of a dead body. The use of Hymenoptera parasitoids in forensic entomology can be relevant to evaluate the time of death. Hymenoptera parasitoids of the larvae and pupae of flies may play an important role in the estimation of the post-mortem period because their time of attack is often restricted to a small, well-defined window of time in the development of the host insect. However, these parasitoids can interfere with the developmental times of colonizing Diptera, and therefore a better understanding of their ecology is needed. The work reported here monitored the presence of adult Hymenoptera parasitoids on decaying pig carcasses in an urban biotope during the summer season (from May to September). Six families and six species of parasitoids were recorded in the field: *Aspilota fuscicornis* Haliday (Braconidae), *Alysia manducator* Panzer, *Nasonia vitripennis* Walker (Pteromalidae), *Tachinaephagus zealandicus* Ashmead (Encyrtidae), *Trichopria* sp. (Diapriidae), and *Figites* sp. (Figitidae). In the laboratory, five species emerged from pupae collected in the field: *Trichopria* sp., *Figites* sp., *A. manducator*, *N. vitripennis*, and *T. zealandicus*. These five species colonize a broad spectrum of Diptera hosts, including those species associated with decomposing carcasses, namely those from the families Calliphoridae, Muscidae, Fanniidae, and Sarcophagidae.

Keywords: *Alysia manducator*, carrion ecology, forensic entomology, *Nasonia vitripennis*, *Tachinaephagus zealandicus*, temperate area

Correspondence: ^a frederickxchr@hotmail.com, ^b jdekeirsschieter@hotmail.com, ^c fverheggen@ulg.ac.be, ^d e.haubruge@ulg.ac.be, *Corresponding author

Editor: Michael Strand was editor of this paper.

Received: 12 December 2011 **Accepted:** 24 May 2012

Copyright : This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 13, Number 32

Cite this paper as:

Frederickx C, Dekeirsschieter J, Verheggen FJ, Haubruge E. 2013. The community of Hymenoptera parasitizing necrophagous Diptera in an urban biotope. *Journal of Insect Science* 13:32. Available online: <http://www.insectscience.org/13.32>

Introduction

As soon as an animal dies, the carcass becomes a food source for many organisms (Grassberger and Frank 2004; Carter et al. 2007). In temperate natural biotopes, the most specialized organisms inhabiting the “cadaver-ecosystem” are insects (Amendt et al. 2004). Necrophagous insects, mainly Diptera and Coleoptera, are attracted to the cadaver, which is then colonized in a relatively predictable sequence called the entomofaunal succession or insect succession (Megnin 1894; Putman 1983; Schoenly and Reid 1987; Marchenko 1988, 2001; Benecke 2004). Study of these insects in a medico-legal context is a component of forensic entomology (Hall 1990; Amendt et al. 2004). Many forensic entomological studies have been conducted using pig carcasses as surrogate human models due to physiological, ethical, legal, and economic reasons (Rodriguez and Bass 1983; Catts and Goff 1992; Anderson and VanLaerhoven 1996; Hart and Whitaker 2005), but few of the studies conducted on pig carcasses have taken place in Europe (Garcia-Rojo 2004; Grassberger and Frank 2004; Wyss and Cherix 2006; Matuszewski et al. 2008; Dekeirsschieter et al. 2011; Matuszewski et al. 2011).

Many published reports are focused on the Diptera community, but very few look at the parasite community (Davies 1999; Amendt et al. 2000; Campobasso et al. 2001; Grassberger and Frank 2003; Schroeder et al. 2003; Amendt et al. 2004; Wang et al. 2008). Predators and parasites are generally considered to be the second most significant group of carrion-frequenting taxa (Goff 2010, 2011). Among these, a special group of parasites, called parasitoids, attack several necrophagous taxa. A parasitoid larvae or pupae feed

exclusively on other arthropods, mainly insects, resulting in the death of the host (Eggleton and Gaston 1990; Amendt et al. 2010). They represent an extremely diverse group, though mainly belonging to Hymenoptera. In Europe, 83 parasitoids species, which attack the larval and pupal stages of synanthropic Diptera, are listed (Fabritius and Klunker 1991). The use of Hymenoptera parasitoids in forensic entomology can be useful to evaluate the time of death (Amendt et al. 2000; Grassberger and Frank 2003; Amendt et al. 2010). The pupal parasitoids of blowflies may play an important role in the estimation of the post-mortem period, because their time of attack is often restricted to a small, well-defined window of time in the development of the host insect (Anderson and Cervenka 2002). This specialized group might also lead to significant problems for forensic entomologists. For example, changes in developmental times for *Lucilia sericata* L. have been observed after attack by the parasitoid *Alysia manducator* Panzer, the result being premature pupation (Holdaway and Evans 1930). Families of Hymenoptera parasitoids of forensic importance include Braconidae, Pteromalidae, and Ichneumonidae (Amendt et al. 2000; Disney and Munk 2004; Turchetto and Vanin 2004). Among them, *Nasonia vitripennis* Walker (Pteromalidae) and *Alysia manducator* Panzer (Braconidae) are the most common parasitoids found on cadavers (Grassberger and Frank 2003; Grassberger and Frank 2004; Turchetto and Vanin 2004).

So far, little information is available on the Hymenoptera post-mortem community in temperate biogeoclimatic countries (Woodcock et al. 2002; Wyss and Cherix 2006; Lefebvre and Gaudry 2009). This paper identifies the Hymenoptera parasitoid community that was identified on large carcasses in a temperate urban biotope during summer.

Materials and Methods

Field site and study periods

This study was conducted during summer 2010 (4 May – 30 September) in an abandoned garden at an urban site located in Belgium. The garden consisted of hazel trees (*Corylus avellana* L.), spruce (*Picea* spp.), and ash trees (*Fraxinus excelsior* L.). The shrub layer was absent. The soil vegetation was scattered, and the herb layer was mainly constituted of nettles (*Urtica dioica* L.) and ivies (*Hedera helix* L.). Regarding the moss layer, there were some sparse areas of *Polytrichum* sp.

The ambient air temperature and humidity were automatically measured once an hour using a data logger (HOBO RH/Temp 8K©; Onset Computer Corporation, www.onsetcomp.com) placed on the lateral side of each cage at a height of 75 cm. The daily mean temperature was calculated on the basis of ambient air temperature recorded at time intervals of 24 hr.

Animal model

Each month, two male piglets, *Sus domesticus* L. (Artiodactyla: Suidae) (5 kg), were killed by penetrative captive bolt (fractured skull) and placed at the experimental sites within

two hours. Piglets were provided by the experimental farm of the Veterinary Medicine Faculty of the University of Liege, Belgium (ethical authorization number: FUSAGx-08-07). Immediately after euthanasia, the pig carcasses were packed in double plastic bags to avoid any insect colonization, before being placed at the experimental site.

Each pig carcass was placed 30 m from each other in a plastic box (50 cm x 95 cm x 50 cm) filled with 30 cm of soil from the site in order to facilitate the collection of pupae samples. This box was placed in a metal mesh cage (180 cm x 90 cm x 90 cm) to avoid scavenging by vertebrate carnivores. Dates of exposure of the piglets were 4 May, 2 June; 30 June; 11 August, and 13 September.

Insect collection and identification

In order to quantify insect colonization on pig carcasses, four yellow traps (plastic containers of 9 cm height and 27 cm diameter) filled with soapy water were placed around each carcass. The distribution of the yellow traps on the ground was as follows: one near the head, one near the dorsal face, one near the anus, and one near the ventral face (Figure 1). The insect traps were removed every week and the collected specimens were conserved in 80% *norvanol D* (ethanol denatured with ether). Only adult stages were included in the count-





Figure 1. Position of yellow traps around each pig carcass. High quality figures are available online.



ing of collected insects during this study.

At the end of each month, pupae present in the soil under the pig carcasses were collected before being transported to the laboratory for rearing. Rearing was conducted under environmentally controlled conditions of $23 \pm 1^\circ\text{C}$ with a daylight regime of 16:8 L: D and 70% RH. Pupae were stored together in plastic containers until either an adult fly or parasitoids emerged. Emerged specimens were killed in 80% *norvanol D*. Pupae from which no adult flies or parasitoids had emerged after eight weeks were dissected for evidence.

Emerged Diptera were identified by their species. However, parasitoid hosts were not identified. Typically, the puparium were very similar in general appearance, being coarctate and light-brown to dark-brown in color, which made identification difficult (Sukontason et al. 2007). Moreover, only one pupal identification key exists. This key identified seven fly species of forensic importance in Thailand, such as *Chrysomya* spp. (Sukontason et al. 2007). Hymenoptera were identified by family, or to genera when it was possible.

Moreover, the subfamilies of Alysiniinae, Pteromalinae, Encyrtinae, Figitinae, and Diapriinae were mounted on insect pins and identified by species. Hymenoptera specimens were determining using different identification keys (Fischer 1971, 1972; Nixon 1980; Fergusson 1986; Delvare and Aberlenc 1989; Goulet and Huber 1993).

Results

Environmental parameters

The mean atmospheric temperatures measured during the decompositional process were 11.0°C for May, 16.6°C for June, 19.9°C for July, 16.2°C for August, and 12.9°C for September (Figure 2). The mean relative humidity was 75.1% for May, 73.2% for June, 71.3% for July, 85.6% for August, and 88.3 % for September (Figure 2).

Hymenoptera specimens collected in the field

The Hymenoptera superfamilies identified on pig carcasses were Ichneumonidea (one family), Chalcidoidea (two families), Cynipoidea (two families), and Proctotrupeoidea (one family) (Table 1). Six families were identified:

Table 1. List of Hymenoptera collected in yellow traps.

Superfamilies	Families	Subfamilies	Species	Abundance (No. of individuals)					
				May	June	July	August	September	Total
Ichneumonoidea	Braconidae	Alysiinae	<i>A. manducator</i>	79	337	85	140	162	803
			<i>A. fuscicornis</i>	-	-	6	-	-	6
		Opiinae	Not identified	11	6	6	-	3	26
Chalcidoidea	Pteromalidae	Pteromalinae	<i>N. vitripennis</i>	3	6	-	1	1	11
		Not identified	Not identified	8	4	4	2	-	18
	Encyrtidae	Encyrtinae	<i>T. zealandicus</i>	-	1	24	18	10	53
Cynipoidea	Figitidae	Figitinae	<i>Figites</i> sp.	3	27	13	30	12	85
	Eucoilidae	Not identified	Not identified	-	1	2	-	-	3
Proctotrupeoidea	Diapriidae	Diapriinae	<i>Trichopria</i> sp.	-	-	3	8	9	20
Total				104	382	143	199	197	1025

were identified: Braconidae, Pteromalidae, Encyrtidae, Figitidae, Eucoilidae, and Diapriidae. The richness of variety in seasonal families was shown to be three families in May, five families in June, six families in July, and five families both in August and September.

Six species of Hymenoptera were identified during the sampling period, and subfamilies of Alysiinae included species of *Alysia manducator* Panzer (803 specimens) and *Aspilota fuscicornis* Haliday (Braconidae) (six specimens). *N. vitripennis* (11 specimens), *Tachinaephagus zealandicus* Ashmead (Encyrtidae) (53 specimens), *Trichopria* sp. (Diapriidae) (20 specimens), and *Figites* sp. (Figitidae) (85 specimens) were also identified. *Figites* sp., *Trichopria* sp., and *A. manducator* were collected all through the summer season. *T. zealandicus* was collected from June to September, however, *A. fuscicornis* was collected only during July.

Of the total individuals (Table 1), 10.15% were collected in May, 37.27% in June, 13.95% in July, 19.41% in August, and 19.22% in September. *A. manducator* was the most abundant species overall (78.34%), fol-

lowed by *Figites* sp. (8.29%) and *T. zealandicus* (5.17%). The remaining families consisted of fewer than 30 collected individuals. In May, *Figites* sp. and *N. vitripennis* were the most abundant species. During June, the predominant species was *Figites* sp., followed by *N. vitripennis*. The most predominant species in July was *T. zealandicus*, followed by *Figites* sp. and *A. fuscicornis*. In August and September, *Figites* sp. was most abundant, followed by *T. zealandicus* and *Trichopria* sp.

Hymenoptera reared from fly pupae

In total, 13,310 Diptera pupae were collected from the soil under carcasses throughout the study (Table 2). Of these, 47.96% were parasitized by Hymenoptera parasitoids, which yielded 6,833 successfully emerged parasitoid specimens. The percentage of parasitism during the summer season was 3.48% in May, 8.99% in June, 8.88% in July, 49.79% in Au-

Table 2. Number of pupae collected, emerged, parasitized, and not emerged in laboratory.

Pupae	No. of pupae					
	May	June	July	August	September	Total
Collected	2241	1079	2715	1456	5819	13310
Emerged	105	603	1117	349	221	2395
Parasitized	78	97	241	725	5243	6384
Unemerged	2058	379	1341	382	355	4515

Table 3. List of Hymenoptera reared from fly pupae.

Superfamilies	Families	Subfamilies	Species	Abundance (No. of individuals)					
				May	June	July	August	September	Total
Ichneumonidea	Braconidae	Alysiinae	<i>A. manducator</i>	77	-	7	592	5084	5760
Chalcidoidea	Pteromalidae	Pteromalinae	<i>N. vitripennis</i>	19	-	-	-	-	19
		Encyrtidae	Encyrtinae	<i>T. zealandicus</i>	-	6	41	425	222
Cynipoidea	Figitidae	Figitinae	<i>Figites</i> sp.	-	95	216	28	11	350
Proctotrupeoidea	Diapriidae	Diapriinae	<i>Trichopria</i> sp.	-	-	10	-	-	10
Total				96	101	274	1045	5317	6833

Table 4. Diptera species emerged in laboratory.

Families	Species	Abundance (No. of individuals)					
		May	June	July	August	September	Total
Calliphoridae	<i>Calliphora vicina</i> Robineau-Desvoidy	105	-	-	6	65	176
	<i>Calliphora vomitoria</i> Linnaeus	-	-	-	-	133	133
	<i>Lucilia sericata</i> Meigen	-	-	9	-	-	9
	<i>Lucilia caesar</i> Linnaeus	-	7	539	337	23	906
	<i>Lucilia silvarum</i> Meigen	-	-	5	1	-	6
Muscidae	<i>Hydrotaea ignava</i> Harris	-	596	579	5	-	1180
Total		105	603	1132	349	221	2410

gust, and 90.10% in September.

Five parasitoid species emerged from pupal samples collected in the field (Table 3). The number of *A. manducator* collected increased dramatically in September. *N. vitripennis* and *Trichopria* sp. were collected only in May and July. *T. zealandicus* and *Figites* sp. were collected from June to September. *A. manducator* was the predominant species (84.30%), followed by *T. zealandicus* (10.16%) and *Figites* sp. (5.12%). Comparatively, parasitism of fly pupae by *N. vitripennis* and *Trichopria* sp. was rare, with both species contributing less than 0.3% of the total parasitism recorded.

Of the Diptera specimens that emerged in the laboratory, two families were identified in association with carcasses: Calliphoridae with five species, and Muscidae with one species (Table 4). All of these species have previously been reported as carrion breeding flies in Europe (Wyss and Cherix 2006; Lefebvre and Gaudry 2009).

Discussion

The objective of this study was to document Hymenoptera, and more precisely Hymenoptera parasitoids of necrophagous Diptera, in an urban site. At this site, the Hymenoptera community was represented by six families and was found to change over time. The make-up of the Hymenoptera community differed between months. The Hymenoptera diversity was highest in July, followed by

June, August, September, and finally May. However, the lowest abundance of Hymenoptera occurred in May, with approximately three times fewer specimens trapped than in June. June had the highest level of caught insects, followed by August, September, and July. Our breeding in the laboratory showed the highest rate of parasitism was observed in September. All previous reports from Europe have been anecdotal, limited to family level or carrion fauna lists (Woodcock et al. 2002; Wyss and Cherix 2006; Lefebvre and Gaudry 2009).

This study identified five species of parasitoids that visited decomposing remains in search of host carrion flies: *A. manducator*, *N. vitripennis*, *Figites* sp., *Trichopria* sp., and *T. zealandicus*. These five species colonized a broad spectrum of Diptera hosts associated with decomposing carcasses, including Calliphoridae, Muscidae, Fanniidae, and Sarcophagidae (Laing 1937; Whiting 1967; Rueda and Axtell 1985; Blanchot 1992; Goulet and Huber 1993; Ferreira De Almeida et al. 2002; Voss et al. 2010).

In the field and laboratory, the most abundant parasitoid species, approximately 78% of the species caught and 84% of the species that emerged, was *A. manducator*. 5,760 pupae were parasitized by *A. manducator* in the field. These females are attracted to decomposing meat (Laing 1937; Blanchot 1992; Reznik et al. 1992). On the carrion, an *A. manducator* female walks over the surface,

stabbing frequently with her ovipositor until a host larva is encountered (Laing 1937; Reznik et al. 1992). Hosts are discovered by contact, and female parasites prefer larger size larvae (Graham-Smith 1919; Blanchot 1992). *A. manducator* is an endoparasitoid, and it lays one egg per host (Reznik et al. 1992; Goulet and Huber 1993). This species is present worldwide, but much more prevalent in temperate regions (Goulet and Huber 1993), and has been collected in an urban site in central Europe in May (Grassberger and Frank 2004). In the British Isles, *A. manducator* is the most common of the parasitic Hymenoptera likely to be seen in carrion (Smith 1986). The seasonal activity of *A. manducator* reported in the present study is in agreement with previous observations in the Paris region, where it was active from May to October (Blanchot 1992). In the present study, the peak of its presence in yellow traps was observed during June (337 specimens).

Three subfamilies are recognized in Figitidae: Anacharitinae, Aspiceratinae, and Figitinae (Goulet and Huber 1993). Aspiceratinae and Figitinae are solitary endoparasitoids of Diptera pupae and early stage larvae, respectively, but the parasitoid emerges from the puparium (Fergusson 1986; Goulet and Huber 1993). Payne and Mason (1971) identified two genera of Figitinae, *Figites* and *Neralsia*, which were collected from pig carcasses. Exposed larvae were generally parasitized; however, *Figites* would also enter the carcass in search of prey. *Neralsia* were only observed parasitizing exposed larvae (Payne and Mason 1971). Six percent of the total caught specimens belonged to the Figitinae subfamily. In the laboratory, 350 pupae were parasitized by one single species of *Figites* sp. These small parasitoids were attracted to carrion where *Lucilia* spp. and *Sarcophaga* spp. larvae were the prevalent species (Payne and Mason

1971). The parasitism of *Figites* sp. in the field corresponded to the time period when *Lucilia* spp. was predominant (June to September) (Payne and Mason 1971).

Encyrtidae is one of the most important chalcidoid families for biological control (Goulet and Huber 1993; Voss et al. 2010). Species are gregarious endoparasitoids of eggs, third instar larvae, and postfeeding or prepupae of several forensically important Diptera (Fanniidae, Muscidae, Calliphoridae) (Goulet and Huber 1993; Ferreira De Almeida et al. 2002; Voss et al. 2010). In 2003, the presence of *T. zealandicus* was detected in northern Italy during September (Turchetto et al. 2003; Turchetto and Vanin 2004). This species, probably native to Australia and New Zealand, has been introduced into various parts of the world in attempts to control pest species of synanthropic Diptera, but no records are available for Europe or the northern regions (Turchetto and Vanin 2010). Following Italy, this is the second recording of *T. zealandicus* in the Palearctic Region (Turchetto et al. 2003).

The family of Diapriidae includes four subfamilies: Belytinae, Ismarinae, Ambositrinae, and Diapriinae (Goulet and Huber 1993). Only Diapriinae contains parasitoid species of necrophagous Diptera (Goulet and Huber 1993). In the field, *Trichopria* sp. made-up approximately 2% of the total specimens caught. In the laboratory, one species of *Trichopria* sp. parasitized only five pupae in July. Moreover, only three specimens were collected in July. These small black insects are endoparasitoids of the immature stages of Diptera (Payne and Mason 1971; Goulet and Huber 1993). In the United Kingdom, three genera of Diapriinae were recorded in the Graham-Smith study on carrion. *Aneurhynchus* and *Psilus* were found in insect-open

carrion, and *Trichopria* in buried carrion (Graham-Smith 1919; Payne and Mason 1971). In France, *Trichopria inermis* Kieffer, a gregarious parasitoid of *L. sericata*, was been observed (Blanchot 1995).

The last species identified in the laboratory was *N. vitripennis*, a gregarious ectoparasitoid of the pupae of several fly species of forensic importance, including blowflies, flesh flies, and houseflies (Whiting 1967; Rueda and Axtell 1985; Blanchot 1992). The attraction of females of this species toward the host can be caused, *in natura*, by decomposing meat (Laing 1937; Blanchot 1992; Frederickx et al. In Press). These wasps are regularly found on carcasses (Blanchot 1995; VanLaerhoven and Anderson 1999; Amendt et al. 2000; Grassberger and Frank 2004; Pohjoismaki et al. 2010) or in bird nests (Whiting 1967; King and Ellison 2005). This species has been recorded worldwide (Braack 1987). *N. vitripennis* was identified in Belgium in 1920 (Mitroiu 2001; Vago 2006), is a cosmopolitan species (Darling and Werren 1990; Yoder et al. 1994), and has been widely investigated in the subjects of genetic, ecological, evolutionary, and developmental research over the last 50 years (Darling and Werren 1990; Grassberger and Frank 2003; Steiner et al. 2006; Gadau et al. 2008). These wasps are commercially supplied and widely used as biological control agents of blowflies in Australia and in the United States (Mandeville et al. 1990; Morgan et al. 1991; Floate et al. 1999; Grassberger and Frank 2003). Only one pupa was parasitized by female *N. vitripennis*. *N. vitripennis* is not considered to be adapted for burrowing, and buried pupae are typically beyond the reach of parasitizing females (Altston 1920; Wylie 1958; Beard 1964; Whiting 1967; Vinson 1976; Voss et al. 2009). In 1950, a higher incidence of parasitism by this species in pupae located on or near

the surface of a carcass, rather than on those buried in the soil was reported (Ullyett 1950). In the present study, no Diptera specimens were collected on the ground.

Forensic interest

Forensic entomology is the application of the study of insect biology to criminal matters and is frequently used to estimate the time that has elapsed since death, or the post-mortem interval (Gennard 2007; Ricciuti 2007; Eberhardt and Elliot 2008). Forensic practitioners have previously postulated the use of parasitoids as a tool in criminal investigations, although the presence of parasitoids at crime scenes has largely been ignored due to their small size and the paucity of biological information available (Grassberger and Frank 2003; Amendt et al. 2007; Voss et al. 2009). Females of Hymenoptera parasitoids usually prefer to parasitize particular instars of their hosts (Reznik et al. 1992). In many cases, the most preferable stages are also the most suitable, creating optimal synchronization (Laing 1937; Vinson 1976; Reznik et al. 1992). *A. manducator* prefer larvae of *Calliphora vicina* that have already finished their feeding but have not yet left the food substrate (Reznik et al. 1992). *N. vitripennis* usually lay eggs in their host one day after the pupation, when the skin of the larva has separated from the inner pupal cuticle (Gunn 2006; Gaudry 2010). However, it has been reported that two day-old pupae were parasitized at a significantly lower rate than pupae exposed for four days (Kaufman et al. 2001).

Considering the abundance of *A. manducator*, *T. zealandicus*, and *Figites sp.*, the use of these species as a reliable forensic indicator for estimating post-mortem interval is promising. In order to estimate post-mortem interval using these species, the calculated developmental time of the parasitoid simply has to be

added to the time of development of the host, therefore providing an extended post-mortem interval timeframe in cases where traditional forensic indicators have completed their development (Grassberger and Frank 2003; Amendt et al. 2010). However, when considering the potential influence, especially of larval parasitoids, it is important to take into account that they can also create significant problems, as seen in the change of developmental times for *L. sericata* after attack of *A. manducator*, which results in a premature pupation (Holdaway and Evans 1930). This problem clearly illustrates the need for further research in this field.

Acknowledgments

Christine Frederickx and Jessica Dekeirsschieter are financially supported by a PhD grant from the Fonds pour la formation à la Recherche dans l'Industrie et l'Agriculture. The voucher Hymenoptera specimens are stored at the entomological conservatory of the Department of Functional and Evolutionary Entomology (Gembloux Agro-Bio Tech, University of Liege, Belgium).

References

Altston A. 1920. The life-history and habits of two parasites of blowflies. *Proceedings of the Zoological Society of London* 3: 195–243.

Amendt J, Krettek R, Niess C, Zehner R, Bratzke H. 2000. Forensic entomology in Germany. *Forensic Science International* 113: 309–314.

Amendt J, Krettek R, Zehner R. 2004. Forensic entomology. *Naturwissenschaften* 91: 51–65.

Amendt J, Campobasso CP, Gaudry E, Reiter C, LeBlanc HN, Hall MJR. 2007. Best practice in forensic entomology - standards and guidelines. *International Journal of Legal Medicine* 121: 90–104.

Amendt J, Zehner R, Johnson DG, Wells J. 2010. Future trends in forensic entomology. In: Amendt J, Campobasso CP, Goff ML, Grassberger M, Editors. *Current concepts in Forensic Entomology*. pp. 353–368. Springer.

Anderson GS, VanLaerhoven SL. 1996. Initial studies on insect succession on carrion in southwestern British Columbia. *Journal of Forensic Sciences* 41: 617–625.

Anderson GS, Cervenka VJ. 2002. Insects associated with the body: their use and analyses. In: Haglund WD, Sorg MH, Editors. *Advances in forensic taphonomy: method, theory and archaeological perspectives*. pp. 173–200. CRC Press.

Beard R. 1964. Parasites of Muscoid Flies. *Bulletin of the World Health Organization* 31: 491–493.

Benecke M. 2004. Forensic entomology: Arthropods and Corpses. In: Tsokos M, Editor. *Forensic Pathology Reviews*. pp. 207–240. Humana Press.

Blanchot P. 1992. Nouveau répertoire bibliographique et nouvelles données biologiques sur les parasites de *Musca domestica* L. (Dipt. Muscidae). *EPHE, Biologie et Evolution des Insectes* 5: 1–54.

Blanchot P. 1995. Inventaire des parasitoïdes de mouches synanthropes recensés en France. *Biologie et Evolution des Insectes* 7-8: 111–119.

- Braack LEO. 1987. Community dynamics of carrion-attendant arthropods in tropical african woodland. *Oecologia* 72: 402–409.
- Campobasso C, Di Vella G, Introna F. 2001. Factors affecting decomposition and Diptera colonization. *Forensic Science International* 120: 18–27.
- Carter DO, Yellowlees D, Tibbett M. 2007. Cadaver decomposition in terrestrial ecosystems. *Naturwissenschaften* 94: 12–24.
- Catts EP, Goff ML. 1992. Forensic Entomology in Criminal Investigations. *Annual Review of Entomology* 37: 253–272.
- Darling C, Werren JH. 1990. Biosystematics of *Nasonia* (Hymenoptera: Pteromalidae): Two New Species Reared from Birds' Nests in North America. *Annals of the Entomological Society of America* 83: 352–370.
- Davies L. 1999. Seasonal and spatial changes in blowfly production from small and large carcasses at Durham in lowland northeast England. *Medical and Veterinary Entomology* 13: 245–251.
- Dekeirsschieter J, Verheggen FJ, Haubruge E, Brostaux Y. 2011. Carrion beetles visiting pig carcasses during early spring in urban, forest and agricultural biotopes of Western Europe. *Journal of Insect Science* 11:173. Available online: <http://www.insectscience.org/11.73/>
- Delvare G, Aberlenc H-P. 1989. *Les insectes d'afrique et d'amérique tropicale- Clés pour la reconnaissance des familles*. CIRAD Departement GERDAT.
- Disney R, Munk T. 2004. Potential use of Braconidae (Hymenoptera) in forensic cases. *Medical and Veterinary Entomology* 18: 442–444.
- Eberhardt TL, Elliot DA. 2008. A preliminary investigation of insect colonisation and succession on remains in New Zealand. *Forensic Science International* 176: 217–223.
- Eggleton P, Gaston KJ. 1990. Parasitoid species and assemblages - convenient definitions or misleading compromises. *Oikos* 59: 417–421.
- Fabritius K, Klunker R. 1991. Die Larven- und Puparienparasitoide von synanthropen Fliegen in Europa. *Angewandte Parasitologie* 32: 1–24.
- Fergusson NDM. 1986. *Charipidae, Ibalidae & Figitidae: Hymenoptera: Cynipoidea*. Royal Entomological Society of London.
- Ferreira De Almeida MA, Pires Do Prado A, Geden CJ. 2002. Influence of temperature on development time and longevity of *Tachinaephagus zealandicus* (Hymenoptera: Encyrtidae), and effects of nutrition and emergence order on longevity. *Biological Control* 31: 375–380.
- Fischer M. 1971. Untersuchungen über die europäischen *Alysiini* mit besonderer Berücksichtigung der Fauna Niederösterreichs (Hymenoptera, Braconidae). *Polskie Pismo Entomologiczne* 41: 19–160.
- Fischer M. 1972. Erste gliederung des paläarktischen *Aspilota*-Arten (Hymenoptera, Braconidae, Alysiinae). *Polskie Pismo Entomologiczne* 42: 323–459.
- Floate K, Khan B, Gibson G. 1999. Hymenopterous parasitoids of filth fly

(Diptera : Muscidae) pupae in cattle feedlots.

Canadian Entomologist 131: 347–362.

Frederickx C, Dekeirsschieter J, Verheggen FJ, Haubruge E. In Press. Host-habitat location by the parasitoid, *Nasonia vitripennis* Walker (Hymenoptera: Pteromalidae). *Journal of Forensic Sciences*.

Gadau J, Niehuis O, Peire A, Werren J, Baudry E, beukeboom L. 2008. The Jewel Wasp–*Nasonia*. In: Hunter W, Kole CS, Editors. *Genome mapping and genomics in animals*. pp. 27–41. Springer.

Garcia-Rojo AM. 2004. A study of the insect succession in carcasses in Alcala de Henares (Madrid administrative region, Spain) using pigs as animal models. *Boletin de la Sociedad Entomologica Aragonesa* 34: 263–269.

Gaudry E. 2010. The Insects Colonisation of Buried Remains. In: Amendt J, Campobasso CP, Goff ML, Grassberger M, Editors. *Current Concepts in Forensic Entomology*. pp. 273–311. Springer.

Gennard DE. 2007. *Forensic Entomology: an Introduction*. John Wiley & Sons, Ltd.

Goff ML. 2010. Early Postmortem Changes and Stages of Decomposition. In: Amendt J, Goff ML, Campobasso CP, Grassberger M, Editors. *Current Concepts in Forensic Entomology*. pp. 1–24. Springer.

Goff ML. 2011. Forensic entomology. In: Mozayani A, Noziglia C, Editors. *The Forensic Laboratory Handbook: Procedures and Practice*, pp. 447–478. Springer.

Goulet H, Huber JT. 1993. *Hymenoptera of the world: an identification guide to families*. Agriculture Canada Publication 1894/E.

Graham-Smith GS. 1919. Further observations on the habits and parasites of common flies. *Parasitology International* 11: 347–384.

Grassberger M, Frank C. 2003. Temperature-related development of the parasitoid wasp *Nasonia vitripennis* as forensic indicator. *Medical and Veterinary Entomology* 17: 257–262.

Grassberger M, Frank C. 2004. Initial study of arthropod succession on pig carrion in a central European urban habitat. *Journal of Medical Entomology* 41: 511–523.

Gunn A. 2006. *Essential Forensic Biology*. John Wiley & Sons, Ltd.

Hall RD. 1990. Medicocriminal entomology. In: Catts EP, Haskell NH, Editors. *Entomology and Death, a Procedural Guide*. pp. 1–8. Forensic Entomology Associates.

Hart AJ, Whitaker AP. 2005. Forensic Entomology. *Antennae* 30: 159–164.

Holdaway FG, Evans AC. 1930. Parasitism a stimulus to pupation: *Alysia manducator* in relation to the host *Lucilia sericata*. *Nature* 125: 598–599.

Kaufman PE, Long SJ, Rutz DA. 2001. Impact of exposure length and pupal source on *Muscidifurax raptorellus* and *Nasonia vitripennis* (Hymenoptera : Pteromalidae) parasitism in a New York poultry facility. *Journal of Economic Entomology* 94: 998–1003.

King B, Ellison J. 2005. Resource quality affects restlessness in the parasitoid wasp

Nasonia vitripennis. *Entomologia Experimentalis et Applicata* 118: 71–76.

Laing J. 1937. Host-finding by insect parasites. 1. Observations on the finding of hosts by *Alysia manducator*, *Mormoniella vitripennis* and *Trichogramma evanescens*. *Journal of Animal Ecology* 6: 298–317.

Lefebvre F, Gaudry E. 2009. Forensic entomology: a new hypothesis for the chronological succession pattern of necrophagous insect on human corpses. *Annales de la Societe Entomologique de France* 45: 377–392.

Mandeville JD, Mullens BA, Yu DS. 1990. Impact of selected pesticides on field population-dynamics of parasitic Hymenoptera (Pteromalidae) in caged-layer poultry manure in Southern California, USA. *Medical and Veterinary Entomology* 4: 261–268.

Marchenko M. 1988. Medico-legal relevance of cadaver entomofauna for the determination of the time of death. *Acta Medicinæ Legalis et Socialis* 38: 257–302.

Marchenko M. 2001. Medicolegal relevance of cadaver entomofauna for the determination of the time of death. *Forensic Science International* 120: 89–109.

Matuszewski S, Bajerlein D, Konwerski S, Szpila K. 2008. An initial study of insect succession and carrion decomposition in various forest habitats of Central Europe. *Forensic Science International* 180: 61–69.

Matuszewski S, Bajerlein D, Konwerski S, Szpila K. 2011. Insect succession and carrion decomposition in selected forests of Central

Europe. Part 3: Succession of carrion fauna. *Forensic Science International* 207: 150–163.

Megnin JP. 1894. *La Faune des Cadavres, Application de l'Entomologie à la Médecine Légale*. Encyclopédie Science, Aide mémoire.

Mitroiu M. 2001. Révision des collections de Chalcidoidea Pteromalidae (Hymenoptera) de l'Institut royal des Sciences naturelles de Belgique, et la découverte de 31 espèces nouvelles pour la Belgique. *Bulletin & Annales de la Societe Royale Belge d'Entomologie* 137: 91–97.

Morgan PB, Berti E, Costa VA. 1991. Life-history of *Spalangia gemina* Boucek (Hymenoptera, Pteromalidae), a fast-breeding microhymenopteran pupal parasitoid of Muscoid flies. *Medical and Veterinary Entomology* 5: 277–281.

Nixon GEJ. 1980. *Diapriidae (Diapriinae): Hymenoptera, Proctotrupoidea*. Royal Entomological Society of London.

Payne JA, Mason WRM. 1971. Hymenoptera associated with pig carrion. *Proceedings of the Entomological Society of Washington* 73: 132–141.

Pohjoismaki JLO, Karhunen PJ, Goebeler S, Saukko P, Saaksjarvi IE. 2010. Indoors forensic entomology: Colonization of human remains in closed environments by specific species of sarcosaprophagous flies. *Forensic Science International* 199: 38–42.

Putman RJ. 1983. *Carrion and Dung: the Decomposition of Animal Wastes*. Hodder.

Reznik SY, Chernoguz DG, Zinovjeva KB. 1992. Host searching, oviposition preferences and optimal synchronization in *Alysia*

manducator (Hymenoptera, Braconidae), a parasitoid of the blowfly, *Calliphora vicina*. *Oikos* 65: 81–88.

Ricciuti E. 2007. *Science 101: Forensics*. HarperCollins.

Rodriguez WC, Bass WM. 1983. Insect activity and its relationship to decay rates of human cadavers in East Tennessee. *Journal of Forensic Sciences* 28: 423–432.

Rueda LM, Axtell RC. 1985. Comparison of hymenopterous parasites of house fly, *Musca domestica* (Diptera: Muscidae), pupae in different livestock and poultry production systems. *Environmental Entomology* 14: 217–222.

Schoenly K, Reid W. 1987. Dynamics of heterotrophic succession in carrion arthropod assemblages- Discrete seres or a continuum of change. *Oecologia* 73: 192–202.

Schroeder H, Klotzbach H, Püschel K. 2003. Insects' colonization of human corpses in warm and cold season. *Legal Medicine* 5: S372–S374.

Smith KGV. 1986. *A manual of forensic entomology*. British Museum (Natural History).

Steiner S, Hermann N, Ruther J. 2006. Characterization of a female-produced courtship pheromone in the parasitoid *Nasonia vitripennis*. *Journal of Chemical Ecology* 32: 1687–1702.

Sukontason KL, Ngern-Klun R, Sripakdee D, Sukontason K. 2007. Identifying fly puparia by clearing technique: application to forensic entomology. *Parasitology Research* 101: 1407–1416.

Turchetto M, Villemant C, Vanin S. 2003. Two fly parasitoids collected during an entomo-forensic: the widespread *Nasonia vitripennis* (Hymenoptera Pteromalidae) and the newly recorded *Tachinaephagus zealandicus* (Hymenoptera Encyrtidae). *Bollettino della Societa Entomologica Italiana* 135: 109–115.

Turchetto M, Vanin S. 2004. Forensic evaluations on a crime case with monospecific necrophagous fly population infected by two parasitoid species. *Aggrawal's Internet Journal of Forensic Medicine and Toxicology* 5: 12–18.

Turchetto M, Vanin S. 2010. Climate change and forensic entomology. In: Amendt J, Campobasso CP, Goff ML, Grassberger M, Editors. *Current Concepts in Forensic Entomology*. pp. 327–351. Springer.

Ulyett GC. 1950. Competition for food and allied phenomena in sheep-blowfly populations. *Philosophical Transactions of the Royal Society of London, Series B, Biological Sciences* 234: 77–174.

Vago J-L. 2006. Révision des collections de Chalcidoidea Pteromalidae (Hymenoptera) de l'Institut royal des Sciences naturelles de Belgique et de la Faculté univeristaire des Sciences agronomiques de Gembloux, et la découverte de 145 espèces nouvelles pour la Belgique. *Bulletin & Annales de la Societe Royale Belge d'Entomologie* 142: 73–99.

VanLaerhoven SL, Anderson GS. 1999. Insect succession on buried carrion in two biogeoclimatic zones of British Columbia. *Journal of Forensic Sciences* 44: 32–43.

Vinson SB. 1976. Host selection by insect parasitoids. *Annual Review of Entomology* 21: 109–133.

Voss SC, Spafford H, Dadour IR. 2009. Hymenopteran Parasitoids of Forensic Importance: Host Associations, Seasonality, and Prevalence of Parasitoids of Carrion Flies in Western Australia. *Journal of Medical Entomology* 46: 1210–1219.

Voss SC, Spafford H, Dadour IR. 2010. Temperature-dependent development of the parasitoid *Tachinaephagus zealandicus* on five forensically important carrion fly species. *Medical and Veterinary Entomology* 24: 189–198.

Wang JF, Li ZG, Chen YC, Chen QS, Yin XH. 2008. The succession and development of insects on pig carcasses and their significances in estimating PMI in south China. *Forensic Science International* 179: 11–18.

Whiting AR. 1967. The biology of the parasitic wasp *Mormoniella vitripennis* (= *Nasonia brevicornis*) (Walker). *The Quarterly Review of Biology* 42: 333–406.

Woodcock BA, Watt AD, Leather SR. 2002. Aggregation, habitat quality and coexistence: a case study on carrion fly communities in slug cadavers. *Journal of Animal Ecology* 71: 131–140.

Wylie HG. 1958. Factors that affect host finding by *Nasonia vitripennis* (Walk.) (Hymenoptera: Pteromalidae). *The Canadian Entomologist* 90: 597–608.

Wyss C, Cherix D. 2006. *Traité d'Entomologie Forensique: Les insectes sur la*

scène de crime. Presses Polytechniques et Universitaires romandes.

Yoder J, Rivers D, Denlinger D. 1994. Water relationships in the ectoparasitoid *Nasonia vitripennis* during larval diapause. *Physiological Entomology* 19: 373–378.