

# Male Genitalia of Neotropical Charaxinae: A Comparative Analysis of Character Variation

Authors: Bonfantti, Dayana, Casagrande, Mirna Martins, and Mielke, Olaf Hermann Hendrik

Source: Journal of Insect Science, 13(35): 1-54

Published By: Entomological Society of America

URL: https://doi.org/10.1673/031.013.3501

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



# Male genitalia of neotropical Charaxinae: A comparative analysis of character variation

Dayana Bonfantti<sup>1a</sup>\*, Mirna Martins Casagrande<sup>1b</sup>, Olaf Hermann Hendrik Mielke<sup>1c</sup>

<sup>1</sup>Departamento de Zoologia; Setor de Ciências Biológicas; Universidade Federal do Paraná; C. P.: 19020; 81531-980, Curitiba, Paraná, Brazil

# Abstract

Charaxinae (Lepidoptera: Nymphalidae) has a worldwide distribution, although it occurs mainly between the tropics. Most species occur in the Neotropics, where three tribes, Preponini, Anaeomorphini, and Anaenini, can be found. Collectively, these three tribes encompass 109 species. Because of its relevance to systematics and taxonomy, the male genitalia of Lepidoptera have been extensively studied. The male genitalia are composed of the last two abdominal segments and their modifications for mating, known as claspers of the bodies. In order to improve upon the systematic classification of the subfamily, 31 species of 13 genera of Neotropical Charaxinae were analyzed. All characters relevant to species and generic taxonomy were analyzed. Most structures showed morphological variations among tribes, genera, and species. These variations demonstrated to be important to Preponini, because the structural patterns of the genitalia allow the separation in two groups, Prepona Boisduval and Archaeoprepona Fruhstorfer, and are in accord with the recent taxonomic classification proposed by Ortiz-Acevado and Willmott (2013), wherein Agrias Doubleday is synonymized in Prepona and Noreppa Rydon within Archaeoprepona. In the same way, Anaeomorpha splendida Rothschild showed considerable differences from Preponini's genera, the tribe in which it was included, confirming the revalidation of the tribe Anaeomorphini (Ortiz-Acevado and Willmott 2013). Substantial variation was found in the genital structures of Anaeini, making it difficult to establish structural patterns for this group. Such structural variation, however, may be very efficient to diagnose species, such as some species of Memphis Hübner and Fountainea Rydon, which can be easily identified through the presence and location of spines on the valva.

# Resumo

Charaxinae (Lepidoptera: Nymphalidae) tem distribuição mundial, embora ocorra principalmente entre trópicos. A maioria das espécies distribui-se na região Neotropical, onde três tribos podem ser encontradas, Preponini, Anaeomorphini, e Anaeini. Coletivamente, estas tribos abrangem 109 espécies. Devido à sua relevância para a sistemática e taxonomia, a genitália masculina de Lepidoptera, formada pelos dois últimos segmentos abdominais, tem sido extensivamente estudada. Com a finalidade de aperfeiçoar a classificação sistemática da subfamília, foram analisadas 31 espécies de 13 gêneros de Charaxinae neotropicais. A maioria das

estruturas apresentou variações morfológicas entre as tribos, gêneros e espécies. Estas variações demostraram ser importantes para Preponini, porque os padrões estruturais das genitálias permitiram que fossem separados em dois grupos, o primeiro *Prepona* Boisduval e o segundo *Archaeoprepona* Fruhstorfer além de estar de acordo com a recente classificação taxonômica proposta por Ortiz-Acevado and Willmott (2013), em que *Agrias* Doubleday esta sinonimizada em *Prepona* e *Noreppa* Rydon em *Archaeoprepona*. Da mesma maneira, *Anaeomorpha splendida* Rothschild apresentou diferenças consideráveis em relação aos gêneros de Preponini, tribo na qual estava incluída, confirmando a revalidação da tribo Aneomorphini (Ortiz-Acevado and Willmott 2013). Variações substanciais nas estruturas genitais de Anaeini foram encontradas, tornando-se difícil estabelecer padrões para esta tribo, embora tais variações estruturais possam ser muito eficientes para diagnosticar espécies, tais como algumas espécies de *Memphis* Hübner e *Fountainea* Rydon, facilmente identificadas através da presença e localização de espinhos na valva.

Keywords: Anaeini, Anaeomorphini, butterflies, morphology, Preponini Correspondence: a dayanabonfantti@gmail.com, b mibras@ufpr.br, c omhesp@ufpr.br, \*Corresponding author Editor: Carla Penz was editor of this paper. Received: 2 August 2011 Accepted: 13 March 2012 Copyright: This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed. ISSN: 1536-2442 | Vol. 13, Number 35

#### Cite this paper as:

Bonfantti D, Casagrande MM, Mielke OHH. 2013. Male genitalia of neotropical Charaxinae: A comparative analysis of character variation. *Journal of Insect Science* 13:35. Available online: <u>http://www.insectscience.org/13.35</u>

#### Introduction

The male genitalia of butterflies has been the subject of several very detailed studies, as its morphology can provide the basis for work of taxonomy, systematic analysis, and phylogenetic analysis (Simonsen 2006a). Some recent work on the comparative morphology of male genitalia has examined genital muscles (Simonsen 2006a, intraspecific structural b) and variations (Caldas 1997; Tóth 2011).

Charaxinae (Lepidoptera: Nymphalidae) are a group of medium to large, robust butterflies. They fly fast and circulate around tree canopies, behaviors that make them difficult to observe. Males and females are easily attracted by bait containing fruit, decomposing animals, feces, urine, and/or sweat (Rydon 1971). Charaxinae butterflies occur mainly between the tropics, primarily inhabiting the Neotropical region (Mielke et al. 2004a).

Recently, this subfamily has undergone drastic changes in its taxonomic classification. Through phylogenetic study of the tribe Preponini, using data from DNA sequences, Ortiz-Acevado and Willmott (2013) classified Preponini as follows: two genera were synonymized, *Noreppa* to *Archaeoprepona* and *Agrias* to *Prepona*; the fifth genus (monotypic), *Anaeomorpha splendida*, had its tribe status revalidated to Anaeomorphini. So, the neotropical Charaxinae contained two tribes, 17 genera, and 109 species (Salazar and Constantine, 2001; Lamas, 2004; Salazar, 2008), and today is made up of three tribes, 15

genera, and 109 species (Ortiz-Acevado and Willmott 2013).

The Charaxinae are characterized by the following synapomorphies: parapatagium sclerotized (Ehrlich 1958); forewings with R<sub>4</sub> and  $R_5$  longer than their common branch;  $R_4$ curved downward near its end, which may be at the costal margin, the apex, or the outer margin of the wing; posterior discal cell closed; anepisternum II present as a distinct sclerite; and mesothoracic pre-episternum well-developed (Ackery 1984). These synapomorphies were corroborated in a series of contributions by Mielke et al. (2004a, b, c), who also found some variation in the way vein R<sub>4</sub> ends in Zaretis itvs itvlus (Westwood) and Prepona claudina annetta (Gray).

With the goal to clarify the classification and systematics of the Neotropical Charaxinae, we have illustrated, compared, and ascertained variation in the male genitalia of 31 species of 13 genera. Some of the characters mentioned have been previously mentioned in other literature, whereas others are new.

## **Materials and Methods**

The species studied and material studied are listed in Table 1. The specimens analyzed are retained at Coleção Entomológica Pe. Jesus Santiago Moure, Departamento de Zoologia, Universidade Federal do Paraná, and Coleção Prof. Dr. Olaf Mielke.

The abdomen of each exemplar was removed and immersed in potassium hydroxide10% for at least 24 hours (depending on the size of the abdomen) to soften the tissues. After that, the abdomen was washed in water.

The genitalia were removed through a slit on the pleura of the 7<sup>th</sup> abdominal segment. After

study, the genital parts were stored in a microvial containing glycerin, which was attached to the same pin as the exemplar.

Drawings were made using a camera lucida attached to a dissecting microscope. The main lines were inked and watercolor was used to mark each sclerite with a different color.

Illustrations were coded as follows: full lines represent the boundaries of morphological structures on the first focal plane; dashed lines represent the boundaries of structures that lay underneath the main structure under study. As with structures represented by dotted lines, structures represented by dashed lines were only shown when they were important for the interpretation of the genital anatomy. Sclerites lacking a well-defined boundary were represented by intermediate colors. Sclerites, and their respective color codes, are as follows: tegumen - yellow; saccus - cadmium red; uncus - navy blue; gnathos - alizarin red; fultura inferior - turquoise; valvae - lemon yellow; penis - green.

In some illustrations, the sclerites were omitted in order to facilitate interpretation. For instance, the fultura inferior was omitted from illustrations showing the lateral aspect of the genitalia. A minimum of nine drawings were produced for each species as follows: lateral, dorsal, and posterior aspects of the genitalia; inner and outer aspects of the valva; and lateral (left and right), dorsal, and ventral aspects of the penis. In the descriptions and comparisons, only the greatest values are given for width and length.

SEM photographs were taken following standard procedures. First, the parts were dehydrated by immersion in decreasing ethanol dilutions (70%, 80%, 90%, and 100%) for 10 minutes, and then immersed for an

additional 10 minutes in ethanol absolute. Second, they were subjected to critical point drying, glued to a stub, and coated.

The main terminology used was adopted from Niculescu (1978) (Figure 1), with adaptations from Petersen (1904), Pierce (1909; 1914), Kuznetsov (1915), Snodgrass (1935), Oiticica (1946), Okagaki et al. (1954), Sibatani et al. (1954), Klots (1956), Ogata et al. (1957), Snodgrass (1957), Srivastava (1965), Casagrande (1979), and Bilotta (1994).

## Results

#### Anaeini

Anaea troglodyta (Fabricius) (Figure 1 A– K). Tegumen triangular in lateral view, anterior projection and appendices angulares wider than long. Arms from tegumen and saccus distinct. Anterior projection from saccus elongated in lateral view; posterior projection shorter than anterior.

Uncus slender, with distal portion descending, in dorsal view fused with tegumen, uncus differing from tegumen by a light sclerotization. Anal tube reduced and entirely membranous. Gnathos open, in lateral view forming two distinct plates, with distal portion flattened on anterior ventral portion, gnathos extended dorsoventrally toward the proximal portion, with spines of various sizes that give the dorsal distal section a serrated aspect.

Valva subtriangular, with inner ends expanded. Fultura inferior forming a semicircle.

Penis: aedeagus sclerotized, with bulbus ejaculatorius projecting from antero-dorsal opening

Coenophlebia archidona (Hewitson) (Figure 2 A-K). Tegumen subtriangular in lateral view, anterior projection wider than long, tegument fold projecting outward from median ventral appendices to region. angulares longer than wide. Arms from tegumen and saccus distinct. Anterior projection from saccus elongate in lateral view: posterior projection absent.

Uncus thick, with distal portion descending, in dorsal view fused with tegumen, uncus differing from tegumen by a light sclerotization. Anal tube developed, with ventral portion sclerotized (subscaphium). Gnathos open, in lateral view with proximal portion entirely fused with tegumen, Cshaped, in posterior view curved with distal portion descending.

Valva subrectangular, with a dorso-proximal notch and a small spine on distal portion. Fultura inferior bilobed, V-shaped, with a deep dorsal notch.

Penis: aedeagus sinuous and completely sclerotized, with bulbus ejaculatorius projecting from antero-dorsal opening.

*Consul electra* (Westwood) (Figure 3 A–K). Tegumen subtriangular in lateral view, anterior projection wider than long, tegument fold projecting outward from anteromedian portion of arms from tegument to their apex, appendices angulares longer than wide. Arms from tegumen and saccus distinct. Anterior projection from saccus elongate in lateral view, about twice as long as wide; posterior projection of saccus present, shorter than anterior projection.

Uncus slender, with distal portion descending, in dorsal view fused with tegumen, uncus differing from tegumen by a light

sclerotization. Anal tube developed, with ventral portion sclerotized (subscaphium). Gnathos open, in lateral view with the distal portion of each gnathos close to each other, with two parts, one dorsal and one ventral distal, ventral part more developed and subrectangular; ventral distal part elongate and with a small descending apical club; in posterior view L-shaped aspect, expanded apically.

Valva rounded, inner tegument expanded over outer tegument on ventral distal portion. Fultura inferior bilobed and linked ventrally with the diaphragma rather than with the valva.

Penis: aedeagus sclerotized, expanded ventrally up to insertion of manica, and bulbus ejaculatorius projecting from antero-dorsal opening.

**Consul fabius drurii** (Butler) (Figure 4 A–K, 33 C, D). Tegumen subtriangular in lateral view; anterior projection and appendices angulares wider than long. Arms from tegumen and saccus distinct. Anterior projection from saccus elongated in lateral view, at least three times longer than wide; posterior projection present, shorter than anterior projection.

Uncus slender, with distal portion descending, in dorsal view fused with tegumen, uncus differing from tegumen by a light sclerotization. Anal tube developed, with ventral portion sclerotized (subscaphium), enveloped by the gnathos ventrally. Gnathos open, in lateral view with two parts: dorsal and ventral-distal, both elongated, ventraldistal part descending, with a slight fold in the tegument. Valva approximately rounded; inner tegument expanded over outer tegument on ventral distal portion and bearing small spines on the harpe. Fultura inferior bilobed.

Penis: aedeagus completely sclerotized, with bulbus ejaculatorius projecting from anterodorsal opening, posterior opening large, originating underneath the manica.

*Fountainea glycerium cratais* (Hewitson) (Figure 5 A–M). Tegumen subtriangular in lateral view; without anterior projection, appendices angulares longer than wide. Arms from tegumen and saccus distinct. Anterior projection from saccus elongated in lateral view, at least three times longer than wide; posterior projection absent.

Uncus slender, apex slender and descending, in dorsal view fused with tegumen, uncus differing from tegumen by a light sclerotization. Anal tube developed, with ventral portion sclerotized (subscaphium) and enveloped by the gnathos ventrally. Gnathos closed and distally forming a round plate, projecting posteriorly.

Valva subrectangular and outer tegument forming a distal spine. Fultura inferior bilobed, with medium ventral notch, and small dorsal spines.

Penis: aedeagus sclerotized, bulbus ejaculatorius projecting from antero-dorsal opening; posterior opening irregular, with deep notches.

*Fountainea halice halice* (Godart) (Figure 6 A–M). Tegumen subtriangular in lateral view, without anterior projection, appendices angulares longer than wide. Arms from tegumen and saccus distinct. Anterior projection from saccus elongated in lateral view, at least twice longer than wide; posterior projection absent.

Uncus slender, with proximal ventral expansion and distal portion descending, in dorsal view fused with tegumen, uncus differing from tegumen by a light sclerotization. Anal tube developed, with ventral portion sclerotized (subscaphium) and enveloped by the gnathos ventrally. Gnathos closed, in lateral view sub rectangular, with anterior and posterior margins sinuous and fused medially

Valva subrectangular, with small spines near dorsal region and a distal spine. Fultura inferior bilobed, formed by two plates joined medially.

Penis: aedeagus sclerotized, with bulbus ejaculatorius projecting from antero-dorsal opening.

*Fountainea nessus* (Latreille) (Figure 7 A– L). Tegumen subtriangular in lateral view, without anterior projection, appendices angulares longer than wide. Arms from tegumen and saccus fused. Anterior projection from saccus elongated in lateral view, at least three times longer than wide; posterior projection absent.

Uncus slender, basal portion enlarged, with distal portion descending, in dorsal view fused with tegumen, uncus differing from tegumen by a light sclerotization. Anal tube developed and entirely membranous. Gnathos open, in lateral view rounded and with distal portion of both gnathos close to each other, differing from the remaining species of *Fountainea*.

Valva subrectangular, bearing distal spine. Fultura inferior large and bilobed. Penis: aedeagus sclerotized, with bulbus ejaculatorius projecting from antero-dorsal opening, posterior opening sinuous.

*Fountainea ryphea phidile* (Geyer) (Figure 8 A–M, 33 A, B). Tegumen subtriangular in lateral view, anterior projection wider than long, appendices angulares longer than wide. Arms from tegumen and saccus distinct. Anterior projection from saccus elongated in lateral view, at least twice as long as wide; posterior projection absent.

Uncus slender, with distal portion descending, in dorsal view fused with tegumen, uncus differing from tegumen by a light sclerotization. Anal tube reduced and entirely membranous. Gnathos closed, in lateral view forming a single round plate, similar to the gnathos of F. glycerium cratais.

Valva subrectangular, bearing distal spine. Fultura inferior wide and semi-circular, with reduced spines on apical dorsal tips.

Penis: aedeagus sclerotized, with bulbus ejaculatorius projecting from antero-dorsal opening.

Hypna clytemnestra forbesi Godman and Salvin (Figure 9 A–L). Tegumen subtraprezoidal in lateral view, anterior projection wider than long, appendices angulares longer than wide. Arms from tegumen and saccus fused. Anterior projection from saccus elongated in lateral view, at least four times longer than wide, apex slender and curved; posterior projection absent.

Uncus thick, dorsal proximal portion crestlike, apex slender and descending, in dorsal view not completely fused with tegumen, uncus with a basal fenestra, with two dorsal distal lobes. Anal tube reduced and entirely

membranous. Gnathos open, in lateral view apostrophe-like, proximal portion underneath tegumen, outer distal portion descending and curved, ventral portion elongated, descending, with distal portions of both gnathos close to each other and enveloped in membranes.

Valva approximately semi-circular. Fultura inferior reduced, V-shaped.

Penis: aedeagus sclerotized and sinuous, with bulbus ejaculatorius projecting from anterodorsal opening.

Hypna clytemnestra huebneri Butler (Figure 10 A–L). Tegumen, subtraprezoidal in lateral view, anterior projection wider than long, appendices angulares longer than wide. Arms from tegumen and saccus distinct, arms from saccus projected dorsally beyond ventral margin of tegumen. Anterior projection from saccus elongated in lateral view, four times longer than wide and descending; posterior projection absent.

Uncus thick, proximal dorsal portion crestlike, apex slender and descending, in dorsal view not completely fused, uncus with a basal fenestra. Anal tube reduced and entirely membranous. Gnathos open, in lateral view apostrophe-like, proximal portion underneath tegumen, outer distal portion descending and curved, elongate in ventral or posterior views, descending, with distal portion of both gnathos close to each other and enveloped in membranes, membranes joined distally.

Valva approximately rounded. Fultura inferior reduced and V-shaped.

Penis: aedeagus sclerotized, sinuous, with bulbus ejaculatorius projecting from anterodorsal opening, posterior opening sinuous. *Memphis acidalia victoria* (H. Druce) (Figure 11 A–L). Tegumen subtriangular in lateral view, without anterior projection, appendices angulares longer than wide. Arms from tegumen and saccus fused. Anterior projection from saccus elongated in lateral view, distal portion enlarged; posterior projection absent.

Uncus slender, ventral margin sinuous, apex slender and descending, in dorsal view fused with tegumen. Anal tube developed and with ventral portion sclerotized (subscaphium). Gnathos open, in lateral view elongated, with proximal portion wider than long, descending distally and with tegument fold projecting outward like a shell, in ventral view with conspicuous suture indicating the fusion between lateral arms of gnathos.

Valva semi-circular, with reduced spines on sacculus, a small notch on dorsal margin and a spine on distal portion. Fultura inferior bilobed.

Penis: aedeagus sclerotized, with bulbus ejaculatorius projecting from antero-dorsal opening, anterior portion curved descending, and posterior portion with sinuous of different shapes and dimensions.

Memphis glauce glauce (C. Felder and R. (Figure 12 **A–K).** Felder) Tegumen subtriangular in lateral view, anterior projection wider than long, tegument fold extended over entire anterior margin, projecting outward, appendices angulares wider than long. Arms from tegumen and saccus distinct. Anterior projection from saccus triangular in lateral view, descending, slender distally, projecting toward the left side; posterior projection present, shorter than anterior projection.

Uncus slender, with ventral margin sinuous, apex descending, in dorsal view fused with tegumen, uncus differing from tegumen by a light sclerotization. Anal tube developed and with ventral portion slcerotized (subscaphium). Gnathos open, in lateral view elongated, proximal portion expanded and distal portion semi-triangular, with median constriction, giving the structure a shovel-like appearance.

Valva subrectangular, each bearing a distal spine; spines reduced on sacculus and costa. Fultura inferior with narrow stripe.

Penis: aedeagus sclerotized, with bulbus ejaculatorius projecting from anterio-dorsal opening, posterior portion with tegument fold near posterior opening.

Memphis hirta (Weymer) (Figure 13 A-K). Tegumen subtriangular in lateral view, anterior projection wider than long. appendices angulares longer than wide. Arms from tegumen and saccus distinct, arm from tegumen with small distal expansions projecting anteriorly. Anterior projection from saccus subrectangular in lateral view, twice as long as wide; posterior projection present, shorter than anterior projection.

Uncus slender, with distal portion descending, in dorsal view fused with tegumen, uncus differing from tegumen by a light esclerotization. Anal tube developed and with ventral portion sclerotized (subscaphium). Gnathos open, in lateral view elongated, proximal portion wider than distal, from posterior view with tegument fold projecting posteriorly on distal portion, shell-like.

Valva semi-circular, each bearing a distal spine, conspicuous spine on dorsal margin and

areas with reduced spines on costa and sacculus. Fultura inferior wide and bilobed.

Penis: aedeagus sclerotized, with bulbus ejaculatorius projecting from antero-dorsal opening.

*Memphis lemnos* (H. Druce) (Figure 14 A– L). Tegumen semi-trapezoidal in lateral view, without anterior projection, tegument fold projecting outward, following anterior margin, appendices angulares longer than wide. Arms from tegumen and saccus fused. Anterior projection from saccus subtriangular in lateral view, slightly longer than wide; posterior projection present, shorter than anterior.

Uncus slender, ventral margin sinuous, apex slender and descending, in dorsal view fused with tegumen, but with proximal lateral portions distinct. Anal tube developed, with ventral portion sclerotized (subscaphium). Gnathos closed, in lateral view elongate, proximal portion wider than distal; distal portion with posterior and anterior projections in posterior view.

Valva subrectangular, bearing distal spine and tufts of setae on inner tegument. Fultura inferior bilobed.

Penis: aedeagus sclerotized, with bulbus ejaculatorius projecting from antero-dorsal opening, posterior-dorsal opening extended to manica.

*Memphis moruus stheno* (Prittwitz) (Figure 15 A–K). Tegumen subtriangular in lateral view, anterior projection wider than long, appendices angulares longer than wide. Arms from tegumen and saccus distinct. Anterior projection from saccus elongated in lateral view, at least three times longer than wide; posterior projection absent.

Uncus slender, ventral margin sinuous and distal portion descending, in dorsal view fused with tegumen, uncus differing from tegumen by a light sclerotization. Anal tube developed, with ventral portion sclerotized (subscaphium). Gnathos closed, in lateral view elongate, proximal portion wider than distal, and with a tegument fold projecting outward.

Valva subrectangular, with small spines on sacculus. Fultura inferior bilobed.

Penis: aedeagus sclerotized, with bulbus ejaculatorius projecting from antero-dorsal opening, posterior-dorsal opening extended to manica.

*Memphis philumena corita* (Fruhstorfer) (Figure 16 A–K). Tegumen subtrapezoidal in lateral view, anterior projection wider than long, tegument fold projecting outward, appendices angulares longer than wide. Arms from tegumen and saccus distinct. Anterior projection from saccus elongated in lateral view, about four times longer than wide; posterior projection present, shorter than anterior.

Uncus slender, with distal portion descending, in dorsal view fused with tegumen, uncus differing from tegumen by a light sclarotization. Anal tube reduced and entirely membranous. Gnathos open, in lateral view elongate, with proximal portion wider than distal; ventral proximal portion underneath tegumen, with distal tegument fold and dorsal distal notch.

Valva subrectangular and unmodified. Fultura inferior bilobed.

Penis: aedeagus sclerotized, with bulbus ejaculatorius projecting from antero-dorsal opening, posterior opening sinuous.

*Memphis polyxo* (H. Druce) (Figure 17 A– K). Tegumen subtrapezoidal in lateral view, anterior projection wider than long, tegument fold on anterior margin, projected outward, appendices angulares longer than wide. Arms from tegumen and saccus distinct. Anterior projection from saccus elongated in lateral view, twice as long as wide; posterior projection absent.

Uncus slender, with proximal portion enlarged and distal portion descending, in dorsal view fused with tegumen, uncus differing from tegumen by a light sclerotization. Anal tube developed, with ventral portion sclerotized (subscaphium). Gnathos closed, in lateral view elongate, proximal portion about twice wider than distal portion, tegument fold projecting outward.

Valva subrectangular, each bearing a distal spine, spines on sacculus and inner tegument small, near costa. Fultura inferior bilobed.

Penis: aedeagus sclerotized and sinuous, with bulbus ejaculatorius projecting from anterodorsal opening, posterior portion with tegument fold, posterior opening sinuous.

*Prozikania suprema* (Schaus) (Figure 18 A– L). Tegumen subtriangular in lateral view, without anterior projection, tegument fold projected outwards on anterior margin, appendices angulares longer than wide. Arms from tegumen and saccus fused. Anterior projection from saccus elongated in lateral view, at least twice longer than wide; posterior projection absent.

Uncus slender, apex slender and descending, in dorsal view fused with tegumen, uncus differing from tegumen by а light sclerotization. Anal tube developed, with ventral portion sclerotized (subscaphium). Gnathos closed, in lateral view elongate, proximal portion twice as wide as distal, outer margin sinuous, distal tegument fold projecting outward.

Valva subrectangular, distally with outer tegument expanded. Fultura inferior wide and forming a semi-circle.

Penis: aedeagus sclerotized, with bulbus ejaculatorius projecting from antero-dorsal opening, posterior opening irregular.

Pseudocharaxes xenocrates punctimarginale (Kaye) (Figure 19 **A–K).** Tegumen subtriangular in lateral view, anterior projection wider than long, tegument fold of anterior margin projected outward, appendices angulares longer than wide. Arms from tegumen and saccus separated, arms from tegumen long, the longest among the species studied in this work. Anterior projection fom saccus semi-triangular in lateral view, about twice as long as wide; posterior projection absent.

Uncus slender, ventral margin sinuous, apex slender and descending, in dorsal view fused with tegument, with triangular fenestra medially. Anal tube developed, with ventral portion sclerotized (subscaphium). Gnathos closed, in lateral view elongate, with distal tegument fold projected outward, giving the ghathos a shell-like appearance.

Valva rectangular, dorsal margin sinuous, spine on distal portion and setae on inner region, setae more numerous than in other species analyzed. Fultura inferior small and semi-circular.

Penis: aedeagus sclerotized, with bulbus ejaculatorius projecting from antero-dorsal opening, posterior opening sinuous.

Siderone nemesis catarina Dottax and Pierre (Figure 20 **A–K).** Tegumen subtriangular view, anterior in lateral than long, projection wider appendices angulares longer than wide. Arms from tegumen and saccus fused. Anterior projection from saccus elongated in lateral view, at least four times longer than wide; posterior projection absent.

Uncus slender, with distal portion descending, in dorsal view fused with tegumen, with triangular fenestra medially. Anal tube developed, with ventral portion sclerotized (subscaphium). Gnathos open, in lateral view elongate, completely enveloped by membranes.

Valva rectangular. Fultura inferior narrow and semi-circular.

Penis: aedeagus sclerotized, with bulbus ejaculatorius projecting from antero-dorsal opening, posterior opening large and sinuous.

Zaretis isidora (Cramer) (Figure 21 A–K). Tegumen subtriangular in lateral view, anterior projection wider than long, appendices angulares as wide as long. Arms from tegumen and saccus distinct. Anterior projection from saccus elongated in lateral view, at least three times longer than wide; posterior projection from saccus absent.

Uncus slender, apex slender and descending, in dorsal view fused with tegumen, distinct from tegumen by light sclerotization. Anal

tube developed, with ventral portion sclerotized (subscaphium). Gnathos open, in lateral view elongate, distal portion of both gnathos close to each other, tegument fold on distal portion shell-like.

Valva rectangular, with inner and outer ends conspicuous. Fultura inferior narrow and semi-circular.

Penis: aedeagus sclerotized, with bulbus ejaculatorius projecting from antero-dorsal opening.

#### Anaeomorphini

Anaeomorpha splendida Rothschild (Figure 22 A–K). Tegumen triangular in lateral view, anterior projection as wide as long, slightly concave on distal portion; appendices angulares wider than long. Arms from tegumen and saccus fused. Anterior projection from saccus subtriangular, in lateral view, wider than long; posterior projection present, subequal to anterior projection, with sharp tip.

Uncus thick, as large as tegumen, apex slender and descending, in dorsal view fused with tegumen, uncus differing from tegumen by a light sclerotization, setae on ventral margin at the beginning of tegument fold, which projects latero-ventrally and has a row of spines near distal portion. Anal tube developed and entirely membranous. Gnathos open, in lateral view with proximal and distal portions covered by the tegumen, in posterior view L-shaped, distal portion longer than proximal.

Valva subtriangular, sharp on distal portion. Fultura inferior rectangular, twice as long as tall, with concave dorsal margin.

Penis: aedeagus sclerotized, with bulbus ejaculatorius projecting from antero-dorsal

#### Preponini

of the posterior opening.

Archaeoprepona amphimachus pseudomeander (Fruhstorfer) (Figure 24 A–L, 33 E). Tegumen rectangular, anterior projection as wide as long, anterior tegument fold projecting outward; appendices angulares wider than long. Arms from tegumen and saccus fused. Anterior projection from saccus subrectangular in lateral view; posterior projection present, shorter than anterior.

Uncus thick, with short setae on ventral distal portion, apex slender, in dorsal view fused with tegumen, uncus differing from tegumen by a light sclerotization. Anal tube reduced and entirely membranous. Gnathos open, in lateral view long, with descending projection, apex shaped like a club, concave, striated.

Valva semi-triangular, distal expansion of inner tegument with short setae, conspicuous spine arising from outer tegument. Fultura inferior bilobed with dorsal median notch.

Penis: aedeagus membranous ventrally, with bulbus ejaculatorius projecting from anterodrosal opening.

Archaeoprepona chromus chromus (Guérin–Méneville) (Figure 24 A–K). Tegumen rectangular in lateral view, anterior projection wider than long, fold from anterior tegument projecting outward, appendices angulares longer than wide. Arms from tegumen and saccus distinct. Anterior projection from saccus elongated dorsoventrally in lateral view; posterior projection shorter than anterior projection, with sharp apex.

Uncus thick, in dorsal view fused with tegumen, uncus differing from tegumen by a light sclerotization, dorsal lobe with conspicuous boundaries. Anal tube reduced and entirely membranous. Gnathos open, in lateral view with proximal portion elongated, distal portion descending and striated.

Valva semi-triangular, distal expansion of tegument, and distal spine. Fultura inferior bilobed, with dorsal and ventral notches.

Penis: aedeagus with ventral portion membranous, bulbus ejaculatorius projecting from antero-ventral opening.

Archaeoprepona demophon muson (Fruhstorfer) (Figure 25 A-K). Tegumen rectangular in lateral view, anterior projection as wide as long, anterior tegument fold projected outward, appendices angulares wider than long. Arms from tegumen and saccus fused. Anterior projection from saccus absent; posterior projection present, developed, subtriangular with distal portion sharp.

Uncus thick, in dorsal view fused with tegumen, uncus differing from tegumen by a light sclerotization, dorsal lobe with conspicuous boundaries and enveloped by numerous spines on lateral portion. Anal tube reduced and entirely membranous. Gnathos open, in lateral view comma-like, striated on medial distal portion.

Valva subtriangular, distal expansions of inner tegument, with spine arising from outer tegument and with long setae. Fultura inferior bilobed and articulated with ventral area of valva.

Penis: aedeagus, ventral portion membranous, medial-anterior region dilated in lateral view,

narrow in dorsal or ventral views, bulbus ejaculatorius projecting from antero-dorsal opening.

Archaeoprepona demophoon andicola (Fruhstorfer) (Figure 26 A–G, 27 A–E). Tegumen rectangular in lateral view, anterior projection wider than long, tegument fold reduced and covering part of the hollow, which distinguishes the arms from tegumen; appendices angulares wider than long. Arms from tegumen and saccus distinct. Anterior projection from saccus subrectangular in lateral view, twice as long as wide; posterior projection, slender distally and descending.

Uncus thick, in dorsal view fused with tegumen, uncus differing from tegumen by a light sclerotization, dorsal lobe with conspicuous boundaries. Anal tube reduced and entirely membranous. Gnathos open, in lateral view comma-like, striated on distal two thirds.

Valva semi-triangular, distal expansion of inner tegument, spine arising from outer tegument. Fultura inferior bilobed and dorsal inner margins rounded.

Penis: aedeagus wider dorsoventrally when compared with other species of the genus, anterior ventral portion rounded, small anterodorsal lobe and median constriction preceding manica, ventral portion membranous, bulbus ejaculatorius projecting from antero-dorsal opening.

Archaeopreponalicomedeslicomedes(Cramer)(Figure28A–L).Tegumenrectangular in lateral view, anterior projectionwider than long, anterior ventral foldprojected outward, appendices angulareswider than long, ventral margin sinuous.

from tegumen and saccus distinct. Anterior projection from saccus subtriangular in lateral view; posterior projection shorter than anterior, diamond-shaped.

Uncus thick, in dorsal view fused with tegument, uncus differing from tegumen by a light sclerotization, dorsal lobe with conspicuous boundaries. Anal tube reduced and entirely membranous. Gnathos open, in lateral view with two flattened, folded plates in lateral view, distal plate descending and striated on outer portion.

Valva subtriangular, distal expansion of inner tegument, and distal spine arising from outer tegument. Fultura inferior bilobed, lateral margins sinuous, a smooth ventral-median notch and deep dorsal median notch.

Penis: aedeagus, ventral portion entirely membranous, bulbus ejaculatorius projecting apically

Archaeoprepona meander castorina (E. Mayr) (Figure 29 A–K). Tegumen rectangular in lateral view, anterior projection wider than long, tegument fold small and projected outward, appendices angulares longer than wide. Arms from tegumen and saccus distinct. Anterior projection of saccus semi-triangular in lateral view, longer than wide; posterior projection present, shorter than anterior projection.

Uncus slender, with expansion from median to distal portion, ventral margin sinuous, apex slender and descending, in dorsal view fused with tegumen, uncus differing from tegumen by a light sclerotization. Anal tube reduced and entirely membranous. Gnathos open with distal portion striated. Valva semi-triangular, distal expansion arising from inner tegument, and distal spine arising from outer tegument. Fultura inferior bilobed.

Penis: aedeagus with a concavity on ventral median portion, membranous ventrally, bulbus ejaculatorius projecting apically.

#### Prepona claudina annetta (Gray)

(Figure 30 A–K). Tegumen semi-trapezoidal in lateral view, anterior projection wider than long, deep depression on posterior ventral portion, appendices angulares wider than long. Arms from tegumen and saccus distinct. Anterior projection from saccus oval in lateral view, longer than wide; posterior projection present, shorter than anterior.

Uncus thick, elevated on proximal dorsal portion, with distal portion descending, in dorsal view fused with tegumen, uncus differing from tegumen by a light sclarotization. Anal tube reduced, entirely membranous and linked to the gnathos. Gnathos open, in lateral view elongate, with long arm and with spiny apical club on, projected dorsally.

Valva triangular. Fultura inferior rectangular, taller than wide, dorsally with two convex areas covered with small spines, tegument fold projecting laterally.

Penis: aedeagus sclerotized dorsally and with ventral sclerotization restricted to median region, bulbus ejaculatorius projects from antero-ventral opening.

**Prepona laertes laertes (Hubner) (Figure 31 A–K).** Tegumen semi-trapezoidal in lateral view, anterior projection wider than long, dorsal margin with small folds, appendices angulares wider than long. Arms from tegumen and saccus distinct. Anterior projection from saccus oval in lateral view; posterior projection absent.

Uncus thick, apex slender and descending, in dorsal view fused with tegumen, conspicuous constriction between uncus and tegumen. Anal tube reduced and entirely membranous. Gnathos open, in lateral view long, with spiny club projected dorsally, and extended beyond uncus.

Valva triangular. Fultura inferior rectangular, taller than wide, with basal lateral folds projected outward.

Penis: aedeagus sclerotized, with tegument fold preceding manica, bulbus ejaculatorius projecting from antero-ventral opening.

**Prepona pylene pylene Hewitson (Figure 32 A–K).** Tegumen semi-trapezoidal, anterior projection wider than long, appendices angulares wider than long. Arms from tegumen and saccus distinct. Anterior projection from saccus oval in lateral view; posterior projection absent.

Uncus thick, with distal portion concave and descending, in dorsal view fused with tegumen, uncus differing from tegumen by the latter through a dorsal process on the proximal portion. Anal tube reduced and entirely membranous. Gnathos open, in lateral view long, with spiny club projecting dorsally, and extended beyond uncus.

Valva triangular. Fultura inferior taller than wide, with lateral fold projecting outward.

Penis: aedeagus sclerotized, bulbus ejaculatorius projecting from antero-ventral opening.

#### Discussion

#### Tegumen

This sclerite is structurally diverse across the group, with differences in at least five areas. Four shapes have been identified in lateral view: (1) semi-trapezoidal, clustering species of Agrias, Prepona, and Hypna, as well as Memphis philumena corita, Memphis lemnos, and Memphis polyxo; (2) rectangular, common to species of Archaeoprepona and Noreppa; (3) triangular, clustering Anaea troglodyta with Anaeomorpha splendida; and (4) subtriangular, common among species of clustering Anaeini, and species of Coenophlebia, Consul. Fountainea. Prozikania, Pseudocharaxes, Siderone, Zaretis, and most species of Memphis.

With a few exceptions, enumerated below, the anterior projection of the tegumen is developed. It is lacking in most species of Fountainea (the anterior projection is found only in Fountainea ryphea phidile, of the four species studied) and a few species of Anaeini (Memphis acidalia victoria, Memphis lemnos, and Prozikania suprema). When present, the anterior projection of the tegumen is often developed, longer than wide or as wide as long. Such projection is reduced or is as wide as long only in Anaeomorpha splendida, Archaeoprepona amphimachus pseudomeander, Archaeoprepna and demophon muson.

Other modifications in the tegumen were the presence or absence of a fold in the tegument on the anterior margin and appendices angulares. The latter are ubiquitous among the species studied and vary in their dimensions, being wider than long, as wide as long, or longer than wide.

#### Saccus

Some differentiation between the ventral arms from the tegumen and the dorsal arms from the saccus was found in 23 of the 31 species analyzed, with special emphasis for *P*. *xenocrates punctimarginale*, with the arms of the tegumen separated and distant from the arms of the saccus. Therefore, the fusion between these two structures has not revealed a suitable character to group species or genera.

Thirty species have a saccus with an anterior projection. The degree of development of this projection varies among species, and six shapes have been identified for it: oval, common among species of Agrias and Prepona; rectangular, a shape found only in Anaea troglodyta; subtriangular, characteristic of Anaeomorpha splendida, Archaeoprepona andicola, demophoon Archaeoprepona licomedes Pseudocharaxes licomedes. xenocrates punctimarginale, and Memphis lemnos; subrectangular, as in Archaeoprepona amphimachus pseudomeander, Archaeoprepona domophon muson. and *Memphis* hirta; triangular, as in Archaeoprepona meander castorina and Memphis glauce glauce; and elongated, species characteristic of of Consul, Fountainea and Hypna, as well as Coenophlebia archidona, Noreppa chromus chromus, Prozikania suprema, Siderone nemesis catarina, Zaretis isidora, Memphis acidalia victoria, Memphis moruus stheno, Memphis phlilumena corita, and Memphis polyxo.

A posterior projection arising from the saccus was present in approximately 50% of the species studied. This projection can be shorter than the anterior projection, a widespread state, or it can be subequal to or longer than the anterior projection, a state present in *Anaeomorpha splendida* and *Archaeoprepona domophon muson*.

#### Uncus

Among the structural differences found in the uncus, we highlight its shape in lateral view, which can be thick (three times longer than wide) or slender (more than three times longer than wide). The tribe Preponini contains only one species with a slender uncus. Archaeoprepona meander meander. All other species in this tribe, as well as the species Hypna and Coenophlebia (Anaeini), have a thick uncus. The uncus is slender in species of Anaea. Consul. Memphis, Fountainea, Siderone, Prozikania, Pseudocharaxes, and Zaretis.

The uncus also varies in the presence or absence of setae on its dorsal portion. The Anaeomorpha uncus of splendida, Archaeoprepona amphimachus pseudomeander, and Prepona laertes laertes have a dorsal seta. Other differences in this structure among species are as follows: the presence or absence of crests (present only in two subspecies of *Hypna clytemnestra*), presence or absence of spines on the dorsal portion (present in Anaeomorpha splendida, Archaeoprepona amphimachus pseudomeander, and Prepona laertes laerte), and the presence or absence of a fenestra. When present, the fenestra may be found either on the central portion (Pseudocharaxes xenocrates punctimarginale and Siderone nemesis catarina) or at the base of the uncus (Hypna clytemnestra forbesi and Hypna clytemnestra huebneri).

#### Anal tube

The degree of sclerotization of the ventral portion was the only difference found in the anal tube. Species that have the ventral

portion of the anal tube sclerotized were considered as having a subscaphium. This character can be used to separate between Anaeini, Anaeomorphini and Preponini. Species of Preponini and Anaeomorphini do not have a subscaphium, whereas species of Anaeini do, with two exceptions, *Fountainea nessus* and *Fountainea ryphea phidile*.

#### Gnathos

The gnathos has two basic shapes in the species studied. It may form a handle enveloping the anal tube, in which case it is called "single" or "close," or it may be divided into two distinct arms, and be called "double" or "open." In our sample, the gnathos is single in species of Anaeini and double in species of and Anaeomorphini Preponini, with some modifications among species.

Five aspects were identified concerning the shape of the gnathos: elongated, present in species of *Agrias* and *Prepona*; shaped as an anvil, unique to *Anaea troglodyta*; covered by the tegument, unique to *Anaeomorpha splendida*; thick, present in *Archaeoprepona* and *Noreppa*; with base wider than distal portion, as in Anaeini, except for *Anaea troglodyta*. Regarding the modifications on the distal portion, they can be smooth, with spiny club, or with distal margin serrated or striated and enveloped by membranes.

#### Valvae

The valvae assume various shapes: triangular, as in Agrias and Prepona; subtriangular, in species of Anaea, Anaeomorpha, and Archaeoprepona; circular, common to the species of Hypna and Consul, as well as Memphis acidalia victoria, Memphis hirta, and Memhis polyxo; rectangular, as in Pseudocharaxes xenocrates punctimarginale,

Siderone nemesis catarina, and Zaretis isidora; and subrectangular, as in all species of Fountainea, Coenophlebia archidona, Memphis philumena corita, Memphis moruus stheno, Memphis lemnos, Memphis glauce glauce, and Prozikania suprema.

Besides the shape, the structure of the valvae may also vary. For instance, the presence of an apical projection is a variable feature within tribes, but stable within genera. In our sample, the valva of all species of *Archaeoprepona*, *Fountainea*, *Consul*, *Prozikania*, *Pseudocharaxes*, and *Memphis* (except for *Memphis moruus stheno*) had an apical projection.

Beyond the characters described above, the following features have been found to be variable: presence or absence of spines on costa (present in *Fountainea halice halice, Memphis glauce glauce, Memphis hirta, Memphis polyxo*); presence or absence of spines on the saccus (present in species of *Memphis*); presence or absence of an apical projection of the valva, presence or absence of setae on the inner tegument of the valva (present only in *Memphis lemnos* and *Pseudocharaxes xenocrates punctimarginale*). These characters can be further investigated, and the location and shape of spines can be detailed.

#### **Fultura inferior**

Differences were found in the fultura inferior of all species. Within this structural diversity, we were able to devise some basic aspects. One is the general shape, with six variations: rectangular, as in all species of *Agrias* and *Prepona*; semi-circular, as in *Anaeomorpha splendida*, *Fountainea ryphea phidile*, *Prozikania suprema*, and *Pseudocharaxes* 

*xenocrates punctimarginale*; elongated, present only in *Coenophlebia archidona*; Vshaped, present among *Hypna* subspecies; narrow, stripe-like, unique to *Memphis hirta*; and bilobed, present in all other species analyzed.

Other characters identified for the fultura inferior were useful for grouping species, such as the presence of absence of tegument folds, uniting species of *Agrias* and *Prepona*. Another such character is the presence or absence of spines and their location.

#### Penis

The penis presents variations in the aedeagus, which can be completely sclerotized or membranous ventrally. A ventrally membranous aedeagus was found in the studied species of *Archaeoprepona* and *Noreppa*.

Another character that varied in the penis was the direction of the projection from the bulbus ejaculatorius. It was ventral in species of *Agrias* and *Prepona*, and also in *Noreppa chromus chromus*. In *Archaeoprepona licomedes licomedes* and *Archaeoprepona meander castorina*, this projection was ventral. It was dorsal in 25 of the 31 species studied.

#### **Final considerations**

The present analysis revealed the importance of conducting detailed morphological analyses that focus on certain tagmas, particularly for groups whose morphology has been neglected, such as the Charaxinae. It is possible that this group has been poorly studied because specimens are rare and consequently they are considered precious. For this reason, their taxonomy has been based on venation and coloration of the wings only. The observation of Rydon (1971) was confirmed, because the differences of the gnathos, club-like and striated-like, were able to differentiate Archaeoprepona and Prepona. According to the results, species of Preponini can be divided into two distinct groups, (including Prepona Prepona claudina annetta) and Archaeoprepona (including Archaeoprepona chromus chromus), thus confirming the arrangement of the genera proposed by Ortiz-Acevedo and Willmott (2013).

Among the characters observed in this study, the following putative synapomorphies support groups within Preponini:

*Archaeoprepona*: Aedeagus with ventral portion membranous and distal portion of gnathos striated;

*Prepona*: Presence of lateral folds in the tegument of the fultura inferior, and gnathos with a club covered with spines.

Morphological characters can resolve taxonomic problems, and can still be more exploited, thus making them more accurate when included in analyses such as female genitalia, color and shape of wings, and even the shape and color of odoriferous scales in males.

The structural modifications of *Anaeomorpha splendida* confirm a tribe, although in some aspects it resembles *Anaea troglodyta*, which deserves attention and further research about the degree of closeness between these two species belonging to different tribes, Anaeomorphini and Anaeini, respectively.

Contrasting with the genitalia of species of Preponini, the gentialia of Anaeini species is variable, making it difficult to establish structural patterns within this group. However, it is possible to distinguish among species based on some genitalic characters. For instance, species of *Memphis* and *Fountainea* are easily separated by the presence and location of spines on the valva. For Comstock (1961), the great variety of forms of gnathos, which despite having a structural pattern as a basal line can provide many features useful for the determination of species.

The results obtained herein reflect detailed dissections and observations of structures from different views (positions) that had not been previously used in the taxonomy of the group. Therefore, they include structures not previously considered. These characters can be used in the future in a phylogenetic analysis or when devising more consistent diagnostic characters for species and genera.

The detailing of such structures can be geared towards the maintenance of the status of Prozikania and Pseudocharaxes as genera, synonymized by Lamas (2004)with Polygrapha, the first as Zikania (nom. nud.), and again used by Salazar (2008). The significant differences in the genitalia of the species of both genera as described and illustrated in this study corroborate the observations of Comstock (1961), Rydon (1971), and Salazar (2008), justifying the maintenance of the genera, despite being monotypical.

#### Acknowledgments

We thank the Centro de Microscopia Eletrônica da Universidade Federal do Paraná for the micro photography, and Conselho Nacional de Desenvolvimento Científico e Tecnológico for the stipends granted to the authors.

#### References

Ackery PR. 1984. Systematic and faunistic studies on butterflies. In: Vane-Wright RI, Ackery PR, Editors. *The Biology of Butterflies*. Academic Press.

Bilotta IG. 1994. Morfologia comparada do abdome das espécies sulbrasileiras de Morphinae (Lepidoptera, Nymphalidae). *Revista Brasileira de Zoologia* 11: 737–748.

Caldas A. 1997. Notes on the male genitalia of the Anaea ryphea – Anaea eurypyle complex (Nymphalidae). *Journal of Lepidopterists' Society* 51: 83–90.

Casagrande MM. 1979. Sobre *Caligo beltrao* (Illiger). IV. Morfologia externa do adulto – abdome (Lepidoptera, Satyridae, Brassolnae). *Revista Brasileira de Biologia* 39: 711–716.

Comstock WP. 1961. *Butterflies of the American Tropics. The Genus Anaea, Lepidoptera, Nymphalidae.* American Museum of Natural History.

Ehrlich PR. 1958. The comparative morphology, phylogeny and classification of the butterflies (Lepidoptera: Papilionoidea). *University of Kansas Science Bulletin* 39: 305–370.

Klots AB. 1956. Lepidoptera. In: Tuxen SL, Editor. *Taxonomist's Glossary of Genitalia in Insects*. Munksgaard.

Kuznetsov NY. 1915. Fauna of Russia and Adjacent Countries, Lepidoptera I. Petrograd.

#### Journal of Insect Science | www.insectscience.org

Lamas G. 2004. Nymphalidae. Charaxinae. Tribe Anaeini. In: Lamas G, Editor. Checklist: Part 4A. Hesperioidea – Papilionoidea. In: Heppner JB, Editor. *Atlas of Neotropical Lepidoptera*, volume 5A. Association for Tropical Lepidoptera/Scientific Publishers.

Mielke CGC, Mielke OHH, Casagrande MM. 2004a. Estudo comparado da morfologia externa de *Zaretis itys itylus* (Westwood) e *Agrias claudina annetta* (Gray) (Lepidoptera, Nymphalidae, Charaxinae) I. Cabeça, apêndices e região cervical. *Revista Brasileira de Zoologia* 21: 657–369.

Mielke CGC, Mielke OHH, Casagrande MM. 2004b. Estudo comparado da morfologia externa de *Zaretis itys itylus* (Westwood) e *Agrias claudina annetta* (Gray) (Lepidoptera, Nymphalidae, Charaxinae) II. Tórax e apêndices. *Revista Brasileira de Zoologia* 21: 421–433.

Mielke CGC, Mielke OHH, Casagrande MM. 2004c. Estudo comparado da morfologia externa de *Zaretis itys itylus* (Westwood) e *Agrias claudina annetta* (Gray) (Lepidoptera, Nymphalidae, Charaxinae) III. Abdome. *Revista Brasileira de Zoologia* 21: 905–912.

Niculescu EV. 1978. L'armature génitale chez les Lépidoptères. *Bulletin de La Société Entomologique de Mulhouse* (Supplement): 1–40.

Ogata M, Okada Y, Okagaki H, Sibatani A. 1957. Male Genitalia of Lepidoptera: Morphology and Nomenclature III. Appendages Pertaining to the Tenth Somito. *Annals of the Entomolgical Society of America* 50: 237–244. Oiticica JF. 1946. Sobre a Morfologia do Penis em Lepidoptera -Boletim do Museu Nacional. *Zoologia* 50: 1–79.

Okagaki H, Sibatani A, Ogata M, Okada Y. 1954. Male Genitalia of Lepidoptera: Morphology and Nomenclature II. Morphology Significance of Sacculus and Furca. *Annals of the Entomolgical Society of America* 48: 439–442.

Ortiz-Acevedo E, Willmott KR. 2013. Molecular systematics of the butterflies tribe Preponini (Nymphalidae: Charaxinae). *Systematic Entomology* 38: 440–449.

Petersen W. 1904. Die Morphologie der Generationsorgane der Schmetterlinge und ihre Bedeutung für die Artbildung. *Mémoires de L'Académie Impériale dês Sciences de St. Pétersbourg* VIII Série XVI, 8: 1–84.

Pierce FN. 1909. *The genitalia of the group Noctuidae of the Lepidoptera of the British Islands*. E.W. Classey.

Pierce FN. 1914. *The genitalia of the British Geometridae*. Liverpool.

Rydon AHB. 1971. The Systematics of the Charaxidae (Lep: Nymphaloidea). *Entomologist's Record and Journal of Variation* 83: 219–233, 311–316, 336–341, 384–388.

Salazar JAE, Constantino LM. 2001. Synthesis of the description of new genera for South America: Rydonia, Annagrapha, Pseudocharaxes, Muyshondtia, Zikania (Lepidoptera, Nymphalidae). *Lambillionea* (supplement 2) 3: 344–370.

Salazar JAE. 2008. Some studies on palpi belonging to neotropical Charaxids and notes

on the wing pattern and behavior of several genera (Lepidoptera: Nymphaloidea, Charaxidae). *Boletin Científico Museo de Historia Natural, Universidad Caldas* 12: 171–205.

Sibatani A, Masami O, Yoshio O, Hiromo O. 1954. Male genitalia of Lepidoptera. Morphology and Nomenclature. I. Divisions of the valvae in Rhopalocera, Phalaenidae (Noctuidae) and Geometridae. *Annals of the Entomological Society of America* 47: 93– 106.

Simonsen TJ. 2006. Glands, muscles and genitalia. Morphological and phylogenetic implications of histological characters in the male genitalia of Fritillary butterflies (Lepidoptera: Nymphalidae: Argynini). *Zoologica Scripta* 35: 231–241.

Simonsen TJ. 2006. The male genitalia segments in fritillary butterflies: Comparative morphology with special reference to the "rectal plate" in *Issoria* (Lepidoptera: Nymphalidae). *European Journal of Entomology* 103: 425–432. Snodgrass RE. 1935. *Principles of insect morphology*. McGrow – Hill Book Company.

Snodgrass RE. 1957. A revised interpretation of the external reproductive organs of male insects. *Smithsonian Miscellaneous Collections* 135: 1–60.

Srivastava KP. 1965. Studies on lemon– butterfly, *Papilio demoleus* L. (Lepidoptera) Part V – skeleton-muscular system of abdomen. *Zoologischer Anzeiger* 177: 217– 236.

Tóth JP, Vargas Z. 2011. Inter-intraspecific variation in the genitalia of the "Melitaea phoebe group" (Lepidoptera, Nymphalidae). *Zoologischer Anzeiger* 250: 258–268.

| Tribe   | Charaxinae Neotropical  |   |
|---|---|---|
| the second se | Species   | Material studied  |
|   | Anaea troglodyta (Fabricius, 1775)  | MEXICO, Yucatan: Piste, 1 male, 15 October 1959, Welling leg., DZ16.619 (DZUP).   |
|   | Coenophlebia archidona (Hewitson, 1860)   | BOLIVIA, Chapare: 400 m a.s.l., 1 male, November 1949, Justus leg., DZ17.158 (DZUP).  |
|   | Consul electra (Westwood, 1850)   | MEXICO, Veracruz: Quintana Roo, 1 male, 15 October 1960, Welling leg., DZ17.157; Tuxtla, S. Tiago, 800 m  |
|   |   | a.s.l., 1 male, 30 October–2 November 1975, Becker leg., DZ19.186 (DZUP).   |
|   | Consul fabius drurii (Butler, 1874)   | male, 6. March 1971, DZ19.091; Paraná: Foz do Iguacu, 1 male, 3 December 1966, DZ18.845 (DZUP).   |
|   | Frank in a local state of (Ileviter 1974)   | BRASIL, Bahia: Campo Formoso, 500 m a.s.l., 1 male, 10 July 1973, Mielke leg., DZ20.302; Pernambuco:  |
|   | Foundationea glycerium cratais (Hewitson, 1874)   | Gravatá, 300 m a.s.l., 1 male, 8 June 1951, Ebert leg., DZ19.945 (DZUP).  |
|   |   | BRASIL, São Paulo: Teodoro Sampaio, Parque Estadual Morro do Diabo, 1 male, 17–19 August 1989, Mielke &   |
|   | Fountainea halice halice (Godart, [1824])   | Lasagrande legs, DZ22.444; Kio de Janeiro: Kio de Janeiro, Lagoinha, 250 m a.s.i., 1 maie, 15 June 1952, Ebert,   |
|   |   | BOLÍVIA, Chulumani: 1400 m a.s.l., 1 male, 22–24 October 1983, Mielke & Casaorande leos. DZ16 485   |
|   | Fountainea nessus (Latreille, [1813])   | (DZUP).   |
|   |   | BRASIL, Santa Catarina: Joinville, 200 m a.s.l., 1male, Miers leg., DZ22.564; Paraná: Londrina, 1 male, 10  |
|   | Fountainea ryphea phidile (Geyer, 1837)   | October 1982, Mielke leg., DZ22.544 ; Mato Grosso: Barra do Bugre, 200 m a.s.l., 1 male, 29 June 1972, Mielke   |
|   |   | & Brown legs, DZ22.554 (DZUP).<br>PRASII Pernambuaa: Limaaira Lunala 6 Juna 1057 Ebert lag. DZ16 577: Gravatá 200 m a s. L. Lunala 24   |
|   | Hypna clytemnestra forbesi Godman & Salvin, 1884  | June 1957, Ebert leg. DZ18 979 (DZ1P)   |
|   | II  | BRASIL, Espírito Santo: Baixo Gandu, 1 male, 20 August 1970, C. & C. T. Elias legs, DZ16.556; Minas Gerais:   |
|   | Hypna ciytemnestra huebneri Butter, 1866  | Parque Estadual do Rio Doce, 200 m a.s.l., 1 male, 27. July .1972, H. & H. D. Hebert legs, DZ18.962 (DZUP).   |
|   |   | Paraná: Londrina, 1 male, 10 September 1985, Mielke & Casagrande legs, DZ22.494; Fênix, 300 m a.s.l., 2 males   |
|   | Memphis acidalia victoria (H. Druce, 18//)  | 29 April 1987, Mielke & Casagrande legs, DZ22.484 e DZ22.534; Foz do Iguaçu, Parque Nacional Iguaçu, 1  |
| Anacini   | Memphis glauce glauce (C. Felder & R. Felder, 1862)   | BRASIL Acre: Santa Rosa do Purus 14 2Km SE 3 males 12 August 2008 Carneiro leg DZ9 294 DZ9 380  |
| , machin  |   | DZ9.516; PERU, Madre de Dios: Parque Nacional del Manu, Pakitza, 340 m a.s.l., 1 male, 4 November 1991.   |
|   |   | Lamas leg., DZ16.506 (DZUP).  |
|   | Memphis hirta (Weymer, 1907)  | BRASIL, Santa Catarina: São Bento do Sul, Rio Vermelho, 850 m a.s.l., 1 male, 15 March 1986, Mielke & Rank  |
|   |   | legs, DZ22.464; Parana: Ponta Grossa, 1 male, September 1948, Woym leg., DZ22.524; Tijucas do Sul, Rincão, 950 m a s. 1. Junela 10 Marsh 1071. Mielka leg. DZ22.514; Tunua Pron. Sr. Luiz Pieldi, 1 mala 16 December 2010.  |
|   |   | 2007 Dolibaina leg. DZ22.514, 1019, 51, Luis Rickii, 1 maie, 10 December  |
|   | Memphis lemnos (H. Druce, 1877)   | BRASIL. Acre: Santa Rosa do Purus, 1 male, 6-8 August 2008, Mielke & Carneiro legs, DZ9.515 (DZUP).   |
|   | Memphis moruus stheno (Prittwitz 1865)  | ARGENTINA, Misiones: Almirante Brown, Reserva Yacutinga, 3 males, 2-5 March 2007, Mielke & Casagrande   |
|   |   | legs, DZ9.943, DZ22.454 e DZ22.504 (DZUP).  |
|   | Memphis philumena corita (Fruhstorfer, 1916)  | BRASIL, Santa Catarina: Joinville, 1 male, 9 October 1971, Mielke leg., DZ22.094; 1 male, 3 September 1973,<br>Mielke leg., DZ21 974: 5 April 1980, 1 male, Mielke leg., DZ19 921 (DZUP).   |
|   | Manufilia and and (II Dense 1874)   | BRASIL, Santa Catarina: Joinville, 200 m a.s.l., 1 male, 8 February 1974, Mielke & Miers legs, DZ16.478; 1  |
|   | Memphis polyxo (11. Druce, 1874)  | male, 19 December 1975, Mielke leg., DZ22.154 (DZUP).   |
|   | Prozikania suprema (Schaus, 1920)   | BRASIL, Minas Gerais: Passa Quatro, Fazenda dos Campos, 1500 m a.s.l., 1 male, 8 August 1911, Zikán leg.,<br>DZ10 009, 1 male, 2 Fabruary 1014, Zikán leg., DZ10 102, 1 male, 22 July, 1016, Zikán leg., DZ16 584 (DZUP)  |
|   |   | Material studies, BRASIL, Mato Grosso: Cáceres, Rio Vermelho, 400 m a.s.l., 1 male, 3 July 1972, Mielke &   |
|   | Pseudocharaxes xenocrates punctimarginale (Kaye, 1918)  | Brown legs, DZ16591; Rondônia: Pimenta Bueno, 1 male, July 1970, DZ18.813 (DZUP).   |
|   |   | BRASIL, Distrito Federal: Brasília, Parque do Gama, 1000 m a.s.l., 1 male, 17 June 1972, Mielke & Brown legs,   |
|   | Siderone nemesis catarina Dottax & Pierre, 2009   | DZ16612; Mato Grosso: Diamantino, Fazendo São João, Alto do Rio Arinos, 1 male, 22 January 1978, Mielke &   |
|   |   | BRASIL, Paraná: Curitiba, Cascatinha, 1 male, 23 April 1966, Mielke leg., DZ21,934; Terra Boa, 650 m a s.l., 1  |
|   | Zaretis isidora (Cramer, 1779)  | male, 10 December 1975, Moure, Mielke & Wedderhoff legs, DZ16563; Ventania, 1000 m a.s.l., 1 male, 8  |
|   |   | September 2007, Mielke leg., DZ22.054 (DZUP).   |
| naeomorphini  | Anaeomorpha splendida Rothschild, 1894  | BRASIL, Rondônia: Candeias do Jamari, Balneário Rio Preto, 1 male, 1–7 September 2001, Mielke &   |
|   |   | Concernando laco (MM54 169 (OM))  |
|   |   | Casagrande legs. OM54.168 (OM).<br>BRASIL, Santa Catarina; Joinville, 1 male, 28 October 1968. Mielke leg., DZ22.144; São Bento do Sul. Rio   |
|   | Anakazannanana amphimachus neudomean dar (Embeterfor 1006)  | Casagrande legs. OM54.168 (OM).<br>BRASIL, Santa Catarina: Joinville, 1 male, 28 October 1968, Mielke leg., DZ22.144; São Bento do Sul, Rio<br>Vermelho, 1 male, DZ22.254; 2 males, 2 December 1969, Ebert leg., DZ22.224 e DZ 22.044; 1 male, 10 January   |
|   | Archaeoprepona amphimachus pseudomeander (Fruhstorfer, 1906)  | Casagrande legs. OM54.168 (OM).<br>BRASIL, Santa Catarina: Joinville, 1 male, 28 October 1968, Mielke leg., DZ22.144; São Bento do Sul, Rio<br>Vermelho, 1 male, DZ22.254; 2 males, 2 December 1969, Ebert leg., DZ22.224 e DZ 22.044; 1 male, 10 January<br>1971, Ebert leg., DZ22.174; Minas Gerais: Caratinga, Estação Ecológica de Caratinga, 400 m a.s.l., 1 male, 29  |
|   | Archaeoprepona amphimachus pseudomeander (Fruhstorfer, 1906)  | Casagrande legs. OM54.168 (OM).<br>BRASIL, Santa Catarina: Joinville, 1 male, 28 October 1968, Mielke leg., DZ22.144; São Bento do Sul, Rio<br>Vermelho, 1 male, DZ22.254; 2 males, 2 December 1969, Ebert leg., DZ22.224 e DZ 22.044; 1 male, 10 January<br>1971, Ebert leg., DZ22.174; Minas Gerais: Caratinga, Estação Ecológica de Caratinga, 400 m a.s.l., 1 male, 29<br>January5 February 2003, Mielke & Casagrande legs, DZ20.622 (DZUP).  |
|   | Archaeoprepona amphimachus pseudomeander (Fruhstorfer, 1906)<br>Archaeoprepona chromus chromus (Guérin-Méneville, [1844])   | Casagrande legs. OM54.168 (OM).<br>BRASIL, Santa Catarina: Joinville, 1 male, 28 October 1968, Mielke leg, DZ22.144; São Bento do Sul, Rio<br>Vermelho, 1 male, DZ22.254; 2 males, 2 December 1969, Ebert leg, DZ22.224 e DZ 22.044; 1 male, 10 January<br>1971, Ebert leg, DZ22.174; Minas Gerais: Caratinga, Estação Ecológica de Caratinga, 400 m a.s.l., 1 male, 29<br>January–3 February 2003, Mielke & Casagrande legs, DZ20.622 (DZUP).<br>COLÓMBIA, Valle do Cauca: Dagua, 1 male, 32 October 1901. Casagrande lag. DZ19633;<br>DEFEUL Modea do Dios: Tambonator Reserva 300 m a c.l. 1 male, 25 October 1901. Casagrande lag. DZ19633;   |
|   | Archaeoprepona amphimachus pseudomeander (Fruhstorfer, 1906)<br>Archaeoprepona chromus chromus (Guérin-Méneville, [1844])   | Casagrande legs. OM54.168 (OM).<br>BRASIL, Santa Catarina: Joinville, 1 male, 28 October 1968, Mielke leg., DZ22.144; São Bento do Sul, Rio<br>Vermelho, 1 male, DZ22.254; 2 males, 2 December 1969, Ebert leg., DZ22.224 e DZ 22.044; 1 male, 10 January<br>1971, Ebert leg., DZ22.174; Minas Gerais: Caratinga, Estação Ecológica de Caratinga, 400 m a.s.l., 1 male, 29<br>January–3 February 2003, Mielke & Casagrande legs, DZ20.622 (DZUP).<br>COLÓMBIA, Valle do Cauca: Dagua, 1 male, January 1967, DZ16.570 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg., DZ19633;<br>BRASIL, Acre: Senador Guiomard Reserva, 621mala 1 male, 29-30 Sentember 2006, Mielke & Casagrande legs.   |
|   | Archaeoprepona amphimachus pseudomeander (Fruhstorfer, 1906)<br>Archaeoprepona chromus chromus (Guérin-Méneville, [1844])<br>Archaeoprepona demophon muson (Fruhstorfer, 1905)  | Casagrande legs. OM54.168 (OM).<br>BRASIL, Santa Catarina: Joinville, 1 male, 28 October 1968, Mielke leg., DZ22.144; São Bento do Sul, Rio<br>Vermelho, 1 male, DZ22.254; 2 males, 2 December 1969, Ebert leg., DZ22.224 e DZ 22.044; 1 male, 10 January<br>1971, Ebert leg., DZ22.174; Minas Gerais: Caratinga, Estação Ecológica de Caratinga, 400 m a.s.l., 1 male, 29<br>January–3 February 2003, Mielke & Casagrande legs, DZ20.622 (DZUP).<br>COLÓMBIA, Valle do Cauca: Dagua, 1 male, January 1967, DZ16.570 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg., DZ19633;<br>BRASIL, Acre: Senador Guiomard, Reserve, 300 m a.s.l., a male, 29-30 September 2006, Mielke & Casagrande leg.,<br>DZ20.402; 1 male, 23–27 July 2008, Mielke & Casagrande legs, DZ19.414; Porto Acre, Reserva Humaită, 2   |
|   | Archaeoprepona amphimachus pseudomeander (Fruhstorfer, 1906)<br>Archaeoprepona chromus chromus (Guérin-Méneville, [1844])<br>Archaeoprepona demophon muson (Fruhstorfer, 1905)  | Casagrande legs. OM54.168 (OM).<br>BRASIL, Santa Catarina: Joinville, 1 male, 28 October 1968, Mielke leg., DZ22.144; São Bento do Sul, Rio<br>Vermelho, 1 male, DZ22.254; 2 males, 2 December 1969, Ebert leg., DZ22.224 e DZ 22.044; 1 male, 10 January<br>1971, Ebert leg., DZ22.174; Minas Gerais: Caratinga, Estação Ecológica de Caratinga, 400 m a.s.l., 1 male, 29<br>January–3 February 2003, Mielke & Casagrande legs, DZ20.622 (DZUP).<br>OCOLÓMBIA, Valle do Cauca: Dagua, 1 male, January 1967, DZ16.570 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande legs, DZ20.402; 1 male, 27–27 July 2008, Mielke & Casagrande legs, DZ19.41; Porto Acre, Reserva Fumaitá, 2<br>males, 28-31. July .2008, Mielke & Casagrande legs, DZ19.994 e DZ19.626 (DZUP).  |
|   | Archaeoprepona amphimachus pseudomeander (Fruhstorfer, 1906)<br>Archaeoprepona chromus chromus (Guérin-Méneville, [1844])<br>Archaeoprepona demophon muson (Fruhstorfer, 1905)  | Casagrande legs. OM54.168 (OM).<br>BRASIL, Santa Catarina: Joinville, 1 male, 28 October 1968, Mielke leg., DZ22.144; São Bento do Sul, Rio<br>Vermelho, 1 male, DZ22.254; 2 males, 2 December 1969, Ebert leg., DZ22.224 e DZ 22.044; 1 male, 10 January<br>1971, Ebert leg., DZ22.174; Minas Gerais: Caratinga, Estação Ecológica de Caratinga, 400 m a.s.l., 1 male, 29<br>January-3 February 2003, Mielke & Casagrande legs, DZ20.622 (DZUP).<br>COLÔMBIA, Valle do Cauca: Dagua, 1 male, January 1967, DZ16.570 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 29-30 September 2006, Mielke & Casagrande legs,<br>DZ0.402; 1 male, 23–27 July 2008, Mielke & Casagrande legs, DZ19.9414; Porto Acre, Reserva Humaitá, 2<br>males, 28–31. July -2008, Mielke & Casagrande legs, DZ19.9414; Porto Acre, Reserva Humaitá, 2<br>Mielse, BC, 1. July -2008, Mielke & Casagrande legs, DZ19.9414; Porto Acre, Reserva Humaitá, 2<br>Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg., DZ20.517;<br>DEVENDENDENDENDENDENDENDENDENDENDENDENDENDE   |
|   | Archaeoprepona amphimachus pseudomeander (Fruhstorfer, 1906)<br>Archaeoprepona chromus chromus (Guérin-Méneville, [1844])<br>Archaeoprepona demophon muson (Fruhstorfer, 1905)<br>Archaeoprepona demophoon andicola (Fruhstorfer, 1904)   | Casagrande legs. OM54.168 (OM).<br>BRASIL, Santa Catarina: Joinville, 1 male, 28 October 1968, Mielke leg., DZ22.144; São Bento do Sul, Rio<br>Vermelho, 1 male, DZ22.254; 2 males, 2 December 1969, Ebert leg., DZ22.24 e DZ 22.044; 1 male, 10 January<br>1971, Ebert leg., DZ22.174; Minas Gerais: Caratinga, Estação Ecológica de Caratinga, 400 m a.s.l., 1 male, 29<br>January–3 February 2003, Mielke & Casagrande legs, DZ20.622 (DZUP).<br>COLÔMBIA, Valle do Cauca: Dagua, 1 male, January 1967, DZ16.570 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg., DZ19633;<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba 1 male, 29–30 September 2006, Mielke & Casagrande legs,<br>DZ20.402; 1 male, 23–27 July 2008, Mielke & Casagrande legs, DZ19.414; Porto Acre, Reserva Humaitá, 2<br>males, 28-31. July .2008, Mielke & Casagrande legs, DZ19.944 (; Porto Acre, Reserva Humaitá, 2<br>males, 28-31. July .2008, Mielke & Casagrande legs, DZ19.944 (; Porto Acre, Reserva Humaitá, 2<br>males, 28-31. July .2008, Mielke & Casagrande legs, DZ19.944 (; Porto Acre, Reserva Humaitá, 2<br>males, 28-31. July .2008, Mielke & Casagrande legs, DZ19.944 (; Porto Acre, Reserva Humaitá, 2<br>males, 28-31. July .2008, Mielke & Casagrande legs, DZ19.944 (; Porto Acre, Reserva Humaitá, 2<br>males, 28-31. July .2008, Mielke & Casagrande legs, DZ19.944 (; Porto Acre, Reserva Humaitá, 2<br>Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg., DZ20.517;<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba, 1 male, 31 January–5 February 2009, Mielke & Casagrande<br>lew. DZ19.9271 (210)  |
|   | Archaeoprepona amphimachus pseudomeander (Fruhstorfer, 1906)<br>Archaeoprepona chromus chromus (Guérin-Méneville, [1844])<br>Archaeoprepona demophon muson (Fruhstorfer, 1905)<br>Archaeoprepona demophoon andicola (Fruhstorfer, 1904)   | Casagrande legs. OM54.168 (OM).<br>BRASIL, Santa Catarina: Joinville, 1 male, 28 October 1968, Mielke leg., DZ22.144; São Bento do Sul, Rio<br>Vermelho, 1 male, DZ22.254; 2 males, 2 December 1969, Ebert leg., DZ22.24 e DZ 22.044; 1 male, 10 January<br>1971, Ebert leg., DZ22.174; Minas Gerais: Caratinga, Estação Ecológica de Caratinga, 400 m a.s.l., 1 male, 29<br>January–3 February 2003, Mielke & Casagrande legs, DZ20.622 (DZUP).<br>COLÓMBIA, Valle do Cauca: Dagua, 1 male, January 1967, DZ16.570 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg., DZ19633;<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba 1 male, 29–30 September 2006, Mielke & Casagrande legs,<br>DZ20.402; 1 male, 23–27 July 2008, Mielke & Casagrande legs, DZ19.414; Porto Acre, Reserva Humaitá, 2<br>males, 28-31. July .2008, Mielke & Casagrande legs, DZ19.994 e DZ19.626 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg, DZ20.517;<br>BRASIL, Acre: Senador Guiomard, Reserve Catuaba, 1 male, 31 January–5 February 2009, Mielke & Casagrande<br>legs, DZ19.972 (DZUP).<br>PERU, Madre de Dios: Parque Manu, Pakitza, 340 m a.s.l., 1 male, 50 totober 1991, Lamas leg., DZ20.517;<br>PERU, Madre de Dios: Parque Manu, Pakitza, 340 m a.s.l., 1 male, 50 totober 1991, Lamas leg., DZ22.114.  |
| Preponini   | Archaeoprepona amphimachus pseudomeander (Fruhstorfer, 1906)<br>Archaeoprepona chromus chromus (Guérin-Méneville, [1844])<br>Archaeoprepona demophon muson (Fruhstorfer, 1905)<br>Archaeoprepona demophoon andicola (Fruhstorfer, 1904)<br>Archaeoprepona licomedes licomedes (Cramer, 1777)  | Casagrande legs. OM54.168 (OM).<br>BRASIL, Santa Catarina: Joinville, 1 male, 28 October 1968, Mielke leg., DZ22.144; São Bento do Sul, Rio<br>Vermelho, 1 male, DZ22.254; 2 males, 2 December 1969, Ebert leg., DZ22.224 e DZ 22.044; 1 male, 10 January<br>1971, Ebert leg., DZ22.174; Minas Gerais: Caratinga, Estação Ecológica de Caratinga, 400 m a.s.l., 1 male, 29<br>January–3 February 2003, Mielke & Casagrande legs, DZ20.622 (DZUP).<br>COLÓMBIA, Valle do Cauca: Dagua, 1 male, January 1967, DZ16.570 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg, DZ19633;<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba 1 male, 29–D3 September 2006, Mielke & Casagrande legs,<br>DZ20.402; 1 male, 23–27 July 2008, Mielke & Casagrande legs, DZ19.414; Porto Acre, Reserva Humairá, 2<br>males, 28-31. July .2008, Mielke & Casagrande legs, DZ19.994 e DZ19.626 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg., DZ20.517;<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba, 1 male, 31 January–5 February 2009, Mielke & Casagrande<br>legs, DZ19.972 (DZUP).<br>PERU, Madre de Dios: Parque Manu, Pakitza, 340 m a.s.l., 1 male, 15 October 1991, Lamas leg., DZ22.114,<br>BRASIL, Acre: Senador Guionard, Reserva Catuaba, 200 m a.s.l., 1 male, 25 September 2004, Mielke & Casagrande<br>legs, DZ19.972 (DZUP).   |
| Preponini   | Archaeoprepona amphimachus pseudomeander (Fruhstorfer, 1906)<br>Archaeoprepona chromus chromus (Guérin-Méneville, [1844])<br>Archaeoprepona demophon muson (Fruhstorfer, 1905)<br>Archaeoprepona demophoon andicola (Fruhstorfer, 1904)<br>Archaeoprepona licomedes licomedes (Cramer, 1777)  | Casagrande legs. OM54.168 (OM).<br>BRASIL, Santa Catarina: Joinville, 1 male, 28 October 1968, Mielke leg., DZ22.144; São Bento do Sul, Rio<br>Vermelho, 1 male, DZ22.254; 2 males, 2 December 1969, Ebert leg., DZ22.224 e DZ 22.044; 1 male, 10 January<br>1971, Ebert leg., DZ22.174; Minas Gerais: Caratinga, Estação Ecológica de Caratinga, 400 m a.s.l., 1 male, 29<br>January–3 February 2003, Mielke & Casagrande legs, DZ20.622 (DZUP).<br>COLÓMBIA, Valle do Cauca: Dagua, 1 male, January 1967, DZ16.570 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg., DZ19633;<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba 1 male, 29–30 September 2006, Mielke & Casagrande legs, DZ20.402; 1 male, 23–27 July 2008, Mielke & Casagrande legs, DZ19.41; Porto Acre, Reserva Humaitá, 2<br>males, 28-31. July .2008, Mielke & Casagrande legs, DZ19.994 e DZ19.626 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg., DZ20.517;<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba, 1 male, 31 January–5 February 2009, Mielke & Casagrand<br>legs, DZ19.972 (DZUP).<br>PERU, Madre de Dios: Parque Manu, Pakitza, 340 m a.s.l., 1 male, 15 October 1991, Lamas leg., DZ22.114,<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba, 200 m a.s.l., 1 male, 15 October 1991, Lamas leg., DZ22.114,<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba, 200 m A.s.l., 1 male, 25 September 2004, Mielke & Miers<br>legs, DZ19.972 (DZUP).   |
| Preponini   | Archaeoprepona amphimachus pseudomeander (Fruhstorfer, 1906)<br>Archaeoprepona chromus chromus (Guérin-Méneville, [1844])<br>Archaeoprepona demophon muson (Fruhstorfer, 1905)<br>Archaeoprepona demophoon andicola (Fruhstorfer, 1904)<br>Archaeoprepona licomedes licomedes (Cramer, 1777)<br>Archaeoprepona mandar metoring (F. New 1922)  | Casagrande legs. OM54.168 (OM).<br>BRASIL, Santa Catarina: Joinville, 1 male, 28 October 1968, Mielke leg., DZ22.144; São Bento do Sul, Rio<br>Vermelho, 1 male, DZ22.254; 2 males, 2 December 1969, Ebert leg., DZ22.224 e DZ 22.044; 1 male, 10 January<br>1971, Ebert leg., DZ22.174; Minas Gerais: Caratinga, Estação Ecológica de Caratinga, 400 m a.s.l., 1 male, 29<br>January-3 February 2003, Mielke & Casagrande legs, DZ20.627 (DZUP).<br>COLÓMBIA, Valle do Cauca: Dagua, 1 male, January 1967, DZ16.570 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg., DZ19633;<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba 1 male, 29-30 September 2006, Mielke & Casagrande leg.,<br>DZ20.402; 1 male, 23-27 July 2008, Mielke & Casagrande legs, DZ19.41; Porto Acre, Reserva Humaitá, 2<br>males, 28-31. July .2008, Mielke & Casagrande legs, DZ19.41; Porto Acre, Reserva Humaitá, 2<br>males, 28-31. July .2008, Mielke & Casagrande legs, DZ19.994 e DZ19.626 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg., DZ20.517;<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba 1 male, 31 January-5 February 2009, Mielke & Casagrand<br>legs, DZ19.972 (DZUP).<br>PERU, Madre de Dios: Parque Manu, Pakitza, 340 m a.s.l., 1 male, 15 October 1991, Lamas leg., DZ22.114,<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba, 200 m a.s.l., 1 male, 2-5 September 2004, Mielke & Miers<br>legs, DZ16513; 1 male, 31 January-5 February 2009, Mielke & Casagrande legs, DZ2.0389 (DZUP).<br>BRASIL, Acre: Senador Guionard, Reserva Catuaba, 200 m a.s.l., 1 male, 2-5 Cotober 1991, Casagrande leg, DZ2.114,<br>BRASIL, Acre: Senador Guionard, Reserva Catuaba, 200 m a.s.l., 1 male, 2-5 September 2004, Mielke & Miers<br>legs, DZ16513; 1 male, 31 January-5 February 2009, Mielke & Casagrande legs, DZ2.0389 (DZUP).<br>BRASIL, Kio de Janeiro: Rio de Janeiro; Rivae, 1 male, September 1995, DZ16.520; Magé, Serra dos Ogãos, 1<br>male 17 Eberuary 104 fa Ebert lego, DZ22 214. Minas Genzie Devane Ended de Dio Devane 2004, O                            |
| Preponini   | Archaeoprepona amphimachus pseudomeander (Fruhstorfer, 1906)<br>Archaeoprepona chromus chromus (Guérin-Méneville, [1844])<br>Archaeoprepona demophon muson (Fruhstorfer, 1905)<br>Archaeoprepona demophoon andicola (Fruhstorfer, 1904)<br>Archaeoprepona licomedes licomedes (Cramer, 1777)<br>Archaeoprepona meander castorina (E. Mayr, 1932)  | Casagrande legs. OM54.168 (OM).<br>BRASIL, Santa Catarina: Joinville, 1 male, 28 October 1968, Mielke leg., DZ22.144; São Bento do Sul, Rio<br>Vermelho, 1 male, DZ22.254; 2 males, 2 December 1969, Ebert leg., DZ22.24 e DZ 22.044; 1 male, 10 January<br>1971, Ebert leg., DZ22.174; Minas Gerais: Caratinga, Estação Ecológica de Caratinga, 400 m a.s.l., 1 male, 29<br>January–3 February 2003, Mielke & Casagrande legs, DZ20.622 (DZUP).<br>COLÔMBIA, Valle do Cauca: Dagua, 1 male, January 1967, DZ16.570 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg., DZ19633;<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba 1 male, 29–30 September 2006, Mielke & Casagrande legs,<br>DZ20.402; 1 male, 23–27 July 2008, Mielke & Casagrande legs, DZ19.414; Porto Acre, Reserva Humaitá, 2<br>males, 28-31. July .2008, Mielke & Casagrande legs, DZ19.914; Porto Acre, Reserva Humaitá, 2<br>males, 28-31. July .2008, Mielke & Casagrande legs, DZ19.994 e DZ19.626 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg., DZ20.517;<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba, 1 male, 31 January–5 February 2009, Mielke & Casagrande<br>legs, DZ19.972 (DZUP).<br>PERU, Madre de Dios: Parque Manu, Pakitza, 340 m a.s.l., 1 male, 15 October 1991, Lamas leg., DZ22.114,<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba, 2000 m a.s.l., 1 male, 2-5 September 2004, Mielke & Miers<br>legs, DZ16513; 1 male, 31 January–5 February 2009, Mielke & Casagrande legs, DZ20.389 (DZUP).<br>BRASIL, Rio de Janeiro: Rio de Janeiro, Gávea, 1 male, September 1955, DZ16.520; Mag&, Serra dos Ógãos, 1<br>male, 17 February 1964, Ebert leg., DZ22.184 (DZ1P).  |
| Preponini   | Archaeoprepona amphimachus pseudomeander (Fruhstorfer, 1906)<br>Archaeoprepona chromus chromus (Guérin-Méneville, [1844])<br>Archaeoprepona demophon muson (Fruhstorfer, 1905)<br>Archaeoprepona demophoon andicola (Fruhstorfer, 1904)<br>Archaeoprepona licomedes licomedes (Cramer, 1777)<br>Archaeoprepona meander castorina (E. Mayr, 1932)  | Casagrande legs. OM54.168 (OM).<br>BRASIL, Santa Catarina: Joinville, 1 male, 28 October 1968, Mielke leg., DZ22.144; São Bento do Sul, Rio<br>Vermelho, 1 male, DZ22.254; 2 males, 2 December 1969, Ebert leg., DZ22.24 e DZ 22.044; 1 male, 10 January<br>1971, Ebert leg., DZ22.174; Minas Gerais: Caratinga, Estação Ecológica de Caratinga, 400 m a.s.l., 1 male, 29<br>January–3 February 2003, Mielke & Casagrande legs, DZ20.622 (DZUP).<br>OCILÓMBIA, Valle do Cauca: Dagua, 1 male, January 1967, DZ16.570 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg., DZ19633;<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba 1 male, 29–30 September 2006, Mielke & Casagrande legs,<br>DZ20.402; 1 male, 23–27 July 2008, Mielke & Casagrande legs, DZ19.414; Porto Acre, Reserva Humaitá, 2<br>males, 28–31. July .2008, Mielke & Casagrande legs, DZ19.944 c DZ19.626 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg, DZ20.517;<br>BRASIL, Acre: Senador Guiomard, Reserve Catuaba, 1 male, 31 January–5 February 2009, Mielke & Casagrande<br>legs, DZ19.972 (DZUP).<br>PERU, Madre de Dios: Parque Manu, Pakitza, 340 m a.s.l., 1 male, 15 October 1991, Lamas leg., DZ22.114,<br>BRASIL, Acre: Senador Guionard, Reserva Catuaba, 200 m a.s.l., 1 male, 55 October 1991, Lamas leg., DZ22.114,<br>BRASIL, Acre: Senador Guionard, Reserva Catuaba, 200 m a.s.l., 1 male, 25 September 2004, Mielke & Miers<br>legs, DZ16513; 1 male, 31 January–5 February 2009, Mielke & Miers<br>legs, DZ16513; I male, 31 January–5 Tebruary 2009, Mielke & Casagrande legs, DZ20.389 (DZUP).<br>BRASIL, Ki de Janeiro, Gávea, 1 male, Spetherber 1958, DZ16.520; Magé, Serra dos Ógãos, 1<br>male, 17 February 1964, Ebert leg., DZ22.214; Minas Gerais: Parque Estadual do Rio Doce, 200 m a.s.l., 1 male,<br>8 September 1972, Ebert leg., DZ21.884 (DZUP).<br>BRASIL, Santa Catarina: Joinville, 100 m a.s.l., 1 male, 25 February 1958, DZ2.334; 1 male, March 1966. Mier  |
| Preponini   | Archaeoprepona amphimachus pseudomeander (Fruhstorfer, 1906)<br>Archaeoprepona chromus chromus (Guérin-Méneville, [1844])<br>Archaeoprepona demophon muson (Fruhstorfer, 1905)<br>Archaeoprepona demophoon andicola (Fruhstorfer, 1904)<br>Archaeoprepona licomedes licomedes (Cramer, 1777)<br>Archaeoprepona meander castorina (E. Mayr, 1932)<br>Perpona claudina annetta (Grav. 1832)   | Casagrande legs. OM54.168 (OM).<br>BRASIL, Santa Catarina: Joinville, 1 male, 28 October 1968, Mielke leg., DZ22.144; São Bento do Sul, Rio<br>Vermelho, 1 male, DZ22.254; 2 males, 2 December 1969, Ebert leg., DZ22.24 e DZ 22.044; 1 male, 10 January<br>1971, Ebert leg., DZ22.174; Minas Gerais: Caratinga, Estação Ecológica de Caratinga, 400 m a.s.l., 1 male, 29<br>January–3 February 2003, Mielke & Casagrande legs, DZ20.622 (DZUP).<br>COLÓMBIA, Valle do Cauca: Dagua, 1 male, January 1967, DZ16.570 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg, DZ19633;<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba 1 male, 29–30 September 2006, Mielke & Casagrande legs,<br>DZ20.402; 1 male, 23–27 July 2008, Mielke & Casagrande legs, DZ19.414; Porto Acre, Reserva Humairá, 2<br>males, 28-31. July .2008, Mielke & Casagrande legs, DZ19.994 e DZ19.626 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg., DZ20.517;<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba, 1 male, 31 January–5 February 2009, Mielke & Casagrande<br>legs, DZ19.972 (DZUP).<br>PERU, Madre de Dios: Parque Manu, Pakitza, 340 m a.s.l., 1 male, 15 October 1991, Lamas leg., DZ22.114,<br>BRASIL, Acre: Senador Guionard, Reserva Catuaba, 200 m a.s.l., 1 male, 25 September 2004, Mielke & Miers<br>legs, DZ16.513; 1 male, 31 January–5 February 2009, Mielke & Casagrande legs, DZ22.114,<br>BRASIL, Acre: Senador Guionard, Reserva Catuaba, 200 m a.s.l., 1 male, 25 September 2004, Mielke & Miers<br>legs, DZ16.513; 1 male, 31 January–5 February 2009, Mielke & Casagrande legs, DZ22.114,<br>BRASIL, Acre: Senador Guionard, Reserva Catuaba, 200 m a.s.l., 1 male, 50 Cotober 1991, Lamas leg., DZ22.114,<br>BRASIL, Acre: Senador Guionard, Reserva Catuaba, 200 m a.s.l., 1 male, 50 September 2004, Mielke & Miers<br>legs, DZ16.05, 13; 1 male, 31 January–5 February 2009, Mielke & Casagrande legs, DZ22.144,<br>BRASIL, Santa Catarina: Joinville, 100 m a.s.l., 1 male, 25 February 1958, DZ26.50; Magé, Serra dos Ógão, 1<br>male, 17 February 1964, E    |
| Preponini   | Archaeoprepona amphimachus pseudomeander (Fruhstorfer, 1906)<br>Archaeoprepona chromus chromus (Guérin-Méneville, [1844])<br>Archaeoprepona demophon muson (Fruhstorfer, 1905)<br>Archaeoprepona demophoon andicola (Fruhstorfer, 1904)<br>Archaeoprepona licomedes licomedes (Cramer, 1777)<br>Archaeoprepona meander castorina (E. Mayr, 1932)<br>Prepona claudina annetta (Gray, 1832)   | Casagrande legs. OM54.168 (OM).<br>BRASIL, Santa Catarina: Joinville, 1 male, 28 October 1968, Mielke leg., DZ22.144; São Bento do Sul, Rio<br>Vermelho, 1 male, DZ22.254; 2 males, 2 December 1969, Ebert leg., DZ22.224 e DZ 22.044; 1 male, 10 January<br>1971, Ebert leg., DZ22.174; Minas Gerais: Caratinga, Estação Ecológica de Caratinga, 400 m a.s.l., 1 male, 29<br>January–3 February 2003, Mielke & Casagrande legs, DZ20.622 (DZUP).<br>COLÓMBIA, Valle do Cauca: Dagua, 1 male, January 1967, DZ16.570 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 29 October 1991, Casagrande leg, DZ19633;<br>BRASIL, Acre: Senador Guiomard, Reserve, 300 m a.s.l., 1 male, 29 October 1991, Casagrande leg, DZ19633;<br>BRASIL, Acre: Senador Guiomard, Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg, DZ20.517;<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg, DZ20.517;<br>BRASIL, Acre: Senador Guiomard, Reserva (Satuaha, 1 male, 31 January–5 February 2009, Mielke & Casagrande legs, DZ19.994 e DZ19.626 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserva (Satuaha, 1 male, 31 January–5 February 2009, Mielke & Casagrande legs, DZ19.972 (DZUP).<br>PERU, Madre de Dios: Parque Manu, Pakitza, 340 m a.s.l., 1 male, 2-5 September 2004, Mielke & Miers<br>legs, DZ19.972 (DZUP).<br>PERU, Madre de Dios: Parque Manu, Pakitza, 340 m a.s.l., 1 male, 2-5 September 2004, Mielke & Miers<br>legs, DZ16513; 1 male, 31 January–5 February 2009, Mielke & Casagrande legs, DZ20.389 (DZUP).<br>BRASIL, Acre: Senador Guionard, Reserva Catuaha, 200 m a.s.l., 1 male, 2-5 September 2004, Mielke & Miers<br>legs, DZ16513; 1 male, 31 January–5 February 2009, Mielke & Casagrande legs, DZ20.389 (DZUP).<br>BRASIL, Acre: Senador Guionard, Reserva Catuaha, 200 m a.s.l., 1 male, 2-5 September 2004, Mielke & Miers<br>legs, DZ16513; 1 male, 31 January–5 February 2009, Mielke & Casagrande legs, DZ20.389 (DZUP).<br>BRASIL, Kio de Janeiro: Rio de Janeiro; Gávea, 1 male, September 1958, DZ16.520; Magé, Serra dos Ogãos, 1<br>male, 17 February 1964, Ebert leg., DZ22.       |
| Preponini   | Archaeoprepona amphimachus pseudomeander (Fruhstorfer, 1906)<br>Archaeoprepona chromus chromus (Guérin-Méneville, [1844])<br>Archaeoprepona demophon muson (Fruhstorfer, 1905)<br>Archaeoprepona demophoon andicola (Fruhstorfer, 1904)<br>Archaeoprepona licomedes licomedes (Cramer, 1777)<br>Archaeoprepona meander castorina (E. Mayr, 1932)<br>Prepona claudina annetta (Gray, 1832)   | Casagrande legs. OM54.168 (OM).<br>BRASIL, Santa Catarina: Joinville, 1 male, 28 October 1968, Mielke leg., DZ22.144; São Bento do Sul, Rio<br>Vermelho, 1 male, DZ22.254; 2 males, 2 December 1969, Ebert leg., DZ22.24 e DZ 22.044; 1 male, 10 January<br>1971, Ebert leg., DZ22.174; Minas Gerais: Caratinga, Estação Ecológica de Caratinga, 400 m a.s.l., 1 male, 29<br>January–3 February 2003, Mielke & Casagrande legs, DZ20.622 (DZUP).<br>COLÔMBIA, Valle do Cauca: Dagua, 1 male, January 1967, DZ16.570 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg., DZ19633;<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba 1 male, 29–30 September 2006, Mielke & Casagrande legs,<br>DZ20.402; 1 male, 23–27 July 2008, Mielke & Casagrande legs, DZ19.414; Porto Acre, Reserva Humaitá, 2<br>males, 28-31. July .2008, Mielke & Casagrande legs, DZ19.914; Porto Acre, Reserva Humaitá, 2<br>males, 28-31. July .2008, Mielke & Casagrande legs, DZ19.914; Porto Acre, Reserva Humaitá, 2<br>males, 28-31. July .2008, Mielke & Casagrande legs, DZ19.914; Casagrande leg, DZ20.517;<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba, 1 male, 31 January–5 February 2009, Mielke & Casagrande<br>legs, DZ19.972 (DZUP).<br>PERU, Madre de Dios: Parque Manu, Pakitza, 340 m a.s.l., 1 male, 15 October 1991, Lamas leg., DZ22.114,<br>BRASIL, Acre: Senador Guionard, Reserva Catuaba, 200 m a.s.l., 1 male, 2-5 September 2004, Mielke & Miers<br>legs, DZ16513; 1 male, 31 January–5 February 2009, Mielke & Casagrande legs, DZ20.389 (DZUP).<br>BRASIL, Rio de Janeiro: Rio de Janeiro, Gávea, 1 male, September 1958, DZ16.520; Magé, Serra dos Ógãos, 1<br>male, 17 February 1964, Ebert leg., DZ21.884 (DZUP).<br>BRASIL, Santa Catarina: Joinville, 100 m a.s.l., 1 male, 25 February 1958, DZ22.334; 1 male, March 1966, Mier<br>leg., DZ10.063; 1 male, 27 March 1970, Mielke leg., DZ22.034; 2 males, 20 November 1970, Mielke leg.,<br>DZ22.194 e DZ2.194; 1 male, 18 January 1971, Ebert leg., DZ16.526; 1 male, 7 March 1986, Miers leg.,<br>DZ20.023 (DZUP).  |
| Preponini   | Archaeoprepona amphimachus pseudomeander (Fruhstorfer, 1906)<br>Archaeoprepona chromus chromus (Guérin-Méneville, [1844])<br>Archaeoprepona demophon muson (Fruhstorfer, 1905)<br>Archaeoprepona demophoon andicola (Fruhstorfer, 1904)<br>Archaeoprepona licomedes licomedes (Cramer, 1777)<br>Archaeoprepona meander castorina (E. Mayr, 1932)<br>Prepona claudina annetta (Gray, 1832)<br>Prepona laertes laertes (Hübner, [1811])   | Casagrande legs. OM54.168 (OM).<br>BRASIL, Santa Catarina: Joinville, 1 male, 28 October 1968, Mielke leg., DZ22.144; São Bento do Sul, Rio<br>Vermelho, 1 male, DZ22.254; 2 males, 2 December 1969, Ebert leg., DZ22.24 e DZ 22.044; 1 male, 10 January<br>1971, Ebert leg., DZ22.174; Minas Gerais: Caratinga, Estação Ecológica de Caratinga, 400 m a.s.l., 1 male, 29<br>January–3 February 2003, Mielke & Casagrande legs, DZ20.622 (DZUP).<br>COLOMBIA, Valle do Cauca: Dagua, 1 male, January 1967, DZ16.570 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg., DZ19633;<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba 1 male, 29–30 September 2006, Mielke & Casagrande legs,<br>DZ20.402; 1 male, 23–27 July 2008, Mielke & Casagrande legs, DZ19.414; Porto Acre, Reserva Humaitá, 2<br>males, 28-31. July -2008, Mielke & Casagrande legs, DZ19.944; Porto Acre, Reserva Humaitá, 2<br>males, 28-31. July -2008, Mielke & Casagrande legs, DZ19.940 e DZ19.626 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg, DZ20.517;<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba, 1 male, 31 January–5 February 2009, Mielke & Casagrand<br>legs, DZ19.972 (DZUP).<br>PERU, Madre de Dios: Parque Manu, Pakitza, 340 m a.s.l., 1 male, 15 October 1991, Lamas leg., DZ22.114,<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba, 200 m a.s.l., 1 male, 2-5 September 2004, Mielke & Miers<br>legs, DZ16513; 1 male, 31 January–5 February 2009, Mielke & Casagrande legs, DZ20.389 (DZUP).<br>BRASIL, Rio de Janeiro: Rio de Janeiro, Gávea, 1 male, September 1958, DZ16.520; Magé, Serra dos Ógãos, 1<br>male, 17 February 1964, Ebert leg., DZ22.144, Minas Gerais: Parque Estadual do Rio Doce, 200 m a.s.l., 1 male,<br>8 September 1972, Ebert leg., DZ21.884 (DZUP).<br>BRASIL, Santa Catarina: Joinville, 100 m a.s.l., 1 male, 25 February 1958, DZ20.323; 1 male, March 1966, Mier<br>leg., DZ10.063; 1 male, 17 March 1970, Mielke leg., DZ21.884 (DZUP).<br>BRASIL, Santa Catarina: Joinville, 100 m a.s.l., 1 male, 25 February 1958, DZ20.234; 1 mal    |
| Preponini   | Archaeoprepona amphimachus pseudomeander (Fruhstorfer, 1906)<br>Archaeoprepona chromus chromus (Guérin-Méneville, [1844])<br>Archaeoprepona demophon muson (Fruhstorfer, 1905)<br>Archaeoprepona demophoon andicola (Fruhstorfer, 1904)<br>Archaeoprepona licomedes licomedes (Cramer, 1777)<br>Archaeoprepona meander castorina (E. Mayr, 1932)<br>Prepona claudina annetta (Gray, 1832)<br>Prepona laertes laertes (Hübner, [1811])   | Casagrande legs. OM54.168 (OM).<br>BRASIL, Santa Catarina: Joinville, 1 male, 28 October 1968, Mielke leg., DZ22.144; São Bento do Sul, Rio<br>Vermelho, 1 male, DZ22.254; 2 males, 2 December 1969, Ebert leg., DZ22.24 e DZ 22.044; 1 male, 10 January<br>1971, Ebert leg., DZ22.174; Minas Gerais: Caratinga, Estação Ecológica de Caratinga, 400 m a.s.l., 1 male, 29<br>January–3 February 2003, Mielke & Casagrande legs, DZ20.622 (DZUP).<br>OCILÓMBIA, Valle do Cauca: Dagua, 1 male, January 1967, DZ16.570 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg, DZ19633;<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba 1 male, 29–30 September 2006, Mielke & Casagrande legs,<br>DZ20.402; 1 male, 23–27 July 2008, Mielke & Casagrande legs, DZ19.414; Porto Acre, Reserva Humaitá, 2<br>males, 28-31. July -2008, Mielke & Casagrande legs, DZ19.994 e DZ19.626 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande legs, DZ20.517;<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba, 1 male, 31 January–5 February 2009, Mielke & Casagrande<br>legs, DZ19.972 (DZUP).<br>PERU, Madre de Dios: Parque Manu, Pakitza, 340 m a.s.l., 1 male, 15 October 1991, Lamas leg., DZ22.114,<br>BRASIL, Acre: Senador Guionard, Reserva Catuaba, 200 m a.s.l., 1 male, 25 September 2004, Mielke & Miers<br>legs, DZ16513; 1 male, 31 January–5 February 2009, Mielke & Miers<br>legs, DZ16513; 1 male, 31 January–5 February 2009, Mielke & Casagrande legs, DZ22.114,<br>BRASIL, Ki de Janeiro, Gávea, 1 male, September 1958, DZ16.520; Magé, Serra dos Ógãos, 1<br>male, 17 February 1964, Ebert leg., DZ22.214; Minas Gerais: Parque Estadual do Rio Doce, 200 m a.s.l., 1 male,<br>8 September 1972, Ebert leg., DZ21.844 (DZUP).<br>BRASIL, Santa Catarina: Joinville, 100 m a.s.l., 1 male, 25 February 1958, DZ22.234; 1 male, March 1966, Mier<br>leg., DZ10.063; 1 male, 27 March 1970, Mielke leg., DZ21.846 (DZUP).<br>BRASIL, Santa Catarina: Joinville, 000 m a.s.l., 1 male, 25 February 1958, DZ22.234; 1 male, March 1966, Mier<br>leg., DZ10.063; 1 male, 27 Mar |
| Preponini   | Archaeoprepona amphimachus pseudomeander (Fruhstorfer, 1906)   Archaeoprepona chromus chromus (Guérin-Méneville, [1844])   Archaeoprepona demophon muson (Fruhstorfer, 1905)   Archaeoprepona demophon andicola (Fruhstorfer, 1904)   Archaeoprepona demophoon andicola (Fruhstorfer, 1904)   Archaeoprepona licomedes licomedes (Cramer, 1777)   Archaeoprepona meander castorina (E. Mayr, 1932)   Prepona claudina annetta (Gray, 1832)   Prepona laertes laertes (Hübner, [1811])   Prepona pylene Hewitson, [1854] | Casagrande legs. OM54.168 (OM).<br>BRASIL, Santa Catarina: Joinville, 1 male, 28 October 1968, Mielke leg., DZ22.144; São Bento do Sul, Rio<br>Vermelho, 1 male, DZ22.254; 2 males, 2 December 1969, Ebert leg., DZ22.24 e DZ 22.044; 1 male, 10 January<br>1971, Ebert leg., DZ22.174; Minas Gerais: Caratinga, Estação Ecológica de Caratinga, 400 m a.s.l., 1 male, 29<br>January–3 February 2003, Mielke & Casagrande legs, DZ20.622 (DZUP).<br>COLÓMBIA, Valle do Cauca: Dagua, 1 male, January 1967, DZ16.570 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg, DZ19633;<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba 1 male, 29–30 September 2006, Mielke & Casagrande legs,<br>DZ20.402; 1 male, 23–27 July 2008, Mielke & Casagrande legs, DZ19.414; Porto Acre, Reserva Humaitá, 2<br>males, 28-31, July .2008, Mielke & Casagrande legs, DZ19.994 e DZ19.626 (DZUP).<br>PERU, Madre de Dios: Tambopata Reserve, 300 m a.s.l., 1 male, 25 October 1991, Casagrande leg, DZ20.517;<br>BRASIL, Acre: Senador Guiomard, Reserva Catuaba, 1 male, 31 January–5 February 2009, Mielke & Casagrande<br>legs, DZ19.972 (DZUP).<br>PERU, Madre de Dios: Parque Manu, Pakitza, 340 m a.s.l., 1 male, 15 October 1991, Lamas leg., DZ22.114,<br>BRASIL, Acre: Senador Guionard, Reserva Catuaba, 200 m a.s.l., 1 male, 25 September 2004, Mielke & Miers<br>legs, DZ16.513; 1 male, 31 January–5 February 2009, Mielke & Casagrande legs, DZ22.114,<br>BRASIL, Acre: Senador Guionard, Reserva Catuaba, 200 m a.s.l., 1 male, 55 September 2004, Mielke & Miers<br>legs, DZ16.513; 1 male, 31 January–5 February 2009, Mielke & Casagrande legs, DZ22.114,<br>BRASIL, Santa Catarina: Joinville, 100 m a.s.l., 1 male, 52 February 2029, Mielke & Miers<br>legs, DZ10.063; 1 male, 27 March 1970, Mielke leg., DZ21.884 (DZUP).<br>BRASIL, Santa Catarina: Joinville, 100 m a.s.l., 1 male, 25 February 1985, NZ26.50; Magé, Serra dos Ógãos, 1<br>male, 17 February 1964, Ebert leg., DZ22.214; Minas Gerais: Parque Estadual do Rio Doce, 200 m a.s.l., 1 male,<br>8 September 1972, Ebert leg., DZ21.844 (DZUP).<br>BRASIL, Santa  |









Journal of Insect Science | www.insectscience.org







Journal of Insect Science | www.insectscience.org





















Journal of Insect Science | www.insectscience.org





view. F. Fultura inferior. G and H Valva: G. Internal. H. External. I–L Penis: I. Right lateral view. J. Left lateral view. K. Dorsal view. L. Ventral view. Scale bar: A and B = 1 cm. C–L = 1 mm. High quality figures are available online.



Journal of Insect Science | www.insectscience.org







Bonfantti et al.









Dorsal view. K. Ventral view. Scale bar: A and B = 1cm. C-K = 1mm. High quality figures are available online.

Journal of Insect Science | www.insectscience.org













**Figure 30.** Prepona claudina annetta (Gray). A. Dorsal. B. Ventral. C–K Genitalia: C. Lateral view. D. Posterior view. E. Dorsal view. F and G Valva: F. Internal. G. External. H–K: Penis: H. Right lateral view. I. Left lateral view. J. Dorsal view. K. Ventral view. Scale bar: A and B = 1 cm. C–K = 1 mm. High quality figures are available online.





