



The Effect of *Beauveria brongniartii* and its Secondary Metabolites on the Detoxification Enzymes of the Pine Caterpillar, *Dendrolimus tabulaeformis*

Authors: Fan, Jinhua, Xie, Yingping, Xue, Jiaoliang, and Liu, Rui

Source: Journal of Insect Science, 13(44) : 1-13

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.013.4401>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



The effect of *Beauveria brongniartii* and its secondary metabolites on the detoxification enzymes of the pine caterpillar, *Dendrolimus tabulaeformis*

Jinhua Fan^a, Yingping Xie^{b*}, Jiaoliang Xue^c, and Rui Liu^d

The School of Life Science, Shanxi University, Taiyuan 030006, China

Abstract

The mortality of pine caterpillar, *Dendrolimus tabulaeformis* Tsai et Liu (Lepidoptera: Lasio-campidae), larvae treated with *Beauveria brongniartii* (Saccardo) Petch (Hypocreales: Clavicipitaceae) conidia and cell-free culture supernatants enriched for the secondary metabolites of the fungus was investigated. In addition, the effects of the treatments on the activities of two insect-related defense response proteins, glutathione *S*-transferase (GST) and esterase (EST), were measured over time. Bioassays were performed using a range of fungal spore (6×10^5 through 6×10^7 spores/mL) and supernatant extract concentrations (5.5–550 $\mu\text{g/mL}$). The results showed that the mortalities of *D. tabulaeformis* larvae were closely related to the concentration of the conidia and the metabolites of *B. brongniartii*. The differences among the treatments all reached a significant level. The activities of the two detoxifying enzymes, GST and EST, in the larvae increased simultaneously post-treatment. After infection with the conidial suspensions, the highest GST activity appeared at 3 days, and the activities of the caterpillars infected with 6×10^6 spores/mL and 6×10^7 spores/mL were significantly higher than in the control. Using α -naphthyl, the highest activity of EST also appeared at 3 days, and the differences for the three different concentrations were significant. A similar trend of change in the EST activity was observed using β -naphthyl. After treatment with the secondary metabolite solution, the highest GST activity appeared at 6 hr, and significant differences were found both for the different durations (2, 4, 6, 12, 24, and 48 hr) and in the three concentration groups. When using α -naphthyl, the EST activity peak appeared at 24 hr, and the differences were significant among the durations of 2, 4, 6, 12, 24, and 48 hr. The effect of the concentration of the secondary metabolite solution notably induced the EST activity in the insects, and a similar result was obtained using β -naphthyl. The data suggest that *B. brongniartii* produces secondary metabolites that disable the immune mechanisms of *D. tabulaeformis*, allowing the fungus to overcome and then kill its host. It was concluded that both the conidial suspensions and the metabolites of *B. brongniartii* were toxic to *D. tabulaeformis* larvae.

Keywords: entomopathogenic fungi, esterase, glutathione S-transferase, immune reaction, toxic
Abbreviations: EST, esterase; GST, glutathione S-transferase; LC₅₀, medium lethal concentration
Correspondence: ^a fanjinhua0329@163.com, ^b xieyingping@eyou.com, ^c xuejl@sxu.edu.cn, ^d liurui124@126.com,
 *Corresponding author
Editor: Nadir Erbilgin was editor of this paper.
Received: 8 March 2012 **Accepted:** 10 October 2012
Copyright : This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.
ISSN: 1536-2442 | Vol. 13, Number 44

Cite this paper as:

Fan J, Xie Y, Xue J, Liu R. 2013. The effect of *Beauveria brongniartii* and its secondary metabolites on the detoxification enzymes of the pine caterpillar, *Dendrolimus tabulaeformis*. *Journal of Insect Science* 13:44. Available online: <http://www.insectscience.org/13.44>

Introduction

The pine caterpillar, genus *Dendrolimus* (Lepidoptera: Lasiocampidae), is one group of the principal pests in coniferous forests worldwide. There are 27 species in China, and, according to the report from the National Forestry Department of China, areas of damage cover more than 2 million hm² each year (Hou 1987). Since the 1970s, entomopathogenic fungi *Beauveria bassiana* and *Beauveria brongniartii* (Saccardo) Petch (Hypocreales: Clavicipitaceae) have been incorporated into an integrated pest management program to control the pine caterpillar (Wang et al. 2010).

The infection mechanism of *Beauveria* in the host insects has been reported (Roberts et al. 2004; Wang et al. 2004; Thomas et al. 2007). The primary route of host invasion is through the external integument via the attachment of the conidia to the cuticle, germination, followed by penetration into the cuticle. Once in the hemocoel, the mycelium ramifies throughout the host, forming yeast-like hyphal bodies or blastospores. Host death is often due to a combination of the action of a fungal toxin, the physical obstruction of blood circulation, nutrient depletion, and the invasion of organs.

During the attack process, the host immune system attempts to resist the fungus, and some

detoxification enzymes inside the insect play a part in protecting the insects from the negative impact of the pathogens and their toxins. When the insects are attacked by these factors, the detoxification enzymes act by regulating the metabolism of hormones, pheromones, and other biologically active substances. General esterase (EST) and glutathione S-transferase (GST) are the most common enzymes involved in the detoxification of penetrated xenobiotics (Zibae et al. 2009b). In past studies, the insect resistance and activity of insect EST and GST induced by certain pesticides have been reported (Terriere 1984; Bakanova et al. 1996; Feyereisen 1999; Serebrov et al. 2006; Zibae et al. 2009a; Zibae et al., 2009b). However, reports are scarce on the changes in the activity of detoxification enzymes when the insects are infected by entomopathogenic fungi and/or the fungal secondary metabolites.

It is known that the entomopathogenic fungal attack of host insects involves both a direct infection by the fungus and the action of its secondary metabolites. Because some secondary metabolites are toxic to insects (Sokolova et al. 1999; Xia et al. 2000; Xia et al. 2001), both the entomopathogenic fungi and their secondary metabolites can induce immunoreactions and changes in the activity of detoxification enzymes in the host. Nevertheless, no data are available on the action and changes in the activity of the detoxifying enzymes of pine caterpillars when infected by

pathogenic fungi and/or exposed to the fungal secondary metabolites.

In a previous study, a new strain, *B. brongniartii*, was isolated from the naturally diseased corpses of *D. tabulaeformis* Tsai et Liu in a pine forest at Chengde, Hebei, China; the fungal secondary metabolites comprised 2-piperridinone, 2-coumaranone, pyrrolo, and certain other toxic components (Fan et al. 2008).

In the present study, the entomopathogenic fungus *B. brongniartii* and the fungal secondary metabolites were employed as pathogens for the pine caterpillar *D. tabulaeformis*, the most destructive pest in the pine forests of northern China. The objective was to determine the changes in the GST and EST activities when the *D. tabulaeformis* larvae were infected with conidial suspensions or were exposed to the secondary metabolites of *B. brongniartii*. This study provides evidence for an understanding of the infection mechanism, the sites of fungal attack, and the immune response of the host insect.

Material and Methods

Insect

The larvae of *D. tabulaeformis* were collected in a pine forest in Chengde (E 117° 51', N 40° 57'), Hebei Province, China. The larvae were reared with fresh pine needles of *Pinus tabulaeformis* Carr. in a rearing room at $27 \pm 1^\circ \text{C}$, with $75 \pm 10\%$ RH, and a 15:9 L:D photoperiod. After completing two generations, the healthy fourth-instar larvae were used for the experiments.

Entomopathogenic fungus

A *B. brongniartii* strain of entomopathogenic fungus was employed in the experiment. We isolated this strain in 2008 from the naturally

infected dead larvae of *D. tabulaeformis* collected in a pine forest in Chengde. Before the experiment, the strain was cultured on potato-dextrose-agar medium for 15 days at $25 \pm 1^\circ \text{C}$, with $75 \pm 10\%$ RH.

Fungal suspension preparation

After culture for 15 days, the fungal conidia were harvested from the surface of the culture medium using a sterile blade. After preparing the conidial suspensions, the conidial concentration was determined using a hemocytometer and was adjusted to 6×10^5 , 6×10^6 , and 6×10^7 spores/mL with 0.1% (v/v) Tween-80 (Kermel, www.chemreagent.com) sterile water solution.

Fungal metabolite preparation

Fungal liquid culture. The harvested conidia were prepared as suspension with a concentration of 1×10^8 spores/mL. One mL aliquot of the conidial suspension was used to inoculate 100 mL liquid medium, which was prepared with 10 g/L peptone, 10 g/L yeast extract, and 10 g/L glucose in a 250 mL conical flask. The fungus was cultured in an incubator (MAXQ 5000, Thermo Scientific, www.thermoscientific.com) at a constant temperature of $25 \pm 1^\circ \text{C}$ and at 265 rpm for 7 days.

Extraction of fungal metabolites. After culturing, crude extracts of the cultured broth were obtained following the method reported by Hu (2006). The fermentation broth was centrifuged (Centrifuge 58042, Eppendorf, www.eppendorf.com) at $10,000 \times g$ for 15 min and then concentrated to 1/5 at 50°C . The concentrated broth was then precipitated with alcohol (final concentration into 70% v/v) (Kermel) for 24 hr. After centrifugation (Centrifuge 58042) at $5,000 \times g$ for 20 min, the supernatant was sequentially extracted with ethyl acetate (Kermel) at a 1:2 ratio.

Lastly, an orange-red powder was obtained after drying at 40° C. This orange-red powder was considered the fungal secondary metabolites used in the experiment.

Bioassays

Using a micro injector (Angle, www.shweiliang.com), the larvae were injected with 5 µl of a range of fungal spore concentrations (6×10^5 – 10^7 spores/mL) and 1 µl of the supernatant extract concentrations (5.5–550 µg/mL). Sixty samples of the larvae were treated separately for each experimental group, and an additional 60 samples of the larvae were synchronously treated with 0.1% (v/v) Tween-80 and/or DMSO (Sigma Aldrich, www.sigmaaldrich.com) (0.5%) alone as the controls. All of the experiments were conducted twice, with five replicates each. The inoculated insects were transferred to rearing chambers (SPX-2051-C, Boxun, www.boxun.com.cn) and cultured for 6 days at $25 \pm 1^\circ$ C with $75 \pm 10\%$ RH and a photoperiod of 15:9 L:D.

Insect mortality

The mortality of the insects was recorded on the 7th day post-treatment. The medium lethal concentration (LC₅₀) was calculated using Probit Analysis of Regression (SPSS 13.0, www-01.ibm.com/software/analytics/spss).

Enzymatic sample preparation

During culturing, the insects in each test group that were inoculated with the conidial suspensions were sampled randomly and daily from the 1st day to the 6th day after the treatment. The metabolite treated insects were sampled randomly at 2, 4, 6, 12, 24, and 48 hr, respectively. The sample number of the insects was 5 insects from each test group at every time point. The insect samples were homogenized using an electric homogenizer (DY89-II, Scientz, www.scientz.com) at 4° C

and $690 \times g$ for 1.5 min. Next, they were cooled in 0.1 mol/L phosphate buffer (PBS, pH 7.5) with 0.3% Triton X-100 (Solarbio, www.solarbio.cn) (tissue: PBS = 1 g:10 mL). The tissue homogenate was then centrifuged (Centrifuge 58042) for 30 min at $15,000 \times g$ and at 4° C. The supernatant, as the enzyme source, was transferred to new tubes and stored (BCD 212DC, Haier, www.haier.com) at -20° C for 2 days only. The tests were conducted twice, with three replicates each, and the enzymatic activities associated with the observed insect resistance were determined.

GST activity determination

The method reported by Oppenorth (1985) was adopted for determining the changes in the GST activity in the test insects; 1-chloro-2, 4-dinitrobenzene was purchased from Sigma Aldrich Company. 195 µL 1-chloro-2, 4-dinitrobenzene (20 mM) were placed in microplate wells, and 5 µl of enzyme solution was added. The optical density value at 340 nm was recorded using a microplate reader (M5, Spectra Max, www.spectramax.com).

EST activity determination

The EST activity of the insects was determined following the method described by Han et al. (1998); α -naphthyl and β -naphthyl were purchased from Sigma Aldrich. A 135 µl aliquot of α -naphthyl acetate (or β -naphthyl) (0.3 mM) and 50 µl fast blue RR salt (Sigma Aldrich) (1 mM) were added to each well of a microplate (Costar 3590, Corning, www.corning.com). The reaction was initiated by the addition of 15 µl of enzyme solution, and the optical density value at 450 nm (M5, Spectra Max) was recorded.

Protein determination

The protein concentrations of the enzyme solution were measured according to the method of Bradford (1976) using bovine serum albu-




Figure 1. The changes in the activity of the two detoxification enzymes *Dendrolimus tabulaeformis* larvae infected with conidial suspension and treated with secondary metabolite of the fungus. (1A) GST activity changes of the larvae treated by the spore suspension; (1B) GST activity changes in larvae treated by the secondary metabolite; (1C) EST activity changes measured under the condition using α -naphthyl in the larvae treated by the spore suspension; (1D) EST activity by using α -naphthyl in larvae treated by secondary metabolite; (1E) EST activity changes measured in the condition using β -naphthyl in the larvae treated by the spore suspension; (1F) Activity of EST by using β -naphthyl in larvae treated by secondary metabolite. High quality figures are available online.

min (Bio-Rad, www.bio-rad.com) as the standard.

Statistical analysis

The data were compared using the univariate analysis of the General Linear Model when significant differences were found at $p = 0.05$ (SPSS 13.0). The differences between samplings were considered statistically significant at a probability of $> 5\%$.

Results

Mortality of larvae infected with the fungal conidia

After inoculation with the conidial suspension of the fungus *B. brongniartii*, the dead larvae were recorded. The mortality data are shown

in Table 1. The larval mortality rate was positively correlated with the conidial concentration in the suspensions. The highest mortality of larvae was $59.67 \pm 3.07\%$ and was found in the group treated with the highest conidial concentration (6.0×10^7 spores/mL). The larval mortalities were $37.00 \pm 2.40\%$ and $23.33 \pm 1.73\%$ in the groups treated with 6.0×10^6 spores/mL and 6.0×10^5 spores/mL, respectively. The differences in the larval mortality were significant among the groups. The LC_{50} was 2.7×10^7 spores/mL (95% confidence interval).

Mortality of larvae treated with the fungal secondary metabolites

The mortality rate of the larvae in the bioassay also positively changed with the concentration of the fungal secondary metabolites (Table 2). In the groups treated with fungal secondary metabolites at three concentrations, 5.5, 55, and 550 $\mu\text{g}/\text{mL}$, the larval mortalities were $29.33 \pm 3.2\%$, $46.00 \pm 1.47\%$, and $71.33 \pm 3.07\%$, respectively. The differences were significant between the groups. The LC_{50} was $2.80 \times 10^2 \mu\text{g}/\text{mL}$ (95% confidence interval).

GST activity alteration of larvae infected with the conidial suspension

The experimental results showed an immunoreaction in the larvae of *D. tabulaeformis* after infection with the conidial suspensions of *B. brongniartii*. The GST activity in the larvae increased from a low level to high level and then decreased again (Figure 1A). The GST activity remained at a very low level at 1 day post-infection. However, at 2 days post-infection, the GST activity increased notably and stayed high until 3 days post-infection. The difference was significant between at 3 days and at the other time points. Afterward, the enzymatic activity declined quickly, and the activity level dropped to the lowest point at 6 days, which was almost the

same as that in the control. Although the GST activity of the insects fluctuated over time, the effect of the conidial concentration in the fungal suspension was marked. For example, in the 6×10^5 spores/mL treatment, there were no significant differences in the GST activities between the control from 1–6 days. A significant difference ($p < 0.05$) was observed in the group treated with 6×10^6 spores/mL. Higher GST activities in the host insect were also found from 1–6 days post-infection in the group infected with 6×10^7 spores/mL, and the difference reached a significant level ($p < 0.05$) compared to the control. Therefore, a higher conidia concentration resulted in the induction of a higher GST activity.

However, it should be noted that with 6×10^7 conidia/mL, the GST activity of the insect reached 0.184 ± 0.013 ($\mu\text{mol}/\text{min}/\text{mg}$ protein) at 3 days, which was the highest value and much higher than other values (Table 3).

EST activity alteration of larvae infected with the conidial suspension

A notable change was also found in the EST activity in *D. tabulaeformis* when the larvae were inoculated with the spore suspension of *B. brongniartii*. The EST activity was measured each day for 1–6 days post-inoculation using α -naphthyl. The results showed that the EST activity fluctuated over time. It initially began to increase, was maintained for 3 days, and then declined at 4 days (Figure 1C). The highest activity was at 3 days and was significantly different from other days ($p < 0.05$). The EST activity was positively correlated with the conidial concentration in the fungal suspension. The larvae were treated with the three concentrations of fungal suspension exhibited higher EST activities that were significant ($p < 0.05$). For example, at 3 days the EST activities were 0.156 ± 0.007 ($\mu\text{mol}/\text{min}/\text{mg}$ protein) in the control, $0.159 \pm$

0.003 ($\mu\text{mol}/\text{min}/\text{mg}$ protein) in the 6×10^5 spores/mL group, 0.184 ± 0.005 ($\mu\text{mol}/\text{min}/\text{mg}$ protein) in the 6×10^6 spores/mL group, and 0.237 ± 0.009 ($\mu\text{mol}/\text{min}/\text{mg}$ protein) in the 6×10^7 spores/mL group (Table 4).

A similar trend in the insect EST activity was observed using β -naphthyl in the experiment (Figure 1E), with a slight increase of the EST activity on the 1st day post-inoculation, which increased for 2 days. The highest EST activity was on the 3rd day; the EST activity started to decline at 4 days, and continued to decrease to the 6th day. The differences between the time points were significant ($p < 0.05$). There was a notable effect of the spore concentration, the EST activities of the treated insects were all higher than in the control section, and the differences were significant ($p < 0.05$). For example, at 3 days the EST activities were 0.207 ± 0.006 ($\mu\text{mol}/\text{min}/\text{mg}$ protein) in the control, 0.214 ± 0.006 ($\mu\text{mol}/\text{min}/\text{mg}$ protein) in the 6×10^5 spores/mL group, 0.227 ± 0.001 ($\mu\text{mol}/\text{min}/\text{mg}$ protein) in the 6×10^6 spores/mL group and 0.266 ± 0.005 ($\mu\text{mol}/\text{min}/\text{mg}$ protein) in the 6×10^7 spores/mL group (Table 5).

GST activity alteration of larvae treated with fungal secondary metabolites

The fungal secondary metabolites altered the GST activity in the *D. tabulaeformis* larvae. Figure 1B shows that GST activity in the treated larvae fluctuated from a low to high level, and then decreased again. The highest activity of GST appeared at 6 hr, and significant differences existed at 2, 4, 6, 12, 24, and 48 hr ($p < 0.05$). The concentration of the secondary metabolite had a notable effect on the GST activity. The high dose (550 $\mu\text{g}/\text{mL}$) group displayed the highest GST activities from 2–48 hr, whereas the activities were the lowest at 5.5 $\mu\text{g}/\text{mL}$ and moderate at

55 µg /mL (Table 6). The differences in the three groups were significant ($p < 0.05$). These results indicated that the secondary metabolites of the pathogenic fungus were the reason for the changes in the insect GST activity.

EST activity alteration of larvae treated with fungal secondary metabolites

Similarly, the secondary metabolites of *B. brongniartii* induced a significant change in the EST activity of the *D. tabulaeformis* larvae. Using α -naphthyl, the EST activity displayed a stable increase during the first period of 2–24 hr post-treatment in all of the test groups, and its peak values appeared at 24 hr. After that, the EST activity declined until 48 hr. The differences in the values for the different time points were significant ($p < 0.05$). The effect of the secondary metabolite concentration in inducing the insect EST activity was marked. For example, at 24 hr the EST activity values were 54.31 ± 3.20 , 90.42 ± 4.33 , and 106.98 ± 5.32 ($\mu\text{mol}/\text{min}/\text{mg}$ protein) for the concentrations of 5.5, 55, and 550 µg/mL, respectively; their differences were significant ($p < 0.05$) (Table 7).

Similar results were obtained using β -naphthyl. The insect EST activity increased from 2 hr to 24 hr post-treatment and declined thereafter (Figure 1F). The differences of the insect EST activities between the time points (2–48 hr) were significant ($p < 0.05$). The concentration effect of the secondary metabolite was also notable. For example, at 24 hr the values of the EST activity were 11.96 ± 0.42 , 12.05 ± 0.47 , and 16.98 ± 0.74 ($\mu\text{mol}/\text{min}/\text{mg}$ protein), for 5.5, 55, and 550 µg/mL, respectively, with the differences being significant ($p < 0.05$) (Table 8).

The EST activity measured in the experiment using α -naphthyl was ten times as high as that

measured in the experiment using β -naphthyl. The EST activity of the *D. tabulaeformis* larvae was stimulated by the fungal metabolites and correlated with the concentration.

Discussion

Entomopathogenic fungi have been studied and applied as important biocontrol agents for certain insect pests. In contrast to the effects of fungal conidia (Lopes et al. 2011; Sabry et al. 2011; Khalid et al. 2012; Nussenbaum et al. 2012), researchers have paid more attention to the role of fungal secondary metabolites on host insects in recent years (Sokolova et al. 1999; Xia et al. 2000; Xia et al. 2001; Zibae et al. 2009a; Zibae et al. 2011). Abdul-Wahid et al. (2012) reported the pathogenicity of both a spore suspension and metabolite solution of two fungi, *Trichoderma harzianum* and *Fusarium solani*, against cockroaches (*Periplaneta americana*). Our study showed that *B. brongniartii* and its metabolites had toxic effects on the pine caterpillar *D. tabulaeformis* larvae, which is in agreement with previous studies. However, in comparison with the conidial infection, it was found that the fungal metabolites were more effective in killing the insects. Moreover, the time until larval mortality was shorter with the metabolites than infection with the conidia. The possible reason was that the conidial infection required more time to complete the series of infection processes, including conidial attachment on the host cuticle, germination, hyphal penetration through the integument, and infection of the internal tissues and organs, whereas the secondary metabolite attack was more direct. Hence, fungal secondary metabolites should be considered for pest management programs in the future.

It is known that certain secondary metabolites of entomopathogenic fungi, including destruxins, cytosporines, and beauverolides, possess functions in impairing the immune response of the host insect, causing death (Gillespie and Claydon 1989; Vilcinskis et al. 1997). Zibae et al. (2011) suggested that the secondary metabolites produced by *B. bassiana* disable several immune mechanisms of *Eurygaster integriceps*, thus helping the fungus to overcome and kill its host. As the enzymatic defense in insects, detoxification enzymes play significant roles in eliminating exotic compounds and maintaining normal physiological functions (Li et al. 2007). During the degradation or sequestration of toxic compounds, the important detoxification enzymes include general ESTs, carboxylesterase, acid phosphatase, alkaline phosphatase, cytochrome P450 monooxygenases, and GSTs (Gordon 1961; Yang et al. 2001). ESTs and GSTs have been reported to be the most important detoxification enzymes that protect insects against the attack from spores and metabolites of entomopathogenic fungi. It is reported that GSTs act in the detoxification process of some insecticides (Flores et al. 2006; Gunasekaran et al. 2011; Sukhirun et al. 2011), whereas ESTs function as hydrolytic enzymes, playing a role in the cleavage of the esters that comprise organic, mineral acids and alcohols or phenols (Michèle et al. 2000). Our study showed that the activities of GST and EST were notably increased when *D. tabulaeformis* larvae were treated with a spore suspension or fungal metabolites. The defense system of the *D. tabulaeformis* larvae reacted to the attack from the conidial infection and the fungal metabolites by resisting oxidative damage and relieving oxidative stress from the toxicants and accelerating toxin metabolism. The trends of GST and EST activities were similar when the larvae were treated with the spores and metabolites. However, the

time required for the alterations was different for the conidial infection and metabolite treatment (Figure 1). This result indicated that the immune reaction of the *D. tabulaeformis* larvae treated with the fungal metabolite was faster than in those infected with the fungal spores, a result different from a previous study. Zibae et al. (2009a) reported that the time required to alter the GST and EST activities of *E. integriceps* adults infected with *B. bassiana* spores was similar to that of adults treated with extracted secondary metabolites of the fungus. This difference was due to the different host insects and the different toxin compositions from different fungi.

The detoxification enzymes of insects have many functions in repairing physiological processes, detoxifying pathogenic products, and metabolizing biologically active compounds. Hence, alterations in their activity may have an impact on the insect to adapt to the environment (Serebrov et al. 2001, 2006). Our study showed that the effect on the detoxification enzyme (EST and GST) activity in *D. tabulaeformis* larvae was closely related to the concentration of the conidia and the metabolites of *B. brongniartii*, and also to the post-treatment time.

In general, our study provides new evidence for understanding the infection mechanism of entomopathogenic fungi and the immune response of the host insect. Our results are significant for applying *B. brongniartii* and its metabolite to control the pest *D. tabulaeformis*.

Acknowledgements

This research was supported by the Shanxi Provincial Key Technologies Research and Development Program (021041 and

2007032009) and the Shanxi Province Forestry Key Program (2010-2012).

References

Abdul-Wahid OA, Elbanna SM. 2012. Evaluation of the insecticidal activity of *Fusarium solani* and *Trichoderma harzianum* against cockroaches; *Periplaneta Americana*. *African Journal of Microbiology Research* 6: 1024–1032.

Bakanova EI, Eremina OY, Roslavtseva SA. 1996. The role of insect microsomal monooxygenases in insecticide degradation. *Agrokhimiya* 10: 145–154.

Bradford M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilising the principle of protein-dye binding. *Analytical Biochemistry* 72: 248–254.

Fan JH, Xie YP, Xue JL, Li BZ. 2008. Isolation and identification of toxins inhibiting *Dentrolimus tabulaeformis* from an antagonistic strain of *Beauveria*. *Acta Microbiologica Sinica* 48: 596–601.

Feyereisen R. 1999. Insect P450 enzymes. *Annual Review of Entomology* 44: 507–533.

Flores AE, Grajales JS, Salas IF, Garica GP, Becerra MHL, Lozano S, Brogdon WG, Black WC, Beaty B. 2006. Mechanisms of insecticide resistance in field populations of *Aedes aegypti* (L.) from Quintana Roo, Southern Mexico. *Journal of the American Mosquito Control Association* 22: 672–677.

Gillespie AT, Claydon N. 1989. The use of entomogenous fungi for pest control and the role of toxins in pathogenesis. *Pesticide Science* 27: 203–215.

Gordon HT. 1961. Nutritional factors in insect resistance to chemicals. *Annual Review of Entomology* 6: 27–54.

Gunasekaran K, Muthukumaravel S, Sahu SS, Vijayakumar T, Jambulingam P. 2011. Glutathione S transferase activity in Indian vectors of malaria: a defense mechanism against DDT. *Journal of Medical Entomology* 48: 561–569.

Han Z, Moores G, Devonshire A, Denholm I. 1998. Association between biochemical marks and insecticide resistance in the cotton aphid, *Aphis gossypii*. *Pesticide Biochemistry and Physiology* 62: 164–171.

Hou TQ. 1987. The pine caterpillars in China. pp. 29-78. Science Press.

Hu FL, Ding XJ, Yang C, Li ZZ, Fan MZ. 2006. Isolation and identification of monoamine oxidase inhibitor from fermentation broth of a strain of *Beauveria*. *Mycosystema* 25: 273–277.

Khalid AH, Mohamed AAA, Ahmed YA, Saad SE, Jin HJ. 2012. Pathogenicity of *Beauveria bassiana* and *Metarhizium anisopliae* against *Galleria mellonella*. *Phytoparasitica* 40: 117–126

Li XZ, Liu YH. 2007. Diet influences the detoxification enzyme activity of *Bactrocera tau* (Walker) (Diptera: Tephritidae). *Acta Entomologica Sinica* 50: 989–995.

Lopes RB, Michereff M, Tigano MS, Neves PMOJ, Lopez EL, Fancelli M, da Silva JP. 2011. Virulence and horizontal transmission of selected Brazilian strains of *Beauveria bassiana* against *Cosmopolites sordidus* under

laboratory conditions. *Bulletin of Insectology* 64: 201–208.

Michèle B, Paul P, Marie-Odile, Soyer-Gobillard. 2000. Esterases in marine dinoflagellates and resistance to the organophosphate insecticide parathion. *International Microbiology* 3: 117–123.

Nussenbaum AL, Lecuona RE. 2012. Selection of *Beauveria bassiana* sensu lato and *Metarhizium anisopliae* sensu lato isolates as microbial control agents against the boll weevil (*Anthonomus grandis*) in Argentina. *Journal of Invertebrate Pathology* 110: 1–7.

Oppenorth FJ. 1985. Glutathione S-Transferase and hydrolytic activity in a tetrachlorvinphos-resistant strain of housefly and their influence on resistance. *Pesticide Biochemistry and Physiology* 11: 176–178.

Roberts DW, St. Leger RJ. 2004. *Metarhizium* spp., cosmopolitan insect-pathogenic fungi: mycological aspects. *Advances in Applied Microbiology* 54: 1–70.

Sabry KH, Abdel-Raheem MA, El-Fatih MM. 2011. Efficacy of the entomopathogenic fungi; *Beauveria bassiana* and *Metarhizium anisopliae* on some insect pests under laboratory conditions. *Egyptian Journal of Biological Pest Control* 21: 33–38.

Serebov VV, Alekseev AA, Glupov VV. 2001. Changes in activity and pattern of hemolymph esterases in larvae of wax moth *Galleria mellonella* L. (Lepidoptera, Pyralidae) during mycosis. *Izvestiya Akademii Nauk Seriya Biologicheskaya* 28: 588–592.

Serebrov VV, Gerber ON, Malyarchuk AA, Martemyanov VV, Alekseev AA, Glupov VV. 2006. Effect of entomopathogenic fungi on

detoxification enzyme activity in greater wax moth *Galleria mellonella* L. (Lepidoptera; Pyralidae) and role of detoxification enzymes in development of insect resistance to entomopathogenic fungi. *Biology Bulletin* 33: 581–586.

Sokolova YY, Sundukov OV. 1999. Inhibition of esterase activity as a property of microsporidial pathogenesis in cricket *Gryllus bimaculatus*. *Parazitologiya* 33(6): 527–537.

Sukhirun N, Pluempanupat W, Bullangpoti V, Koul O. 2011. Bioefficacy of *Alpinia galanga* (Zingiberaceae) rhizome extracts, (E)-p-acetoxycinnamyl alcohol, and (E)-p-coumaryl alcohol ethyl ether against *Bactrocera dorsalis* (Diptera: Tephritidae) and the impact on detoxification enzyme activities. *Journal of Economic Entomology* 104: 1534–1540.

Terriere LC. 1984. Induction of detoxication enzymes in insects. *Annual Review of Entomology* 29: 71–78.

Thomas MB, Read AF. 2007. Can fungal biopesticides control malaria? *Nature Reviews Microbiology* 5: 377–383.

Wang B, Li ZZ, Yao J. 2010. Development status, tendency and strategy of fungal insecticides industry. *Journal of Anhui Agricultural Science* 38: 6269–6270, 6361.

Wang JJ, Cheng WX, Ding W, Zhao ZM. 2004. The effect of the insecticide dichlorvos on esterase activity extracted from the psocids, *Liposcelis bostrychophila* and *L. entomophila*. *Journal of Insect Science* 4:23. Available online: <http://www.insectscience.org/4.23/>

Xia Y, Clarkson JM, Charnley AK. 2001. Acid phosphatases of *Metarhizium anisopliae*

during infection of the tobacco hornworm *Manduca sexta*. *Archives of Microbiology* 176: 427–434.

Xia Y, Dean P, Judge AJ, Gillespie JP, Clarkson JM, Charnley AK. 2000. Acid phosphatases in the haemolymph of the desert locust, *Schistocerca gregaria*, infected with the entomopathogenic fungus *Metarhizium anisopliae*. *Journal of Insect Physiology* 46: 1249–1257.

Yang XQ, Gao XW, Zheng BZ. 2001. Comparison of the activity of the enzymes related to insecticide resistance in *Trialeurodes vaporariorum* and *Bemisia tabaci*. *Chinese Journal of Pesticide Science* 3: 38–43.

Zibae A, Bandani AR, Talaei-Hassanlouei R, Malagoli D. 2011. Cellular immune reactions of the sunn pest, *Eurygaster integriceps*, to the entomopathogenic fungus, *Beauveria bassiana* and its secondary metabolites. *Journal of Insect Science* 11: 138. Available online: <http://www.insectscience.org/11.138/>

Zibae A, Bandani AR, Tork M. 2009a. Effect of the entomopathogenic fungus, *Beauveria bassiana*, and its secondary metabolite on detoxifying enzyme activities and acetylcholinesterase (AChE) of the Sunn pest, *Eurygaster integriceps* (Heteroptera: Scutellaridae). *Biocontrol Science and Technology* 19: 485–498.

Zibae A, Jalali Sendi J, Alinia F, Ghadam-yari M. 2009b. Diazinon resistance in different selected strains of *Chilo suppressalis* Walker (Lepidoptera; Pyralidae), Rice Striped Stem Borer, in the north of Iran. *Journal of Economic Entomology* 102: 1189–1196.

Table 1. Toxicity of the spore suspension on the larvae of *Dendrolimus tabulaeformis*.

Conidial concentration in the suspension (spores/mL)	Days post-treatment	Number of insects	Mortality (r _c /%)	LC ₅₀ (spores/mL)
Control	7	60	3.67 ± 1.07a	2.7 × 10 ⁷
6.0 × 10 ⁵	7	60	23.33 ± 1.73b	
6.0 × 10 ⁶	7	60	37.00 ± 2.40c	
6.0 × 10 ⁷	7	60	59.67 ± 3.07d	

PROBIT: $(p) = -3.871 + 0.5217X$. Data (mean ± SE) in the same column followed by a common letter do not differ significantly ($p > 0.05$) according to univariate analysis of General Linear Model.

Table 2. Toxicity of the secondary metabolites on the larvae *Dendrolimus tabulaeformis*.

Concentration of the secondary metabolite (µg/mL)	Days post-treatment	Number of insects	Mortality (r _c /%)	LC ₅₀ (µg/mL)
Control	7	60	8.33 ± 2.00a	280
5.5	7	60	29.33 ± 3.20b	
55	7	60	46.00 ± 1.47c	
550	7	60	71.33 ± 3.07d	

PROBIT: $(p) = -0.527 + 0.002X$. Data (mean ± SE) in the same column followed by a common letter do not differ significantly ($p > 0.05$) according to univariate analysis of General Linear Model.

Table 3. The GST activity (µmol/min/mg protein) in the *Dendrolimus tabulaeformis* larvae infected with the conidial suspension.

Spore concentration (spores/mL)	Days post-treatment					
	1	2	3	4	5	6
Control group	0.011 ± 0.002a	0.020 ± 0.005a	0.032 ± 0.004a	0.012 ± 0.001a	0.009 ± 0.001a	0.004 ± 0.002a
6.0 × 10 ⁵	0.017 ± 0.005ab	0.023 ± 0.001ab	0.043 ± 0.004ab	0.014 ± 0.004ab	0.015 ± 0.003ab	0.012 ± 0.003ab
6.0 × 10 ⁶	0.019 ± 0.007b	0.036 ± 0.008b	0.050 ± 0.010b	0.024 ± 0.003b	0.017 ± 0.009b	0.016 ± 0.008b
6.0 × 10 ⁷	0.051 ± 0.008c	0.053 ± 0.002c	0.184 ± 0.013c	0.076 ± 0.002c	0.056 ± 0.010c	0.038 ± 0.007c
Univariate (time)	b	c	d	bc	b	a

Results are the mean ± SE of three repetitions. The same letter in each column indicates no significant difference ($p > 0.05$, Duncan Univariate).

Table 4. The EST activity (µmol/min/mg protein) for α-naphthyl in the *Dendrolimus tabulaeformis* larvae infected with the conidial suspension.

Spore concentration (spores/mL)	Days post-treatment					
	1	2	3	4	5	6
Control group	0.042 ± 0.005a	0.103 ± 0.004a	0.156 ± 0.007a	0.085 ± 0.002a	0.064 ± 0.002a	0.038 ± 0.002a
6.0 × 10 ⁵	0.074 ± 0.002b	0.147 ± 0.003b	0.159 ± 0.003b	0.154 ± 0.005b	0.110 ± 0.005b	0.040 ± 0.008b
6.0 × 10 ⁶	0.083 ± 0.002c	0.166 ± 0.004c	0.184 ± 0.005c	0.164 ± 0.004c	0.148 ± 0.002c	0.053 ± 0.004c
6.0 × 10 ⁷	0.084 ± 0.003d	0.193 ± 0.006d	0.237 ± 0.009d	0.184 ± 0.004d	0.150 ± 0.008d	0.066 ± 0.004d
Univariate (time)	b	d	e	d	c	a

Results are the mean ± SE of three repetitions. The same letter in each column indicates no significant difference ($p > 0.05$, Duncan Univariate).

Table 5. The EST activity (µmol/min/mg protein) for β-naphthyl in the *Dendrolimus tabulaeformis* larvae infected with the conidial suspension.

Spore concentration (spores/mL)	Days post-treatment					
	1	2	3	4	5	6
Control group	0.092 ± 0.013a	0.109 ± 0.008a	0.207 ± 0.006a	0.116 ± 0.010a	0.106 ± 0.008a	0.097 ± 0.003a
6.0 × 10 ⁵	0.101 ± 0.012b	0.132 ± 0.004b	0.214 ± 0.006b	0.153 ± 0.003b	0.142 ± 0.009b	0.138 ± 0.007b
6.0 × 10 ⁶	0.103 ± 0.009c	0.152 ± 0.006c	0.227 ± 0.001c	0.188 ± 0.003c	0.147 ± 0.010c	0.145 ± 0.004c
6.0 × 10 ⁷	0.104 ± 0.002d	0.211 ± 0.001d	0.266 ± 0.005d	0.208 ± 0.008d	0.168 ± 0.006d	0.146 ± 0.003d
Univariate (time)	a	d	f	e	c	b

Results are the mean ± SE of three repetitions. The same letter in each column indicates no significant difference ($p > 0.05$, Duncan Univariate).

Table 6. The GST activity (µmol/min/mg protein) in the *Dendrolimus tabulaeformis* larvae treated with the secondary metabolites.

Concentration of the metabolite (µg/mL)	Hours post-treatment					
	2	4	6	12	24	48
Control group	0.039 ± 0.010a	0.131 ± 0.020a	0.378 ± 0.011a	0.131 ± 0.022a	0.039 ± 0.011a	0.014 ± 0.008a
5.5	0.085 ± 0.009b	0.195 ± 0.015b	0.497 ± 0.012b	0.195 ± 0.035b	0.085 ± 0.017b	0.065 ± 0.021b
55	0.261 ± 0.016c	0.204 ± 0.019c	0.652 ± 0.014c	0.289 ± 0.043c	0.204 ± 0.021c	0.077 ± 0.009c
550	0.289 ± 0.016d	0.257 ± 0.020d	0.944 ± 0.030d	0.708 ± 0.002d	0.573 ± 0.006d	0.257 ± 0.004d
Univariate (time)	b	c	f	e	d	a

Results are the mean ± SE of three repetitions. The same letter in each column indicates no significant difference ($p > 0.05$, Duncan Univariate).

Table 7. The EST activity ($\mu\text{mol}/\text{min}/\text{mg}$ protein) for α -naphthyl in the *Dendrolimus tabulaeformis* larvae treated with the secondary metabolites.

Concentration of the metabolite ($\mu\text{g}/\text{mL}$)	Hours post-treatment					
	2	4	6	12	24	48
Control group	11.24 \pm 1.01a	11.52 \pm 0.82a	11.91 \pm 0.76a	25.89 \pm 0.41a	41.87 \pm 6.25a	12.17 \pm 0.49a
5.5	11.51 \pm 1.60b	12.62 \pm 3.81b	13.32 \pm 0.37b	31.59 \pm 1.69b	54.31 \pm 3.20b	14.58 \pm 5.88b
55	25.19 \pm 2.01c	25.57 \pm 4.63c	26.32 \pm 3.28c	40.21 \pm 2.12c	90.42 \pm 4.33c	41.75 \pm 9.19c
550	29.39 \pm 3.12d	30.62 \pm 4.03d	31.14 \pm 2.37d	43.28 \pm 0.36d	106.98 \pm 5.32d	57.95 \pm 0.93d
Univariate (time)	a	b	c	e	f	d

Results are the mean \pm SE of three repetitions. The same letter in each column indicates no significant difference ($p > 0.05$, Duncan Univariate).

Table 8. The EST activity ($\mu\text{mol}/\text{min}/\text{mg}$ protein) for β -naphthyl in the *Dendrolimus tabulaeformis* larvae treated with the secondary metabolites

Concentration of the metabolite ($\mu\text{g}/\text{mL}$)	Hours post-treatment					
	2	4	6	12	24	48
Control group	3.53 \pm 0.31a	3.73 \pm 0.24a	5.22 \pm 0.33a	6.54 \pm 0.90a	10.92 \pm 0.38a	4.26 \pm 0.53a
5.5	3.73 \pm 0.48b	3.93 \pm 0.12b	6.77 \pm 0.29b	8.64 \pm 1.31b	11.96 \pm 0.42b	6.73 \pm 0.20b
55	4.56 \pm 0.62c	5.63 \pm 0.09c	8.25 \pm 1.00c	9.92 \pm 0.62c	12.05 \pm 0.47c	8.27 \pm 1.10c
550	7.53 \pm 0.90d	9.01 \pm 0.27d	9.49 \pm 0.87d	15.72 \pm 1.49d	16.98 \pm 0.74d	13.16 \pm 1.03d
Univariate (time)	a	b	c	e	f	d

Results are the mean \pm SE of three repetitions. The same letter in each column indicates no significant difference ($p > 0.05$, Duncan Univariate).