

## **Cloning, Expression Analysis, and Molecular Modeling of the Gamma-Aminobutyric Acid Receptor Alpha2 Subunit Gene from the Common Cutworm, *Spodoptera litura***

Authors: Zuo, Hongliang, Gao, Lu, Hu, Zhen, Liu, Haiyuan, and Zhong, Guohua

Source: Journal of Insect Science, 13(49) : 1-13

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.013.4901>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



## Cloning, expression analysis, and molecular modeling of the gamma-aminobutyric acid receptor $\alpha 2$ subunit gene from the common cutworm, *Spodoptera litura*

Hongliang Zuo<sup>a</sup>, Lu Gao<sup>b</sup>, Zhen Hu<sup>c</sup>, Haiyuan Liu<sup>d</sup>, and Guohua Zhong<sup>e\*</sup>

Laboratory of Insect Toxicology, South China Agricultural University, Guangzhou 510642, China

### Abstract

Intensive research on the molecule structures of the gamma-aminobutyric acid (GABA) receptor in agricultural pests has great significance to the mechanism investigation, resistance prevention, and molecular design of novel pesticides. The GABA receptor  $\alpha 2$  (*SIGABAR $\alpha 2$* ) subunit gene in *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae) was cloned using the technologies of reverse transcription PCR and rapid amplification of cDNA ends. The genomic DNA sequence of *SIGABAR $\alpha 2$*  has 5164 bp with 8 exons and 7 introns that were in accordance with the GT-AG splicing formula. The complete mRNA sequence of *SIGABAR $\alpha 2$*  was 1965 bp, with an open reading frame of 1500 bp encoding a protein of 499 amino acids. The GABA receptor is highly conserved among insects. The conserved regions include several N-glycosylation, O-glycosylation, and phosphorylation sites, as well as 4 transmembrane domains. The identities that *SIGABAR $\alpha 2$*  shared with the GABA receptor  $\alpha 2$  subunit of *Spodoptera exigua*, *Heliothis virescens*, *Chilo suppressalis*, *Plutella xylostella*, *Bombyx mori* ranged from 99.2% to 87.2% at the amino acid level. The comparative 3-dimensional model of *SIGABAR $\alpha 2$*  showed that its tertiary structure was composed of 4 major  $\alpha$ -helices located at the 4 putative transmembrane domains on one side, with some  $\beta$ -sheets and 1 small  $\alpha$ -helix on the other side. *SIGABAR $\alpha 2$*  may be attached to the membrane by 4  $\alpha$ -helices that bind ions in other conserved domains to transport them through the membrane. The results of quantitative real time PCR demonstrated that *SIGABAR $\alpha 2$*  was expressed in all developmental stages of *S. litura*. The relative expression level of *SIGABAR $\alpha 2$*  was the lowest in eggs and increased with larval growth, while it declined slightly in pupae and reached the peak in adults. The expressions of *SIGABAR $\alpha 2$*  in larvae varied among different tissues; it was extremely high in the brain but was low in the midgut, epicuticle, Malpighian tube, and fat body.

**Keywords:** developmental expression, GABA receptor, gene structure, three-dimensional model, tissue distribution  
**Abbreviations:** **GABA**, gamma-aminobutyric; **qRT-PCR**, quantitative real time PCR; **RACE**, rapid amplification of cDNA ends; **RT-PCR**, reverse transcription PCR; **SIGABAR $\alpha$ 2**, the mRNA of *Spodoptera litura* gamma-aminobutyric receptor  $\alpha$ 2 subunit; **SIGABAR $\alpha$ 2**, the protein of *Spodoptera litura* gamma-aminobutyric receptor  $\alpha$ 2 subunit; **TM**, transmembrane

**Correspondence:** <sup>a</sup> [zhzuo@163.com](mailto:zhzuo@163.com), <sup>b</sup> [gaolu0424@163.com](mailto:gaolu0424@163.com), <sup>c</sup> [huzhlll@163.com](mailto:huzhlll@163.com), <sup>d</sup> [zhanghutuo@163.com](mailto:zhanghutuo@163.com), <sup>e</sup> [guohuazhong@scau.edu.cn](mailto:guohuazhong@scau.edu.cn), \*Corresponding author.

**Editor:** Craig Coates was editor of this paper.

**Received:** 8 October 2011 | **Accepted:** 28 February 2013

**Copyright:** This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

**ISSN:** 1536-2442 | Vol. 13, Number 49

**Cite this paper as:**

Zuo H, Gao L, Hu Z, Liu H, Zhong G. 2013. Cloning, expression analysis, and molecular modeling of the gamma-aminobutyric acid receptor alpha2 subunit gene from the common cutworm, *Spodoptera litura*. *Journal of Insect Science* 13:49. Available online: <http://www.insectscience.org/13.49>

## Introduction

Gamma-aminobutyric acid (GABA) is an inhibitory neurotransmitter in vertebrates and invertebrates that can suppress neurotransmission, hyperpolarize postsynaptic membrane, decrease ion inflow, and slow down cellular metabolism, a process that results in the neuron of postsynaptic membrane being brought into a protective inhibitory condition (Zhao et al. 2005; Ju et al. 2010). The GABA receptor in insects has been intensively explored, and remarkable progress has been made in recent years (Darlison et al. 2005; Casida and Tomizawa 2008). It is an oligomeric protein, its molecule weight is 220–400 kD, and it is composed of 5 different subunits:  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$ , and  $\rho$ . Every subunit has 4 transmembrane domains (TM1, TM2, TM3, and TM4), and the 5 subunits are linked with the hydrophobic ion channel by the TM2 domains to form a chloride ion channel (Chang and Weiss 2000; Ozoe and Akamatsu 2001; Johnston 2005). In terms of gene structure, several kinds of gene sequence of GABA receptors have been cloned from insects whose complete genome sequence have been identified, such as *Drosophila melanogaster*, *Bombyx mori*, *Tribolium castaneum*, and *Anopheles gambiae* (Chen et al. 1994; Miyazaki et al. 1995; Holt et al. 2002; Yu et al. 2010).

The GABA receptor is a major target of a series of important insecticides, such as cyclopentadiene and abamectin. However, resistance to these pesticides is getting more serious in several pest species (Wang and Wu 2007; Chen et al. 2011). The resistance to pesticides was mainly caused by the mutation of target genetic structures in pests (Zeineb et al. 2008; He et al. 2009). It was shown that the mutation of the GABA receptor in *Anopheles funestus*, *Drosophila simulans*, and *Sogatella furcifera* can significantly enhanced their resistance to the traditional pesticides (Gaelle et al. 2005; Lu et al. 2009; Nakao et al. 2010; Wondji et al. 2011). At present, the structures of the GABA receptor have been isolated from only a few species, for instance, *Spodoptera exigua* (Shang et al. 2009), *Plutella xylostella* (Yuan et al. 2010), and *Laodelphax striatellus* (Narusuye et al 2007). The vast majority of GABA receptors in different agricultural pests remain to be identified. Therefore, intensive research on the molecule structures of the GABA receptor in agricultural insects has great significance for research of pesticides, resistance prevention, and novel pesticide invention.

The common cutworm, *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae), is a prevalent agricultural pest in China, causing

**Table 1.** Primers used in the experiment.

Primers	Sequences(5'-3')
Degenerate Primers	SIGARDF: GTGCTCTCYATCAGCTCC
	SIGARDR: GATGTCCCCGATSGTGTA
3'RACE Specific Primer	SIGAR3EF1: ATCCGCAATTCATCATCTT
	SIGAR3EF2: CTATCACCGCTTCTTGTC
3'-Full RACE Core Set Ver. 2.0 (TAKARA)	3'RACE Inner Primer: CGCGGATCCTCCACTAGTGATTCACTATAGG
	3'RACE Outer Primer: TACCGTCGTTCCACTAGTGATTT
5'RACE Specific Primer	SIGAR5ER: CATTGGACAAGAAGCGGTGAT
	SIGAR5ER1: AGAATGATGAATGCCGATG
	SIGAR5ER2: GGGTCTGTCCAAAACCTGC
5' RACE System for Rapid Amplification of cDNA Ends, version 2.0 (Invitrogen)	AUAP Primer: GGCCACGCGTCGACTAGTAC
	UAP Primer: CUACUACUACUAGGCCACGCGTCGACTAGTAC
Specific Primers of Genomic DNA	SIGARDNAF: CACGGGCCCGTCGGCTCCCTGGT
	SIGARDNAR: CTTATTAATAAACTAATTTAATCC
Specific Primers of qRT-PCR	SIGARRTF: GCTGTTACAGAAAATGGCTGCT
	SIGARRTR: TCTACTCGGTGGTGGTGGTCT
Specific Primers of Reference Gene	SIACRTF: CGTCCCATCTACGAAGGTT
	SIACRTR: AGCGGTGGTGGTGAAAGAG

enormous economic losses due to its wide range, rapid reproduction, and resistance to pesticides (Huang et al. 2006). Chemical pesticides are still the main method for controlling *S. litura* all over the world (Ahmad et al. 2008). However, the resistance of *S. litura* to traditional pesticides is getting more and more serious (Shad et al. 2010). Analysis of genetic structure and expression pattern of the GABA receptor in *S. litura* may provide a new strategy to suppress the resistance of pests and increase the efficiency of pesticides. In this study, cDNA of gamma-aminobutyric acid receptor  $\alpha 2$  subunit in *S. litura* (named as *SIGABAR $\alpha 2$* ) of a laboratory susceptible strain was amplified via RT-PCR, and development- and tissue-specific expression patterns were analyzed using quantitative real-time PCR (qRT-PCR).

## Materials and Methods

### Insect

*S. litura* of a laboratory susceptible strain were reared with artificial diet under conditions of  $26 \pm 1^\circ \text{C}$ , 70–80% RH, and 16:8 L:D in a rearing room. The artificial diet contained 100 g of soybeans meal, 80 g of wheat bran, 8 g of casein, 8 g of vitamin C, 2 g of sorbic acid, 0.2 g of choline chloride, 0.2 g of inositol, and 0.2 g of cholesterol in 1 L volume.

### RNA preparation and synthesizing of first-strand cDNA

Fifth instar larvae of *S. litura* were ground in mortar with liquid nitrogen. Total RNA was extracted utilizing the Trizol Reagent (Omega, <http://omegabiotek.com/>) according to the protocol recommended by the manufacturer. 2  $\mu \text{L}$  total RNA was used as template to synthesize first-strand cDNA by PrimerScript Reverse Transcriptase and oligod(T)<sub>18</sub> (TaKaRa, [www.takara-bio.com](http://www.takara-bio.com)).

### Partial sequence amplification of *SIGABAR $\alpha 2$*

Degenerate primers SIGARDF and SIGARDR (Table 1) were designed according to the sequences of the GABA receptor in *B. mori* (NM001099824.1), *S. exigua* (EF535530.1), *P. xylostella* (FJ665610.1), and *Heliothis virescens* (AF006189.1) as reported on NCBI ([www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)). A partial sequence of *SIGABAR $\alpha 2$*  was amplified from first-strand cDNA of 5th instar larvae by SIGARDF and SIGARDR. Thermal cycling was done using the following protocol for touchdown PCR:  $94^\circ \text{C}$  for 3 min; 10 cycles at  $94^\circ \text{C}$  for 30 sec, annealing temperature decrease from  $55$  to  $50^\circ \text{C}$  for 30 sec, 2 cycles at each annealing temperature,  $72^\circ \text{C}$  for 1 min; another 20 cycles at  $94^\circ \text{C}$  for 30 sec,  $50^\circ \text{C}$  for 30 sec,  $72^\circ \text{C}$  for 1 min, and an additional polymerization step at  $72^\circ \text{C}$  for 10 min. Ampli-

fied fragments were purified by 1.5% agarose gels, ligated into pMD 20-T vector (TaKaRa), transformed into *Escherichia coli* of DH5 $\alpha$  to clone, and then positive clones were sequenced (Invitrogen, [www.invitrogen.com](http://www.invitrogen.com)).

### Rapid amplification of cDNA ends (RACE) of *SIGABAR $\alpha$ 2*

Specific primers SIGAR3EF1, SIGAR3EF2, SIGAR5ER, SIGAR5ER1, and SIGAR5ER2 (Table 1) were designed according to the partial sequence of *SIGABAR $\alpha$ 2*, cloned as below to amplify the cDNA ends of *SIGABAR $\alpha$ 2* using Nested PCR. For 3' end amplification, according to the instructions of 3'-Full RACE Core Set Version 2.0 (TaKaRa), the first round PCR used primers SIGAR3EF1 and Outer Primer with the following protocol: 94° C for 3 min; 30 cycles at 94° C for 30 sec, 55° C for 30 sec, 72° C for 2 min; and an additional polymerization step at 72° C for 10 min. The second round of PCR used primers of SIGAR3EF2 and Inner Primer, with the first round PCR product as the template. The thermal cycling protocol was as follows: 94° C for 3 min; 30 cycles at 94° C for 30 sec, 60° C for 30 sec, 72° C for 2 min; and an additional polymerization step at 72° C for 10 min. For 5' end amplification, the 5' RACE cDNA of *S. litura* was synthesized using primer SIGAR5ER according to the manufacturer's instructions of 5' RACE System for Rapid Amplification of cDNA Ends, version 2.0 (Invitrogen). Subsequently, the first round PCR used primers SIGAR5ER1 and Abridged Anchor Primer with the following protocol: 94° C for 2 min; 30 cycles at 94° C for 30 sec, 55° C for 30 sec, 72° C for 1 min; and an additional polymerization step at 72° C for 10 min. The second round PCR used primers SIGAR5ER2 and UAUP Primer, with the first round PCR product as the template. The thermal cycling protocol was as follows: 94° C for 2 min; 30 cycles at 94° C for 30 sec, 60° C for

30 sec, 72° C for 1 min; and an additional polymerization step at 72° C for 7 min. The products of 3' RACE and 5' RACE were purified from 2.0% agarose gels and ligated into the T-vector (TaKaRa), then transformed into *Escherichia coli* of DH5 $\alpha$  to clone, and positive clones were sequenced (Invitrogen).

### Genomic DNA isolation, and *SIGABAR $\alpha$ 2* DNA sequence amplification

Total genomic DNA was isolated from the adult *S. litura* according to the instruction of E.Z.N.A. Insect DNA Kit (OMEGA). Specific primers of genomic DNA SIGADNAF and SIGADNAR (Table 1) were designed according to the complete sequence of *SIGABAR $\alpha$ 2*. PCR reaction was done using TaKaRa LA Taq, which is particularly applicable to long sequence amplification. The PCR reaction was performed with the following protocol: 94° C for 3 min; 30 cycles at 94° C for 30 sec, 60° C for 30 sec, 72° C for 6 min; and an additional polymerization step at 72° C for 10 min. The amplified DNA that were isolated from the adult *S. litura* were purified by 1.5% agarose gels, then ligated into pMD 20-T vector (TaKaRa) and transformed into *E. coli* of DH5 $\alpha$  to clone; then, positive clones were sequenced.

### Analysis of sequence structure and amino acid of *SIGABAR $\alpha$ 2*

Sequence similarity and analysis for conserved domains were performed using BLAST programs on NCBI. Amino acid sequences were derived and analyzed using EditSeq of DNASTAR 7.1 ([www.dnastar.com](http://www.dnastar.com)). Signal peptide was analyzed by online software SignalP 3.0 Server (<http://www.cbs.dtu.dk/services/SignalP/>) (Dyrlov et al. 2004). Transmembrane domains prediction was performed with online software TMHMM Server v. 2.0 (<http://www.cbs.dtu.dk/services/TMHMM/>).

N-glycosylation, O-glycosylation, and phosphorylation sites was analyzed by NetNGlyc 1.0 Server, DictyOGlyc 1.1 Server, and NetPhos 2.0 Server (Center for Biological Sequence Analysis, [www.cbs.dtu.dk](http://www.cbs.dtu.dk)) respectively (Julenius et al. 2005). The secondary structure of SIGABAR $\alpha$ 2 was analyzed using the online software PHD secondary structure prediction method ([http://npsa-pbil.ibcp.fr/cgi-bin/secpred\\_phd.pl](http://npsa-pbil.ibcp.fr/cgi-bin/secpred_phd.pl)). Multiple sequence alignments were performed by Multalin version 5.4.1 (<http://multalin.toulouse.inra.fr/multalin/multalin.html>). A phylogenetic tree was constructed using MEGA 5.05 ([www.megasoftware.net](http://www.megasoftware.net)) with the neighbor joining method and reconstructed with 1000 replicate bootstrap analysis.

### Homology modeling of SIGABAR $\alpha$ 2

The comparative modeling of the tertiary structure of SIGABAR $\alpha$ 2 was performed using the module of Build Homology Models of Discovery Studio 2.0 (Accelrys, [www.accelrys.com](http://www.accelrys.com)). The template model that was applied for comparative modeling was searched in the 3D structure of Protein Data Bank (RCSB, [www.rcsb.org/pdb](http://www.rcsb.org/pdb)). Ten models were predicted, and the model that attained the highest score was chosen as the final template model of SIGABAR $\alpha$ 2. The amino acid rationality of the constructed model was analyzed by Profile-3D and Ramachandran plots.

### Quantitative real-time PCR

The total RNA was extracted from eggs, 1st, 2nd, 3rd, 4th, 5th, and 6th instar larva, prepupae, 1st, 7th, and 14th day pupae, and adults. The brain, malpighian tube, midgut, epicuticle, and fat body of *S.litura* 6th instar larvae were extracted to examine relative transcription levels of SIGABAR $\alpha$ 2 mRNA in different developmental stages and tissues using qRT-PCR. The reaction was performed with BIO-

RAD CFX96 Real-Time PCR Detection System (Bio-Rad, [www.bio-rad.com](http://www.bio-rad.com)) following the manufacturer's recommendations. The final volume of the 25  $\mu$  L reaction system contained 2  $\mu$  L cDNA (less than 100ng), 12.5  $\mu$  L SYBR *Premix Ex Taq*<sup>TM</sup> (TaKaRa), 8.5  $\mu$  L ddH<sub>2</sub>O, 1  $\mu$  L of forward primer (10  $\mu$ M), and 1  $\mu$  L of reverse primer (10  $\mu$ M). qRT-PCR primers for SIGABAR $\alpha$ 2 (SIGARRTF and SIGARRTR, Table 1) and *S.litura*  $\beta$ -actin protein (SIACRTF and SIACRTR, Table 1) were designed according to the complete sequence of SIGABAR $\alpha$ 2 and  $\beta$ -actin cloned from *S.litura* respectively, and synthesized by the technical qRT-PCR primer design and synthesis company TaKaRa. The optimized real-time PCR protocol consisted of an initial step at 95° C for 30 sec followed by 40 cycles at 95° C for 5 sec, 60° C for 30 sec, and 15 sec at 72° C for extension and plate reading. After the cycling protocol, melting curves were obtained by increasing the temperature from 70° C to 95° C (0.4° C/sec) to denature the double-stranded DNA. All samples of real-time PCR were replicated 3 times, the dates were recorded, and data were analyzed by Bio-Rad CFX Manager. Quantification of the transcript level of SIGABAR $\alpha$ 2 was conducted according to the  $2^{-\Delta\Delta C_t}$  method (Livak and Schmittgen 2001), and statistical difference was determined followed by Duncan's Multiple Ranges Test method.

## Results

### Cloning and analysis of SIGABAR $\alpha$ 2 complete sequence

A 370 bp fragment was amplified using degenerate primers SIGARDF and SIGARDR from the cDNA of 5th instar *S.litura*. The 5' end (378 bp) and 3' end (1399 bp) of SIGABAR $\alpha$ 2 was amplified using RACE technique and the specific primers SIGAR3RF1,

SIGAR3RF2, SIGAR5RR1, and SIGAR5RR2, respectively, which was designed based on the partial sequence of *SIGABARα2*. The complete sequence of *SIGABARα2* was spliced according to the partial 5' end and 3' end sequence by EditSeq of DNASTAR 7.1 and searched on the NCBI ([www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)) at amino acid level. The search result demonstrated that it shared 99.2% identity with GABA receptor α2 subunit gene from *S. exigua* (*SeGABARα2*, EF535530.1). It was named as *SIGABARα2*, and the GenBank accession number is JN792582.

The analysis of sequence structure and amino acids indicated that the full length of *SIGABARα2* had 1966 bp that consisted of a 76 bp 5' end untranslated region, a 389 bp 3' end untranslated region, and a 1500 bp open reading frame (Figure 1). The *SIGABARα2* consisted of 499 amino acids, which had a computed molecular mass of 55.53 kD and predicted isoelectric point of 8.84. The amino acid sequence had a predicted N-terminal signal peptide consisting of 29 amino acids (Figure 2a), 2 N-glycosylation sites (Figure 2b), 1 O-glycosylation site (Figure 2c), 30 phosphorylation sites (Figure 2d), a highly conserved dicysteine-loop (Cys-loop), and 4 transmembrane domains (TM1, 253Y-275L; TM2, 284V-306S; TM3, 316D-338M; and TM4, 464I-481Y in *SIGABARα2*) (Figure 2e). Prediction of the *SIGABARα2* secondary structure demonstrated that the amino acid sequence contained 15.43% α-helix, 30.46% extended strand, and 54.11% random coil (Figure 3).

### Homology modeling and rationality analysis of *SIGABARα2*

The amino acid sequence of *SIGABARα2* was blasted in the protein database. The search results indicated that the amino acid sequence

of *SIGABARα2* shared the highest homology (42%) with glutamate-gated chloride channel in *Caenorhabditis elegans* (protein database ID: 3RHW) (Hibbs and Gouaux 2011). The 3D structure of *SIGABARα2* was constructed using the homology tertiary model of 3RHW (Figure 4a). The entire tertiary structure of *SIGABARα2* was like a shuttle. It was mainly composed of 4 α-helices (252G to 270W, 279T to 303A, 313S to 340K, and 458I to 781Y) (Figure 4b) on one side and some β-sheets (67V to 249R) on the other side. The β-turns and irregular curls were distributed in the entire model randomly. Interestingly, the 4 α-helices were located at the 4 putative transmembrane domains, and β-sheets were located after the N-terminal signal peptide. Analysis of Profile-3D showed that the Verify Score of the amino acid index was mostly above zero except for amino acids at the C terminal (Figure 4c), which indicated that most amino acids of *SIGABARα2* were located at rational position in the tertiary structure. Figure 4d shows 13 amino acids (red plots) in the disallowed region and 486 (97.4%) amino acids (green plots) in the allowed region. The results of Profile-3D and Ramachandran plots demonstrated that the tertiary structure of *SIGABARα2* that was simulated by Build Homology Models of Discovery Studio 2.0 was logical.

### Analysis of multiple sequence alignments and the phylogenetic tree

Multiple alignments of known amino acid sequences of GABA receptors were analyzed, and the results indicated that the sequences were highly conserved among different species of insects (Figure 5). The amino acid sequence of *SIGABARα2* has the highest conservation degree among Lepidoptera, with 99.2% to *S. exigua*, 97.6% to *H. virescens*, 95.4% to *Chilo suppressalis*, 92.8% to *P. xylostella*, and 87.2% to *B. mori*. It also shares

remarkable homology among Diptera, with 86.0% to *Culex quinquefasciatus*, 84.4% to *A. gambiae*, 84.0% to *Anopheles funestus*, 83.65% to *Musca domestica*, 82.0% to *Lucilia cuprina*, 82.0% to *Drosophila simulans*, 81.7% to *Aedes aegypti*, and 81.6% to *D. melanogaster*. It has a relatively low identity to Coleoptera and Homoptera, with 83.6% to *T. castaneum* and 78.2% to *L. striatellus*, respectively. After comparing homological amino acid sequences of GABA receptors among different insects, it can be concluded that the highly conserved positions were located at 2 N-glycosylation sites (N42 and N237), 1 O-glycosylation site (456S), 2 phosphorylation sites (404S and 461Y), 1 dicysteine-loop (Cys-loop), and 4 transmembrane domains (TM1, TM2, TM3, and TM4) in SIGABAR $\alpha$ 2. The phylogenetic tree showed that the sequences of amino acid of GABA receptors had the nearest genetic distance in the same order (Figure 6). Lepidoptera, Diptera, Coleoptera, and Homoptera were well segregated from each other. The amino acid sequence of SIGABAR $\alpha$ 2 had the nearest genetic distance to the GABA receptor of *S. exigua* and the farthest genetic distance to the GABA receptor of *L. striatellus*. The genetic distances of the insects between Lepidoptera and Diptera were generally close.

### Genomic character of SIGABAR $\alpha$ 2

The genomic DNA of SIGABAR $\alpha$ 2 was amplified using the specific primers SIGARDNAF and SIGARDNAR from the total DNA of *S. litura* adults. The sequencing results demonstrated that the genomic DNA of SIGABAR $\alpha$ 2 had a size of 5164 bp (Genbank accession number: JN794059) with 7 introns in it (Figure 7). These introns were located at 170 - 1531, 1924 - 2542, 2627 - 3093, 3232 - 3484, 3770 - 4010, 4193 - 4196, and 4439 - 4620 sites on the genomic DNA sequence of SIGABAR $\alpha$ 2 respectively. All of these 7 introns

were in accordance with the GT-AG splicing formula.

### Development- and tissue-specific expression patterns of SIGABAR $\alpha$ 2

The development- and tissue-specific expression patterns of SIGABAR $\alpha$ 2 were detected using qRT-PCR. The results demonstrated that SIGABAR $\alpha$ 2 was expressed in every development stage of *S. litura* (Figure 8a). The relative expression level of SIGABAR $\alpha$ 2 in eggs was the lowest among the various development stages and increased with the larval growth. The relative expression level in 1st, 2nd, 3rd, 4th, 5th, and 6th instar larvae was 5.74, 9.25, 10.17, 13.29, 14.37, and 14.70-fold higher than in eggs, respectively. However, the relative expression level in prepupae and 1st, 7th, and 14th day pupae was 13.84, 13.48, 13.10, and 13.69-fold higher than in eggs, respectively, which were a bit lower than in 6th instar. The highest relative expression level was in adults, which was 16.35-fold higher than in eggs. The relative expression level was remarkably dissimilar in the different tissues of *S. litura* larvae (Figure 8b). The lowest relative expression level tissue was the fat body. The relative expression level in the Malpighian tube, midgut, epicuticle, and brain was 2.57, 8.88, 10.93, and 79.34-fold higher than in the fat body, respectively.

### Discussion

The GABA receptor has been an important target for insecticides. It became the new focus of insect toxicology research because of the emergence of abamectin and fipronil, which target the GABA receptor in insects (Whiting 2003; Li et al. 2006). However, resistance to these insecticides appeared in many kinds of insects; for example, the sensitivity to insecticides was reduced 100-fold in a resistant *D. melanogaster* population, the re-



sistance being caused by a point mutation (Ala 302 to Ser) within the subunit gene *Rdl* of the GABA receptor (Ffrench-Constant et al. 1993; Anthony et al. 1998). A similar mutation was found for the resistant mechanism to fipronil in *P. xylostella* (Zhou et al. 2006). Therefore, further study on the molecular structure characters of GABA receptors is important to prevent resistance to these insecticides and develop novel, efficient, and safe insecticides that target the GABA receptor.

Studies of the GABA receptor subunits are still very limited, except for the RDL subunit. In the present study, the complete sequence of the GABA receptor  $\alpha 2$  subunit in *S. litura* was cloned using the technology of RT-PCR and RACE. The amino acid of SIGABA $\alpha 2$  shares a high degree of conservation with those in other insects, especially those in Lepidoptera, which has the same N-terminal signal peptide at the initial position of the amino acid sequence. The different degrees of identity of GABA receptors among insects may be related to the genetic diversity. Multiple alignments of amino acid sequences of GABA receptors in different insects demonstrated that the highly conserved region was mainly present in positions after the N-terminal signal peptide of the amino acid sequence, such as the N-glycosylation, O-glycosylation, phosphorylation, and 4 transmembrane domains. Combining the result of multiple alignments of amino acid sequences and the simulated 3D structure of SIGABA $\alpha 2$ , it is likely that the  $\alpha 2$  subunit of GABA receptors in insects shares a similar tertiary structure that is embedded into the membrane by 4 transmembrane domains ( $\alpha$ -helixes in the tertiary structure of SIGABA $\alpha 2$ ) that bind ions in other conserved domains ( $\beta$ -sheets in the tertiary structure of SIGABA $\alpha 2$ ) to transport them through the membrane. The DNA sequence information of SIGABA $\alpha 2$  may help

to understand the feedback and regulation mechanism in some physiological process of SIGABA $\alpha 2$  in *S. litura*.

The expression of SIGABA $\alpha 2$  showed a distinct developmental- and tissue-specific pattern in *S. litura*. The expression level of SIGABA $\alpha 2$  was the lowest in the eggs, which was probably due to the incomplete development of the nervous system in the embryo. It increased rapidly with larval growth, which is correlated with the rapid growth and complete development of tissues and the nervous system in the 4th, 5th, and 6th instar larvae. Perhaps the low expression level of SIGABA $\alpha 2$  may be the reason why 1st, 2nd, and 3rd instar larvae are more sensitive to abamectin than 4th, 5th, and 6th instar larvae of *S. litura*. The relative expression level was reduced slightly in the pupal stage compared with 5th and 6th instar larvae and adults, which may be a result of the metamorphic development in the pupae when the nervous system was in the transformative process and the transportation of signals from the external was retarded. The tissue-specific pattern of SIGABA $\alpha 2$  indicated that the GABA receptor was mainly expressed in the brain of the larvae, and was extremely low in the Malpighian tube, midgut, epicuticle, and fat body compared to the brain. These results are in accordance with the distribution of the nervous system being mainly concentrated in the brain, while just a few neurocytes are distributed in other tissues, especially in the fat body. The characters of expression of the GABA receptor in *S. litura* suggest that the best period for controlling *S. litura* using pesticides targeting the GABA receptor is in the early instars. These results may be beneficial to develop some novel, efficient, and safe insecticides targeting the GABA receptor and to create new methods to restrain the GABA receptor in

pests, such as RNA interference and pest-resistant transgenic plants.

### Acknowledgements

The authors gratefully acknowledge Mr. Lloyd D. Crossman for his kind manuscript correction. The work was supported by a grant from the China National Nature Science Foundation (30971944).

### References

- Ahmad M, Sayyed AH, Saleem MA, Ahmad M. 2008. Evidence for field evolved resistance to newer insecticides in *Spodoptera litura* (Lepidoptera: Noctuidae) from Pakistan. *Crop Protection* 27(10): 1367–1372.
- Anthony N, Unruh T, Ganser D, Ffrench-Constant R. 1998. Duplication of the Rdl GABA receptor subunit gene in an insecticide-resistant aphid, *Myzus persicae*. *Molecular and General Genetics* 260(2 - 3): 165–175.
- Casida JE, Tomizawa M. 2008. Insecticide interactions with  $\gamma$ -aminobutyric acid and nicotinic receptors: predictive aspects of structural models. *Journal of Pesticide Science* 33(1): 4–8.
- Chang YC, Weiss DS. 2000. Functional domains of GABA receptors. *Trends in Neurosciences* 16: 127–139.
- Chen R, Belelli D, Lambert JJ, Peters JA, Reyes A, Lan NC. 1994. Cloning and functional expression of a *Drosophila*  $\gamma$ -aminobutyric acid receptor. *Proceedings of the National Academy of Sciences of the United States of America* 91(13): 6069–6073.
- Chen XL, Yuan LZ, Du YZ, Zhang YJ, Wang JJ. 2011. Cross-resistance and biochemical mechanisms of abamectin resistance in the western flower thrips, *Frankliniella occidentalis*. *Pesticide Biochemistry and Physiology* 101(1): 34–38.
- Darlison MG, Pahal I, Thode C. 2005. Consequences of the evolution of the GABA (A) receptor gene family. *Cellular and molecular neurobiology* 25(3-4): 607–624.
- Dyrlov BJ, Nielsen H, Von Heijne G, Brunak S. 2004. Improved prediction of signal peptides: SignalP 3.0. *Journal of Molecular Biology* 340(4): 783–795.
- Ffrench-Constant RH, Steichen JC, Rocheleau TA, Aronstein K, Roush RT. 1993. A single-amino acid substitution in a  $\gamma$ -aminobutyric acid subtype A receptor locus is associated with cyclodiene insecticide resistance in *Drosophila* populations. *Proceedings of the National Academy of Sciences of the United States of America* 90(5): 1957–1961.
- Gaelle LG, Hamon A, Berge JB, Amichot M. 2005. Resistance to fipronil in *Drosophila simulans*: influence of two point mutations in the RDL GABA receptor subunit. *Journal of Neurochemistry* 92(6): 1295–1305.
- He L, Gao XW, Wang JJ, Zhao ZM, Liu NN. 2009. Genetic analysis of abamectin resistance in *Tetranychus cinnabarinus*. *Pesticide Biochemistry and Physiology* 95(3): 147–151.
- Hibbs RE, Gouaux E. 2011. Principles of activation and permeation in an anion-selective Cys-loop receptor. *Nature* 474(7349): 54–60.
- Holt RA, Mani SG, Halpern A, et al. 2002. The genome sequence of the malaria mosquito

*Anopheles gambiae*. *Science* 298(5591): 129–149.

Huang SJ, Han ZJ. 2006. Mechanisms for multiple resistances in field populations of common cutworm, *Spodoptera litura* (Fabricius) in China. *Pesticide Biochemistry and Physiology* 87(1): 14–22.

Johnston GAR. 2005. GABA (A) receptor channel pharmacology. *Current Pharmaceutical Design* 11(15): 1867–1885.

Julenius K, Molgaard A, Gupta R, Brunak S. 2005. Prediction, conservation analysis, and structural characterization of mammalian mucin-type O-glycosylation sites. *Glycobiology* 15(2): 153–64.

Ju XL, Fusazaki S, Hishinuma H, Qiao XM, Ikeda L, Qzoe Y. 2010. Synthesis and structure-activity relationship analysis of bicyclophosphorothionate blockers with selectivity for housefly-aminobutyric acid receptor channels. *Pest Management Science* 66(9): 1002–1010.

Li JK, Xu HH, Jiang DX. 2006. Insect pest control based on  $\gamma$ -aminobutyric acid. *World pesticides* 28(1): 29–31, 36.

Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the  $2^{-\Delta\Delta CT}$  method. *Methods* 25(4): 402–408.

Lu WC, He L, Xue CH, Li M, Wang JJ. 2009. Advances in research of gamma-aminobutyric acid receptors of insects. *Chinese Bulletin of Entomology* 46(1): 52–158.

Miyazaki M, Matsumura F, Beeman RW. 1995. DNA sequence and site of mutation of the GABA receptor of cyclodiene-resistant red

flour beetle, *Tribolium castaneum*. *Comparative Biochemistry and Physiology B: Biochemistry and Molecular Biology* 111B(3): 399–406.

Nakao T, Naoi A, Kawahara N, Hirase K. 2010. Mutation of the GABA receptor associated with fipronil resistance in the whitebacked planthopper, *Sogatella furcifera*. *Pesticide Biochemistry and Physiology* 97(3): 262–266.

Narusuye K, Nakao T, Abe R, Nagatomi Y, Hirase K, Ozoe Y. 2007. Molecular cloning of a GABA receptor subunit from *Laodelphax striatella* (Fallen) and patch clamp analysis of the homo-oligomeric receptors expressed in a *Drosophila* cell line. *Insect Molecular Biology* 16(6): 723–733.

Ozoe Y, Akamatsu M. 2001. Molecular interactions of non-competitive antagonists with ionotropic  $\gamma$ -aminobutyric acid receptors: studies into species difference. *ACS Symposium Series* 774(Agrochemical Discovery): 256–268.

Shad SA, Sayyed AH, Saleem MA. 2010. Cross-resistance, mode of inheritance and stability of resistance to emamectin in *Spodoptera litura* (Lepidoptera: Noctuidae). *Pest Management Science* 66(8): 839–846.

Shang QL, Liang P, Gao XW. 2009. Cloning, developmental and tissue-specific expression of  $\gamma$ -aminobutyric acid(GABA) receptor alpha 2 subunit gene in *Spodoptera exigua* (Hübner). *Pesticide Biochemistry and Physiology* 93(1): 1–7.

Wang LH, Wu YD. 2007. Cross-resistance and biochemical mechanisms of abamectin resistance in the B-type *Bemisia tabaci*. *Journal of Applied Entomology* 131(2): 98–103.

Whiting PJ. 2003. The GABAA receptor gene family: New opportunities for drug development. *Current Opinion in Drug Discovery & Development* 6(5): 648–657.

Wondji CS, Dabire RK, Tukur Z, Irving H, Djouaka R, Morgan JC. 2011. Identification and distribution of a GABA receptor mutation conferring dieldrin resistance in the malaria vector *Anopheles funestus* in Africa. *Insect Biochemistry and Molecular Biology* 41(7): 484–491.

Yuan GR, Gao WY, Yang YH, Wu YD. 2010. Molecular cloning, genomic structure, and genetic mapping of two Rdl-orthologous genes of GABA receptors in the diamondback moth, *Plutella xylostella*. *Archives of Insect Biochemistry and Physiology* 74(2): 81–90.


Yu LL, Cui YJ, Lang GJ, Zhang MY, Zhang CX. 2010. The ionotropic  $\gamma$ -aminobutyric acid receptor gene family of the silkworm, *Bombyx mori*. *Genome* 53(9): 688–697.

Zeineb ES, Bruno L, Gaelle LG, Alain H. 2008. RNA editing regulates insect gamma-aminobutyric acid receptor function and insecticide sensitivity. *NeuroReport* 19(9): 939–943.


Zhao XL, Yeh JZ, Salgado, VL, Narahashi T. 2005. Sulfone metabolite of fipronil blocks  $\gamma$ -aminobutyric acid- and glutamate-activated chloride channels in mammalian and insect neurons. *Journal of Pharmacology and Experimental Therapeutics* 314(1): 363–373.

Zhou XM, Wu QJ, Hu MY, Zhang YJ, Zhu GR, Xu BY. 2006. Cloning and expression characteristics of cDNA fragment of GABAA receptor gene in diamondback moth, *Plutella*


*xylostella*. *Archives of Insect Biochemistry and Physiology* 33(2): 300–305.




**Figure 1.** cDNA nucleotide sequence and deduced amino acid sequence of *Spodoptera litura* SIGABAR $\alpha$ 2. Predicted N-terminal signal peptides were double framed; 2 N-glycosylation sites were treble underlined; 1 O-glycosylation site was double underlined; 30 phosphorylation sites were single underlined; 1 dicysteine-loop was single framed; 4 transmembrane domains were shaded. High quality figures are available online.




**Figure 2.** Analysis of the amino acid sequence of *Spodoptera litura* SIGABAR $\alpha$ 2. (a) Signal P prediction. (b) N-glycosylation site prediction. (c) O-glycosylation site prediction. (d) Phosphorylation site prediction. (e) Transmembrane domains prediction. High quality figures are available online.




**Figure 3.** Secondary structure prediction of the amino acid sequence of *Spodoptera litura* SIGABAR $\alpha$ 2. Random coils are shown in purple, alpha helices are shown in blue, extended strands are shown in red. High quality figures are available online.



**Figure 4.** Prediction of the tertiary structure of *Spodoptera litura* SIGABAR $\alpha$ 2. (a) Comparative modeling of the 3D structure of SIGABAR $\alpha$ 2.  $\alpha$ -helixes are shown in red,  $\beta$ -sheets are shown in blue,  $\beta$ -turns are shown in green, and irregular curls are shown in gray. (b) Comparative second structure in the tertiary model of SIGABAR $\alpha$ 2.  $\alpha$ -helixes are shown in red bars,  $\beta$ -sheets are shown in blue arrows, and irregular curls are shown in gray bars. (c) Profile-3D analysis on the 3D model of SIGABAR $\alpha$ 2. (d) Ramachandran plot analysis of SIGABAR $\alpha$ 2. Credible amino acids are marked with green plots and incredible amino acids are marked with red. High quality figures are available online.




**Figure 5.** Multiple alignment of the amino acid sequence deduced from *Spodoptera litura* SIGABAR $\alpha$ 2 with other GABA receptors. Highly conserved N-phosphorylation, O-glycosylation sites, and phosphorylation sites are marked by black squares, diamonds, and rotundity, respectively; The Cys-loop was marked by a black triangle; transmembrane domains were marked by black bars. The sequences were obtained from the GeneBank database, and the GeneBank accession numbers are as follow: *Spodoptera exigua*: EF535530.1, *Heliothis virescens*: AF006189.1, *Plutella xylostella*: FJ655610.1, *Bombyx mori*: NM001099824.1, *Chilo suppressalis*: HM566200.1, *Aedes aegypti*: U28803.1, *Anopheles funestus*: JF460792.1, *Anopheles gambiae*: XM001688723.1, *Culex quinquefasciatus*: XM001850045.1, *Lucilia cuprina*: AF024647.1, *Musca domestica*: AB177547.2, *Drosophila simulans*: AY017266.1, *Drosophila melanogaster*: U02042.1, *Laodelphax striatellus*: AB253526.1, *Tribolium castaneum*: NM001114337.1. High quality figures are available online.



**Figure 6.** Phylogenetic tree of the GABA receptor in some insects. High quality figures are available online.



**Figure 7.** Schematic model of the genomic DNA of *Spodoptera litura* SIGABAR $\alpha$ 2. The lines at the 2 sides of the model indicate the 5' and 3' UTRs; black and white regions indicate exons and introns respectively; small arrows locating the positions of PCR primers for the amplifying of genomic sequence of SIGABAR $\alpha$ 2. High quality figures are available online.



**Figure 8.** The relative expression levels of *Spodoptera* SIGABAR $\alpha$ 2. (a) The relative expression levels of SIGABAR $\alpha$ 2 in different developmental stages of *S. litura*. (b) The relative expression levels of SIGABAR $\alpha$ 2 in different tissue of *S. litura* larva. Each bar represents the mean  $\pm$  SD of 3 independent assays. The same letters above each bar indicate the expression level no significant difference in the different stages by DMRT ( $p = 0.05$ ). High quality figures are available online.