

Copper Resistance Selection and Activity Changes of Antioxidases in the Flesh Fly *Boettcherisca peregrina*

Authors: Wu, Guoxing, Gao, Xi, Zhu, Jiaying, Hu, Cui, and Ye, Gongyin

Source: Journal of Insect Science, 14(116) : 1-10

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.014.116>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Copper resistance selection and activity changes of antioxidases in the flesh fly *Boettcherisca peregrina*

Guoxing Wu^{1,2a}, Xi Gao^{2b}, Jiaying Zhu^{3c}, Cui Hu^{1d}, Gongyin Ye^{1e*}

¹State Key Laboratory of Rice Biology, Key Laboratory of Agricultural Entomology, Ministry of Agriculture, Institute of Insect Sciences, Zhejiang University, Hangzhou 310058, China

²College of Plant Protection, Yunnan Agricultural University, Kunming 650201, China

³Key Laboratory of Forest Disaster Warning and Control of Yunnan Province and Key Laboratory of Southwest Mountain Forest Resources Conservation and Utilization of Ministry of Education, Southwest Forestry University, Kunming 650224, China

Abstract

Natural populations of *Boettcherisca (Sarcophaga) peregrina* Robineau-Desvoidy (Diptera: Sarcophagidae) were maintained for 20 generations and reared either on unpolluted diet or on polluted diet containing copper at a median lethal concentration (LC₅₀) determined every five generations. This resulted in two reliable strains: the relative susceptible strain (S) and the copper-resistant strain (R). The metal accumulation, growth and development, reproduction, and antioxidant enzymes were analyzed in the two strains. The results showed that compared with the S strain, the R strain showed increased metal accumulation and fecundity of female adults. Regardless of whether larvae were fed on diet with or without Cu²⁺, the R strain showed higher activity of superoxide dismutase and glutathione S-transferase than the S strain, although without statistical significance. Moreover, the activity of superoxide dismutase and glutathione S-transferase increased when *B. peregrina* larvae were exposed to Cu²⁺ at 100 µg/g but decreased when they were exposed to Cu²⁺ at 800 µg/g. Larval catalase activity in the R strain was higher than in the S strain when larvae were fed on diet with or without Cu²⁺, although these differences were significant only at the 100 µg/g concentration. Moreover, the activity of catalase decreased when larvae were exposed to experimental Cu²⁺. Beyond all expectations, larval glutathione reductase activity was not significantly different between the two strains but changed slightly when larvae were exposed to experimental Cu²⁺. These results indicate that copper resistance in *B. peregrina* larvae is mediated by superoxide dismutase, catalase, and glutathione S-transferase. These results also help in establishing a physiological link between antioxidant activity and the resistance level of *B. peregrina* to copper.

Keywords: superoxide dismutase, catalase, glutathione S-transferase, development, reproduction

Correspondence: ^a wugxl@163.com, ^b chonchon@163.com, ^c jyzhu001@gmail.com, ^d chu@zju.edu.cn, ^e gyye@zju.edu.cn, *

Corresponding author

Editor: Nannan Liu was editor of this paper.

Received: 7 October 2012 **Accepted:** 7 March 2013 **Published:** 1 September 2014

Copyright: This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 14, Number 116

Cite this paper as:

Wu G, Gao X, Zhu J, Hu C, Ye G. 2014. Copper resistance selection and activity changes of antioxidases in the flesh fly *Boettcherisca peregrina*. *Journal of Insect Science* 14(116). Available online: <http://www.insectscience.org/14.116>

Introduction

Heavy metal pollution has become a global environmental problem and severely threatens biological diversity and human health. Because insects form an important group with global biological diversity (Sun et al. 2007), much attention has been paid to the potential effects of heavy metal pollution on insects. One of the important indicators of heavy metal pollution in insects are antioxidases, such as superoxide dismutase (SOD), catalase, peroxidase, glutathione peroxidase (GSH-Px), glutathione S-transferase (GST), and glutathione reductase (GR), that remove reactive oxygen species generated by insects exposed to heavy metals (Zamam et al. 1994, Ahmad 1995, Pardini 1995, Migula and Glowacka 1996, Stone et al. 2002, Wilczek et al. 2003, Li et al. 2005, Wang et al. 2006). Like other heavy metals that are required in trace amounts to maintain homeostasis, copper is also one of the micronutrients essential for insects although an excess dietary intake of copper can be toxic in some circumstances. Many of the toxic effects of copper, such as increased lipid peroxidation in cell membranes and DNA damage, are related to its role in the generation of oxygen free radicals (Kadiiska et al. 1992, Bremner 1998, Schümann et al. 2002). In insects, induction of reactive oxygen species by copper alters the activity of antioxidant enzymes (Korsloot et al. 2004, Migula et al. 2004, Wang et al. 2006). Thus far, not much is known about the relationships between antioxidases and copper resistance levels in insects. Therefore, we developed a copper-resistant strain of the flesh fly *Boettcherisca* (*Sarcophaga*) *peregrina* Robineau-Desvoidy (Diptera: Sarcophagidae) and compared the changes effected by exposure to Cu^{2+} between copper-resistant and susceptible flesh fly strains.

Flesh flies have been models to study various aspects of insects, such as physiology, biochemistry, development, and reproduction, among others. Larvae of these flies feed on carrion or feces and cause myiasis in livestock and humans (Braverman et al. 1994, Iqbal et al. 2011). *Boettcherisca peregrina* is one flesh fly species that has been studied widely because of its use in forensic entomology (Sukontason et al. 2010). As a model insect, *B. peregrina* could explain the cytotoxic effects caused by metal pollution on insects. Previous studies have reported the effect of copper on the activity of SOD, catalase, and peroxidase (Wang et al. 2006), the development and reproduction (Wu et al. 2007), and the ultrastructure of midgut and Malpighian tubules (Wu et al. 2009) in *B. peregrina* larvae. Here we report the selection of a copper-resistant strain and changes in antioxidase activity in the flesh fly *B. peregrina*. These results will be helpful to understand the relationship between the activity of antioxidases and the level of copper resistance in *B. peregrina*.

Materials and Methods

Insects

Boettcherisca peregrina was maintained in an artificial climate chamber ($25 \pm 1^\circ\text{C}$, photoperiod of 14:10 L:D) for five years in the laboratory. Larvae were fed on wheat bran:water:porcine liver mixed at a ratio of 3:5:6, and adults were fed on water and sucrose.

Toxicity determination

One-day-old flesh fly larvae were transferred to a glass vial containing 100 g artificial diet supplemented with the following Cu^{2+} concentrations: 50, 100, 200, 400, 800, 1,600, and 3,200 $\mu\text{g/g}$ of artificial diet (Wu et al. 2009). The control group was fed on artificial diet

without Cu^{2+} . Three replicates of about 30 larvae each were used for each Cu^{2+} concentration and the control. The number of dead individuals in each treatment was counted when the larvae pupated. Regression equations, LC_{50} , and confidence interval were calculated by using a data processing system (DPS) for practical statistics (Tang and Feng 2002).

Selection of fly strains

One *B. peregrina* population fed on unpolluted diet was maintained in the laboratory for 20 generations (F_{20}) and resulted in a copper-susceptible strain (S). A copper-resistant strain (R) of *B. peregrina* was created by rearing one-day-old larvae on diet containing Cu^{2+} at LC_{50} concentrations (median lethal concentration) determined every five generations. Individuals surviving the treatment were screened and used for the next generation. Such selection was continued for 20 generations, resulting in the R strain.

Accumulation of Cu^{2+} in larvae and its effects on growth and development

Groups of 300 newly hatched larvae (within 8 hr) in the 20th generation (F_{20}) were fed on diets containing Cu^{2+} at concentrations of 0, 100, and 800 $\mu\text{g/g}$. Each group was reared in a glass bottle, and each concentration was replicated three times. After four days of treatment, 30 larvae from each group were picked randomly, washed with distilled water, and starved for 24 hr. They were then dried on paper towels and weighed on an electronic scale (AB204-E, Mettler Toledo, www.mt.com). Each treatment was divided into four groups with one group of larvae treated with xylene:ethanol (1:1) solution to measure larval body length by using a vernier caliper after stretching. The second group of larvae was used to determine tissue metal content by using an atomic absorption

spectrophotometer (AAnalyst100, Perkin Elmer, www.perkinelmer.com) after digestion with 1 mL mixed acid ($\text{HClO}_4:\text{HNO}_3 = 1:5$ v/v) (Wu et al. 2007). The third group was used to determine enzyme activity, and the remaining larvae in the fourth group were allowed to pupate, emerge, and mate to determine egg production by individual females.

Enzyme activity measurement

To measure enzyme activity, larvae were first washed with the appropriate buffer solution, mixed with ice-cold buffer (1 mL buffer was added to 0.5 g larvae), and homogenized on ice. The homogenate was then centrifuged at $10,000 \times g$ for 10 min at 4°C , and the supernatant was used as the enzyme preparation.

The SOD activity was determined as described previously by McCord and Fridovich (1969) and Deng and Yuan (1991). Briefly, about 10 μL enzyme preparation was added to 4.5 mL Tris-HCl (50 mM; pH 8.2), mixed with 10 μL 45 mM pyrogallol, and homogenized immediately. The homogenate was transferred to a 1 cm cuvette to measure the optical density (OD) at 325 nm every 30 sec, maintaining the auto oxidation rate around 0.07 OD/min. One activity unit was defined as the amount of enzyme required to inhibit 50% auto oxidation in 1 min in 1 mL enzyme preparation. The SOD activity and specific activity were then calculated by using Equations [1] and [2].

SOD activity (U/mL) =

$$\frac{\frac{0.070 - \Delta A_{325\text{nm}} / \text{min}}{0.070} \times 100\%}{50\%} \times \text{reaction volume} \frac{\text{dilution factor}}{\text{sample volume}}$$

[1]

Specific activity (U/mg protein) =

$$\frac{SOD \text{ activity } (U / mL)}{\text{Protein concentration } (mg / mL)}$$

[2]

Catalase activity was measured according to the method described by Barbehenn (2002). The reaction solution contained 665 μ L phosphate buffer (66 mM; pH 7.0), 25 μ L enzyme preparation, and 10 μ L 3% H₂O₂. The OD was measured continuously for 5 min every 30 sec at 240 nm. Catalase activity was expressed as the amount of H₂O₂ reduced per mg protein in 1 min. The extinction coefficient was 39.4 M⁻¹ · cm⁻¹ (Aebi 1984).

Activity of GR was measured by using Bergmeyer’s method (Bergmeyer 1963) with slight modifications. Briefly, about 3 mL reaction mix was prepared that contained 0.1 mM phosphate buffer (pH 7.8), 1 mM Na₂EDTA, 1 mM oxidized glutathione (GSSG), 0.2 mM NADPH-Na₄, and 140 μ L enzyme preparation. Absorbance at 340 nm was measured continuously for 5 min by using a UV spectrophotometer.

The GST was measured as described by Habig et al. (1974). About 50 μ L enzyme preparation was mixed with 1.93 mL 0.1 M phosphate buffer (pH 7.6) and 100 μ L 0.05 M reduced glutathione, incubated at 25°C for 5 min, and then mixed with 20 μ L 0.01 M 1-chloro-2,4-dinitrobenzene. The OD was then measured at 340 nm within 5 min. The extinction coefficient was 9.6 mM⁻¹ · cm⁻¹.

Protein concentration was determined according to Bradford (1976) by using Coomassie Brilliant blue G₂₅₀. A standard curve was prepared with bovine serum albumin.

Data analysis

Data were analyzed by analysis of variance (ANOVA) in the DPS software (Tang and Feng 2002) followed by Duncan’s multiple comparison method to compare within treatments. Levels of significance at $P < 0.05$ were considered as significant and at $P < 0.01$ as highly significant , whereas $P > 0.05$ was considered as not significant.

Results

Selection for copper resistance

To select copper-resistant *B. peregrina*, larvae were fed on diet containing Cu²⁺ at an LC₅₀ concentration, which was determined once every five generations. Copper resistance in *B. peregrina* larvae developed slowly. For the R strain, LC₅₀ values are listed in Table 1. The LC₅₀ in F₂₀ of the R strain was only 1.64-fold higher than that in F₀ and 1.68-fold higher than that in the S strain, which fed on unpol- luted diet for 20 generations.

Cu²⁺ accumulation in larvae

Larvae of the S and R strains fed on diet containing Cu²⁺ at 800 μ g/g accumulated more Cu²⁺ than those fed on diet with Cu²⁺ at 100 μ g/g ($P < 0.01$). On both diets, R-strain larvae accumulated more Cu²⁺ than S-strain larvae. A significant difference in Cu²⁺ accumulation was observed between R- and S-strain larvae fed on diet with Cu²⁺ at 800 μ g/g ($P < 0.01$), but when they were fed on diet with Cu²⁺ at 100 μ g/g, the difference was not statistically significant ($P > 0.05$) (Fig. 1).

Table 1. Resistance development of *B. peregrina* larvae to Cu²⁺.

Generation	Regression equation	Correlation coefficient (r)	LC ₅₀ (95% confidence interval) (μ g/g)	RR
F ₀	y = -1.72 + 2.49 x	0.96	494.89 (411.03 ~ 622.45)	n/a
F ₅	y = -1.88 + 2.51 x	0.96	553.52 (455.04 ~ 711.53)	1.12
F ₁₀	y = -2.17 + 2.57 x	0.96	612.58 (498.52 ~ 805.09)	1.24
F ₁₅	y = -2.94 + 2.79 x	0.97	700.21 (561.91 ~ 951.60)	1.41
F ₂₀	y = -2.43 + 2.55 x	0.95	812.99 (699.22 ~ 982.88)	1.64

RR values represent resistance rates at the LC₅₀ relative to F₀ generation; n/a, not applicable.

Effects of Cu²⁺ on larval development

As shown in Tables 2 and 3, the R and S strains were significantly different from each other in body weight ($P < 0.01$) and length ($P < 0.01$) when larvae were fed on Cu²⁺-free diet. After a four-day treatment with Cu²⁺, no significant difference in body weight ($P > 0.05$) was observed between the two strains at 100 µg/g, but a significant difference was observed at 800 µg/g. However, the body lengths were signifi-




Figure 1. Cu²⁺ accumulation in R and S strains of *B. peregrina* larvae. Values are means ± standard deviation. Same lower-case letters represent no significant difference after exposure to same concentration of Cu²⁺ ($P < 0.05$) and different letters represent significant differences ($P < 0.05$) (Duncan's multiple range test); fw, fresh weight; S, susceptible strain; R, copper-resistant strain.

Table 2. Body weight of R- and S-strain larvae of *B. peregrina* after exposure to Cu²⁺.

Cu ²⁺ concentration (µg/g)	Body weight (mg per larva)	
	S	R
0	65.45 ± 3.07a	47.33 ± 6.09b
100	59.47 ± 4.57a	54.87 ± 3.03a
800	42.96 ± 2.25a	31.00 ± 2.77b

Values are means ± standard deviation. Same lower-case letters in a row represent no significant difference ($P > 0.05$); different letters represent significant differences ($P < 0.05$) (Duncan's multiple range test). S, susceptible strain; R, copper-resistant strain.

Table 3. Body length of R- and S-strain larvae of *B. peregrina* after exposure to Cu²⁺.

Cu ²⁺ concentration (µg/g)	Body length (cm)	
	S	R
0	1.59 ± 0.03a	1.36 ± 0.07b
100	1.47 ± 0.15a	1.26 ± 0.12a
800	1.28 ± 0.19a	1.20 ± 0.10a

Values are means ± standard deviation. Same lower-case letters in a row represent no significant difference ($P > 0.05$); different letters represent significant differences ($P < 0.05$) (Duncan's multiple range test). S, susceptible strain; R, copper-resistant strain.

Table 4. Adult female fecundity in R and S strains of *B. peregrina* after exposure to Cu²⁺.

Cu ²⁺ concentration (µg/g)	Eggs per ♀	
	S	R
0	79.4 ± 5.42a	81.3 ± 7.96a
100	80.8 ± 6.41a	89.6 ± 7.29a
800	47.4 ± 7.47a	53.4 ± 5.91a

Values are means ± standard deviation. Same lower-case letters in a row represent no significant difference ($P > 0.05$) (Duncan's multiple range test). S, susceptible strain; R, copper-resistant strain.

cantly different at both concentrations ($P < 0.01$). Interestingly, body weight decreased at low concentrations of Cu²⁺ (100 µg/g) in the S strain but increased slightly in the R strain. Similar body weights between the two strains at 100 µg/g ($P < 0.05$) suggest the adaptation of the R strain to the low Cu²⁺ concentration.

Effects of Cu²⁺ on adult reproduction

The R and S strains showed no significant difference in adult egg production ($P > 0.05$) when larvae were fed on Cu²⁺-free diet (Table 4). After Cu²⁺ treatment during the larval stage, adult egg production declined significantly in the R and S strains at high Cu²⁺ concentrations (800 µg/g) ($P < 0.01$). However, treatment of larvae with low Cu²⁺ concentrations (100 µg/g) did not cause a significant difference between the R and S strains ($P < 0.05$).

Effects of Cu²⁺ on larval enzyme activity

As shown in Table 5, SOD activity of the R strain was higher than that of the S strain

Table 5. Enzyme activity in R and S strains of *B. peregrina* larvae.

Cu ²⁺ concentration (μg/g)	SOD activity (U/mg protein)		CAT activity (μmol/min/mg protein)		GR activity (U/mg protein)		GST activity (μmol/min/mg protein)	
	S	R	S	R	S	R	S	R
0	183.62 ± 24.03a	193.56 ± 12.76a	34.56 ± 3.74a	49.58 ± 11.82a	0.028 ± 0.004a	0.030 ± 0.002a	248.34 ± 34.67a	264.42 ± 43.42a
100	190.41 ± 19.08a	201.09 ± 22.86a	23.19 ± 4.75b	33.83 ± 3.81a	0.028 ± 0.004a	0.030 ± 0.007a	268.91 ± 35.63a	329.97 ± 50.74a
800	108.25 ± 11.06a	125.41 ± 10.64a	20.69 ± 6.36a	28.98 ± 5.76a	0.036 ± 0.004a	0.034 ± 0.003a	171.20 ± 21.17a	179.39 ± 15.18a

Values are means ± standard deviation. Same lower-case letters in a row in an enzyme activity represent no significant difference ($P > 0.05$) and different letters represent significant differences ($P < 0.05$) (Duncan's multiple range test). SOD, superoxide dismutase; CAT, catalase ; GR, glutathione reductase; GST, glutathione S-transferase; S, susceptible strain; R, copper-resistant strain.

when larvae were fed on diet with or without Cu²⁺, although without statistical significance ($P > 0.05$). The SOD activity was enhanced after larvae of both strains were continuously fed for four days on diet with Cu²⁺ at 100 μg/g but was suppressed when larvae were fed on diet containing Cu²⁺ at 800 μg/g.

Larval catalase activity was higher in the R strain than in the S strain when larvae were fed on diet with or without Cu²⁺, although these differences were significant ($P < 0.05$) only at the 100 μg/g concentration (Table 5). After four days of Cu²⁺ treatment, larval catalase activity was suppressed significantly in both strains ($P < 0.05$) in a dose-dependent manner; the higher the Cu²⁺ concentration the lower the catalase activity. At the low Cu²⁺ concentration (100 μg/g), the activity of catalase was significantly different between the two strains ($P < 0.05$), whereas at the high Cu²⁺ concentration (800 μg/g), the catalase activity was not significantly different between the two strains ($P > 0.05$).

Larval GR activity was similar in the R and S strains when larvae were fed on diet with or without Cu²⁺ (Table 5). Even after a four-day Cu²⁺ treatment, larval GR activity in both strains showed no significant change ($P > 0.05$).

Similar to SOD activity, larval GST activity was higher in the R strain than in the S strain when larvae were fed on diet with or without

Cu²⁺, but without statistical significance ($P > 0.05$) (Table 5). Compared with larval GST activity on Cu²⁺-free diet, GST activity in both strains increased after a four-day treatment with Cu²⁺ at 100 μg/g, although there was no significant difference. However, after a four-day treatment with Cu²⁺ at 800 μg/g, larval GST activity was significantly suppressed in the R and S strains ($P < 0.05$) compared with their larval GST activity in the Cu²⁺-free treatment.

Discussion

Changes in the activity of antioxidant enzymes are important to tolerate copper accumulation in insects (Korsloot et al. 2004, Migula et al. 2004, Wang et al. 2006). However, such antioxidant enzyme activity has not been reported in relative copper-resistant insect strains. In the present study, regardless of the presence or absence of Cu²⁺ in the diet of *B. peregrina* larvae, the R strain had higher SOD, catalase, and GST activity than the S strain. Moreover, the activity of SOD and GST increased when *B. peregrina* larvae were exposed to Cu²⁺ at 100 μg/g but decreased when larvae were exposed to Cu²⁺ at 800 μg/g. Our results differ from those of Wang et al (2006), who reported that the activity of SOD and catalase in *B. peregrina* was significantly inhibited with increasing Cu²⁺ concentrations. These discrepancies support the notion that patterns in antioxidative enzyme activity are species specific and

correlate to the levels of metal pollution or metal loads in the insect's body (Migula et al. 2004).

In contrast, larval GR activity was not significantly different between the R and S strains, and the activity of GR slightly changed when larvae were exposed to experimental Cu^{2+} . This finding is similar to the reported GR activity in *Phyllobius betulae* F. (Coleoptera: Curculionidae) (Migula et al. 2004).

At homeostatic conditions, SOD produces hydrogen peroxide by rapidly dismutating $\text{O}_2^{\bullet-}$ ($2\text{O}_2^{\bullet-} + 2\text{H}^+ \rightarrow \text{H}_2\text{O}_2 + \text{O}_2$) (Richter and Schweizer 1997, Wolin and Mohazzab-H 1997), which is a superoxide anion radical predominantly produced in the respiratory chain of mitochondria by auto oxidation of reduced components (Winyard et al. 1994). In the presence of H_2O_2 , which is an oxidizing environment, Cu^{2+} reacts with reduced glutathione (GSH) to produce Cu^+ and a thiyl radical, GS^\bullet , which reacts with GS^- to result in GSSG^\bullet . The latter is a strongly reducing molecule that reacts rapidly with oxygen to yield $\text{O}_2^{\bullet-}$ (Brouwer and Brouwer-Hoexum 1998). In our study, the increased SOD activity in *B. peregrina* indicates its critical role in converting $\text{O}_2^{\bullet-}$ into H_2O_2 to mitigate the damaging effects exerted by excess $\text{O}_2^{\bullet-}$.

In *B. peregrina*, an increase in the amounts of H_2O_2 also results in increased catalase activity, which is required to decompose H_2O_2 (Sohal et al. 1995). In general, H_2O_2 is degraded to H_2O by two enzyme systems, catalase and glutathione peroxidase (GSH-Px) (Korsloot et al. 2004), although differences between organisms have been observed. For example, GSH-Px plays an important role in mammals but is not present in nematodes and insects (Orr and Sohal 1994, Beckmann and Ames 1997). This indicates that the reaction

$\text{H}_2\text{O}_2 + 2\text{GSH} \rightarrow \text{GSSG} + 2\text{H}_2\text{O}$ catalyzed by GSH-Px and the GSH-regenerated reaction $\text{GSSG} + \text{NADPH} + \text{H}^+ \rightarrow 2\text{GSH} + \text{NADP}^+$ catalyzed by GR do not occur in insects. This could explain the low GR activity that was not significantly different between the R and S strains of *B. peregrina*.

The enzyme GST plays an active role in the detoxification of endogenous and exogenous compounds and is ubiquitously distributed in the biota. Increased GST activity was reported in the carabid beetle *Pterostichus oblongopunctatus* F. (Coleoptera: Curculionidae) collected from metal-polluted areas (Stone et al. 2002). In the western honey bee, *Apis mellifera* L. (Hymenoptera: Apidae), Smirle and Winston (1988) emphasized the role of GST in defense against the cytotoxic action of metals, and similar results were observed in cadmium-treated red wood ants *Formica aquilonia* Yarrow (Hymenoptera: Formicidae) (Migula 1997). Changes in GST activity in the carabid beetle *Poecilus cupreus* L. also depended on the metals used and their doses to detoxify cadmium or zinc (Wilczek et al. 2003). This holds true for GST activity in *B. peregrina* larvae fed on diet with Cu^{2+} . Compared with the larval GST activity in the Cu^{2+} -free treatment, GST activity in the R and S strains of *B. peregrina* increased after a four-day exposure to Cu^{2+} at 100 $\mu\text{g/g}$ and significantly decreased after a four-day exposure to Cu^{2+} at 800 $\mu\text{g/g}$. Moreover, larval GST activity was higher in the R strain than in the S strain indicating that copper resistance in *B. peregrina* may be linked to GST activity.

In conclusion, results of the present study showed that increased resistance to Cu^{2+} in the R strain resulted in enhanced fecundity and Cu^{2+} accumulation compared with the S strain. Copper resistance in *B. peregrina* larvae was mediated by SOD, catalase, and GST rather

than GR. Antioxidative enzyme activity was correlated to the levels of metal exposure or the metal loads in the body. These factors should therefore be considered in the design of experiments to investigate antioxidative enzyme activity in *B. peregrina*.

Acknowledgements

This research was supported by the National Natural Science Foundation of China (Grant No. 30230070 and 30960221). We thank the College of Environmental and Resource Sciences of Zhejiang University for allowing us to use the atomic absorption spectrophotometry.

References

- Aebi, H. 1984.** Catalase in vitro, pp. 121–126. In L. Packer (ed.), *Methods in enzymology*. Academic Press, Orlando, FL.
- Ahmad, S. 1995.** Oxidative stress from environmental pollutants. *Archives of Insect Biochemistry and Physiology* 29: 135–137.
- Barbehenn, R. V. 2002.** Gut-based antioxidant enzymes in a polyphagous and a graminivorous grasshopper. *Journal of Chemical Ecology* 28: 1329–1347.
- Beckmann, K. B., and B. N. Ames. 1997.** Oxidants, antioxidants, and aging, pp. 201–246. In J. G. Scandalios (ed.), *Oxidative stress and the molecular biology of antioxidant defenses*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Bergmeyer, H. U. 1963.** *Methods of enzymatic analysis*. Academic Press, New York, NY, and London, United Kingdom.
- Bradford, M. M. 1976.** A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* 72: 248–254.
- Braverman, I., I. Dano, D. Saah, and B. Gapany. 1994.** Aural myiasis caused by flesh fly larva, *Sarcophaga haemorrhoidalis*. *Journal of Otolaryngology* 23: 204–205.
- Bremner, I. 1998.** Manifestations of copper excess. *The American Journal of Clinical Nutrition* 67: 1069–1073.
- Brouwer, M., and T. Brouwer-Hoexum. 1998.** Biochemical defence mechanisms against copper-induced oxidative damage in the blue crab, *Callinectes sapidus*. *Archives of Biochemistry and Biophysics* 351: 257–264.
- Deng, B. Y., and Q. S. Yuan. 1991.** Improved method of pyrogallol self-oxidation for determination of superoxide dismutase. *Progress in Biochemistry and Biophysics* 18: 163.
- Habig, W. H., M. J. Pabst, and W. B. Jakoby. 1974.** Glutathione S-transferases: the first enzymatic step in mercapturic acid synthesis. *The Journal of Biological Chemistry* 249: 7130–7139.
- Iqbal, J., P. R. Hira, M. M. Marzouk, F. Al-Ali, F. Shelahi, N. Khalid, N. Wyatt, and M. J. Hall. 2011.** Pressure sores and myiasis: flesh flies (Diptera: Sarcophagidae) complicating a decubitus ulcer. *Annals of Tropical Medicine and Parasitology* 105: 91–94.
- Kadiiska, M. B., P. M. Hanna, L. Hernandez, and R. P. Mason. 1992.** In vivo evidence of hydroxyl radical formation after

acute copper and ascorbic acid intake: electron spin resonance spin-trapping investigation. *Molecular Pharmacology* 42: 723–729.

Korsloot, A., C.A.M. van Gestel, and N. M. van Straalen. 2004. *Environmental stress and cellular response in arthropods*. CRC Press, Boca Raton, FL.

Li, L. J., X. M. Liu, Y. P. Guo, and E. B. Ma. 2005. Activity of the enzymes of the antioxidative system in cadmium-treated *Oxya chinensis* (Orthoptera: Acridoidae). *Environmental Toxicology and Pharmacology* 20: 412–416.

McCord, J. M., and I. Fridovich. 1969. Superoxide dismutase. An enzymic function for erythrocuprein (hemocuprein). *The Journal of Biological Chemistry* 244: 6049–6055.

Migula, P. 1997. Molecular and physiological biomarkers in insects as the tools in environmental hazard assessment. *Acta Phytopathologica et Entomologica Hungarica* 32: 231–243.

Migula, P., and E. Glowacka. 1996. Heavy metals as stressing factors in the red wood ants (*Formica polystena*) from industrially polluted forests. *Fresenius' Journal of Analytical Chemistry* 354: 653–659.

Migula, P., P. Laszczyca, M. Augutyniak, G. Wilczek, K. Rozpedek, A. Kafel, and M. Woloszyn. 2004. Antioxidative defence enzymes in beetles from a metal pollution gradient. *Biologia Bratislava* 59: 645–654.

Orr, W. C., and R. S. Sohal. 1994. Extension of life span by overexpression of superoxide

dismutase and catalase in *Drosophila melanogaster*. *Science* 263: 1128–1130.

Pardini, R. S. 1995. Toxicity of oxygen from naturally occurring redox-active pro-oxidants. *Archives of Insect Biochemistry and Physiology* 29: 101–118.

Richter, C., and M. Schweizer. 1997. Oxidative stress in mitochondria, pp.169–200. In J. G. Scandalios (ed.), *Oxidative stress and the molecular biology of antioxidant defenses*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.

Schümann, K., H. G. Classen, H. H. Dieter, J. König, G. Multhaup, M. Rückgauer, K. H. Summer, J. Bernhardt, and H. K. Biesalski. 2002. Hohenheim consensus workshop: copper. *European Journal of Clinical Nutrition* 56: 469–483.

Smirle, M. J., and M. L. Winston. 1988. Detoxifying enzyme activity in worker honey bees: an adaptation for foraging in contaminated ecosystems. *Canadian Journal of Zoology* 66: 1938–1942.

Sohal, R. S., A. Agarwal, S. Agarwal, and W. C. Orr. 1995. Simultaneous overexpression of copper- and zinc-containing superoxide dismutase and catalase retards age-related oxidative damage and increases metabolic potential in *Drosophila melanogaster*. *The Journal of Biological Chemistry* 270: 15671–15674.

Stone, D., P. Jepson, and R. Laskowski. 2002. Trends in detoxification enzymes and heavy metal accumulation in ground beetles (Coleoptera: Carabidae) inhabiting a gradient of pollution. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* 132: 105–112.

Sukontason, K., N. Bunchu, T. Chaiwong, K. Moophayak, and K. L. Sukontason. 2010. Forensically important flesh fly species in Thailand: morphology and developmental rate. *Parasitology Research* 106: 1055-1064.

Sun, H. X., Y. Liu, and G. R. Zhang. 2007. Effects of heavy metal pollution on insects. *Acta Entomologica Sinica* 50(2): 178–185. (In Chinese.)

Tang, Q. Y., and M. G. Feng. 2002. *DPS[®] data processing system for practical statistics*. China Science Press, Beijing, China. (In Chinese.)

Wang, H., G. X. Wu, G. Y. Ye, C. Hu, and J. A. Cheng. 2006. Accumulation of cuprum and cadmium and their effects on the antioxidant enzymes in *Boettcherisca peregrina* exposed to cuprum and cadmium. *Journal of Zhejiang University (Agriculture & Life Science)* 32(1): 77–81. (In Chinese.)

Wilczek, G., P. Kramarz, and A. Babczyńska. 2003. Activity of carboxylesterase and glutathione S-transferase in different life-stages of carabid beetle (*Poecilus cupreus*) exposed to toxic metal concentrations. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* 134: 501–512.

Winyard, P. G., C. J. Morris, V. R. Winrow, M. Zaidi, and D. R. Blake. 1994. Free radicals in the inflammatory response: redox regulation of gene transcription and proteolytic tissue destruction, pp. 361–383. In C. A. Rice-Evans and R. H. Burdonm (eds.), *Free radical damage and its control*. Elsevier Science BV, Amsterdam, The Netherlands.

Wolin, M. S., and K. M. Mohazzab-H. 1997. Mediation of signal transduction by

oxidants, pp. 21–48. In J. G. Scandalios (ed.), *Oxidative stress and the molecular biology of antioxidant defenses*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.

Wu, G. X., X. Gao, G. Y. Ye, C. Hu, and J. A. Cheng. 2007. Effects of dietary copper on the growth, development and reproduction of *Boettcherisca peregrina* (Diptera: Sarcophagidae) in the parental generation and first filial generation. *Acta Entomologica Sinica* 50(10): 1042–1048. (In Chinese.)

Wu, G. X., G. Y. Ye, C. Hu, and J. A. Cheng. 2009. Ultrastructural alterations in the midgut and Malpighian tubules of *Boettcherisca peregrina* exposure to cadmium and copper. *Ecotoxicology and Environmental Safety* 72: 1137–1147.

Zamam, K., R. S. MacGill, J. E. Johnson, S. Ahmad, and R. S. Pardini. 1994. An insect model for assessing mercury toxicity: effect of mercury on antioxidant enzyme activities of the housefly (*Musca domestica*) and the cabbage looper moth (*Trichoplusia ni*). *Archives of Environmental Contamination and Toxicology* 26: 114–118.