



Viability of *Cabralea canjerana* Extracts to Control the South American Fruit Fly, *Anastrepha fraterculus*

Authors: Magrini, Flaviane Eva, Specht, Alexandre, Gaio, Juliano, Girelli, Cristiane Priscila, Miguez, Ignacio, et al.

Source: Journal of Insect Science, 14(47) : 1-13

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.014.47>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Viability of *Cabralea canjerana* extracts to control the South American fruit fly, *Anastrepha fraterculus*

Flaviane Eva Magrini^{1a}, Alexandre Specht^{1,2b}, Juliano Gaio^{1c}, Cristiane Priscila Girelli^{1d}, Ignacio Miguez^{3e}, Horacio Heinzen^{3f}, Valdirene Camatti Sartori^{1g}, and Veronica Cesio^{3h}

¹Universidade de Caxias do Sul – Instituto de Biotecnologia, Laboratório de Controle de Pragas, Caxias do Sul, RS, Brasil 95070-560

²Embrapa Cerrados, BR 020 Km 18 Planaltina, DF, Brasil 73310-970

³Universidad de la Republica, Facultad de Química, Laboratorio de Química de Productos Naturales, Montevideo, Uruguay (11800)

Abstract

Several representatives of Meliaceae contain biologically active compounds that are toxic to insects with few negative effects on the environment and humans. Our study evaluated the activity of ethyl acetate and ethanol extracts from the fruit and seeds of *Cabralea canjerana* (Vellozo) Mart (Sapindales: Meliaceae) on *Anastrepha fraterculus* (Wiedemann) (Diptera: Tephritidae). Limonoids and triterpenes were detected in fruit and seed extracts. Each extract was added to an artificial diet at three concentrations and tested after 24, 48, and 72 hr of extract application. Ethyl acetate extracts were the most active ones and showed the effect of both dose and time elapses after application on the insects. The highest toxic effect on *A. fraterculus* adults was from ethyl acetate extracts from fruit, followed by extracts from seeds. These extracts showed anti-feedant activities. Extract solutions sprinkled on fruits of *Carica papaya* (L.) (Brassicales: Caricaceae) caused oviposition repellency and negatively affected the biological development of *A. fraterculus*. Ethyl acetate extracts highly hampered oviposition, but seed extracts showed lesser oviposition deterrence. The fruit and seed extracts diminished pupal viability. Particularly, the ethyl acetate fruit extract caused malformed adults. The sex ratio was also affected, resulting in female predominance for the fruit extract, while the seed extract showed a dose-dependent effect. Low doses caused male abundance, but at higher concentrations the effect was reversed. These encouraging results showed that the *C. canjerana* extracts have great potential as new tools to be used in integrated pest management programs to protect fruits against *A. fraterculus*.

Keywords: antifeedant activity, toxic activity, oviposition deterrence

Correspondence: ^a flavimagrini@hotmail.com, ^b alexandre.specht@embrapa.br, ^c jgaio@ucs.br, ^d cpgirelli@ucs.br, ^e nacho0412@gmail.com, ^f heihoracio@gmail.com, ^g vc Sartori@ucs.br, ^h veronicacesio@gmail.com, *Corresponding author

Editor: Thomas Miller was editor of this paper.

Received: 30 May 2012 **Accepted:** 30 May 2013 **Published:** 10 April 2014

Copyright: This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 14, Number 47

Cite this paper as:

Magrini FE, Specht A, Gaio J, Girelli CP, Miguez I, Heinzen H, Sartori VC, Cesio V. 2014. Viability of *Cabralea canjerana* extracts to control the South American fruit fly, *Anastrepha fraterculus*. *Journal of Insect Science* 14:47. Available online:

<http://www.insectscience.org/14.47>

Introduction

The South American fruit fly, *Anastrepha fraterculus* (Wiedemann) (Diptera: Tephritidae), is a key pest of many fruits in Argentina, Uruguay, and the southern and southeastern regions of Brazil (Malavasi et al. 2000). Larvae of this fly develop in fruits of more than 67 native and cultivated hosts (Zucchi 2000) and may compromise up to 100% of fruit production (Carvalho 2006). The control of fruit fly adults and larvae is normally carried out with toxic baits. Unfortunately, the chemical compounds (Kovaleski et al. 2000) in these baits not only kill the pest insects, but also harm the applicators and the environment. Residues from these baits contaminate the soil and consequently the food that grows on it, cause biological unbalance, and determine the selection of resistant populations (Whalon et al. 2008). Alternatively, in organic orchards, essential oils, plant extracts, and soaps may be used (Steffens and Schmutterer 1982; Stark et al. 1990; Prokopy and Powers 1995; Van-Randen and Roitberg 1998a, b; Salles and Rech 1999; Singh 2003; Khan et al. 2007; Ali et al. 2011; Efrom et al. 2011).

Hundreds of limonoids have been isolated from Meliaceae (Connolly 1983; Viegas Jr. 2003). These bioactive compounds act in different ways against different insect orders (Mordue and Nisbet 2000). Among the main known Meliaceae are the neem, *Azadirachta indica* (Schmutterer 1990; Bostid 1992); the chinaberry tree, *Melia azedarach* (Carpinella et al. 2002); cedar, *Cedrela fissilis* (Ambrozin et al. 2006); andiroba, *Carapa guianensis* (Lavie et al. 1973; Qi et al. 2004); *Dysoxylum malabaricum* (Govindachari et al. 1994, 1995); *Sandoricum koetjape* (Powell et al. 1991); representatives of *Trichilia* (Roel et al. 2000; Matos et al. 2009; Baatile et al. 2011); and *Cabralea canjerana* (Vellozo) Mart.

(Sapindales: Meliaceae) (Rao et al. 1975; Schmeda-Hirschmann et al. 1992; Braga et al. 2006; Sarria et al. 2011).

Cabralea canjerana is a perennial Meliaceae tree that occurs from Costa Rica to northern Argentina (Pennington and Styles 1975; Barreiros and Souza 1986). Several therapeutic properties of this plant have been exploited, mainly by indigenous populations (Bueno et al. 2005). From *C. canjerana* stems and seeds, limonoids of the gedunin group, mexicanolids (Rao et al. 1975), and dammarane triterpenes have been described (Rao et al. 1975; Braga et al. 2006; Sarria et al. 2011). Schmeda-Hirschmann et al. (1992) described the insecticidal activity of hexane and ethanol extracts of leaves and fruits of *C. canjerana* against nymphs of *Rhodnius neglectus*, one of the vectors of the Chagas disease. Recently, Sarria et al. (2011) described the effects of triterpenes and ocotilone limonoids, cabraleadiol, angolensate, and 3-methyl- β deacetylfissinolide isolated from ethanolic extracts of fruits and seeds against the fall armyworm, *Spodoptera frugiperda*. Our study aimed to evaluate the activity of ethanolic and ethyl acetate extracts obtained from fruits and seeds of *C. canjerana* on the ingestion and oviposition by adults and immatures of *A. fraterculus*.

Materials and Methods

Vegetal material

Fruits of *C. canjerana* were collected in December 2009 from trees growing within the campus of the University of Caxias do Sul (UCS), Caxias do Sul, Brazil (29°9'46"S and 51°8'52"W). The material was identified and classified by the plant taxonomist Dr. Ronaldo Adelfo Wasun at the Museum of Natural Sciences of the UCS. The voucher specimen (in

HUCS # 35647) was deposited in the herbarium of this museum.

Plant extract preparation and identification of compounds

The seeds were manually separated from the fruits (Cañigueral et al. 1998). The pericarp was cut into small pieces (1 cm³), weighed (100 g), and homogenized in hexane (1:3) at room temperature for 30 min with magnetic stirring. Then, the solution was filtered and the hexane extract was discarded. The remaining fruits were extracted sequentially with ethyl acetate and ethanol (1:10) after 48 hr stirring for each solvent. All extracts were filtered with Watman filter paper # 1. The filtrate was evaporated under reduced pressure. Seed extracts were prepared the same way as the fruit extracts. Ethyl acetate fruit and seed extracts were evaluated phytochemically by thin layer chromatography (Merck aluminium TLC silica gel 60 F254, Merck Millipore, www.merckmillipore.com) using the following mobile phase: chloroform:methanol (97:3) with different dyeing reagents: Dragendorff, copper sulfate, vanillin sulfuric acid, sugars-specific dyeing reagent (diphenylamine:aniline:phosphoric acid/acetone), UV, fluorescence, and anisaldehyde (Anonymous 1974; Wagner and Bladt 1996).

Typically, 15 µL of 1 mg/mL extract solutions were seeded in a thin layer chromatography and developed to 10 cm. As both extracts were seeded at the same concentration, rough visual comparison of the relative amounts of the compounds present in each extract could be performed.

Insects and rearing methodology

The *A. fraterculus* used in the bioassays were obtained from insects reared in the Pest Control Laboratory, Institute of Biotechnology,

University of Caxias do Sul. The specimens were maintained at 25 ± 3°C, 70 ± 10% RH, and a 12:12 L:D photoperiod. Adults were confined in cages (30 x 30 x 40) lined with cheesecloth, and larvae were reared on papaya, *Carica papaya* (L.) (Brassicales: Caricaceae), substrate. Other details of the methodology followed Salles (1992) and Machota Jr. et al. (2010).

Antifeedant activity of extracts on adults

The tests were free choice tests. Females and males of *A. fraterculus* that were 15–20 days old were deprived of food for 12 hr. Three pairs were placed in each cylindrical plastic cage (9 cm diameter x 10 cm height). Each cage was considered as a repetition, and each treatment consisted of seven repetitions. A solution of commercial hydrolyzed protein currently employed as an artificial feed attractant for fruit flies (Bio Anastrepha 3%, Bio Controle, www.biocontrole.com.br) (Scoz et al. 2006) was used as nourishing solution both in the control and the treatment. In each treatment, the extracts were included at 5 mg/mL as well as Ponceau red dye at 1g L⁻¹ concentration level (Sigma-Aldrich, www.sigmaaldrich.com) (Cruz et al. 1997; Scoz et al. 2004) to test flies' positive feedings. The test solutions were offered through a roll of cotton inserted into a 4 mL container (Scoz et al. 2004). The antifeedant activity was assessed 24 hr after the beginning of the experiment, and the percentage of antifeedant effect was calculated using the Obeng-Ofori (1995) formula:

$$AF = [(Nc - Nt) / (Nc + Nt)] \times 100.$$

AF = percentage of antifeedant effect

Nc = number of insects on the control after the exposure.

Nt = number of insects on the treatment after the exposure.

Toxic effect of the extracts to adults

This assay was performed as described in the previous section, but the choice possibility was eliminated. The extracts were used at the concentrations of 1, 2.5, and 5 mg/mL in the different treatments, with the addition of 1 g/L⁻¹ Ponceau red dye to the hydrolyzed protein (3%) solution. Only hydrolyzed protein with dye was used to rear the control group in separate cages. The number of dead insects was assessed 24, 48, and 72 hr after the beginning of the experiment. The determination of the lethal concentration (LC₅₀) and lethal time (TL₅₀) was performed using a Probit analysis (Finney 1971). When insect mortality in the control group was over 10%, the mortality values were corrected using Abbott's formula (1925).

Effect of extracts on oviposition

Ten *A. fraterculus* couples, 15 to 20 days old, kept in wooden cages covered with voile fabric (30 x 30 x 30 cm) were assayed in each treatment as well as in control trials. The reported result for each experiment is the average of five repetitions.

Typically, in a no choice assay, one papaya fruit (*C. papaya*) was placed in each cage. The fruits were sprayed with the extract diluted with water and tween at 1, 2.5, and 5 mg/mL concentration level. Only water and tween were sprayed on the control fruits. The fruits were divided in four, and each set of fruits was offered to the flies at different times after spraying (1, 24, 48, and 72 hr). The fruits were exposed to insects for 24 hr, then transferred and stored individually in plastic pots (1000 mL) containing expanded and crushed vermiculite and covered with plastic sheets wrapped in voile fabric. After 15 days, the fruit leftovers were removed, the vermiculite was sifted, and the number of pupae per fruit was investigated.

Effect of the extracts on fruit fly development

Besides the number of insects per fruit, certain morphological parameters of the pupae and adults were also evaluated. The shape of the pupae was characterized as normal or larviform (Hallman and Zhang 1997). Then, the specimens, individualized by fruit, were kept in pots containing moistened vermiculite until the emergence of the adults. In the first days after emergence, the adults were euthanized in a freezer (-17°C) to later identify their sex (Zucchi et al. 2000) and to calculate the sex ratio (Silveira Neto et al. 1976). The percentage of morphological deformities was also assessed in them.

Statistical Analysis

The experimental data were analyzed using one-way ANOVA, whose means were compared by Duncan's test ($P \leq 0.05$) using the transformed square-root average ($x + 0.5$). The means and standard deviations presented in the tables represent the original values.

Results

The phytochemical study was performed following standard procedures (Wagner and Bladt 1996) using thin layer chromatography as a tool to characterize the "digital fingerprint" of the extracts under study. Different dyeing reagents (Draggendorf, copper sulfate/phosphoric acid, vanillin/sulfuric acid, UV at 254 and 365 nm) were employed to obtain the chromatographic profiles for specific groups of secondary metabolites contained in the ethyl acetate fruit and seed extracts. The reported phytochemicals isolated from *C. canjerana* were dammarane type triterpenoids and phenolics, whose presence was confirmed in both extracts. After color development with copper sulfate and vanillin sulfuric acid reagents, their profiles were different (Figure 1).

The major compounds presented at $R_f = 0.65$ and $R_f = 0.4$ revealed yellow and green with vanillin sulfuric acid respectively, indicating the possible presence of lignans and oxidized terpenes in fruits. Lignans were not identified in the seed extract.

All extracts of *C. canjerana* affected the survival of fruit fly adults (Figure 2). The effect was more important when flies were exposed to fruits that were treated 48 to 72 hr before the beginning of the experiment. Even in the lack of a choice, flies did not feed the diet supplemented with extracts for at least 24 hr after the first exposure to the trial sample. However, we found that the flies in the control group fed immediately after being released into the cages. Both time and dose-dependent effects were observed in the mortality of *A. fraterculus* adults (Figure 2). The most significant effects on mortality were associated with the ethyl acetate extracts of both fruits and seeds. They caused the highest mortality among fruit flies without statistical significance between them. Nevertheless, the smallest value for the median lethal concentration (LC_{50}) was observed for the ethyl acetate seeds extract, but the lowest lethal time average (TL_{50}) was observed for ethyl acetate fruit extract (Table 1). These differences in pharmacokinetics as well as in the pharmacodynamics of the toxic action could be due to differences in the chemical profile of the extracts. On the other hand, the mortality induced by the ethanol extracts of fruits (1 mg/mL) and seeds (1 and 2.5 mg/mL) was less than 10% (Figure 2).

Three extracts showed good antifeedant activity towards *A. fraterculus* adults (Figure 3). The ethyl acetate fruit and seed extracts as well as the ethanolic fruit extract showed 66–57% antifeedant activity, whereas the ethanolic seed extract extracts had a 44% antifeedant


activity. This antifeedant effect could be related to the toxicity observed for the ethyl acetate extracts. As the insects died, the antifeedant effect calculation shifts to higher values.

All extracts, at the evaluated concentrations, inhibited oviposition, especially in the early hours. There was also a dose-dependent effect observed. At higher concentrations, significant inhibition of oviposition was observed, even 72 hr after application of the extract, except for the ethanol seeds extract. However, at the highest concentration level, only the ethanol seeds extract completely repelled flies' oviposition in the first 24 hr (Table 2).

Besides the effects described and quantified above, a significant increase in the percentage of larviform pupae from papaya fruits sprayed with fruit extracts of *C. canjerana* was observed, especially at higher concentrations and during the first 24 hr (Table 3).

The pupal viability (Table 4) of the insects from the treatments differed from that observed in the control, especially for the papaya fruit sprayed with *C. canjerana* fruits and seeds ethyl acetate extracts. When the highest concentration of this extract was used, it caused approximately 50% fruit fly mortality, and the effect lasted 72 hr. This residual effect was independent of the applied doses.


The different extracts influenced the proportion of individuals of each sex in different ways. Overall, the sex ratio (Figure 4) was skewed. Females prevailed in the treatments using higher concentrations of the extracts obtained with ethyl acetate. Males, by contrast, were more abundant when the ethanolic extract of fruits was used. These effects were kept throughout the 72-hr period.



The ethyl acetate seed extract caused the highest abnormalities ratio in emerging adults for all the extracts tested. Interestingly, ethanolic seed extracts caused no abnormalities at all. On the other hand, both fruit extracts caused adult abnormalities when compared to the control. Nevertheless, no direct relationship was observed between this occurrence and either concentration or time after spraying (Figure 5). Abnormalities included incomplete emergence, lack of wings, stunted wings, no characteristic wing color, constricted body, oval body, and lack of antennae, mouthparts, and atrophied abdominal bulge.

Discussion

The phytochemical evaluation revealed qualitative differences between the chromatographic profiles of the fruit and seed extracts, which may explain the different biological results obtained for the same bioassay. The colors observed with the vanillin/sulfuric



acid reagent suggests the presence of both triterpenes and lignan type compounds, particularly the yellow spot detected with the latter that was not seen using copper sulfate-phosphoric acid, whereas the reddish-purple color that developed in these cases (Figure 1) indicated the presence of limonoids and triterpenoids, which are the most common type of bioactive compounds reported for different representatives of Meliaceae (Regnault-Roger et al. 2004; Hu et al. 2011), including *Cabralea* species, where the presence of dammarane-type triterpenes in branches and stems was highlighted (Braga et al. 2006).

Cabralea canjerana ethyl acetate fruit and seed extracts caused higher mortality rate to *A. fraterculus* than commercial neem oil (Botton et al. 2003; Efrom et al. 2011), but the sublethal effects were similar to those described previously (Van-Randen and Roitberg 1998a). Given that the mortality rate was directly correlated with the concentration of the extracts and the exposure period (Figure 2,

Table 1. Mean lethal concentration LC_{50} and mean lethal time (TL_{50}) of fruit and seed extracts of *Cabralea canjerana*, made with ethyl acetate and ethanol, incorporated to the artificial diet* of *Anastrepha fraterculus* adults.

Vegetal Structure – solvent	CL_{50} (mg/mL)	Variation	TL_{50} (hours)	Variation
Fruit ethyl acetate	4.18 b**	3.71–4.76	45.36 c	28.01–59.29
Fruit ethanol	4.81 b	4.27–5.58	71.72 b	66.21–79.92
Seed ethyl acetate	2.73 c	0.35–2.44	62.94 b	57.51–70.21
Seed ethanol	7.46 a	6.05–10.53	95.00 a	81.85–126.54

*Hydrolyzed protein (Bio Anastrepha 3%). ** Means followed by distinct letters in each column are not significantly different according to the Duncan's test $P \leq 0.05$.




Figure 3. Mean percent antifeedant activity on *Anastrepha fraterculus* adults offered an artificial diet containing extracts in the concentration of 5 mg/mL and choice of (1) fruit ethyl acetate; (2) fruit ethanol; (3) seed ethyl acetate; (4) seed ethanol. High quality figures are available online.

Table 1), and therefore with volume ingested, we suggest that mortality of *A. fraterculus* was due to a cumulative process. It should be noted that, even in the absence of choice, flies within an experiment initially stopped eating, and just after 24 hr they restarted to feed on the fruit.

The antifeedant activity of the various extracts on *A. fraterculus* (Figure 3) adults indicates that, like other Meliaceae, including neem (Jacobson 1989; Schmetterer 1995; Senthil Nathan et al. 2005, 2006; Coria et al. 2008; Alouani et al. 2009; Masood et al. 2009), *C. canjerana* also has various compounds that act as insect antifeedant agents. After ingestion of the test diet containing the extracts, the flies remained at the bottom of the cage in a lethargic state, not responding to external stimuli. This type of formulation, combining a feed attractant with the ethyl acetate seed extract, is an interesting alternative to control *Anastrepha* flies.

Many of the biological properties (larviform pupae, pupal viability, oviposition, number of eggs, deformed adults, sex distribution) described in our study are related to the well-

Table 2. Mean number and standard error (SE) of *Anastrepha fraterculus* pupae per fruit (*Carica papaya*) pulverized with three concentrations of extracts of fruits and seeds of *Cabralea canjerana*, obtained with ethyl acetate and ethanol.

Vegetal Structure	Concentration	Periods			
		1 hr	24 hr	48 hr	72 hr
Fruit ethyl acetate	1	5.00* ± 0.00cB**	2.00 ± 3.24cBC	9.20 ± 3.20cdB	30.40 ± 3.55aA
	2.5	2.80 ± 0.56cAB	14.40 ± 2.69bA	9.00 ± 0.95cdA	7.00 ± 0.81cdA
	5	1.00 ± 0.20cC	9.60 ± 1.66bA	5.50 ± 1.27dB	5.60 ± 2.39cdB
Fruit ethanol	1	40.00 ± 6.07aA	23.50 ± 1.55aA	22.00 ± 1.18bA	8.00 ± 1.06cdB
	2.5	11.67 ± 0.20bA	13.50 ± 0.97aA	16.40 ± 0.97bA	7.20 ± 1.24cdB
	5	14.40 ± 1.70abA	17.80 ± 4.10aA	16.00 ± 1.64bcA	7.00 ± 1.41cdB
Seed ethyl acetate	1	33.33 ± 1.52abA	28.75 ± 5.06aA	28.25 ± 7.24aA	32.33 ± 6.50aA
	2.5	16.00 ± 3.11abA	17.00 ± 2.11aA	16.00 ± 2.12bcA	27.20 ± 6.80aA
	5	10.50 ± 1.43bA	8.75 ± 2.55bA	9.40 ± 2.55cdA	14.80 ± 6.50bcA
Seed ethanol	1	5.40 ± 0.67cB	21.67 ± 3.21aA	19.67 ± 1.53bA	24.67 ± 5.10aA
	2.5	2.50 ± 0.42cB	20.00 ± 4.58aA	16.00 ± 3.68bcA	19.40 ± 0.23bA
	5	0.00 ± 0.00cC	16.50 ± 3.08bA	29.33 ± 4.72aA	17.80 ± 1.75bAB
Control	-	39.53 ± 1.08aA	29.66 ± 0.88aA	31.58 ± 0.60aA	35.58 ± 1.38aA

*Mean number of five replicates per treatment with ten adult couples of *Anastrepha fraterculus* aged 10 to 15 days. **Means followed by the same letters are not significantly different (Duncan's test $P \leq 0.05$; lowercase letters for differences between extracts and concentrations; capital letters for differences between time intervals).

known ecdisone-like properties of meliaceae triterpenoids and limonoids. These compounds are found mainly in the ethyl acetate extracts from *C. canjerana* due to their physicochemical properties. Ethanol extracts were less active, as they were obtained after the ethyl acetate extraction of the vegetal material, and little if any triterpenoid aglycones were left to be extracted.

The oviposition deterrence (Table 2) induced by the extracts of fruits and seeds of *C. canjerana* in the no choice treatment was similar to that obtained in other studies with other Meliaceae species, where a reduction in the number of tephritid pupae was obtained with

Table 3. Percentage and standard error (SE) of *Anastrepha fraterculus* larviform pupae per fruit (*C. papaya*) pulverized with different concentrations of *Cabralea canjerana* ethyl acetate and ethanol fruit and seed extracts.

Vegetal Structure	Concentration	Periods			
		1 hr	24 hr	48 hr	72 hr
Fruit ethyl acetate	1	20.00* ± 1.06cA**	10.05 ± 1.32 bB	3.52 ± 0.07bC	3.70 ± 0.05cC
	2	45.00 ± 0.56aA	15.12 ± 1.08bB	5.55 ± 0.16bC	8.07 ± 0.12bcB
	5	33.35 ± 0.97bA	29.81 ± 0.75aA	12.01 ± 0.20aB	16.67 ± 0.18bB
Fruit ethanol	1	9.06 ± 0.10dA	4.48 ± 0.18cB	3.00 ± 0.15bB	2.75 ± 0.09cB
	2.5	9.67 ± 0.12dA	3.79 ± 0.07cB	4.09 ± 0.22bB	4.41 ± 0.19cB
	5	35.45 ± 1.98bA	15.55 ± 0.50bB	16.64 ± 0.82aB	18.01 ± 0.98aB
Seed ethyl acetate	1	0.00 ± 0.00cA	0.00 ± 0.00dA	0.00 ± 0.00cA	1.04 ± 0.03dA
	2.5	1.61 ± 0.15cA	1.00 ± 0.02dA	1.00 ± 0.05cA	0.00 ± 0.00dA
	5	0.00 ± 0.00cA	0.00 ± 0.00dA	1.00 ± 0.05cA	0.00 ± 0.00dA
Seed ethanol	1	0.00 ± 0.00cA	0.00 ± 0.00dA	0.00 ± 0.00cA	0.00 ± 0.00dA
	2.5	0.00 ± 0.00cA	0.00 ± 0.00dA	0.00 ± 0.00cA	0.00 ± 0.00dA
	5	-	0.00 ± 0.00dA	0.00 ± 0.00cA	0.00 ± 0.00dA
Control	-	0.00 ± 0.00cA	0.00 ± 0.00dA	0.00 ± 0.00cA	0.00 ± 0.00dA

*Mean number of five replicates per treatment with ten adult couples of *Anastrepha fraterculus* aged 10 to 15 days. **Means followed by the same letters are not significantly different (Duncan's test $P \leq 0.05$; lowercase letters for differences between extracts and concentrations; capital letters for differences between time intervals).

Table 4. Mean percentage and standard error (SE) of pupal viability of *Anastrepha fraterculus* from fruits (*Carica papaya*) pulverized with three concentrations of extracts of fruits and seed of *Cabralea canjerana*, obtained with ethyl acetate and ethanol.

Vegetal Structure	Concentration (mg/mL)	Periods			
		1 hr	24 hr	48 hr	72 hr
Fruit ethyl acetate	1	100.00* ± 0.00aA**	52.14 ± 8.31bC	65.18 ± 5.20bB	66.16 ± 5.13bcB
	2.5	40.28 ± 2.89cB	62.5 ± 13.26bA	48.00 ± 10.84cB	31.78 ± 1.84dC
	5	50.00 ± 0.00cA	53.14 ± 8.11bA	50.00 ± 2.20bcA	42.25 ± 4.65dA
Fruit ethanol	1	51.65 ± 3.89bcA	69.96 ± 5.27abA	60.95 ± 5.79bA	60.71 ± 5.05cA
	2.5	63.60 ± 3.89bB	78.33 ± 3.94abA	58.09 ± 2.36bB	60.78 ± 7.76cB
	5	62.16 ± 0.76bA	69.73 ± 3.04abA	72.62 ± 2.80bA	60.05 ± 0.82cA
Seed ethyl acetate	1	85.19 ± 5.24abA	60.00 ± 5.24bB	87.50 ± 3.61abA	66.81 ± 4.87bcAB
	2.5	68.89 ± 1.83bA	59.22 ± 2.02bAB	68.32 ± 3.46bA	50.67 ± 2.71cB
	5	92.81 ± 2.18aA	61.11 ± 2.02bB	73.46 ± 0.52bB	77.77 ± 3.79abAB
Seed ethanol	1	65.03 ± 4.25bA	60.07 ± 5.76bA	67.85 ± 5.28bA	62.75 ± 3.24bcA
	2.5	98.00 ± 1.25aA	48.85 ± 1.25cC	69.23 ± 0.05bBC	91.67 ± 4.81aA
	5	-	61.35 ± 2.74bA	62.12 ± 3.35bA	80.15 ± 5.95abA
Control	-	98.65 ± 0.83aA	95.07 ± 1.75aA	98.83 ± 0.87aA	96.31 ± 1.30aA

*Mean number of five replicates per treatment with ten adult couples of *Anastrepha fraterculus* aged 10 to 15 days. ** Means followed by the same letters are not significantly different (Duncan's test $P \leq 0.05$; lowercase letters for differences between extracts and concentrations; capital letters for differences between time intervals).

neem extracts (Chen et al. 1996; Masood et al. 2009; Ali et al. 2011).

The extracts inhibited the number of hatching eggs and/or turned eggs and larvae unviable after oviposition for at least 72 hr, effects that were also described by Van-Randen and Roitberg (1998b). The high percentage of larviform pupae in the treatments (Table 3), especially in insects that fed on the ethyl acetate fruit extract treatment, indicates that the triterpenoids and limonoids interfered with the hardening of the cuticle and the stabilization of protein structures (Cruz 2000). These results are similar to the ones obtained with azadirachtin (1 mg/L), where its ingestion by newly-hatched larvae of *C. capitata* completely prevented adult emergence (Vinuela et al. 2000).

The lowest pupal viability (Table 4) was found in the treatment with the ethyl acetate fruit extract. Pupal viability decrease has been reported in several studies that found a reduction in the survival of tephritids that had fed on meliceae extracts (Steffens and Schmutterer 1982; Van-Randem and Roitberg 1998b; Vinuela et al. 2000; Mahfusa et al. 2007).

The variations in sex ratio (Figure 4) associated with the different extracts indicate that




Figure 4. Sex ratio of flies that emerged from fruits of *Carica papaya* pulverized with different concentrations of *Cabralea canjerana* fruits and seed extracts, obtained with ethyl acetate and ethanol, in four distinct periods. High quality figures are available online.

these compounds acted differentially on individuals of each sex. As demonstrated before, several Meliaceae, especially neem, act on the hormone systems of insects (Rembold 1995; Mordue (Luntz) and Nisbet 2000) and therefore have different effects on males and females. This is the first study that addresses the differential effects of extracts on the immature stages and the sex ratio of fruit flies. These results indicate that more studies should be conducted detailing the effects of different substances and extracts on the development of males and females, especially on their reproductive systems.

Besides altering the viability and the number of larviform pupae, the extracts of fruits and seeds used in this study were responsible for




Figure 5. Mean number of flies with deformities that emerged from *Carica papaya* fruits pulverized with different concentrations of fruit and seed extracts of *Cabralea canjerana*, obtained with ethyl acetate and ethanol, in four distinct periods. High quality figures are available online.

an increase in the percentage of deformed adults (Figure 5), especially the ethyl acetate seed extract. The abnormalities found in adults were consistent with those described in other studies involving Meliaceae and tephritids (Steffens and Schmutterer 1982; Singh 2003; Silva et al. 2011).

The results of this study highlight toxic and deterrent properties (antifeedant and anti-oviposition) of extracts from *C. canjerana* to which *A. fraterculus* is vulnerable. These effects were observed for at least 72 hr. The different modes of action of the extracts of fruits and seeds of *C. canjerana* on *A. fraterculus*, particularly the antifeedant activity and oviposition deterrence for at least 72 hr, indicate their possible use in the integrated pest management of this fruit fly.

Acknowledgements

We thank the Fundação de Amparo a Pesquisa do Rio Grande do Sul (FAPERGS) and the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), for their financial support.

References

Abbott WS. 1925. A method of computing the effectiveness of an insecticide. *Journal of Economic Entomology* 18: 265–267.

Ali H, Ahmad S, Hassan G, Amin A, Hussain Z, Naeem M. 2011. Bioefficacy of different plant extracts against melon fruit fly in bitter gourd. *Pakistan Journal of Weed Science Research* 17:143–149.

Alouani A, Rehim N, Soltani N. 2009. Larvicidal activity of a neem tree extract (Azadirachtin) against mosquito larvae in the

Republic of Algeria. *Jordan Journal of Biological Sciences* 2:15–22.

Ambrozin ARP, Leite AC, Bueno FC, Vieira PC, Fernandes JB, Bueno OC, Silva MFGF, Pagnocca FC, Hebling MJA, Bacci Jr M. 2006. Limonoids from andiroba oil and *Cedrela fissilis* and their insecticidal activity. *Journal of the Brazilian Chemical Society* 17: 542–547.

Anonymous. 1974. *Dyeing reagents for thin layer and paper chromatography*. Darmstadt: Merck.

Baatile M, Komane EI, Olivier AMV. 2011. *Trichilia emetica* (Meliaceae) – A review of traditional uses, biological activities and phytochemistry. *Phytochemistry Letters* 4: 1–9.

Barreiros HDS, Souza DSE. 1986. Notas Geográficas e Taxonômicas sobre *Cabralea canjerana* (Vell.) Mart. no Brasil (Meliaceae). *Revista Brasileira de Biologia* 46: 17–26.

Bostid RFR. 1992. *Neem. A tree for solving global problems*. National Academy Press.

Botton M, Scoz PL, Garcia MS, Colleta VD. 2003. Novas alternativas para o monitoramento e controle de *Anastrepha fraterculus* (Wiedmann, 1830) (Diptera: Tephritidae) em frutíferas temperadas. 6^o Encontro Nacional de Fruticultura de Clima Temperado 163–170.

Braga PAC, Soares MS, Silva FMGF, Vieira PC, Fernandes JB, Pinheiro AL. 2006. Dammarane triterpenes from *Cabralea canjerana* (Vell.) Mart. (Meliaceae): Their chemosystematic significance. *Biochemical Systematics and Ecology* 34: 282–290.

- Bueno NR, Castilho RO, Costa RB, Pott A, Pott VJ, Scheidt GN, Batista MS. 2005. Medicinal plants used by the Kaiowá and Guarani indigenous populations in the Caarapó Reserve, Mato Grosso do Sul, Brazil. *Acta Botânica Brasilica* 19: 39–44.
- Cañigueral S, Vila R, Wichtl M. 1998. *Plantas medicinales y drogas vegetales para infusión y tisana*. OEMF Internacional srl.
- Carpinella MC, Ferrayoli C, Valladares G, Defago M, Palacios S. 2002. Potent limonoid insect antifeedant from *Melia azedarach*. *Bioscience, Biotechnology and Biochemistry* 66: 1731–1736.
- Carvalho RS. 2006. Biocontrole de moscas-das-frutas: histórico, conceitos e estratégias. *Bahia Agrícola* 7: 14–17.
- Chen CC, Dong YJ, Cheng LL, Hou RF. 1996. Deterrent effect of neem seed kernel extract on oviposition of the oriental fruit fly (Diptera: Tephritidae) in Guava. *Journal of Economic Entomology* 89: 462–466.
- Connolly JD. 1983. Chemistry of the Meliaceae and Cneoraceae. In: Waterman PG, Grunden MF, Editors. *Chemistry and Chemical Taxonomy of the Rutales*. pp. 175–213. Academic Press.
- Coria C, Almirom W, Valladares G, Carpinella C, Ludueña F, Defago M. 2008. Larvicide and oviposition deterrent effects of fruit and leaf extracts from *Melia azedarach* L. on *Aedes Aegypti* (L.) (Diptera: Culicidae). *Bioresource Technology* 99: 3066–3070.
- Cruz IBM, Humeres E, Oliveira AK. 1997. Toxicity of fenthion to *Anastrepha fraterculus* (Wied.) (Diptera: Tephritidae): dose response analyses. *Anais da Sociedade Entomológica do Brasil* 26: 471–479.
- Cruz IBM, Nascimento JC, Taufer M, Oliveira AK. 2000. Morfologia do aparelho reprodutor e biologia do desenvolvimento. In: Malavasi A, Zucchi RA, Editors. *Moscas-das-frutas de importância econômica no Brasil*. pp. 55–66. Ribeirão Preto.
- Efrom CFS, Redaelli LR, Meirelles RN, Ourique CB. 2011. Laboratory evaluation of phytosanitary products used for control of the South American fruit fly, *Anastrepha fraterculus*, in organic farming. *Crop Protection* 30: 1162–1167. doi:10.1016/j.cropro.2011.05.007.
- Finney DJ. 1971. *Probit analysis*, 3rd edition. Cambridge University Press.
- Govindachari TR, Suresh G, Krishna Kumari GN. 1994. Triterpenoids from *Dysoxylum malabaricum*. *Phytochemistry* 37: 1127–1129.
- Govindachari TR, Narasimhan NS, Suresh G, Partho PD, Gopalakrishnan G, Krishna Kumari GN. 1995. Structured-related insect antifeedant and growth regulating activities of some limonoids. *Journal of Chemical Ecology* 21: 1585–1601.
- Hallman GJ, Zhang QH. 1997. Inhibition of fruit fly (Diptera: Tephritidae) development by pulsed electric field. *Florida Entomologist* 80: 239–248.
- Hu JF, Fan H, Wang LJ, Wu SB, Zhao Y. 2011. Limonoids from the fruits of *Meliatoosendan*. *Phytochemistry Letters* 4: 292–297.
- Jacobson M. 1989. *Focus on Biochemical Pesticides, I. The Neem Tree*. CRC Press.

- Khan M, Hossain MA, Islam MS. 2007. Effects of neem leaf dust and a commercial formulation of a neem compound on the longevity, fecundity and ovarian development of the melon fly, *Bactrocera cucurbitae* (Coquillett) and the oriental fruit fly, *Bactrocera dorsalis* (Hendel) (Diptera: Tephritidae). *Pakistan Journal of Biological Sciences* 10: 3656–3661.
- Kovaleski A, Sugayama RL, Malavasi A. 2000. Controle químico em macieiras. In: Malavasi A, Zucchi RA, Editors. *Moscas-das-frutas de importância econômica no Brasil*. pp. 135–141. Ribeirão Preto.
- Lavie D, Levy EC, Zelnik R. 1973. Constituents of *Carapa guianensis* Aubland and their biogenetic relationship. *Bioorganic Chemistry* 2: 59–64.
- Machota Jr R, Bortoli LC, Tolotti A, Botton M. 2010. Técnica de criação de *Anastrepha fraterculus* (Wied., 1830) (Diptera: Tephritidae) em laboratório utilizando hospedeiro natural. *Embrapa Uva e Vinho, Boletim de Pesquisa e Desenvolvimento* 15.
- Mahfuza K, Aftab MH, Saidul MI. 2007. Effect of Neem leaf dust and commercial formulation of a Neem compound on the longevity, fecundity and ovarian development of melon fly, *Bactrocera cucurbitae* (Coquillett) and oriental fruit fly, *Bactrocera dorsalis* (Hendel) (Diptera: Tephritidae). *Pakistan Journal of Biological Sciences* 10: 3656–3661.
- Malavasi A, Zucchi RA, Sugayama RL. 2000. Biogeografia. In: Malavasi A, Zucchi RA, Editors. *Moscas-das-frutas de importância econômica no Brasil*. pp. 93–98. Ribeirão Preto.
- Masood KK, Rashid MMU, Abdullah K. 2009. Effect of Neem derivatives on infestation, settling and oviposition of melon fruit fly (*Bactrocera cucurbitae* Coq.) (Tephritidae: Diptera). *Pakistan Entomology* 31: 11–15.
- Matos AP, Nebo L, Vieira PC, Fernandes JB, Silva MFGF, Rodrigues RR. 2009. Constituintes químicos e atividade inseticida dos extratos de frutos de *Trichilia elegans* e *T. Catigua* (Meliaceae). *Química Nova* 32: 1553–1556.
- Mordue (Luntz) JA, Nisbet AJ. 2000. Azadirachtin from the neem tree *Azadirachta indica*: its action against insects. *Anais da Sociedade Entomológica do Brasil* 29: 615–632.
- Obeng-Ofori D. 1995. Plant oils as grain protectants against infestations of *Cryptolestes pusillus* and *Rhyzopertha dominica* in stored grain. *Entomologia Experimentalis et Applicata* 77: 133–139.
- Pennington TD, Styles BD. 1975. *A generic monograph of Meliaceae*. Blumea.
- Powell RG, Mikolajczak KN, Zilkowski BW, Mantus EK, Cherry D, Clardy J. 1991. Limonoid antifeedants from seed of *Sandoricum koetjape*. *Journal of Natural Products* 54: 241–246.
- Prokopy RJ, Powers PJ. 1995. Influence of neem seed extract on oviposition and mortality of *Conotrachelus nenuphar* (Col., Curculionidae) and *Rhagoletis pomonella* (Diptera, Tephritidae) adults. *Journal of Applied Entomology* 119: 63–65.

- Qi SH, Wu DG, Zhang S, Luo XD. 2004. Constituents of *Carapa guianensis* Aubl. (Meliaceae). *Pharmazie* 59: 488–490.
- Rao MM, Meshulam H, Zelnik R, Lavie D. 1975. Structure and stereochemistry of limonoids of *Cabralea eichleriana*. *Phytochemistry* 14: 1071–1075.
- Regnault-Roger C, Philogène BJR, Vincent C. 2004. *Biopesticidas de origen vegetal*. Edição Mundi-Prensa.
- Rembold H. 1995. Biological effects of neem and their modes of action: growth and metamorphosis. In: Schmutterer H, Editors. *The neem tree: source of unique natural products for integrated pest management, medicine, industry and other purposes*. pp. 177–194. VCH.
- Roel AR, Vendramim JD, Frighetto, RTS, Frighetto N. 2000. Atividade tóxica de extratos orgânicos de *Trichilia pallida* (Swartz) (Meliaceae) sobre *Spodoptera frugiperda* (J.E. Smith). *Anais da Sociedade Entomológica do Brasil* 29: 799–808.
- Salles LAB. 1992. Metodologia de criação de *Anastrepha fraterculus* (Wied., 1830) (Diptera:Tephritidae) em dieta artificial em laboratório. *Anais da Sociedade Entomológica do Brasil* 21: 479–486.
- Salles LAB, Rech NL. 1999. Efeito de extratos de nim (*Azadirachta indica*) e cinamomo (*Melia azedarach*) sobre *Anastrepha fraterculus* (Wied.) (Diptera:Tephritidae). *Revista Brasileira de Agrociência* 5: 225–227.
- Sarria AL, Soares MS, Matos AP, Fernandes JB, Vieira PC, Silva MFGF. 2011. Effect of triterpenoids and limonoids isolated from *Cabralea canjerana* and *Carapa guianensis* (Meliaceae) against *Spodoptera frugiperda* (J. E. Smith). *Z.Naturforsch* 66: 245–250.
- Schmeda-Hirschmann G, Rojas de Arias AA. 1992. Screening method for natural products on triatomine bugs. *Phytotherapy Research* 6: 68–73.
- Schmutterer H. 1990. Properties and potential of natural pesticides from the neem tree, *Azadirachta indica*. *Annual Review of Entomology* 35: 271–297.
- Schmutterer H. 1995. *The neem tree: source of unique natural products for integrated pest management, medicine, industry and other purposes*. VCH.
- Scoz PL, Botton M, Garcia MS. 2004. Controle químico de *Anastrepha fraterculus* (Wied.) (Diptera: Tephritidae) em laboratório. *Ciência Rural* 34: 1689–1694.
- Scoz PL, Botton M, Garcia MS, Pastori LP. 2006. Avaliação de atrativos alimentares e armadilhas para o monitoramento de *Anastrepha fraterculus* (wiedemann, 1830) (Diptera: tephritidae) na cultura do pessegueiro (*Prunus persica* (L.) Batsh). *Idesia (Chile)* 24:7–13.
- Senthil Nathan S, Kalaivani K, Muruga K. 2005. Effects of neem limonoids on the malaria vector *Anopheles stephensi* Liston (Diptera: Culicidae). *Acta Tropica* 96: 47–55.
- Senthil Nathan S, Savith G, George DK, Narmadha A, Suganya L, Chung PG. 2006. Efficacy of *Melia azedarach* L. extract on the malarial vector *Anopheles stephensi* Liston (Diptera: Culicidae). *Bioresource Technology* 97: 1316–1323.

Silva MA, Alvarenga CD, Bezerra-Silva GCD, Mastrangelo T, Lopes-Mielezrski GN, Giustolin T. 2011. Toxic effects of neem seed cake on the larval-pupal (prepupal) stage of Mediterranean fruit fly (Diptera: Tephritidae). *Fruits* 66: 363–369.

Silveira Neto S, Nakano O, Bardin D, Villa Nova NA. 1976. *Manual de ecologia dos insetos*. Agronômica Ceres.

Singh S. 2003. Effects of aqueous extract of neem seed kernel and azadirachtin on the fecundity, fertility and post-embryonic development of the melon fly, *Bactrocera cucurbitae* and the oriental fruit fly, *Bactrocera dorsalis* (Diptera: Tephritidae). *Journal of Applied Entomology* 127: 540–547.

Stark JD, Vargas RI, Thalman RH. 1990. Azadirachtin effects on metamorphosis, longevity and reproduction of three tephritid fruit fly species (Diptera). *Journal of Economic Entomology* 83: 2168–2174.

Steffens RJ, Schmutterer H. 1982. The effect of a crude methanolic neem (*Azadirachtin indica*) seed kernel extract on metamorphosis and quality of adults of the Mediterranean fruit fly, *Ceratitidis capitata* Wied. (Diptera, Tephritidae). *Zeitschrift für Angewandte Entomologie* 94: 98–103.

Van-Randen EJ, Roitberg BD. 1998a. Effect of a neem (*Azadirachta indica*) – based insecticide on oviposition deterrence, survival, behavior and reproduction of adult western cherry fruit fly (Diptera: Tephritidae). *Ecology and Behavior* 91: 123–131.

Van-Randen EJ, Roitberg BD. 1998b. The effect of a neem (*Azadirachta indica*) based insecticide on survival and development of juvenile western cherry fruit fly (*Rhagoletis*

indifferens) (Diptera: Tephritidae). *The Canadian Entomologist* 130: 869–876.

Viegas Jr C. 2003. Terpenos com atividade inseticida: uma alternativa para o controle químico de insetos. *Química Nova* 26: 390–400.

Vinuela E, Adán A, Smagghe G, González M, Medina MP, Budia F, Vogt H, Estal P. 2000. Laboratory effects of ingestion of azadirachtin by two pests (*Ceratitidis capitata* and *Spodoptera exigua*) and three natural enemies (*Chrysoperla carnea*, *Opius concolor* and *Podisus maculiventris*). *Biocontrol Science and Technology* 10: 65–177.

Zucchi RA. 2000. Espécies de *Anastrepha*, sinónimas, plantas hospedeiras e parasitóides. In: Malavasi A, Zucchi R.A, Editors. *Mosca-das-frutas de importância econômica no Brasil: conhecimento básico e aplicado*. pp. 41-48. Ribeirão Preto.

Wagner H, Bladt S. 1996. *Plant Drug Analysis*. 2nd edition. Springer.

Whalon ME, Mota-Sanchez D, Hollingworth RM. 2008. *Global Pesticide Resistance in Arthropods*. CABI.