

Interaction of Omega, Sigma, and Theta Glutathione Transferases with p38b Mitogen-Activated Protein Kinase from the Fruit Fly, Drosophila melanogaster

Authors: Wongtrakul, J., Janphen, K., Saisawang, C., and Ketterman, A.J.

Source: Journal of Insect Science, 14(60) : 1-13

Published By: Entomological Society of America

URL: https://doi.org/10.1673/031.014.60

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Interaction of Omega, Sigma, and Theta glutathione transferases with p38b mitogen-activated protein kinase from the fruit fly, Drosophila melanogaster

J. Wongtrakul^{1a}, K. Janphen^{1b}, C. Saisawang^{2c}, and A.J. Ketterman^{2d*}

¹Research Institute for Health Sciences (RIHES), Chiang Mai University, P.O.BOX 80 CMU, Chiang Mai, Thailand 50200 ²Institute of Molecular Biosciences, Mahidol University, Nakhon Pathom, Thailand 73170

Abstract

Glutathione S-transferases (GSTs) are a diverse family of phase II detoxification enzymes found in almost all organisms. Besides playing a major role in the detoxification of xenobiotic and toxic compounds, GSTs are also involved in the regulation of mitogen activated protein (MAP) kinase signal transduction by interaction with proteins in the pathway. An *in vitro* study was performed for Theta, Omega, Sigma GSTs and their interaction with MAP kinase p38b protein from the fruit fly *Drosophila melanogaster* Meigen (Diptera: Drosophilidae). The study included the effects of all five Omega class GSTs (DmGSTO1, DmGSTO2a, DmGSTO2b, DmGSTO3, DmGSTO4), all five Theta class GSTs (DmGSTT1, DmGSTT2, DmGSTT3a, DmGSTT3b, DmGSTT4), and one Sigma class glutathione transferase on the activity of *Drosophila* p38b, including the reciprocal effect of this kinase protein on glutathione transferase activity. It was found that DmGSTT2, DmGSTT3b, DmGSTO1, and DmGSTO3 activated p38b significantly. Substrate specificities of GSTs were also altered after co-incubation with p38b. Although p38b activated DmGSTO1, DmGSTO2a, and DmGSTT2, it inhibited DmGSTT3b and DmGSTO3 activity toward xenobiotic and physiological substrates tested. These results suggest a novel link between Omega and Theta GSTs with the p38b MAP kinase pathway.

Abbreviations: CDNB, 1-chloro-2,4-dinitrobenzene; CuOOH, cumene hydroperoxide; DCNB, 1,2-dichloro-4-nitrobenzene; EA, ethacrynic acid; GST, glutathione S-transferases, HED, hydroxyethyldisulfide; JNK, Jun kinase; MAPK, mitogen activated protein kinase; PNBC, p-nitrobenzyl chloride; PNPB, p-nitrophenethyl bromide; 4-HNE, 4-hydroxynonenal.

Correspondence: a jwongtrakul@yahoo.com, b kanokwanjan@yahoo.com, c biotbug@yahoo.com, d albertketterman@yahoo.com, *Corresponding author

Editor: David Schooley was editor of this paper.

Received: 3 July 2012 Accepted: 22 October 2012 Published: 1 May 2014

Copyright: This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 14, Number 60

Cite this paper as:

Wongtrakul J, Janphen K, Saisawang C, Ketterman AJ. 2014. Interaction of Omega, Sigma, and Theta glutathione transferases with p38b mitogen-activated protein kinase from the fruit fly, *Drosophila melanogaster*. *Journal of Insect Science* 14(60). Available online: http://www.insectscience.org/14.60

Introduction

S-transferases (GSTs, Glutathione EC 2.5.1.18) are multifunctional enzymes involved in detoxification and excretion of physiological and toxic substances (Haves et al. 2005). Several studies indicated that in addition to the enzymatic functions, GSTs also have roles such as regulation of mitogenactivated protein kinase (MAPK) pathways. A number of reports have shown that GST enzymes can interact and modulate several proteins in MAPK pathways, e.g. JNK, TRAF2, and ASK1 (Adler et al. 1999; Cho et al. 2001; Wang et al. 2001; Wu et al. 2006; Laborde 2010). Mammalian GSTs have been categorized into at least seven classes; Alpha, Mu, Pi, Omega, Sigma, Theta, and Zeta (Frova 2006). By extending the mammalian nomenclature, insect GSTs have been grouped into six classes: Delta, Epsilon, Omega, Sigma, Theta, and Zeta (Enavati et al. 2005). Studies on the insect GSTs have mainly focused on their roles in conferring insecticide resistance, e.g., Delta and Epsilon GSTs (Hemingway 2000; Enayati et al. 2005). The Omega, Sigma, and Theta classes have been much less studied, and their non-enzymatic roles have not been elucidated. There is accumulating evidence that suggests GSTs in Omega, Sigma, and Theta classes have important roles in oxidative stress (Singh et al. 2001; Burmeister et al. 2008; Jaramillo-Gutierrez et al. 2009; Piaggi et al. 2010; Nair and Choi 2011; Kim et al. 2012), which also suggests additional roles in cell signaling in response to oxidative stress.

Cellular behavior in response to extracellular stimuli is mediated via intracellular signaling pathways, such as the MAPK pathways. Several MAPK families have been reported, including extra-cellular signal-regulated kinase, Jun kinase (JNK/SAPK), and p38b

MAPK (Ono and Han 2000). The p38b MAPK pathway plays a major role in apoptosis, differentiation, survival, proliferation, development, inflammation, and other stress responses. The p38b MAPK has a role in a number of important diseases, such as cardiovascular diseases and inflammatory diseases, including cancers. Therefore, modulating p38b MAPK may represent a novel therapeutic target for those diseases. This study, using a Drosophila model, focuses on the p38b kinase and possible interactions between GST enzymes and p38b in modulating p38b signal transduction. Several studies have found that GSTs were involved in p38b signaling. It was reported that GST mu directly interacts with ASK1 and modulates p38b MAPK signaling (Cho et al. 2001; Dorion et al. 2002). In addiregulator tion. GST mu is а of dexamethasone-induced apoptosis by inhibition of Bim through down regulation of p38b MAPK and up regulation of NF-KB p50 (Hosono et al. 2010). GST mu expression also has been shown to regulate p38b phosphoryla-It was found that hypertension tion. susceptible C57BL/6 VSM cells, vascular smooth muscle cells, possessed lower levels of GSTM1 compared to VSM cells from a resistant strain and had significantly more p38b mitogen-activated protein kinase phosphorylation after H₂O₂ exposure (Yang et al. 2009). H₂O₂ treatment and knockdown of GSTM1 in the resistant VSM cells also showed increased phosphorylation of p38b kinase.

Previously, we reported a GST-p38b interaction with insect specific GST Delta class, (Wongtrakul et al. 2012), therefore we would like to further extend the study with Omega, Theta, and Sigma GSTs from *Drosophila melanogaster* Meigen (Diptera: Drosophilidae). The study included all five Omega class GSTs (DmGSTO1, DmGSTO2a,

DmGSTO2b, DmGSTO3, DmGSTO4), all GSTs (DmGSTT1, five Theta class DmGSTT2, DmGSTT3a, DmGSTT3b, DmGSTT4), and the one Sigma class GST. Four of the GSTs are splice variants from their respective genes, DmGSTO2a and DmGSTO2b from Omega 2 gene and DmGSTT3a and DmGSTT3b from the Theta 3 gene (Saisawang et al. 2012). These three classes are expressed in all known eukaryotes. Using an in vitro kinase assay and two different p38b substrates, ATF2 and Jun, we examined whether substrate specificity changes occurred for p38b upon GST interaction. We also studied kinase effects on GST activity toward both xenobiotic and physiological substrates. Our data showed interaction effects occurred between several GST isoforms and p38b MAPK.

Materials and Methods

Recombinant protein expression

The expression and purification of all proteins were carried out as previously described (Bukhtiyarova et al. 2004; Udomsinprasert et al. 2004; Saisawang et al. 2012). For ATF2, a HiTrap Chelating HP (GE Healthcare, www.gehealthcare.com) charged with NiCl₂ was used according to the manufacturer's instruction. The purified proteins were stored in 50% glycerol, 10 mM DTT, 50 mM phosphate buffer pH 6.5 for GSTs or 50 mM Tris-HCl pH 8.0 for the kinase pathway proteins at -20°C.

GST substrate specificity

Substrate specificity of DmGST enzymes toward five xenobiotic substrates (1-chloro-2,4dinitrobenzene (CDNB), 1,2-dichloro-4nitrobenzene (DCNB), ethacrynic acid (EA), p-nitrophenethyl bromide (PNPB), and pnitrobenzyl chloride (PNBC)) were measured in a 96-well plate in a spectra MR Microplate spectrophotometer (Dynex Technologies, <u>www.dynextechnologies.com</u>), using the appropriate pH and λ_{max} (Habig et al. 1974). Specific activity is reported as µmol/min/mg protein.

Effect of p38b interaction on GST activity

The effects of p38b interaction on GST activity toward some model substrates (4hydroxynonenal (4-HNE) (Bruns et al. 1999), cumene hydroperoxide (CuOOH) (Hiratsuka et al. 1997), and hydroxyethyldisulfide (HED) (Collinson and Grant 2003)) were performed. For 4-HNE, GST activities were measured spectrophotometrically in 1-mL quartz cuvettes in a UV-visible spectrophotometer (Thermo Scientific, www.thermoscientific.com). DmGST and p38b were desalted prior to co-incubation at room temperature using 1:1 molar ratio for 5 minutes, then GST activity was measured in comparison with the activity in the absence of p38b.

Kinase assay

All DmGST and MAPK pathway proteins were desalted to remove DTT and glycerol prior to the kinase assay. Kinase activity assays were performed using the ADP Quest Assay (DiscoveRx, www.discoverx.com). The p38b kinase activity was monitored using SynergyTM HT plate reader (Biotek, www.biotek.com) operating in kinetic mode, and fluorescence intensity measurements (530 nm excitation and 590 nm emission) were made every 10 seconds for 2 minutes. Briefly, p38b (4µg) was combined using 1:1 molar ratio of GST and 100 µM ATP in ADP assay buffer. The solution was incubated for 3 minutes at room temperature before addition of 10 µg ATF2 or Jun, reagent A and reagent B, respectively. All reactions were performed in a final volume of 100 µL. Fluorescence intensity values were plotted against time to



A Low Molecular Weight Calibration Kit was used for SDS electrophoresis (Amersham Biosciences, GE Life Science, <u>www.gelifesciences.com</u>). Values are shown in kilodaltons. M: molecular weight marker; lane 1: purified DmGSTO1; lane 2: purified DmGSTO2a; lane 3: purified DmGSTO2b; lane 4: purified DmGSTO3; lane 5: DmGSTO4; lane 6: DmGSTS1; lane 7: DmGSTT1; lane 8: DmGSTT2; lane 9: DmGSTT3a; lane 10: DmGSTT3b; lane 11: DmGSTT4. High quality figures are available online.

DmGSTs	Substrates					
	CDNB	DCNB	PNBC	PNPB	EA	CuOOH
DmGSTT1	8.33 ± 0.188	0.045 ± 0.001	ND	0.062 ± 0.003	ND	3.80 ± 0.236
DmGSTT2	4.98 ± 0.211	< 0.018	0.194 ± 0.038	ND	ND	ND
DmGSTT3a	0.406 ± 0.056	ND	ND	0.673 ± 0.084	ND	2.18 ± 0.217
DmGSTT3b	0.030 ± 0.005	0.009 ± 0.001	0.311 ± 0.014	0.271 ± 0.005	0.029 ± 0.011	1.03 ± 0.034
DmGSTT4	ND	ND	ND	ND	ND	ND
DmGSTS1	1.85 ± 0.061	ND	ND	ND	ND	-
DmGSTO1	0.001 ± 0.0003	ND	ND	ND	ND	(<u>2</u>)
DmGSTO2a	0.003 ± 0.0015	ND	ND	ND	ND	-
DmGSTO2b	ND	ND	ND	ND	ND	-
DmGSTO3	0.079 ± 0.0009	ND	0.0064 ± 0.0006	0.0078 ± 0.0013	ND	-
DmGSTO4	0.003 ± 0.0003	ND	ND	ND	ND	-

Specific activity is shown as µmol/min/mg protein. Results are means ± SD for at least three separate assays. ND is not detected. The substrate concentrations used were: CDNB, 1 mM; DCNB, 1 mM; PNBC, 1.2 mM; PNPB, 0.1 mM; EA, 0.2 mM; and CuOOH, 1.5 mM.

obtain a slope. The correlation between fluorescence intensity and time was linear. The background reaction rate was measured in a reaction lacking enzyme. Enzyme activities were obtained by subtracting the slope of background from the experimental sets. All experiments were performed in triplicate, and the activities were calculated using Graphpad Prism 4.0 (www.graphpad.com).

Results

Substrate profiling

The recombinant GSTs used in this report were purified to greater than 90% homogeneity as shown by SDS-PAGE (Figure 1). GSTs display differences in substrate specificities in conjugating chemical substances. Therefore, six different substrates, CDNB, DCNB, EA, PNPB, PNBC, and CuOOH, were chosen to study for the enzymatic activities. Table 1

summarizes the results of the GST activity profiling under standard assay conditions. Only DmGSTT3b showed activity towards EA. It also had the greatest activity toward PNBC. All of the enzymes were most active with CDNB as substrate, except DmGSTT4 and DmGSTO2b, which showed no CDNB activity at the concentrations tested. Unlike other Omega DmGSTs, DmGSTO3 appears to have a broader substrate specificity displaying activity toward PNBC and PNPB. The remaining Theta enzymes also showed broader substrate specificity with the tested substrates compared to the Omega GSTs. DmGSTT3a had the greatest activity toward PNPB, which is the substrate preference of Theta class, compared to DmGSTT3b and DmGSTT1, which had 2.48-fold and 10.8fold less activity than DmGSTT3a, respectively. Although DmGSTT1 had the lowest toward PNPB activity compared to DmGSTT3a and DmGSTT3b, it demonstrated the greatest activity for DCNB, as well as ac-

tivity for CDNB. Unlike DmGSTT3a, the splice variant DmGSTT3b had detectable activity toward DCNB. DmGSTT2 can be from discriminated DmGSTT1 and DmGSTT3 by its inability to utilize PNPB. DmGSTT2 also had the ability to utilize PNBC and possessed high CDNB conjugating activity, approximately 60% of DmGSTT1. DmGSTT1, DmGSTT3a, and DmGSTT3b can also metabolize cumene hydroperoxide, suggesting possible physiological roles in oxidative stress product metabolism. The peroxidase activity of DmGSTT1 was 2-fold greater than DmGSTT3a, whereas the activity of DmGSTT3b was less than DmGSTT3a by approximately 2-fold. DmGSTT2 and DmGSTT4 had no peroxidase activity.

GST specific activity changes upon interaction with p38b

Four substrates were used to study whether p38b affected substrate specificity of GST. CDNB is a classical GST xenobiotic substrate,



and three different model substrates were used: CuCOOH for general peroxidase activity, HED for thiol transferase activity, and 4-HNE, a physiological substrate originating from lipid peroxidation. For conjugating activity of GSTs toward CDNB, it was found that p38b increased DmGSTO1, DmGSTO2a, and DmGSTT2 activities 58%, 26%, and 14%, respectively (Figure 2). Intriguingly, p38b decreased CDNB conjugating activity of DmGSTT3b by approximately 84%. Peroxidase and thiol transferase activities toward





CuCOOH and HED in the presence and absence of p38b were furthered studied for The-Theta enzymes and Omega enzymes. It was found that peroxidase activity of Theta GSTs was changed, ranging from 0% to 7% (data not shown), whereas thiol transferase activity change ranged from 0.2% to 14% (Figure 2). DmGSTO3 demonstrated the greatest percent change, approximately 14%. In addition, 4-HNE substrate, a lipid peroxidation product, was also tested with DmGSTS1. The activity of DmGSTS1 did not change upon preincubation with p38b (data not shown). Therefore, p38b showed varying effects on the GST activity for CDNB and thiol transferase activity for both Omega and Theta class GSTs.

p38b kinase substrate specificity changes upon GST interaction

Kinase activity of p38b toward ATF2 and Jun substrates in the presence and absence of Drosophila Theta, Omega, and Sigma GSTs was determined. class For Theta GSTs, DmGSTT1, DmGSTT3a, and DmGSTT4 did not affect kinase activity of p38b toward both ATF2 and Jun substrates (Figures 3 and 4). It was found that DmGSTT2 enzyme significantly increased p38b activity toward ATF2 approximately 3.1-fold. Moreover, the enzyme also increased p38b activity toward Jun approximately 4.3-fold. Unlike DmGSTT3a, the other spliced variant of the same gene, DmGSTT3b, which possesses an extra 40 amino acids at the N-terminus, showed a 3.4fold increase in p38b kinase activity toward ATF2. Interestingly, the interaction between DmGSTT3b and p38b also increased kinase activity for Jun 7.4-fold, which was the greatest activation of p38b observed. Kinase activity of p38b in the presence of the five Omega GSTs was slightly increased toward ATF2, ranging from 1.2 to 1.5-fold; although, none of these proteins significantly activated p38b kinase. DmGSTO2a, DmGSTO2b, and DmGSTO4 did not alter p38b activity toward Jun substrate, whereas DmGSTO1 and DmGSTO3 significantly increased p38b activity approximately 2.0 and 1.6-fold. Unlike the Theta and Omega enzymes, DmGSTS1 did not affect the p38b activity, which remained unchanged when tested with both kinase substrates.

Discussion

Using standard GST assays, the substrate panel demonstrated that Drosophila Theta class GSTs had broader conjugating activities compared to the Omega class. With the exception of DmGSTT4, the enzymes tested preferred CDNB as the substrate in transferase reactions. DmGSTT1 and DmGSTT3a utilized PNPB but not PNBC, although both are reportedly Theta class GST substrates. For comparison, the mosquito AcGSTT1 utilized both PNBC and PNPB with 10-fold more activity toward PNBC compared to PNPB (Wongtrakul et al. 2010). DmGSTT1 had lower PNPB activity compared to AcGSTT1 (approximately 11.8-fold less), but greater activity compared to CDNB (approximately 1.34-fold greater). In contrast, although DmGSTT1 utilized DCNB as a substrate, AcGSTT1 did not. The xenobiotic substrate selectivity of Omega and Sigma GSTs was more restricted. With the exception of DmGSTO2b and DmGSTO3, the enzymes acted only as CDNB-conjugating enzymes. Similar substrate selectivity was reported for AcGSTO1 (Wongtrakul et al. 2010). The DmGSTO3 enzyme appeared to show broader substrate specificity compared to the other Omega GSTs because it showed activity towards both PNBC and PNPB, which are Theta class substrates. The GSTs in the present study appeared to show little or no peroxidase activity for the tested substrates either in this or a previous study (Saisawang et al. 2012).

This appears to be in contrast to the previous reports for these GST classes and demonstrates that GST isoforms from the same class may have quite different substrate specificity (Singh et al. 2001; Burmeister et al. 2008; Jaramillo-Gutierrez et al. 2009; Piaggi et al. 2010; Nair and Choi 2011; Kim et al. 2012).

DmGSTO1 significantly increased the activation of p38b toward Jun (Figure 4). In addition, p38b also increased DmGSTO1 activity toward CDNB (Figure 2). These data show that DmGSTO1 interacts with p38b, and the effects have an impact on both enzymes. DmGSTO2a was reported to play several roles in a *Drosophila* model of Parkinson's disease (although referred to as DmGSTO1a in the report) (Kim et al. 2012). Overexpression of DmGSTO2a suppressed phenotypes caused by the *parkin* gene, resulting in loss of function, e.g., collapsed thorax, downturned wing muscle degeneration and donaminergic

function, e.g., collapsed thorax, downturned wing, muscle degeneration, and dopaminergic neuron loss. The enzyme was shown to be involved in MAPK signaling, as the expression of DmGSTO2a inhibited activation of JNK in *parkin* mutants of *Drosophila*. The overexpressed DmGSTO2a blocked the activation of JNK and apoptosis in the *parkin* mutants,



which also protected against the degeneration of indirect flight muscle. In our results, both DmGSTO2a and DmGSTO2b appeared not to interact with p38b for either ATF2 or Jun, suggesting that these GSTs were not involved with the p38b pathway at this level.

We observed that DmGSTO3 significantly increased p38b activity toward Jun (Figure 4). In addition, p38b also decreased DmGSTO3 thiol transferase activity by approximately 15% (Figure 2). This observation suggests the specific interactions of both proteins, the kinase and the GST, may affect the respective physiological roles. In adult flies, the DmGSTO3 transcript was strongly expressed in the regions associated with digestion and detoxification, namely the crop, gut, and tubules (Walters et al. 2009). In embryo, expression of DmGSTO3 was only detected in stages 13–16, the stages of gut development. Although reportedly involved with oxidative stress, the DmGSTO3 shows little or no activity for oxidative stress substrates. A Fox-like transcription factor binding motif is found in proximity to the DmGSTO3 gene, and this binding motif is associated with signaling pathway and cell fate decisions in development (Li et al. 2008). Down regulation of DmGSTO3 in dcr-2 mutant Drosophila flies appeared to correlate with decreased survival under oxidative stress as well as a decreased life span (Lim et al. 2011). This raises the possibility that DmGSTO3 may have similar roles as the human GSTO1. Human GSTO1 has been shown to modulate ryanodine receptors, which are intracellular calcium channels (Dulhunty et al. 2001). It was shown some of the modulation was dependent upon Omega enzyme thiol activity. Omega enzyme activity was also found to be important for glutathionvlation of ATP synthase β subunit for activity of F₁F₀-ATP synthase activity (Kim et al. 2012). These data show the Omega GSTs

possess a substrate/protein interaction specificity extending beyond p38b kinase interactions, as well as each Omega GST showing unique preferences.

Comparing DmGSTT1, DmGSTT2, DmGSTT3a, and DmGSTT3b showed an amino acid identity ranging from 40% to 85% and an amino acid similarity ranging from 58% to 85% (Figure 5). Accordingly, these isoenzymes have distinct substrate profiles (Table 1). DmGSTT1 had substrate preferences for CDNB and cumene hydroperoxide. DmGSTT2 had no detectable activity with PNPB, which is a highly specific substrate for Theta GST, whereas both DmGSTT3a and DmGSTT3b had high activity. This is most likely due to the differences in their hydrophobic substrate binding site (H-site) conformations. It was found that p38b affected DmGSTT3b activity by reducing the CDNB activity 80% while showing no effect on GST activity for HED (Figure 2). This suggests the binding of p38b induces conformational changes in the GST that differentially affect substrate specificity. Similar effects for kinase-GST interactions were also reported for JNK decreasing activity for several studied GSTs toward CDNB, DCNB, and PNBC (Udomsinprasert et al. 2004). In contrast, DmGSTT3b activated p38b kinase activity significantly toward both ATF2 and Jun substrates compared to the control (Figures 3 and 4). A possible region of GST-p38b interaction might be the N-terminus of GST, as DmGSTT3b had 40 amino acids extra at the N-terminus compared to DmGSTT3a, with the remainder of the two proteins being identical (Figure 5). The extra amino acids of DmGSTT3b increased p38b activity compared to the DmGSTT3a-p38b interaction 3.1and 6.9-fold for ATF2 and Jun, respectively. This suggests that the N-terminus of GST may be involved in p38b interaction, which also

has been reported for a Delta class GST. DmGSTD11a has a 21 amino acid extension at the N-terminus, which significantly affected p38b activity (Wongtrakul et al. 2012). Evidence for an additional N-terminus role is that DmGSTT3b activity can be detected for DCNB, PNBC, and EA, whereas no detectable activity was observed for DmGSTT3a (Table 1). Our study also suggests a role for GSTT2 in MAPK cell signaling (Figures 3 and 4). DmGSTT2 similarly affected p38b activity, increasing it toward both ATF2 and Jun substrates. The mechanisms for the activation of p38b by DmGSTT2, as well as the DmGSTT2 CDNB conjugating activity increases induced by p38b, remain to be elucidated. Alignments of the four sequences of the GSTs that interacted with p38b, T2, T3b, O1, and O3, and those from the same GST class that appeared to not affect p38b activity, did not yield any clues as to the amino acids that were involved in the proteinprotein interactions (Figure 5). The interactions therefore must be determined by tertiary changes that occur with the sequence variations between the enzymes.

In summary, employing several substrates for p38b MAPK as well as for DmGSTs, we observed that GSTs from Theta and Omega, but not Sigma, classes demonstrated protein interp38b. actions with These interactions increased p38b kinase activity toward its transcription factor substrates ATF2 and Jun. This has been reported previously for several Delta class GSTs interacting with p38b and showing reciprocal effects on substrate specificity (Wongtrakul et al. 2012). The p38b-GST protein interactions would cause conformational changes affecting substrate preference for both the GST and the p38b kinase. The physiological functions of these interactions are areas for future research.

Acknowledgments

This work was supported by the Thailand Research Fund (TRF) to J. Wongtrakul and funding from Mahidol University to A. J. Ketterman.

References

Adler V, Yin Z, Fuchs SY, Benezra M, Rosario L, Tew KD, Pincus MR, Sardana M, Henderson CJ, Wolf CR, Davis RJ, Ronai Z. 1999. Regulation of JNK signaling by GSTp. *EMBO Journal* 18: 1321-1334.

Bruns CM, Hubatsch I, Ridderström M, Mannervik B, Tainer JA. 1999. Human glutathione transferase A4-4 crystal structures and mutagenesis reveal the basis of high catalytic efficiency with toxic lipid peroxidation products. *Journal of Molecular Biology* 288: 427-439.

Bukhtiyarova M, Northrop K, Chai X, Casper D, Karpusas M, Springman E. 2004. Improved expression, purification, and crystallization of p38alpha MAP kinase. *Protein Expression and Purification* 37: 154-161.

Burmeister C, Lüersen K, Heinick A, Hussein A, Domagalski M, Walter RD, Liebau E. 2008. Oxidative stress in *Caenorhabditis elegans*: protective effects of the Omega class glutathione transferase (GSTO-1). *FASEB Journal* 22: 343-354.

Cho SG, Lee YH, Park HS, Ryoo K, Kang KW, Park J, Eom SJ, Kim MJ, Chang TS, Choi SY, Shim J, Kim Y, Dong MS, Lee MJ, Kim SG, Ichijo H, Choi EJ. 2001. Glutathione S-transferase Mu modulates the stressactivated signals by suppressing apoptosis

signal-regulating kinase 1. *Journal of Biological Chemistry* 276: 12749-12755.

Collinson EJ, Grant CM. 2003. Role of yeast glutaredoxins as glutathione S-transferases. *Journal of Biological Chemistry* 278: 22492-22497.

Dorion S, Lambert H, Landry J. 2002. Activation of the p38 signaling pathway by heat shock involves the dissociation of glutathione S-transferase mu from Ask1. *Journal of Biological Chemistry* 277: 30792-30797.

Dulhunty A, Gage P, Curtis S, Chelvanayagam G, Board P. 2001. The glutathione transferase structural family includes a nuclear chloride channel and a ryanodine receptor calcium release channel modulator *Journal of Biological Chemistry* 276: 3319-3323.

Enayati AA, Ranson H, Hemingway J. 2005. Insect glutathione transferases and insecticide resistance. *Insect Molecular Biology* 14: 3-8.

Frova C. 2006. Glutathione transferases in the genomics era: new insights and perspectives. *Biomolecular Engineering* 23: 149-169.

Habig WH, Pabst MJ, Jakoby WB. 1974. Glutathione S-transferases. The first enzymatic step in mercapturic acid formation. *Journal of Biological Chemistry* 249: 7130-7139.

Hayes JD, Flanagan JU, Jowsey IR. 2005. Glutathione transferases. *Annual Review of Pharmacology and Toxicology* 45: 51-88.

Hemingway J. 2000. The molecular basis of two contrasting metabolic mechanism of

insecticide resistance. *Insect Biochemistry and Molecular Biology* 30: 1009-1015.

Hiratsuka A, Yamane H, Yamazaki S, Ozawa N, Watabe T. 1997. Subunit Ya-specific glutathione peroxidase activity toward cholesterol 7-hydroperoxides of glutathione S-transferases in cytosols from rat liver and skin. *Journal of Biological Chemistry* 272: 4763-4769.

Hosono N, Kishi S, Iho S, Urasaki Y, Yoshida A, Kurooka H, Yokota Y, Ueda T. 2010. Glutathione S-transferase M1 inhibits dexamethasone-induced apoptosis in association with the suppression of Bim through dual mechanisms in a lymphoblastic leukemia cell line. *Cancer Science* 101: 767-773.

Jaramillo-Gutierrez G, Rodrigues J, Ndikuyeze G, Povelones M, Molina-Cruz A, Barillas-Mury C. 2009. Mosquito immune responses and compatibility between *Plasmodium* parasites and anopheline mosquitoes. *BMC Microbiology* 9: 154.

Kim K, Kim SH, Kim J, Kim H, Yim J. 2012. Glutathione s-transferase omega 1 activity is sufficient to suppress neurodegeneration in a *Drosophila* model of Parkinson disease. *Journal of Biological Chemistry* 287: 6628-6641.

Laborde E. 2010. Glutathione transferases as mediators of signaling pathways involved in cell proliferation and cell death. *Cell Death and Differentiation* 17: 1373-1380.

Li HM, Buczkowski G, Mittapalli O, Xie J, Wu J, Westerman R, Schemerhorn BJ, Murdock LL, Pittendrigh BR. 2008. Transcriptomic profiles of *Drosophila melanogaster* third instar larval midgut and

responses to oxidative stress. *Insect Molecular Biology* 17: 325-339.

Lim DH, Oh CT, Lee L, Hong JS, Noh SH, Hwang S, Kim S, Han SJ, Lee YS. 2011. The endogenous siRNA pathway in *Drosophila* impacts stress resistance and lifespan by regulating metabolic homeostasis. *FEBS Letters* 585: 3079-3085.

Nair PM, Choi J. 2011. Identification, characterization and expression profiles of *Chironomus riparius* glutathione S-transferase (GST) genes in response to cadmium and silver nanoparticles exposure. *Aquatic Toxicology* 101: 550-560.

Ono K, Han J. 2000. The p38 signal transduction pathway: activation and function. *Cellular Signalling* 12: 1-13.

Piaggi S, Raggi C, Corti A, Pitzalis E, Maschera MC, Saviozzi M, Pompella A, Casini A. 2010. Glutathione transferase omega 1-1 (GSTO1-1) plays an anti-apoptotic role in cell resistance to cisplatin toxicity. *Carcinogenesis* 31: 804-811.

Saisawang C, Wongsantichon J, Ketterman AJ. 2012. A preliminary characterization of the cytosolic glutathione transferase proteome from *Drosophila melanogaster*. *Biochemical Journal* 442: 181-190.

Singh SP, Coronella JA, Beneš H, Cochrane BJ, Zimniak P. 2001. Catalytic function of *Drosophila melanogaster* glutathione Stransferase DmGSTS1-1 (GST-2) in conjugation of lipid peroxidation end products. *European Journal of Biochemistry* 268: 2912-2923.

Udomsinprasert R, Bogoyevitch MA, Ketterman AJ. 2004. Reciprocal regulation of glutathione S-transferase spliceforms and the *Drosophila* c-Jun N-terminal Kinase pathway components. *Biochemical Journal* 383: 483-490.

Walters KB, Grant P, Johnson DL. 2009. Evolution of the GST omega gene family in 12 *Drosophila* species. *Journal of Heredity* 100: 742-753.

Wang T, Arifoglu P, Ronai Z, Tew KD. 2001. Glutathione S-transferase P1-1 (GSTP1-1) inhibits c-Jun N-terminal kinase (JNK1) signaling through interaction with the C terminus. *Journal of Biological Chemistry* 276: 20999-21003.

Wongtrakul J, Pongjaroenkit S, Leelapat P, Nachaiwieng W, Prapanthadara LA, Ketterman AJ. 2010. Expression and characterization of three new glutathione transferases, an epsilon (AcGSTE2-2), omega (AcGSTO1-1), and theta (AcGSTT1-1) from *Anopheles cracens* (Diptera: Culicidae), a major Thai malaria vector. *Journal of Medical Entomology* 47: 162-171.

Wongtrakul J, Sukittikul S, Saisawang C, Ketterman AJ. 2012. Mitogen-activated protein (MAP) kinase p38b interaction with delta class glutathione transferases from the fruit fly *Drosophila melanogaster*. *Journal of Insect Science* 12:107. Available online: www.insectscience.org/12.107

Wu Y, Fan Y, Xue B, Luo L, Shen J, Zhang S, Jiang Y, Yin Z. 2006. Human glutathione S-transferase P1-1 interacts with TRAF2 and regulates TRAF2-ASK1 signals. *Oncogene* 25: 5787-5800.

Yang Y, Parsons KK, Chi L, Malakauskas SM, Le TH. 2009. Glutathione S-Transferasem1 regulates vascular smooth muscle cell

proliferation, migration, and oxidative stress. *Hypertension* 54: 1360-1368.