

Resistance and cross-resistance in populations of the leafrollers, *Choristoneura rosaceana* and *Pandemis pyrusana*, in Washington apples

Authors: Dunley, John E., Brunner, Jay F., Doerr, Michael D., and Beers, E. H.

Source: Journal of Insect Science, 6(14) : 1-7

Published By: Entomological Society of America

URL: https://doi.org/10.1673/2006_06_14.1

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Resistance and cross-resistance in populations of the leafrollers, *Choristoneura rosaceana* and *Pandemis pyrusana*, in Washington apples

John E. Dunley, Jay F. Brunner, Michael D. Doerr and E. H. Beers

Washington State University, Tree Fruit Research and Extension Center, 1100 N. Western Ave., Wenatchee, WA 98801

Abstract

Insecticide bioassays of the leafrollers, *Choristoneura rosaceana* (Harris), and *Pandemis pyrusana* Kearfott (Lepidoptera: Tortricidae), were used to investigate resistance and cross-resistance between azinphosmethyl and other insecticides. Comparisons of field-collected populations with susceptible laboratory colonies of both leafroller species were made in 1996–97, prior to registration and field introduction of several of insecticides, and were re-tested in 2000–2001 following several years of use in the field. Insecticides tested included azinphosmethyl, chlorpyrifos, methyl parathion, tebufenozide, methoxyfenozide, spinosad, indoxacarb, acetamiprid, *Bacillus thuringiensis*, and azadirachtin. Azinphosmethyl-susceptible laboratory colonies were used for comparison to field populations. Resistance to azinphosmethyl was found in all populations of *C. rosaceana* (5.2–26.8 fold) and *P. pyrusana* (8.4–24.9 fold) collected from commercial orchards. Cross-resistance between azinphosmethyl and the insect growth regulators tebufenozide and methoxyfenozide was found in all but one population of the two leafroller species. No cross-resistance was found to chlorpyrifos. Some of the populations tested were cross-resistant to spinosad and indoxacarb, but the responses to these materials were more variable.

Keywords: insecticide resistance, organophosphate, insect growth regulator, benzoylhydrazine

Abbreviations: LCR Lethal concentration ratio, LC₅₀ Lethal concentration (50% of the population)

Correspondence: dunleyj@wsu.edu, jfb@wsu.edu, mdoerr@wsu.edu, ebeers@wsu.edu

Received: 25.3.2005 | Accepted: 7.12.2005 | Published: 25.8.2006

Copyright: Creative Commons Attribution 2.5 (<http://creativecommons.org/licenses/by/2.5>)

ISSN: 1536-2442 | Volume 2006, Number 14

Cite this paper as:

Dunley JE, Brunner JF, Doerr MD, Beers EH. 2006. Resistance and cross-resistance in populations of the leafrollers, *Choristoneura rosaceana* and *Pandemis pyrusana*, in Washington apples. 7pp. *Journal of Insect Science* 6:14, available online: insectscience.org/6.14

Introduction

Leafrollers are among the most destructive lepidopteran pests of tree fruits in Washington, second only to codling moth, *Cydia pomonella*. Although a number of leafroller species occurs in Washington, two species, *Choristoneura rosaceana* (Harris), and *Pandemis pyrusana* Kearfott, are the most important in commercial orchards. Damage from larval feeding on fruit can occur around bloom, during mid-summer, and just prior to harvest (Beers et al. 1993).

Leafrollers, and a wide spectrum of other apple pests, have been controlled using the organophosphate insecticides for over four decades. Organophosphate insecticides have played a key role in pest management during that time, especially after the phase-out of organochlorine insecticides. Organophosphate insecticides initially had a very wide spectrum of activity, including control of most primary and secondary insect pests of apple, as well as tetranychid mites. Efficacy against various pests declined over time with greater or less rapidity, until lepidopteran pests were the primary remaining targets. By the mid-1990s, only a few organophosphate insecticides were still widely used for codling moth and leafroller control, including azinphosmethyl, chlorpyrifos, and methyl parathion (Beers et al. 1996).

The organophosphate-based apple integrated pest management system was relatively stable for many years; however, regulatory action on organophosphate insecticides (Food Quality Protection Act of 1996) brought pressure for change in the fundamental approach to apple integrated pest management. In addition, there has been increasing evidence for organophosphate resistance in the codling moth, a key pest (Dunley and Welter 2000; Knight et al. 1994; Varela et al. 1993), along with reports of decreasing efficacy against leafrollers (Brunner 1996). The introduction of pheromone mating disruption represented a major change in apple integrated pest management and resulted in significant reductions in the use of organophosphate insecticides (Thomson et al. 2001, Brunner et al. 2002). The successful implementation of mating disruption of codling moth has promoted the investigation of alternative chemistries for lepidopteran pest control that are potentially less toxic to natural enemies, including *Bacillus thuringiensis*, insect growth regulators, neem-based insecticides, neonicotinyls, and

fermentation products. This change in approach has greatly diversified the array of control tactics available to the producer, and has provided greater opportunities for biological control.

The purpose of this study was to determine the status of susceptibility of the two primary leafroller pests to organophosphate insecticides, and examine field populations of leafrollers for resistance and cross-resistance to new insecticides.

Material and Methods

Rearing and Bioassay Methods

Laboratory colonies of *C. rosaceana* and *P. pyrusana* were collected initially from field populations occurring on apple. The *P. pyrusana* colony was collected from Yakima, WA, in 1985, and the *C. rosaceana* colony was collected from Mattawa, WA, in 1990. These two colonies have been reared continuously since their collection on a pinto bean diet using the method of Shorey and Hale (1965) under constant growth room conditions of $23 \pm 2^{\circ}\text{C}$, with a photoperiod of 16:8 L:D.

C. rosaceana and *P. pyrusana* field populations were collected either from the overwintering or summer generation larvae. Larvae were reared on pinto bean diet in the laboratory through to neonates of the first generation. Field populations were collected from two apple blocks in the Tree Fruit Research and Extension Center in Wenatchee Washington (TF1 and TF2); the Wenatchee Valley College orchard in East Wenatchee (WV); Stemilt Hill (SH), south of Wenatchee, on the west side of the Columbia River; Columbia River Orchards (CRO), 20 miles south of Wenatchee on the east side of the Columbia River; Quincy Washington (QC); Mattawa Washington (MA1 and MA2); Brewster Washington (BR); Bridgeport Washington (AR); an orchard on the Snake River, near Pasco, Washington (BO); and Milton-Freewater, Oregon (MF1, MF2, and MF3).

In 2000–01, more populations from a wider geographic distribution were tested. Three products tested in 1996–97 were not retested because of changes in use. Chlorpyrifos use in apple had been restricted to the pre-bloom period, methyl parathion registration had been withdrawn completely, and tebufenozide use had been largely replaced by methoxyfenozide. Indoxacarb, a newer material in the registration process, was added.

A leaf disk bioassay was used to expose leafroller larvae to insecticide residues. Treatments (insecticide concentrations) were prepared by dilution of the formulated insecticide in 500 ml water. A wetting agent, Latron B-1956 (Dow AgroSciences, www.dowagro.com/ (<http://www.dowagro.com/>)) (2 ml) was added to the stock solution. Serial dilutions were made from the stock solution, with 4 to 8 concentrations per insecticide. The control was water plus the wetting agent.

Leaves were collected from untreated apple trees (*Malus domestica* Borkhausen) at the Washington State University Tree Fruit Research and Extension Center, Wenatchee. Whole leaves were dipped in the various insecticide concentrations and allowed to air dry. Disks (2.3 cm diameter) were then cut from each leaf, and four leaf disks were placed in a small covered Petri dish (Falcon 1006, 50 x 9 mm, Becton-Dickinson Labware, www.bd.com/ (<http://www.bd.com/>)). Five 1- to 2-day-old leafroller larvae were placed at random on the leaf disks in each dish. A total of 10 dishes (50 larvae) were prepared per insecticide concentration. Petri dishes were placed in a plastic container and held at $23 \pm 2^\circ\text{C}$ and 16:8 (L:D). Larvae were examined after 7 d and mortality recorded, except for the two azadirachtin compounds, for which mortality was recorded after 14 days. Failure to respond to gentle probing with a camel's hair brush was classified as dead.

Statistical Analysis

Probit regression parameters (slope and LC) were estimated using the probit option of POLO-PC (LeOra 1987). Significant differences in LC₅₀ between leafroller species were determined by non-overlapping 95% fiducial limits. Comparisons

with susceptible populations among leafroller species were made using a lethal concentration ratio tests (LCR) (Robertson and Preisler 1992); ratios that had confidence limits not encompassing 1.0 were considered significantly different ($\alpha = 0.05$).

Results

The two laboratory colonies of *C. rosaceana* and *P. pyrusana* did not differ significantly in their responses to the organophosphate insecticides azinphosmethyl and methyl parathion, although *P. pyrusana* was more susceptible to chlorpyrifos than *C. rosaceana* (Table 1). There were no differences between the species for the insecticides with insect growth regulator activity. *P. pyrusana* was more susceptible to indoxacarb than *C. rosaceana*, but the species did not differ in responses to spinosad, acetamiprid, or *B. thuringiensis* (Table 1).

In the 1996–97 samples, the first generation for both populations of field-collected *C. rosaceana* (MF1 and MA1) had significantly higher LC₅₀ values to azinphosmethyl (9.2- to 10-fold) compared to the susceptible laboratory colony (Table 2). The field-collected populations had LC₅₀ values for chlorpyrifos that did not differ from that of the laboratory colony and neither population was cross-resistant to azinphosmethyl (Table 2). The field populations were significantly more tolerant of tebufenozide compared to the laboratory colony. LCRs for methoxyfenozide (12.8- and 26.3-fold) were higher than those for tebufenozide, suggesting a higher degree of cross-resistance (Table 2). Tebufenozide and methoxyfenozide were cross-resistant to azinphosmethyl even though they had not been used against *C. rosaceana* in the field.

Table 1. LC₅₀ values of OP-susceptible laboratory colonies of *C. rosaceana* and *P. pyrusana* to various insecticides, 1996–1997

	<i>C. rosaceana</i> - lab colony			<i>P. pyrusana</i> - lab colony		
		mg AI/liter			mg AI/liter	
			95% FL			95% FL
Insecticide	Slope ± SE	LC ₅₀	lower - upper	Slope ± SE	LC ₅₀	lower - upper
Organophosphates						
Azinphosmethyl	4.4±0.7	4.90	3.9 - 6.0	2.8±0.5	7.20	5.2 - 9.2
Chlorpyrifos	2.6±0.5	2.60	1.5 - 3.6	2.1±0.3	0.70	0.5 - 1.0
Methyl parathion	4.1±0.3	3.50	2.0 - 5.2	4.0±0.3	2.60	1.1 - 4.2
IGRs						
Tebufenozide	2.5±0.5	12.40	7.1 - 17.8	1.7±0.3	6.30	3.3 - 9.2
Methoxyfenozide	2.1±0.2	0.45	0.32 - 0.6	1.9±0.3	0.33	0.2 - 0.5
Azadirachtin ¹	1.6±0.5	15.20	4.6 - 26.2	1.2±0.3	6.10	0.5 - 14.2
Azadirachtin ³	1.7±0.3	5.40	2.2 - 8.7			
Miscellaneous						
Spinosad	2.9±0.4	0.28	0.21 - 0.4	2.6±0.6	0.18	0.09 - 0.2
Indoxacarb	2.2±0.5	1.90	1.2 - 2.5	1.5±0.2	0.26	0.1 - 0.5
Acetamiprid	2.7±0.9	45.90	9.8 - 59.8	1.3±0.4	99.50	48.5 - 220.1
<i>Bacillus thuringiensis</i>	2.0±0.3	198.10	101.4 - 287.4	1.6±0.5	278.30	39.2 - 474.1

Table 2. LC₅₀ and LCR values of laboratory (susceptible) and G₁ of field-collected populations of *C. rosaceana*, 1996–97

Insecticide	Year	<i>C. rosaceana</i>		mg AI/liter		LCR 95% CL ²		Significance
		Colony	n	Slope ± SE	LC ₅₀	lower - upper	lower - upper	
Azinphosmethyl	1997	LAB	400	4.4±0.7	4.90	3.9 - 6.0	----	
		MF1	400	1.4±0.3	45.30	23.5 - 67.0	9.2	*3
		MA1	400	1.6±0.2	49.10	24.4 - 65.6	10.0	*
Chlorpyrifos	1997	LAB	400	2.6±0.5	2.60	1.5 - 3.6	----	
		MF1	400	2.2±0.4	4.30	2.6 - 5.8	1.6	ns
		MA1	400	1.7±0.3	2.40	1.3 - 3.6	0.9	ns
Tebufenozide	1997	LAB	300	2.5±0.5	12.40	7.1 - 17.8	----	
		MF1	300	2.0±0.4	31.60	16.8 - 46.4	2.5	*
		MA1	300	1.8±0.3	34.00	20.6 - 53.2	2.7	*
Methoxyfenozide	1996	LAB	400	2.1±0.2	0.45	0.32 - 0.6	----	
		MF1	400	1.8±0.4	5.70	3.3 - 8.7	12.8	*
		MA1	400	0.7±0.3	11.80	3.2 - 8.6	26.3	*
Spinosad	1997	LAB	400	2.9±0.4	0.28	0.21 - 0.4	----	
		MF1	400	1.5±0.2	0.31	0.21 - 0.4	1.1	ns
		MA1	400	1.9±0.3	0.13	0.09 - 0.2	0.5	*

There was no difference in the tolerance between field populations of *C. rosaceana* to spinosad and the laboratory colony, but a weak (2-fold) negatively-correlated cross-resistance to spinosad was found in one *C. rosaceana* population (MA1) (Table 2).

All of the field (commercial orchard) populations of *C. rosaceana* were significantly resistant to azinphosmethyl relative to the laboratory colony (Table 3). Resistance levels were 5 to 27 fold, and resistance levels from the Mattawa and Milton-Freewater areas (MA and MF collections) were higher than found in 1997. The highest resistance levels were found in the Milton-Freewater area, which had a history of severe leafroller problems and liberal organophosphate use (J.F. Brunner unpublished data). In general, populations from the southern part of the region tended to be more resistant than those from the northern region (AR, BR).

All of the populations tested were resistant to methoxyfenozide and were also cross-resistant to azinphosmethyl (Table 3). However, there was a significant increase in the LC₅₀ estimate for the laboratory population in 2001, 2.0 mg AI/liter, from 0.45 mg AI/liter in 1996. Thus, the LCR values did not increase in correspondence to the higher levels of azinphosmethyl resistance.

None of the azinphosmethyl-resistant field populations showed resistance to spinosad, with LC₅₀ estimates ranging from 0.2 to 0.6 mg AI/liter (Table 3). Nevertheless, three of the eight populations had significantly different LCRs relative to the susceptible population. For these three populations, the LC₅₀s were lower than the susceptible population, indicating the possibility of

a negatively correlated cross-resistance, although the differences were small, similar to that found in 1997.

Only two *C. rosaceana* populations were tested with indoxacarb (Table 3). One, the TF2 population, was not significantly different the susceptible laboratory colony. The other, AR, had an extremely high LC₅₀ estimate, and correspondingly high LCR. High mortality of larvae of this population could not be obtained with the rates used, resulting in a very flat slope for the probit line. While these are very preliminary data they confirm reports of high *C. rosaceana* resistance in Canada (Smirle et al. 2002) and are an indicator of potential problems for the use of this material against *C. rosaceana*. The AR population was only moderately resistant (6.2-fold) to azinphosmethyl.

Of the five *P. pyrusana* field populations tested, all were resistant to azinphosmethyl, including the TF1 and WV populations, which had not received any organophosphate insecticides for five years prior to collection (Table 4). Resistance levels ranged from 8.4- to 24.9-fold, with the highest level found in the CRO population.

The only field-collected *P. pyrusana* population tested was resistant to methoxyfenozide, and was cross-resistant to azinphosmethyl (Table 4). Two of three field populations were slightly resistant to spinosad (1.9- to 2.5-fold) and they appeared to be cross-resistant to azinphosmethyl. The one population that was an exception was TF1, which had not received organophosphate insecticides for several years. Unlike *C. rosaceana*, all significant LCRs were positive for spinosad (>1). Response of *P. pyrusana* to indoxacarb was variable, with only

Table 3. LC₅₀ and LCR values for G₁ of field-collected populations of *C. rosaceana* in comparison to a susceptible laboratory population, 2000–01

						mg AI/liter			
		<i>C. rosaceana</i>				LC ₅₀ 95% FL		LCR 95% CL ²	
Insecticide	Year	Colony	n	Slope ± SE	LC ₅₀	lower - upper	LCR ¹	lower - upper	Significance
Azinphosmethyl	2001	LAB	300	3.9±0.9	6.7	4.3 - 8.6	----	----	----
	2001	BO	300	2.3±0.4	70.6	38.4 - 106.0	10.5	6.5 - 16.9	*3
	2001	MF2	300	2.5±0.6	167.8	98.8 - 266.1	25.0	2.8 - 220.2	*
	2001	MF3	300	2.7±0.5	179.7	127.2 - 255.3	26.8	17.8 - 39.9	*
	2001	BR	300	2.2±0.4	34.7	18.6 - 51.0	5.2	3.0 - 8.7	*
	2001	MA2	300	2.9±0.5	129.9	88.1 - 178.3	19.4	12.8 - 28.9	*
	2001	AR	300	2.8±0.6	41.3	21.3 - 57.7	6.2	3.8 - 9.8	*
Methoxyfenozide	2001	LAB	300	2.3±0.4	2.0	1.2 - 2.7	----	----	
	2001	MF2	300	2.2±0.6	16.0	9.1 - 31.0	8.2	4.7 - 14.2	*
	2001	BR	300	2.7±0.7	9.7	4.6 - 15.4	5.0	2.9 - 8.5	*
	2001	MA2	300	1.6±0.3	11.4	6.1 - 3.0	5.8	3.2 - 10.5	*
	2001	AR	300	1.4±0.2	10.9	5.7 - 30.3	5.6	3.0 - 10.4	*
Spinosad	2001	LAB	300	4.8±1.0	0.5	0.35 - 0.6	----	----	
	2001	BO	300	2.4±0.6	0.5	0.017 - 0.8	1.0	0.6 - 1.8	ns
	2001	TF2	210	2.6±0.5	0.2	0.13 - 0.3	0.4	0.2 - 0.6	*
	2001	MF2	300	2.5±0.4	0.5	0.34 - 0.7	1.0	0.6 - 1.5	ns
	2001	MF3	300	2.4±0.4	0.6	0.41 - 0.8	1.2	0.8 - 1.9	ns
	2001	BR	300	3.0±0.7	0.3	0.17 - 0.4	0.6	0.4 - 0.9	*
	2001	MA2	300	4.6±1.1	0.5	0.33 - 0.7	1.1	0.7 - 1.6	ns
	2001	MA3	150	2.7±0.7	0.3	0.12 - 0.4	0.5	0.3 - 0.9	*
	2001	AR	300	2.3±0.4	0.4	0.22 - 0.5	0.7	0.4 - 1.1	ns
Indoxacarb	2000	LAB	300	1.8±0.2	1.2	0.7 - 1.7	----	----	
	2000	TF2	300	1.4±0.2	1.4	0.08 - 2.3	1.2	0.7 - 2.1	ns
	2000	AR	300	0.4±0.2	5161 ⁴	----	4,440	1.1 - 1.8 x 10 ⁷	*

Table 4. LC₅₀ and LCR values for G₁ of field-collected populations of *P. pyrusana* in comparison to a susceptible laboratory population, 2000–01

					mg AI/liter				
		<i>P. pyrusana</i>			LC ₅₀	95% FL		LCR 95% CL ²	
Chemical	Year	Colony	n	Slope ± SE	LC ₅₀	lower - upper	LCR ¹	lower - upper	Significance
Azinphosmethyl	2001	LAB	300	5.5±1.0	5.0	3.7 - 6.2	----		
	2000	TF1	300	2.4±0.5	69.3	36.5 - 100.3	13.9	9.2 - 21.4	*3
	2001	SH	300	3.4±0.5	42.0	32.5 - 52.5	8.4	6.1 - 11.8	*
	2001	WV	240	2.3±0.5	88.9	56.2 - 123.1	17.8	11.6 - 27.6	*
	2000	CRO	300	2.3±0.4	124.4	89.3 - 172.7	24.9	17.4 - 36.2	*
Methoxyfenozide	2000	QC	300	4.0±0.8	46.9	34.3 - 46.9	9.4	6.7 - 13.4	*
	2001	LAB	300	2.6±0.4	1.0	0.8 - 1.3	----		
Spinosad	2001	SH	150	2.3±0.5	5.9	2.9 - 10.0	5.7	3.4 - 9.5	*
	2001	LAB	300	1.7±0.3	0.37	0.3 - 0.6	----		
	2001	TF1	150	3.0±0.8	0.43	0.4 - 0.6	1.2	0.6 - 2.3	ns
	2001	SH	300	2.5±0.4	0.70	0.7 - 1.0	1.9	1.1 - 3.4	*
	2001	WV	265	2.3±0.5	0.94	0.9 - 1.4	2.5	1.3 - 4.9	*
Indoxacarb	2001	LAB	300	3.6±0.7	0.38	0.4 - 0.5	----		
	2001	TF1	300	3.0±0.6	0.36	0.4 - 0.5	1.0	0.5 - 1.4	ns
	2001	CRO	300	2.1±0.4	1.18	1.2 - 1.6	3.2	2.0 - 5.2	*
	2001	QC	300	2.6±0.6	0.30	0.3 - 0.4	0.8	0.5 - 1.3	ns

one population having a significant LC₅₀ higher than the laboratory colony and with an LCR greater than one. This is the same population (CRO) had the highest azinphosmethyl LC₅₀.

Discussion

Information on resistance levels and cross-resistance is an important factor in structuring a resistance management program for leafrollers. The prevalence of azinphosmethyl resistance in both leafroller species reflects the decades of exposure, and indicates its current unsuitability as a leafroller management tool. While cross-resistance would be likely with other organophosphate insecticides, the lack of

cross-resistance to the organophosphate insecticide chlorpyrifos (1997) is consistent with the results of Dunley and Welter (2000), who found negatively-correlated cross-resistance in codling moth. The same would likely have been true of methyl parathion, which was relied on for leafroller control through the 1980's and also has been found to have a negative correlation to azinphosmethyl resistance in codling moth (Dunley and Welter 2000). Since methyl parathion is no longer registered for use in tree fruit and chlorpyrifos is restricted to prebloom use only, their negatively correlated cross-resistance is unavailable as tools in a resistance management program.

Cross-resistance between azinphosmethyl and the benzoylhydrazine insect growth regulators (tebufenozide and methoxyfenozide) means they must be used with caution, and only in a resistance management program. Such cross-resistance has been noted elsewhere (Sauphanor and Bouvier 1995; Carrière et al. 1996; Waldstein et al. 1999; Wearing 1998; Pree et al. 2001; Ahmad et al. 2002; Ahmad and Hollingworth 2004). The insect growth regulators are an important class of chemistry for use in mating-disruption based programs, and their conservation as tactics is vital. A clear demonstration of cross-resistance can provide convincing evidence to producers of the necessity of practicing resistance management (French-Constant and Roush 1990). Too often there is a tendency for producers to make a one-for-one switch of insecticides after resistance becomes apparent placing greater selection pressure on the new product.

The insect pest management program for Washington apples has more tools for chemical control of key pests than at any time in the last 40 years. The variety of insecticide choices available to control apple pests makes apple integrated pest management decisions more complex, but also provides opportunities for reducing the historical reliance on broad-spectrum neurotoxic insecticides. Additionally, some of the new chemistries are just as effective as the organophosphates, but may provide more stability to apple integrated pest management due to selectivity and better conservation of biological control agents. This opportunity for improving integrated pest management could be squandered if new insecticides are not effectively managed by following good resistance management practices. The information presented here shows that new insecticides, such as methoxyfenozide, can face the potential of reduced efficacy even before they are introduced. While spinosad does not appear to have been influenced by some pre-existing cross-resistance to organophosphate insecticides in leafrollers, it is difficult to predict how long it will remain effective if over-used. While it might be possible to select a susceptible leafroller population for resistance to spinosad or some other new insecticide in the laboratory this requires a great deal of effort and the ability of such an approach to predict the rate of resistance development is unclear. Therefore it seems necessary to conduct surveys of field populations of leafrollers on a regular basis as a way of tracking changes in resistance levels.

Apple producers in Washington will need to adopt a new level of sophistication when it comes to managing new insecticides. While the newer insecticides generally have narrower spectrums of activity than older products, many still affect more than one pest. We are recommending to Washington apple producers that an insecticide, or insecticide class, not be used against the same pest in two consecutive generations. If a product is used to control a pest at one time in the growing season it would not be used later in the season (next generation) against the same pest. If the product is also effective against another pest it could be used unless the application timing would expose the first target pest to selection. For example, it is possible to apply a product like methoxyfenozide against codling moth in the spring and then against *C. rosaceana* in the summer. However, this latter application would also impact the second codling moth generation and thus be counter to resistance management practices. While new insecticides provide more choices for management of apple pests, strategies to preserve those choices will limit the options in order to provide long-term stability to integrated pest management programs.

Acknowledgements

This project was funded in part by the Washington Tree Fruit Research Commission and Dow AgroSciences. The authors gratefully acknowledge the help of Lyla Lampson for leafroller field collections, Kathleen Pierre for colony maintenance, and many summer students for field and laboratory work.

References

- Ahmad M, Hollingworth RM, Wise JC. 2002. Broad-spectrum insecticide resistance in obliquebanded leafroller *Choristoneura rosaceana* (Lepidoptera: Tortricidae) from Michigan. *Pest Management Science* 58: 834-838.
- Ahmad M, Hollingworth RM. 2004. Synergism of insecticides provides evidence of metabolic mechanisms of resistance in the obliquebanded leafroller *Choristoneura rosaceana* (Lepidoptera: Tortricidae). *Pest Management Science* 60: 465-473.
- Beers, EH, Dunley JE, Brunner JF, Grove GG, Williams KM, Mayer DF, Parker R, Witmer G, Schreiber A, Daniels C, Maxwell T, Roberts, S. 1996. *1996 Crop Protection Guide for Tree Fruits in Washington*. Washington State University Cooperative Extension, Pullman, WA.
- Beers EH, Brunner JF, Willett MJ, Warner GM, editors. 1993. *Orchard Pest Management: A Resource Book for the Pacific Northwest*. Good Fruit Grower.

- Brunner JF. 1996. Management of leafrollers in Washington apple orchards. *Proceedings of the Washington State Horticultural Association* 91: 257-263.
- Brunner JF, Welter S, Calkins C, Hilton R, Beers EH, Dunley JE, Unruh T, Knight A, Van Steenwyk R, VanBuskirk P. 2002. Mating disruption of codling moth: a perspective from the Western United States. *IOBC-WPRS Bulletin* 25: 911-19.
- Carrière Y, Deland JP, Roff DA. 1996. Obliquebanded leafroller (Lepidoptera: Tortricidae) resistance to insecticides: among-orchard variation and cross-resistance. *Journal of Economic Entomology* 89: 577-582.
- Dunley JE, Welter SC. 2000. Correlated insecticide cross-resistance in azinphosmethyl resistant codling moth. *Journal of Economic Entomology* 93: 955-962.
- French-Constant RH, Roush RT. 1990. Resistance detection and documentation: the relative roles of pesticidal and biochemical assays. In: Roush RT, Tabashnik BE, editors. *Pesticide Resistance in Arthropods*, pp. 4-38. New York: Chapman and Hall.
- Knight AL, Brunner JF, Alston D. 1994. Survey of azinphosmethyl resistance in codling moth (Lepidoptera: Tortricidae) in Washington and Utah. *Journal of Economic Entomology* 87: 285-292.
- LeOra Software. 1987. *POLO-PC: A user's guide to probit or logit analysis*.
- Pree DJ, Whitty KJ, Pogoda MK, Bittner LA. 2001. Occurrence of resistance to insecticides in populations of the obliquebanded leafroller from orchards. *The Canadian Entomologist* 133: 93-103.
- Robertson JL, Preisler HK. 1992. *Pesticide bioassays with arthropods*. CRC Press.
- Sauphanor B, Bouvier JC. 1995. Cross-resistance between benzoylureas and benzoylhydrazines in the codling moth, *Cydia pomonella* L. *Pesticide Science* 45: 369-375.
- Shorey HH, Hale RL. 1965. Mass-rearing of the larvae of nine noctuid species on a simple artificial medium. *Journal of Economic Entomology* 58: 522-524.
- Smirle MJ, Lowery DT, Zurowski CL. 2002. Resistance and cross-resistance to four insecticides in populations of obliquebanded leafroller (Lepidoptera: Tortricidae). *Journal of Economic Entomology* 95: 820-825.
- Thomson D, Brunner JF, Gut LJ, Judd G, Knight A. 2001. Ten years implementing codling moth mating disruption in the orchards of Washington and British Columbia. Starting right and managing for success! In: Witzgall P, Kienzle J, editors. *IOBC-WPRS Bulletin Vol. 24(2)*, pp. 23-20. Monfavet, France.
- Varela LG, Welter SC, Jones VP, Brunner JF, Riedl H. 1993. Monitoring and characterization of insecticide resistance in codling moth (Lepidoptera: Tortricidae) in four western states. *Journal of Economic Entomology* 86: 1-10.
- Waldstein DE, Reissig WH, Scott JG, Straub RW. 1999. Susceptibility of obliquebanded leafroller (Lepidoptera: Tortricidae) populations from commercial apple orchards and an unsprayed habitat in New York to tebufenozide. *Journal of Economic Entomology* 92: 1251-1255.
- Wearing CH. 1998. Cross-resistance between azinphos-methyl and tebufenozide in the greenheaded leafroller, *Planotortrix octo*. *Pesticide Science* 54: 203-211.